

Lifetime positron annihilation spectroscopy and photo-inactivated bacteria

L V Elnikova¹

¹*A. I. Alikhanov Institute for Theoretical and Experimental Physics,
B. Cheremushkinskaya st. 25, Moscow 117218, Russia*

Combined positron and visible light irradiations in the photodynamical therapy (PDT) applications are analyzed. Objectives and goals of PDT are killing or irreversible oxidative damage of pathogenic cells, or rather their cell walls, cell membranes, peptides, and nucleic acids by photo-activated oxygen of photosensitizer injected into the target cell during light irradiation. In this paper, the arguments for involving of lifetime positron annihilation spectroscopy to control of photodamaging cells in the course of the PDT procedure are given on the examples of gram-positive and gram-negative bacteria basing on a brief survey of the literature.

PACS numbers:

INTRODUCTION

One of the most topical applications in use of affect irradiation onto living organisms might be photodynamical therapy (PDT) combined with positron annihilation methods.

PDT utilizes light of any waveband in combination with a photosensitizing agent to induce a phototoxic reaction which results with damage or death of pathogenic cells. Antibacterial PDT, directed against bacterial and yeast cells, has most demonstrative evidence in dermatology [1]. At present, PDT is a main competitor for the antibiomatic cure. If penicillin, discovered by Fleming in 1928, was not spread out in medicine, PDT would not be forgotten at a long date.

Photosensitizing reactions in PDT are the processes, in which absorption of light by a photosensitizer (or a dye) induces chemical changes in the outer wall at the surface of several types of bacterial and yeast cells, increases their permeability, and allows significant amounts of photosensitizer to be accumulated at the level of the cytoplasmic membrane. Two types of such reactions may carry out, either via radical mechanism (type I) or energy migration to produce reactive singlet oxygen (type II). The target cells of bacteria are classified to gram-positive (Gram (+), e.g. murein sacculus) and gram-negative (Gram (-), e.g. *Escherichia coli* (*E. coli*)), they differ in structure and thickness of the peptidoglycan layer. The positive charge of the dye appears to promote a tight electrostatic interaction with negatively charged sites at the outer surface of any species of bacterial cells.

Singlet oxygen $^1\text{O}_2$ was first observed in 1924. In 1931, Kautsky first proposed that $^1\text{O}_2$ might be a reaction intermediate in dye-sensitized photooxygenation [2]. Singlet oxygen is metastable states of triplet oxygen (O_2) with more high energy is less stable than triplet oxygen (O_2). The energy difference between the lowest energy of singlet state and the lowest energy of triplet state is about 11400 K ($T_e(a^1\Delta_g \leftarrow X_3\Sigma_g^-) = 0.98 \text{ eV} (=94.2 \text{ kJ/mol})$) and corresponds to the transition in near-infrared at $\sim 1270 \text{ nm}$.

The theory of molecular orbital predicts three low-lying excited singlet states of triplet molecular oxygen $\text{O}_2(X_3\Sigma_g^-)$: $\text{O}_2(a^1\Delta_g)$, $\text{O}_2(a'^1\Delta'_g)$, and $\text{O}_2(b^1\Sigma_g^+)$, which differ in only spins and occupation of antibonding degenerated π_g orbitals. Amid these states, namely $\text{O}_2(a^1\Delta_g)$ is called "singlet oxygen" as non-degenerated and more long-living state. Because of the differences in the electron shells, singlet and triplet oxygen differ in their chemical properties. Lifetime of singlet oxygen in vacuum equals 72 min and significantly decreases in dependence on a media. Singlet oxygen reacts with many kinds of biological molecules such as DNA, proteins and lipids [3].

Some notions to develop PAS as a new noninvasive technique for the detection of molecular damage by UV radiation have issued in [4] and references therein. The experiments on UV irradiated mouse's [4] and cancer-diseased [5] skin strikingly demonstrate the perspectives of positron annihilation spectroscopy (PAS) in medicine.

There were observed, that the long-living o -Ps is trapping into the 10-15 nm depth of the outer cell layer [4]. These scales imply to be comparable to penetration of photoinactivated dye oxygen into the outer layer of gram-negative bacteria, which is of just appropriate to porine and lipopolysaccharide size.

These data motivate the consideration the perspectives of positrons and Ps in the PDT use. To future experimental confirmations and medical applications, we rest here at the evaluation of positronium states for photoirradiated *E. coli* and the porphyrin photosensitizer, 4 N-methyl-pyridinium (meso) (T_4MPyP), in the presence of the Tris-EDTA agent.

TRIAL GRAM (-) BACTERIUM IN THE PDT PROCEDURE

Consider the photosensitization of *E. coli* with visible light of 250 W tungsten lamps for T_4MPyP due to the PDT action (Figures 1, 2). The presence of the Tris-EDTA agent [1], [6] in the system might be useful to enhancing of the membrane permeability and facilitating




FIG. 1: Pathway of type I and type II reactions of a light absorbing photosensitizer. After light activating of the ground state of a photosensitizer (PS), activated form of PS* can follow two alternative pathways via reactive singlet oxygen (1O_2), hydrogen peroxide, hydroxyl radical (type II) or organic substrate (S) (type I). The intermediates react rapidly with their surroundings: cell wall, cell membrane, peptides, nucleic acids, [1].




FIG. 2: Schematic representations of the arrangement of the cell walls of gram-negative (A) and gram-positive (B) bacteria. Gram (-) bacteria cell wall consists of a thin, inner wall composed of 2-3 layers of peptidoglycan (2-3 nm thick), a periplasmic space and an outer lipid bilayer (7 nm). The outer membrane contains phospholipids, lipoproteins, lipopolysaccharides and proteins like porins (A). Gram (+) bacteria cell wall appears as a 15-80 nm thick cell wall composed of up to 100 interconnecting layers of peptidoglycan (B). Teichoic acids are interwoven in the peptidoglycan layers. Some have a lipid attached (lipoteichoic acid). Also proteins are ingrained in the peptidoglycan layers, [1].

the penetration of phototoxic molecules to the cytoplasmic membrane. The addition of Tris-EDTA to Gram (-) bacteria removes the divalent cations (e.g. Ca^{2+} , Mg^{2+} ions) which are present in large numbers to stabilize adjacent negative charged lipopolysaccharides molecules at the outer membrane.

Typically, cells were incubated with 8.4 μM porphyrin

solution (1 mL) for 5 min at 37°C, cell pellets were washed once with 5 mM phosphate buffer, pH=7.4, and treated with 2% aqueous sodium dodecyl sulphate (SDS) in order to disrupt the cells and obtain the incorporation of the porphyrin in a monomeric state into the surfactant micelles.

Absorbed maximum of T_4MPyP is lying at 424 nm, the extinction coefficient equals to 194 $M^{-1}cm^{-1}$, and respective quantum yield of singlet oxygen is 0.74 [6].

In the 12 mL'th Pyrex test tube, there are 10^6 cell/mL. The cell survival has controlled due to the standard procedure [6]. During irradiation time 1, 5, and 10 min, 8.4 μM 'th T_4MPyP kills 0.0, 3.1, and 4.5 the *E. coli* cells respectively.

IRRADIATED CELLS AND POSITRONIUM STATES

Biochemical analysis performed on irradiated cells suggest that the cytoplasmic membrane is an important target of the photo-process [7]. To reach this membrane from the outside, singlet oxygen have to destroy an outer cell layer (Figure 2). In particular, the result of its damage should be phase transformations or the pore formation of the outer lipidic layer of a cell. In this respect, DPPC [8] and stratum corneum [9] have been studied with the Doppler-broadening and PALS techniques (without specific therapeutic applications).


Consider the main results of photo-destructive reactions of photosensitizers in the bacterial layer structure and chemical agents, and the positronium formation after that.

Role of singlet oxygen

Singlet oxygen is essentially used in PTD [1], in the considered scheme, it is produced by meso-Tetra(4-N-methyl-pyridyl)porphin (Figure 3). T_4MPyP is cationic porphyrin charged from +2 to +4. After neutralization of a whole outer layer with chemical or biological agents, one may approximately consider a target cell as neutral before immobilizing a dye (a photosensitizer) and light irradiation.

As it has been is known, molecular triplet oxygen in the ground state [15], [16] is a quencher for the Ps formation due to the mechanism of dissociative trapping, where the energies and the trapping cross sections are $E_{maxI} = 6.2$ eV, $E_{maxII} = 8.3$ eV, $\sigma_{maxI} = \sigma_{maxII} = 1.3 \cdot 10^{-18}$ cm^2 .

Photodynamic reactions based on primary porphyrin photoreduction by the oxidizing compound are an example of a type I mechanism where molecular oxygen plays the role of an electron acceptor [10]:







FIG. 3: The chemical structure of meso-Tetra(4-N-methylpyridyl)porphyrin (T4MPyP).

where P , $^1P^*$ and $^3P^*$ are photosensitizer molecules in the ground, excited singlet and triplet states, and D is a substrate of photooxygenation. In the alternative "type II" mechanism, the primary event is the energy transfer from triplet photosensitizers to dioxygen with population of its singlet $^1\Delta_g$ state (1O_2); then, singlet oxygen oxidizes appropriate substrates [10]:



Phosphorescence kinetics after irradiation is described by the equation [10]

$$L(t) = \frac{k_{gen}[^3P]_0[O_2]}{\tau_r(1/\tau_t - 1/\tau_\Delta)} [e^{-t/\tau_\Delta} - e^{-t/\tau_t}], \quad (3)$$

where k_{gen} is the rate constant of energy transfer from $^3P^*$ to O_2 resulting in 1O_2 generation, τ_r is the 1O_2 radiative lifetime; τ_Δ is the real 1O_2 lifetime, τ_t is the $^3P^*$ lifetime; $[^3P]_0$ is the concentration of triplet photosensitizer molecules just after irradiation; $[O_2]$ is the dioxygen concentration in solution.

For some agents, there was found (references in [10]), that the quenching of photosensitized 1O_2 phosphorescence obeyed the Stern-Volmer equation, which may be simplified to the next form:

$$\tau_\Delta = 1/k_Q C_{max}, \quad (4)$$

where C_{max} the molar concentration of a neat solvent used as a quencher. In different solvents, τ_Δ lies in the range from 10 to 250 μs .

For example, under excitation by Nd:Yag laser, 532 nm, photosensitized singlet oxygen phosphorescence in air-saturated D_2O for the tetra-(*p*-sulphophenyl) porphyrin photosensitizer has the peak achieved in $\sim 10 \mu s$ after the laser flash.

In [10] and references therein, there is proven phosphorescence emitted by solvated 1O_2 molecules in H_2O , by comparing the 1O_2 spectra with those in a gas phase.

The quantum yield φ_{ph} corresponds to an equation

$$\varphi_{ph} = \varphi_\Delta \varphi_r = \varphi_\Delta t_\Delta / t_r, \quad (5)$$

where φ_Δ is the quantum yield of 1O_2 generation by a photosensitizer, φ_r is the 1O_2 phosphorescence quantum yield, t_r is the 1O_2 radiative lifetime. The zero-time

intensity (I_0 , in quanta per second) is calculated by extrapolation of the exponential phosphorescence decay to zero time [10]:

$$I_0 = I_{las} \varphi_\Delta / t_r, \quad (6)$$

where I_{las} is the number of quanta absorbed by a photosensitizer during a laser flash.

The total phosphorescence yield in photosensitizer solutions is significantly less than 100%. The phosphorescence quantum yields in hydrogen-atom containing solvents are 0.02 % and less. The lowest φ_{ph} was found in water, the natural medium of living organisms.

The experimental observation of the 1O_2 fluorescence in the case of a mouse with implanted cancer tumor loaded with Photofrin II and tetra methyl pyridyl porphyrin (TMPP, $10^{-4}M$) [11] in water phosphate buffer. The author of [11], Parker, used the Lexel Model 700L dye laser providing an output wavelength in the range of 600 to 700 nm and fixed at 630 nm, its an average power of was 20 mW. In the TMPP solution, photosensitized phosphorescence at 1270 nm was detected. Parker received $\tau_\Delta = 3.2 \mu s$ and $\tau_r = 2.4 \mu s$.

Review data, received before 1998 on another agents, photosensitizers and target objects one may find once more in [10].

The rate constants of 1O_2 interaction with biomolecules can be measured using chemical 1O_2 traps ([12], [13], [14]) or by quenching 1O_2 phosphorescence (in biological quenchers, such as glycine, tryptophan and so on, k_q is of the order of 10^7 - 10^8 (see references in [10])). Rate constants of the 1O_2 quenching by nucleotides, saccharides and organic acids approximately equal to 10^4 - 10^7 .

So, the quantum yield of singlet oxygen may serve as a criterion of a cell damage.

From the other hand, the oxygen molecule is a very effective quenching agent for positronium formation [16], owing to two unpaired electrons of the ground-state, molecular O_2 exhibits both spinflip and nonspin flip quenching.

Paul, Lee and Celitans [17], [18], [19] concluded, that in a liquid phase, the above mentioned quenched effect (the decreasing of τ_2 in a positronium spectra) is described by the quenching constant [15]

$$\sigma_q v = (\sigma_q v)_{gas} + K_{PsO_2}. \quad (7)$$

The first term in the right side of (7) means the temperature-independent conversion constant in a gas phase (the subscript "gas"). Second one denotes the constant for formation of the PsO_2 : K_{PsO_2} molecule, which equals to $K_{PsO_2}^0 e^{-E/RT}$, where E is an activation energy of a medium molecule, R is the universal gas constant and T is the absolute temperature.

PAL spectra of SDS

SDS, indispensable in the given PDT process, should be aggregated in a micellar phase due to its phase diagram. In [20], the micellar SDS contribution in PS lifetime spectra was studied. Lifetime spectra were analyzed in terms of four components ascribed to *p*-Ps, free e^+ and *o*-Ps in the aqueous (*o*-Ps^{aq}) and organic (*o*-Ps^{org}) sub-phases, in increasing order of the lifetimes τ_i , $i = 1, 2, 3, 4$.

Quantitatively, the molar reaction rate constant (k_{diff}) is given by Smoluchowsky equation

$$k_{diff} = 4\pi DRN_A/1000, \quad (8)$$

where N_A is the Avogadro number, $R = R_{Ps} + R_{core}$ and $D = R_{Ps} + R_{mic}$ are the sum of reaction radii and diffusion coefficients of the reactants respectively (see ref. in [20]). Each diffusion coefficient may be expressed as a function of the hydrodynamic radii of the reactants, R_h : $D = \frac{k_B T}{6\pi\eta R_h}$, where k_B is the Boltzmann constant, T is the absolute temperature, and η is the viscosity of the medium. The trapping processes from water to organic substrates *o*-Ps^{org} are given as [21]:

$$2\gamma \xleftarrow{\lambda_3^0} o - Ps^{aq}(+micelle) \xrightarrow{k(t)C_{mic}} o - Ps^{org} \xrightarrow{\lambda_4^0} 2\gamma. \quad (9)$$

With no addition of NaCl [21], the total intensity associated to *o*-Ps^{aq} is $I_{tot} = I_3 + I_4 = 22 \pm 0.1$ %. By comparison with $I_3^0 = 27$ % for pure water, this is due to some slight inhibition induced by the sulphate polar heads. In the presence of NaCl, I_{tot} is slightly lower and constant, at 21.4 ± 0.1 %.

The appeared results declared that τ_1 and τ_2 were constant at 178 ± 21 and 402 ± 4 ps, respectively. The value of τ_1 is significantly higher than the intrinsic *p*-Ps lifetime in vacuum, $\tau_s = 125$ ps ($\lambda_s = 8$ ns⁻¹, $\lambda_3^0 = 1/1.795$ ns⁻¹ at 303 K, $\tau_4 = 3.87 \pm 0.05$ ns [21]).

Evaluation of spectral contributions of lipopolysaccharides and porins

Consider the next appropriate results. One of Gram (+) bacteria [9], stratum corneum, was studied with PALS to qualify the pore diameters. More specifically, Yucatan miniature pig stratum corneum, separated with heat, was irradiated with polycarbonate (of positron annihilation lifetime 2.103 ± 0.076 ns) due to sandwich scheme with the ²²Na source 0.51 MeV before samples ($15 \times 15 \times 0.5$ mm) isolated by the enveloping aluminium foil.

As it has been generally known, the thickness of an outer layer of Gram (+) bacteria is 15-80 nm (Figure 2). There, pore diameters (from 0.54 to 0.6 nm) in cyclodextrins and polycarbonats were measurable with PALS by means of an application of the well-known Tao-Eldrup formula.

In contrast to stratum corneum, a Gram (-) bacterium has an outer layer of the 10-15 nm thickness, and shows low level of photosensitivity, perhaps due to highly organized outer wall, including murein and a trilamellar structure outside of the peptidoglycan layer [6]. Therefore, to increase the permeability of Gram (-) bacteria, the biological agent such as T₄MPyP is involving in PDT procedures. So, the logarithmic decrease in the overall survival of *E. coli* cells irradiated in the presence of the 8.4 μM T₄MPyP porphyrin is 4.5 after 10 min irradiation.

The reader may find the detailed description of cell killing experiments on Gram (-) *E. coli*, for instance, in [6] and [22].

There, pores may distinct with cylindrical geometry, and expected to be of the same character sizes, as those in a Gram (+) bacterium. And this fair illustration of the Gram (+) bacterial PALS application might be spreadable out onto other bacterial species.

SUM OF COMPONENTS OF EXPECTED SPECTRA OF AQUEOUS *E. COLI*- T₄MPYP

It known that the cytoplasm of an organic cell (a main target of the photo-process [7]) consists of water (75-85%), protein with a mean molecular mass of 10-20%, lipids with a mean molecular mass 2-3%, and other compounds (2%) [10]. Only four amino acids (tryptophan, histidine, methionine and cysteine) actively interact with singlet oxygen, therefore ? protein macromolecule contains 60 residues of active amino acids. The cytoplasm density is known to be about 1 g/sm³. Molar concentrations of water, proteins and lipids in the cytoplasm are 44, 0.004 and 0.04 M respectively. The mean k_q values for active aminoacids is about $2 \cdot 10^7$ M⁻¹s⁻¹. So, the total rate of ¹O₂ deactivation has been calculated as follows [10]: $\sum_g (k_g C_g) = 5 \cdot 10^6$ s⁻¹, where g is a component index. Hence, $\tau_\Delta \approx 200$ ns. The quenching activities of protein amino acids greatly exceed those of lipids and water, water contributes 2-3% into overall quenching. For a cytoplasm membrane $\tau_\Delta \approx 40$ ns [10].

The diffusion length of singlet oxygen in different species is estimated as follows

$$l_\Delta \simeq 1/\sqrt{6D\tau_\Delta}, \quad (10)$$

where D is the ¹O₂ diffusion coefficient presented for the discussed medium in [23]. So, in H₂O, a lipid membrane and a cytoplasm, $l_\Delta = 1900, 2200$ and 90 Å respectively.

Comparison the lifetime spectral contributions of species is shown in Table 1.

NOTES ON APPARATUS REALIZATION

There are several features of the PDT-positron combined irradiation.

TABLE I: Contribution of different agents into lifetime spectra

Medium	τ_3 <i>o</i> -Ps, ns	τ_{Δ} , ns $^1\text{O}_2$
water	1.8 [24]	1...4 [10]
SDS	1.795 [21]	–
T ₄ MPyP	–	3200 [11] (TMPP)
<i>E. coli</i> outer layer	–	50 [10]
<i>E. coli</i> cytoplasmic membrane	–	40 [10]
<i>stratum corneum</i> cytoplasmic membrane	2.1–2.2 [9]	~40 [10]

Typical PALS experiments with living systems ([9], [25] etc) are carried out in frames of the sandwich type configuration of a sample with the positron source (for example, ^{22}Na). However, such a construction does not allow to study a living system in an equivalent to a living media an aqueous aggregation, and also causes troubles in delivery of light from a laser or a lamp. An alternative bulk construction schemes of a samples in ampules, applicable for liquids on the whole, are described, for instance, in [26].

CONCLUSIONS

The above arguments have proved the possibility of control with positron spectroscopy of states of outer membrane layers in bacterial cells irradiated in PDT. The next conclusions have been made.

1. As the lifetime order of magnitudes for *o*-Ps and $^1\text{O}_2$ is comparable and different for an each medium (with the exception of water), hence, one may measure the Ps lifetime during the photoinactivation by singlet oxygen during the PDT procedure.

2. The scenario involving the reactions of singlet oxygen of PDT into PALS might be directly tested on the Gram (+) bacterial system [9] and spread out onto Gram (-) bacteria.

3. There is announced that for both Gram (+) and Gram (-) bacteria, the quenching effect of singlet oxygen will be impaired depending of the cell death.

Acknowledgments

The author thanks M. G. Strakhovskaya for her introduction in PDT and joint collaboration. In this text,

the review by Lingjie Zhao "Free Radical and Radiation Biology" (Graduate Program 2001) was used.

References

- [1] Maisch T, Szeimies R–M, Jori G and Abels C 2004 *Photochem. Photobiol. Sci.* **3** 907
- [2] Kearns D R 1971 *Chemical Reviews* **71** 395
- [3] Briviba K, Klotz L O and Sies H 1997 *Biol. Chem.* **378** 1259
- [4] Jean Y C, Chen H, Liu G and Gadzia J E 2007 *Radiat. Phys. and Chem.* **76** 70
- [5] Jean Y C, Li Y, Liu G, Chen H, Zhang J and Gadzia J E 2006 *Applied Surface Science* **252** 3166
- [6] Villanueva A, Bertaloni G and Jori G 1995 *J. Braz. Chem. Soc.* **71** 123
- [7] Bertolini G, Rossi F, Valduga G, Jori G and Van Lier J 2006 *FEMS Microbiology Letters* **71** 149
- [8] Jean Y C and Hancock A J 1982 *J. Chem. Phys* **77** 5836
- [9] Itoh Y, Shimazu A, Sadzuka Y, Sonobe T, Itai S 2008 *Int. J. of Pharmaceutics* **358** 91
- [10] Krasnovsky A A, Jr. 1998 *Membr. Cell Biol.* **12** 665
- [11] Parker J G 1987 *IEEE Circuits and Devices* *??gazine* 10
- [12] Foote C 1976 *Free Radicals in Biology* (New York ? San Francisco ? London: Academic Press)
- [13] Lindig B A and Rodgers M A J 1981 *Photochem. Photobiol.* **33** 627
- [14] Matheson I B C, Etheridge R D, Kratoch N R and Li J 1975 *J. Photochem. Photobiol.* **21** 165
- [15] Goldanskii V I 1968 *Fizicheskaya khimia positrona i positronia* (Moscow: Nauka)
- [16] Ferrell R A 1958 *Phys. Rev.* **110** 1355
- [17] Paul D A 1959 *Canad. J. Phys.* **37** 1059
- [18] Lee J, Celitans G J 1965 *J. Chem. Phys.* **42** 437
- [19] Lee J, Celitans G J 1966 *J. Chem. Phys.* **44** 2506
- [20] Bockstahl F, Pachoud E, Duplâtre G and Billard I 2000 *Chemical Physics* **256** 307
- [21] Bockstahl F, Duplâtre G 1999 *Phys. Chem. Chem. Phys.* **1** 2767
- [22] Minnock A, Vernon D I, Schofield J, Griffiths J, Parish J H and Brown S B 2000 *Antimicrob Agents Chemother.* **44** 522
- [23] Fu Y and Kanofsky J R, Jr. 1995 *Photochem. Photobiol.* **62** 692
- [24] Duplâtre G, Talamoni J, Abbé J Ch and Haessler A 1984 *Radiat. Phys. Chem.* **23** 531
- [25] Uchiyama Y, Ito K, Li H-L, Ujihara Y, Jean Y C 1996, *J. of Radioanalyt. and Nucl. Chem.* **211** 111
- [26] Duplâtre G, Al-Shukri L M and Haessler A 1980 *J. of Radioanalytical Chem.* **55** 199