

Classical description of quantum randomness using stochastic gauge systems

Michel Feldmann *

Abstract

We present a classical probability model appropriate to the description of quantum randomness. This tool, that we have called *stochastic gauge system*, constitutes a contextual scheme in which the Kolmogorov probability space depends upon the experimental setup, in accordance with quantum mechanics. Therefore, the probability space behaves like a gauge parameter. We discuss the technical issues of this theory and apply the concept to classically emulate quantum entangled states and even ‘super-quantum’ systems. We exhibit bipartite examples leading to maximum violation of Bell-CHSH inequalities like EPR pairs or exceeding the Tsirelson bound like PR-boxes, as well as tripartite cases simulating GHZ or W-states. We address also the question of partially correlated systems and multipartite entanglements. In this model, the classical equivalent of the entanglement entropy is identified with the Kullback-Leibler divergence. Hence, we propose a natural generalisation of this function to multipartite systems, leading to a simple evaluation of the degree of entanglement and determining the bounds of maximum entanglement.

PACS 03.65.Ud (Entanglement and quantum non-locality)

Contents

1	Introduction	2
2	Simulation of bipartite quantum systems	3
2.1	Bell-type systems	3
2.1.1	Contextuality and Bell’s inequalities	3
2.1.2	Partial measurement	4
2.1.3	Total correlation	5
2.2	Classical evolution of a Bell-type system	5
2.2.1	Ignition states and projection function	5
2.2.2	Gauge probability distributions	7
2.2.3	Classical collapse	7
2.2.4	General 2-setting totally correlated systems	9
2.3	Simulation of the EPR spin experiment	9
2.3.1	Two settings	10
2.3.2	Three settings and Bell’s inequality violation	10
2.3.3	Four settings and CHSH inequality violation	10
2.3.4	Regular settings	11
2.3.5	Continuous settings	13
3	Simulation of multipartite entangled systems	13
3.1	Classical analogues of quantum systems	13
3.1.1	Local consistency	14
3.1.2	Degrees of freedom	14
3.2	Randomness and entropy	15
3.2.1	Entropy of measurement	15
3.2.2	Entropy of entanglement	15
3.2.3	Multipartite entanglements	17
3.2.4	Entanglement scheme	19
3.2.5	Information inequalities	19
3.3	Classical multipartite collapse	20
3.3.1	Collapse mechanism	20
3.3.2	One-step collapse	22

*Electronic adress: michel.feldmann@polytechnique.org

3.3.3	Multi-step collapse	23
3.4	Typical examples	23
3.4.1	One-region systems	24
3.4.2	Bipartite systems (separable, entangled, PR-box)	24
3.4.3	Tripartite entangled systems (W and GHZ states)	25
4	Conclusion	31

1 Introduction

While quantum mechanics is one of the pillars of modern physics, its very foundations remain controversial. A crucial point is that probability in quantum theory obeys different rules than do the classical probabilities, as emphasized by Richard Feynman in the 1950's. Later, E. T. Jaynes noted that quantum probabilities are strikingly similar to the Bayesian probabilities. One controversial issue concerns 'local realism' highlighted by the well-known EPR paradox [1, 2]: Nowadays, it is widely accepted that local realism does not hold in microphysics. This 'spooky' property is derived from a famous theorem by Bell [3, 4] complemented by Kochen and Specker [5] and supported by a number of experimental verifications [6]. As a result, a huge literature has come to light, aiming to reconcile this break of local realism, if not with common sense, at least with a form of logic. Currently, a number of different approaches are competing, often at the interface between physics and philosophy (see for instance references [7–12]). However, following Jaynes [13] and among several authors, [14–23] we have warned about a flaw in the proof of Bell's theorem [24]. In order to falsify Bell's claims, we have exhibited a very simple classical counterexample with just three dice [25] based on the present model. In our opinion, this proves that the concept of quantum non-locality remains actually unfounded. Conversely, if quantum mechanics is really compatible with classical local realism, quantum phenomena can be classically emulated, at least to some extent. That is the purpose of the present paper.

We have constructed classical tools, namely, consistent parametric probability distributions, that we have called *stochastic gauge systems*. This model constitutes a contextual probability scheme in which the Kolmogorov probability space depends upon the experimental setup, in accordance with quantum mechanics. Hence, the recourse to non-locality is not necessary. By contrast, the flaw in Bell's theorem is to postulate the existence of an absolute probability space whatever the settings, in opposition to both theoretical and experimental evidences. Beyond the EPR paradox, we suggest that classical simulation of quantum phenomena may provide some insights in several fundamental questions.

Our underlying picture is the following: We propose that the classical equivalent of a stationary quantum system is a spatially extended object in dynamic equilibrium. Therefore, in the absence of perturbation, its evolution is completely deterministic and there is no room for randomness. This analogy suggests that quantum collapse can be compared with a classical break of equilibrium. The stability of a classical equilibrium is similar to the resistance to decoherence of a quantum state. Based on Bell's theorem, it is generally assumed that the notion of entanglement is strictly of quantum nature [26, 27]. By contrast, we argue that the classical analogue of quantum entanglement is the property of *contextual dependence*, better underscored when the break of equilibrium of a spatially extended object can be triggered from two distant locations separated by a boundary. We have found that this boundary behaves like an ignition front where the *contextual gauge probability trial* is performed. Therefore, the *contextual entropy* should be located on this boundary. This resembles the holographic hypothesis suggested by 't Hooft [28] and Susskind [29] in black hole physics.

We will first deal with the technical issues of our gauge probability model before addressing the question of classical simulation of quantum entangled systems. We will then describe the classical Bell-type model and will exhibit a number of examples with several settings and total correlation between two regions. Specially, we will compute exactly some typical bipartite configurations leading to maximum violation of Bell or CHSH inequalities. We will next extend the model to general locally consistent systems, including non quantum devices like PR-boxes and multipartite systems like GHZ and W tripartite states. We also deal with partial correlation and define the concept of classical entanglement entropy. The entropy of a classical entangled system is not extensive contrary to the entropy of a separable system. In thermodynamics, this supports the existence of long range interactions [30, 31]. In this context, the classical entanglement entropy should be identified with the Kullback-Leibler divergence. We propose natural generalizations, including the definition of classical 'multipartite entanglement entropies' and a concept of 'entanglement scheme' to characterize complex multipartite entanglement.

2 Bipartite systems

In classical physics, suppose that a spatially extended object is in equilibrium and that this equilibrium can be broken from two distant locations. Equilibrium is a global concept: Break of equilibrium from one location signifies break of equilibrium for the whole object and consequently break of equilibrium at the second location as well. This means *instantaneous correlation* at a distance between the two locations. Nevertheless, this does not imply *instantaneous physical effect* since the relaxation towards a new equilibrium will require a delay. A trivial picture is the evolution of a stationary huge soap bubble (possibly in dynamic equilibrium, for instance in uniform rotation). We compare this phenomenon with what happens in quantum EPR experiment. For this purpose, we identify the spatial extension of the quantum system with the locus of all possible regions of measurement. The quantum system in unitary evolution is considered as a stationary object in dynamic equilibrium. The measurement of one parameter in one location interrupts the unitary evolution or breaks the equilibrium. The final state is a new dynamic equilibrium. We claim that the behaviour of similar extended objects in equilibrium may be viewed as a form of *classical entanglement*.

2.1 Bell-type systems

Consider a pair of quantum *entangled* entities $\{\mathcal{E}_0, \mathcal{E}_1\}$ (e.g., particles). The pair, or its global wave function, will be compared with a *classical extended object* stretching out on two space regions \mathcal{R}_0 and \mathcal{R}_1 . We will first focus on totally correlated entities (as defined in Sec. 2.1.3) referred to as Bell-type systems.

In region \mathcal{R}_0 , we suppose that an observer \mathcal{O}_0 selects freely a setting u_0 (e.g., an angle of polarization), element of a given set Θ_0 , and measures on \mathcal{E}_0 a random dichotomic observable x_0 , element of a dyadic set $\mathbf{X} = \{0, 1\}$, (or a spin $s_0 = 2x_0 - 1$). This measurement breaks the dynamic equilibrium of the whole extended object. Similarly, in region \mathcal{R}_1 , a second observer \mathcal{O}_1 selects independently a setting $u_1 \in \Theta_1$ and measures on \mathcal{E}_1 a random dichotomic observable $x_1 \in \mathbf{X}$, (or a spin $s_1 = 1 - 2x_1$). For the sake of simplicity, we will assume that $\Theta_0 = \Theta_1 = \Theta$.

The spins are well defined physical observables and thus the interchange of the numbering in the pair $\{\mathcal{E}_0, \mathcal{E}_1\}$ interchanges the binary digits ‘0’ and ‘1’. More generally, we will call *bit reversal* the exchange of ‘0’ and ‘1’ in \mathbf{X} . Due to symmetry between \mathcal{E}_0 and \mathcal{E}_1 , physical Bell-type systems are often invariant by bit reversal.

Note that for an ensemble of runs, the binary strings $\{x_0\}$ and $\{x_1\}$ can be viewed as digital signals emitted from two distant ports \mathcal{R}_0 and \mathcal{R}_1 .

2.1.1 Contextuality and Bell’s inequalities

As soon as the system $\{\mathcal{E}_0, \mathcal{E}_1\}$ interacts with the settings u_0 or/and u_1 , *instantaneously*, the initial equilibrium is broken. Therefore, there is some conditional probability $P(x_0; x_1 | u_0; u_1)$ that \mathcal{O}_0 will observe (after a delay) the outcome x_0 and \mathcal{O}_1 the outcome x_1 . In fact, the function $P(x_0; x_1 | u_0; u_1)$ (or $P(\mathbf{x} | \mathbf{u})$ in short) describes the symmetry of the extended object. It will be convenient to have a special name for the entries $(\mathbf{x} | \mathbf{u})$.

Definition (Target). *We will name global target or simply target each entry $(\mathbf{x} | \mathbf{u})$. Let $\mathfrak{T} = \{(\mathbf{x} | \mathbf{u})\}$ be the set of all possible targets.*

For the purpose in hand, we can identify \mathfrak{T} with a *coarse-grained description* of the potential *future* equilibrium of the extended object. According to Bell’s theorem [3], Bell’s inequalities hold, but only when the theorem is valid, i.e., in *non contextual* systems. By contrast, if the conditional probability $P(x_0; x_1 | u_0; u_1)$ depends upon the settings u_0 and u_1 , Bell’s inequalities can be violated.

Actually, there is a number of formulations of Bell’s inequalities. We have shown [24] that basically, these inequalities are nothing else than the well known triangle inequalities with respect to a convenient metrics, namely the *Hamming distance* [32] $d_H(\{x_0\}, \{x_1\})$ between the digital signals $\{x_0\}$ and $\{x_1\}$ emitted from the two ports \mathcal{R}_0 and \mathcal{R}_1 for an ensemble of runs. A similar metrics was proposed by Santos [33] using Boolean logic. Define the mean Hamming distance for one run as

$$d(u_0, u_1) \stackrel{(\text{def})}{=} \mathbf{E}[d_H(x_0, x_1)],$$

where $\mathbf{E}[\cdot]$ stands for the expectation value with respect to $P(\mathbf{x} | \mathbf{u})$. Clearly, we have

$$d(u_0, u_1) = P(1; 0 | u_0; u_1) + P(0; 1 | u_0; u_1) \leq 1.$$

Let $\theta_0, \theta_1, \theta_2 \in \Theta$ be three different settings: Then, Bell’s inequalities simply read:

$$d(\theta_0, \theta_2) \leq d(\theta_0, \theta_1) + d(\theta_1, \theta_2). \quad (1)$$

In other words, Bell's inequalities imply the existence of a metrics, $d(\theta, \theta')$ between the elements θ, θ' of Θ . The original Bell's inequalities make use of the expectation value $M(u_0, u_1) = \mathbf{E}[s_0 \cdot s_1]$. In non contextual systems [24] (and also in contextual systems invariant by bit reversal), we have,

$$d(u_0, u_1) = (1/2)[1 + M(u_0, u_1)]. \quad (2)$$

Thus, starting from the following formulation of the triangle inequality,

$$|d(\theta_0, \theta_1) - d(\theta_0, \theta_2)| \leq d(\theta_1, \theta_2),$$

we obtain the well known original Bell's inequality,

$$|M(\theta_0, \theta_1) - M(\theta_0, \theta_2)| \leq 1 + M(\theta_1, \theta_2). \quad (3)$$

Another popular formulation involving four different settings, $\theta_0, \theta_1, \theta_2$ and θ_3 , is the CHSH inequality [34], which reads,

$$|M(\theta_0, \theta_1) - M(\theta_0, \theta_3)| + M(\theta_3, \theta_1) + M(\theta_3, \theta_2) \leq 2. \quad (4)$$

In non contextual systems, these inequalities are again a simple consequence of the triangle inequality [33], given that all mean Hamming distances $d(\theta, \theta')$ are less or equal to unity.

These inequalities will hold true *for any settings* if it possible to define a triangle or a quadrilateral in the Hamming metrics *at one and the same time*, i.e., if it is possible to prepare different experiments with different settings governed by the same probability space. Clearly, this implies that $P(x_0; x_1|u_0; u_1)$ does not depend on the contextual parameters u_0 and u_1 . In this classical framework, violation of Bell's inequalities is simply a criterion of *contextual dependency* of the probability space and does not in the least imply instantaneous effect at a distance nor violation of *local realism*.

2.1.2 Partial measurement, nonsignaling correlations, local consistency and causal horizon

Quantum mechanical collapse may be viewed as a break of equilibrium of the wave function. When the particle \mathcal{E}_1 interacts with the setting u_1 in region \mathcal{R}_1 , the whole system decoheres and the pair of particles is no more entangled. Then, possibly after a delay, the wave function splits into two local wave functions. However, in region \mathcal{R}_0 , this splitting is not observable and the observer \mathcal{O}_0 is not aware of what happens in \mathcal{R}_1 ; he independently selects his own setting u_0 and proceeds to the measurement of x_0 , while the local marginal probability $\text{Prob}(x_0|u_0)$ of x_0 given u_0 is not affected, as long as both the setting u_1 and the outcome x_1 are ignored. In other words, the local probability $\text{Prob}(x_0|u_0)$ depends on the knowledge of the observer \mathcal{O}_0 and is thus an observer-dependent concept. This property characterizes quantum *partial measurements* in two different locations, when the global Hilbert space is regarded as a tensor product of two partial subspaces. This means that partial measurement is not a way to communicate and is thus compatible with space-like separation of the two regions. Adopting the point of view of the observers, we will call *local consistency* this *nonsignaling* correlation [35–37]. This property is also encountered in a number of situations in relativity, cosmology and black hole physics, where quantum entanglement holds between two regions separated by a so called *causal horizon*, and even in some classical laboratory analogues of black holes [38, 39]. On the contrary, it is rarely met in usual classical physics, e.g., a soap bubble is not locally consistent. Nevertheless, as we shall show in the following sections, it is quite easy to design *ad hoc* classical devices, like dice games [25], which fulfil this criterion.

Consider a system with K discrete settings, $\Theta = \{\theta_0, \theta_1, \dots, \theta_{K-1}\}$. When local consistency holds, the partial probability in one region, say \mathcal{R}_0 , does not depends on the setting u_1 in the second region \mathcal{R}_1 . Let $u_0 = \theta_k$ be selected in region \mathcal{R}_0 we have:

$$\forall u_1 \in \Theta : \text{Prob}(x_0|u_0 = \theta_k) = P(x_0; 0|\theta_k, u_1) + P(x_0; 1|\theta_k; u_1) \quad (5)$$

and similarly for the second port. When accounting for the normalization relation

$$\text{Prob}(x_i = 0|u_i = \theta_k) + \text{Prob}(x_i = 1|u_i = \theta_k) = 1$$

for $i = 0, 1$, Eq.(5) provides a number of $2(K - 1)$ relations between the target probabilities $P(\mathbf{x}|\mathbf{u})$ for each setting θ_k and $2K(K - 1)$ for all settings (Table 1).

For a continuous ensemble of settings, when $P(x_0; x_1|u_0; u_1)$ is continuous and differentiable, we have for $i = 0, 1$,

$$\frac{\partial \text{Prob}(x_i|u_i)}{\partial u_{1-i}} = 0. \quad (6)$$

2.1.3 Total correlation

In Bell-type experiment the outcomes $x_0 = x_1$ are identical when the same setting $u_0 = u_1 = \theta \in \Theta$ is selected in the two regions. Then:

$$P(1; 0|\theta; \theta) = P(0; 1|\theta; \theta) = 0. \quad (7)$$

This provides a number of 2 additional relations between the target probabilities $P(\mathbf{x}|\mathbf{u})$ for each setting θ and $2K$ for all settings (Table 1). We will call this property *total correlation*.

For *totally correlated and locally consistent systems*, it follows from Eqs. (5), (7) that for $\xi \in \mathbf{X}$ and $\theta \in \Theta$

$$\text{Prob}(x_0 = \xi|u_0 = \theta) = \text{Prob}(x_1 = \xi|u_1 = \theta) = P(\xi; \xi|\theta; \theta). \quad (8)$$

Note that ‘total correlation’ is different from ‘maximal entanglement’. Totally correlated and maximally entangled bipartite systems are generally referred to as ‘EPR pairs’. We will meet later examples of bipartite maximally entangled systems which are not totally correlated (Table 20 and 30). We will discuss the concept of maximum entanglement in Sec. 3.2.5.

2.2 Classical evolution of a Bell-type system

As soon as the system interacts with the settings u_0 and/or u_1 , the initial equilibrium is broken and the system relaxes towards a future equilibrium. Violation of Bell’s inequalities proved that a simple random trial is unable to describe the contextual correlations $P(x_0; x_1|u_0; u_1)$. Hence, we address the following issue: Is it nevertheless possible to implement the joint conditional probability $P(\mathbf{x}|\mathbf{u})$ in classical physics? In other words, is it possible to classically simulate a quantum collapse?

We claim that the answer is definitively *yes*: We are going to show that any *locally consistent* and *totally correlated* Bell-type system relaxation can be achieved by cascading (1) propagation with finite velocity towards a so-called *ignition point*, (2) random trial at this ignition point and (3) backwards propagation, again with finite velocity. We will next identify the locus of all possible ignition points with a *causal horizon*.

2.2.1 Ignition states, ignition set and projection function

In this paragraph, we will construct a convenient outcome set, that we will call *ignition set*, which meets the requirements of contextual systems. We will call its elements *ignition states*. Let us first consider discrete systems.

Discrete settings Consider a finite ensemble of K settings, $\theta_0, \theta_1, \dots, \theta_{K-1}$. In Sec. 2.1.1, we have compared the set of targets $\mathfrak{T} = \{\mathbf{x}|\mathbf{u}\}$ with a coarse-grained description of the potential future equilibrium of the extended object. Now, we will similarly construct a convenient *fine-grained description*¹ as follows: In region \mathcal{R}_i define the *local target* $(x_i|u_i)$. When $x_i = \xi$ and $u_i = \theta_k$, we write $(x_i|u_i) = (\xi|\theta_k)_i$. Each local target, e.g. $(\xi|\theta_k)_0$ in region \mathcal{R}_0 , can be regarded as the union of all (global) targets compatible with $x_0 = \xi$ and $u_0 = \theta_k$.

$$(x_0|u_0) = (\xi|\theta_k)_0 = \bigcup_{x_1=0}^1 \bigcup_{u_1=0}^{K-1} (x_0; x_1|u_0; u_1).$$

Conversely

$$(x_0; x_1|u_0; u_1) = (x_0|u_0) \cap (x_1|u_1).$$

According to Eq.(8), we have $\text{Prob}(\xi|\theta_k) = P(\xi; \xi|\theta_k; \theta_k)$. When total correlation holds, the marginal probabilities are equal in the two regions and the local targets $(\xi|\theta_k)_0$ and $(\xi|\theta_k)_1$ are identical. Let $\mathfrak{T}_{\text{loc}} = \{(\xi|\theta_k)\}$ be the set of local targets (relevant for both regions). The cardinality of $\mathfrak{T}_{\text{loc}}$ is $2K$.

Let $\mathcal{J} = \{j\}$ be a set of integers. We aim to construct a sequence of $\text{card}(\mathcal{J})$ fine grains. Define an application of $\Theta \times \mathcal{J} \rightarrow \mathbf{X} = \{0, 1\}$ or rather an application $\Pi(k, j): \llbracket 0, K-1 \rrbracket \times \mathcal{J} \rightarrow \mathbf{X} = \{0, 1\}$

$$\xi = \Pi(k, j). \quad (9)$$

We will call *projection function* the function $\Pi(k, j)$. Now, we construct the sequence of $\text{card}(\mathcal{J})$ fine grains $\hat{\lambda}_j$ by the following intersection of local targets:

$$j \in \mathcal{J} \mapsto \hat{\lambda}_j \stackrel{(\text{def})}{=} \bigcap_{k=0}^{K-1} \left(\Pi(k, j)|\theta_k \right). \quad (10)$$

¹This designation describes a method but does not fit very well, because the number of useful fine grains is often less than the number of coarse grains!

Therefore, the maximum number of distinct fine grains is 2^K and then $\mathfrak{J} = \llbracket 0, 2^K - 1 \rrbracket$. Let

$$j = \sum_{k=0}^{K-1} j_k 2^k$$

be the binary expansion of an integer j , ($0 \leq j \leq 2^K - 1$) and j_k the coefficient of 2^k . A convenient definition of the projection function is

$$\Pi(k, j) \stackrel{(\text{def})}{=} j_k. \quad (11)$$

For reasons explained in Sec. 2.2.3, we will call *ignition state* each fine grain $\hat{\lambda}_j$. Clearly, the ignition states included in a given global target $(x_0; x_1 | u_0; u_1)$ are also included in the local targets $(x_0 | u_0)$ and $(x_1 | u_1)$ and conversely. Therefore, we have

$$\hat{\lambda}_j \in (x_0; x_1 | u_0; u_1) \Leftrightarrow \hat{\lambda}_j \in (x_0 | u_0) \cap (x_1 | u_1).$$

It is also convenient to define the index subset,

$$\mathbf{I}(\xi, k) \stackrel{(\text{def})}{=} \{j \mid \Pi(k, j) = \xi\}. \quad (12)$$

Clearly, we have $\mathbf{I}(0, k) + \mathbf{I}(1, k) = \mathfrak{J}$. For simplicity, we will often identify k with θ_k , j with $\hat{\lambda}_j$ and use the same symbol,

$$\mathbf{I}(\xi, \theta_k) \stackrel{(\text{def})}{=} \mathbf{I}(\xi, k) ; \Pi(\theta_k, \lambda_j) \stackrel{(\text{def})}{=} \Pi(k, j).$$

In addition define the index subset,

$$\mathbf{J}(\mathbf{x}, \mathbf{u}) \stackrel{(\text{def})}{=} \{j \mid \hat{\lambda}_j \in (\mathbf{x}, \mathbf{u})\} = \mathbf{I}(x_0, u_0) \cap \mathbf{I}(x_1, u_1). \quad (13)$$

Finally, let $\mathbf{\Lambda}$ be the set of all ignition states,

$$\mathbf{\Lambda} \stackrel{(\text{def})}{=} \{\hat{\lambda}_0, \hat{\lambda}_1, \dots, \hat{\lambda}_{\text{card}(\mathfrak{J})-1}\}.$$

We will call this ensemble *ignition set*. While the maximum value of $\text{card}(\mathfrak{J})$ is 2^K , we shall see that only $2K$ ignition states are necessary and thus, this set is strongly redundant for large K . A more convenient set \mathfrak{J} of cardinality just $2K$ is the set of ‘double-plateau’ integers \mathbf{D}_K defined as the integers of $\llbracket 0, 2^K - 1 \rrbracket$ whose binary expansion is a chain of identical bits except for a maximum of one jump. For instance, $\mathbf{D}_3 = \{\overline{011}, \overline{111}, \overline{110}, \overline{100}, \overline{000}, \overline{001}\} = \{3, 7, 6, 4, 0, 1\}$ and $\mathbf{D}_4 = \{\overline{0011}, \overline{0111}, \overline{1111}, \overline{1110}, \overline{1100}, \overline{1000}, \overline{0000}, \overline{0001}\} = \{3, 7, 15, 14, 12, 8, 0, 1\}$. For reasons explained in Sec. 2.3.5 this particular order in \mathbf{D}_K , beginning with $K/2$ zeros for K even or $(K-1)/2$ zeros for K odd, will be referred to as the *trigonometric order*. For $r \in \llbracket 0, 2K - 1 \rrbracket$, this order defines a *double-plateau* function $D_K(r)$,

$$D_K : \llbracket 0, 2K - 1 \rrbracket \rightarrow \llbracket 0, 2^K - 1 \rrbracket : r \mapsto j = D_K(r) \quad (14)$$

For example, we have $D_4(0) = 3$ or $D_4(6) = 0$.

Note that the ignition states $\hat{\lambda}_j$ have no definite probability for $K > 1$. Therefore, in general, they cannot be regarded as hidden variables. Nevertheless, we will define in the next section (Sec. 2.2.2) a number of *gauge* probability distributions $g_k(\hat{\lambda}_j)$ on $\mathbf{\Lambda}$, each compatible with $P(\mathbf{x}, \mathbf{u})$. For instance, for $K = 3$, only 6 ignition states are necessary: The ignition states can be pictured by dice sides, while each probability distribution corresponds to a particular biased die [25].

Continuous settings The discrete projection function Eq. (9) is not appropriate for continuous settings. We have thus to guess a convenient *projection function* $\Pi(\theta, \lambda) : \Theta \times \mathbb{R} \rightarrow \mathbf{X} = \{0, 1\}$. The fine-grained description for continuous settings is next defined by:

$$\lambda \in \mathbb{R} \mapsto \hat{\lambda} = \bigcap_{\theta \in \Theta} (\Pi(\theta, \lambda) | \theta). \quad (15)$$

We will use such a projection function in Sec. 2.3.5. More generally, the structure of the ignition set should reflect the system symmetry.

Number of settings	K
Number of targets $(\mathbf{x} \mathbf{u})$	$4K^2$
Number of local targets $\text{card}(\mathfrak{T}_{\text{loc}})$	$2K$
Constraints of normalization	K^2
Constraints of local consistency	$2K(K-1)$
Constraints of total correlation	$2K$
Degrees of freedom	K^2
Number of gauge distributions $g_k(\lambda_j)$	K
Rank of the gauge equations	$2K$

Table 1: Degrees of freedom in general bipartite Bell-type systems with K possible settings. Due to total correlation, the K^2 degrees of freedom are shared between the two regions

2.2.2 Gauge probability distributions

For ease of exposition and simplicity, we will focus on finite systems with $K \in \mathbb{N}$ different settings.

We have emphasized that the ignition states $\hat{\lambda}_j$ have no definite probability. Now, we are going to define a number of K parametric probability distributions $g_k(\lambda_j)$ on the ignition set $\mathbf{\Lambda}$, labelled k from 0 to $K-1$. Of course, each distribution has to be compatible with the target probability $P(\mathbf{x}|\mathbf{u})$. We regard each target as an union of ignition states. Therefore, the probability of each target $P(\mathbf{x}|\mathbf{u})$ will be identified with the sum of the probabilities of its ignition states. There are a number of consistent possibilities and two families of solutions.

A first family is obtained from region \mathcal{R}_0 : Let $u_0 = \theta_k$ and let p_{kj} be the unknown probability of $\hat{\lambda}_j$ in the probability distribution labelled k . For each setting $u_1 \in \Theta$ and each outcome pair x_0, x_1 , we write one compatibility condition:

$$\forall u_1, \forall x_0, \forall x_1 : \sum_{j \in \mathbf{J}(\mathbf{x}, \mathbf{u})} p_{kj} = P(x_0; x_1 | u_0 = \theta_k; u_1) \quad (16)$$

where $\mathbf{J}(\mathbf{x}, \mathbf{u})$ is defined by Eq.(13). From Eq.(12), we derive

$$\mathbf{J}(x_0, x_1, u_0, u_1) = \mathbf{I}(x_0, u_0) \cap \mathbf{I}(x_1, u_1).$$

We have clearly

$$\forall u_1 \in \Theta : \mathbf{I}(x_0, u_0) = \mathbf{I}(x_0, u_0) \cap [\mathbf{I}(0, u_1) \cup \mathbf{I}(1, u_1)],$$

and thus, local consistency, Eq.(5), is automatically encoded. Similarly, we have

$$\mathbf{J}(0, 1, \theta_k, \theta_k) = \mathbf{I}(0, \theta_k) \cap \mathbf{I}(1, \theta_k) = \emptyset,$$

and thus, total correlation is also secured.

Accounting for local consistency and total correlation, each parametric probability distribution labelled k is defined by a linear system of $4K - 2(K-1) - 2 = 2K$ independent equations and $\text{card}(\mathfrak{T})$ unknowns. There is generally a set of positive solutions. A second family of solutions could be obtained similarly from region \mathcal{R}_1 , by interchanging the indices 0 and 1. Select one solution p_{kj} and define

$$g_k(\hat{\lambda}_j) \stackrel{(\text{def})}{=} p_{kj} \quad (17)$$

Clearly, we have for any u_1 :

$$\sum_{j \in \mathfrak{T}} g_k(\hat{\lambda}_j) = \sum_{x_0=0}^1 \sum_{x_1=0}^1 P(x_0; x_1 | \theta_k; u_1) = 1.$$

We will call the function $g_k(\hat{\lambda}_j)$ *gauge probability distribution*.

2.2.3 Classical collapse

We aim to describe the random system relaxation by use of the ignition set $\mathbf{\Lambda} = \{\hat{\lambda}\}$ and the K probability distributions $g_k(\hat{\lambda})$. The process is the following: In each region, the observer selects his own setting u_0 or u_1 . These settings are transmitted with finite velocity towards a so-called *ignition point* \mathcal{I} . Only *one setting* u_0 or u_1 , labelled θ_{igni} is kept, e.g., the first received setting. Now, at \mathcal{I} , we perform a trial in the ignition set $\mathbf{\Lambda}$ using the only probability distribution $g_{\text{igni}}(\hat{\lambda}_j)$ to draw a single ignition state $\hat{\lambda}_{j_{\text{igni}}}$. Let $j_{u_0} = \Pi(u_0, \hat{\lambda}_{j_{\text{igni}}})$ and $j_{u_1} = \Pi(u_1, \hat{\lambda}_{j_{\text{igni}}})$. These

coefficients, j_{u_0} and j_{u_1} , are transmitted with finite velocity towards the end regions \mathcal{R}_0 and \mathcal{R}_1 respectively, where the final outcomes are $x_0 = j_{u_0} = \Pi(u_0, \hat{\lambda}_{j_{\text{igni}}})$ and $x_1 = j_{u_1} = \Pi(u_1, \hat{\lambda}_{j_{\text{igni}}})$.

Conversely, to compute the probability $P(\mathbf{x}|\mathbf{u})$ we have to collect all ignition states $\hat{\lambda}_j$ within the target $(\mathbf{x}|\mathbf{u})$:

$$P(x_0; x_1 | u_0; u_1) = \sum_{\lambda_j \in (\mathbf{x}|\mathbf{u})} g_{\text{igni}}(\hat{\lambda}_j) \quad (18)$$

where the label of g_{igni} is such that θ_{igni} is any one of the two settings, u_0 or u_1 , and $\lambda_j \in (x_0; x_1 | u_0; u_1)$ means that $x_0 = \Pi(u_0, \hat{\lambda}_j)$ and $x_1 = \Pi(u_1, \hat{\lambda}_j)$.

Now the proof is a straightforward verification of the identity of Eq.(16) and Eq.(18). By construction, Eq.(7) is automatically satisfied and the choice of u_0 or u_1 for θ_{igni} to select the distribution g_{igni} at the ignition point \mathcal{I} does not affect the overall probability $P(x_0; x_1 | u_0; u_1)$.

Ignition front and causal horizon Note first that the process implies the existence of an ignition point located between the two end regions. Then, the locus of all possible ignition points draws a form of frontier between \mathcal{R}_0 and \mathcal{R}_1 , whereas no transfrontier communication is possible between the two regions. Such an *ignition front* is not recognized in quantum mechanical collapse since this process is generally assumed to be superluminal and non local [40], but it could be *light cones*, *light-sheets* [41] or *null surfaces*. In black holes physics, it looks like the *event horizon*. We will thus identify this ignition front with a *causal horizon* [42].

Gauge probability distributions There are two possible probability distributions for the trial at \mathcal{I} , defined by the settings of the two end regions. The real process, if any, is not observable and the choice between these two distributions is indifferent. Therefore, the probability distributions $g_k(\hat{\lambda})$ behave like global *gauge parameters*: The choice of one particular distribution correspond to a gauge selection at \mathcal{I} and this choice does not affect the observable result. This *gauge invariance* is fundamental. In our opinion, it is the core of the EPR paradox. We will call *stochastic gauge system* the ensemble of the ignition set $\mathbf{\Lambda} = \{\hat{\lambda}\}$ and a number of parametric distributions $g_k(\hat{\lambda})$.

Analogy with the holographic hypothesis The parametric trial is performed on the ignition front, i.e., on the causal horizon and not in regions \mathcal{R}_0 and \mathcal{R}_1 . This meets a suggestion by 't Hooft and Susskind in black hole physics, namely, the *holographic hypothesis* [28, 29, 41], that space of quantum states in a region must be associated with the two-dimensional boundary rather than the volume of the system. Note that this assumption might be checked by experiments on entangled entities more easily than in black holes.

Consistency conditions Due to gauge invariance, the parametric probability distributions $g_k(\lambda_j)$ are not independent [24]. Suppose that the observers \mathcal{O}_0 and \mathcal{O}_1 select respectively the settings u_0 and $u_1 \in \Theta$. Let g_{u_0} and g_{u_1} be the gauge distributions associated with u_0 and u_1 respectively. For each run, the ignition state can be obtained by a parametric trial on $\mathbf{\Lambda}$ with respect to either g_{u_0} or g_{u_1} . Then, Eq.(5) implies that for $i = 0, 1$ and $x = 0$ or 1 ,

$$\text{Prob}(x_i | u_i) = \sum_{\hat{\lambda}_j \in (x_i | u_0)} g_{u_0}(\hat{\lambda}_j) = \sum_{\hat{\lambda}_j \in (x_i | u_1)} g_{u_1}(\hat{\lambda}_j) \quad (19)$$

Clearly, these conditions mean that $\text{Prob}(x_i | u_i)$ is well defined, i.e., that gauges distributions are compatible with local consistency. Conversely, when the *consistency conditions*, Eq.(19), are fulfilled, using Eq.(18), the probability distributions $g_k(\lambda_j)$ define a single random system $P(x_0; x_1 | u_0; u_1)$. An alternative formulation with expectation values reads,

$$\mathbf{E}(x_i | u_i) = \sum_{\hat{\lambda}_j \in (1 | u_0)} g_{u_0}(\hat{\lambda}_j) = \sum_{\hat{\lambda}_j \in (1 | u_1)} g_{u_1}(\hat{\lambda}_j) \quad (20)$$

We will give some examples, but beforehand, let us summarize our results so far:

Summary (Emulation of Bell-type systems). *Any Bell-type system can be emulated by a stochastic gauge system composed of an auxiliary outcome set $\mathbf{\Lambda} = \{\hat{\lambda}_j\}$, called ignition set, and a set of gauge distributions $g_k(\hat{\lambda}_j)$. One distribution g_k is associated with each setting θ_k . A setting u_i is freely selected in each end region \mathcal{R}_i and transmitted with finite velocity towards a so-called ignition point \mathcal{I} , where each setting is associated with its gauge distribution. Only one of the two distributions, g_{igni} , is used at \mathcal{I} . This choice is called gauge selection. A single trial is performed at \mathcal{I} to draw a unique ignition state $\hat{\lambda}_j$. This result is transmitted with finite velocity backwards the end regions. The final outcomes x_i are next computed by a projection function $\Pi(u_i, \hat{\lambda}_j)$. Gauge selection at \mathcal{I} does not affect the probability of the final outcomes. The ignition point is located at*

the boundary of the two regions. Therefore, if the two regions are separated by a causal horizon, the gauge probability system is located on the horizon.

In classical systems, we will call this random process *classical collapse*.

2.2.4 General 2-setting totally correlated systems

Consider first a Bell-type system with just $K = 2$ settings, θ_0 and θ_1 . When accounting for local consistency and total correlation, it is easily shown that the general system depends on 4 parameters, $q_i, i = 1, 2, 3, 4$ (Table 2), subjected to the following constraints: $0 \leq q_i \leq 1$; $q_3 \geq q_1, q_2$; $q_4 \geq q_1, q_2$; and $q_1 + q_2 \geq q_3, q_4$. The linear system Eq.(16) admits $2K = 4$ unknowns for 4 independent equations. The gauge distributions $g_k(\lambda_j)$ are given in Table 3. Suppose that the observers choose two different settings, e.g., $u_0 = \theta_0$ and $u_1 = \theta_1$. Then, at the ignition point \mathcal{I} , any distribution of the working set can be selected. The trial with respect to this particular distribution gives an ignition state λ_j . Finally, since $u_0 = \theta_0$, we get $x_0 = 0$ or 1 depending on whether j is even or odd. Similarly, $x_1 = 0$ or 1 according to $j < 2$ or $j \geq 2$. Gauge invariance means that all statistical properties of x_0 and x_1 do not depend on the gauge selection, because they can be computed directly from $P(x_0, x_1 | u_0, u_1)$.

Note that, even in this simple case, the concept of distance between θ_0 and θ_1 is inconsistent in contextually dependent systems because the two setting pairs (θ_0, θ_1) and (θ_1, θ_0) cannot be defined at one and the same time. Thus, generally, $d(\theta_0, \theta_1) \neq d(\theta_1, \theta_0)$. This violation of the metric identity $d(\theta_0, \theta_1) = d(\theta_1, \theta_0)$ can be regarded as a criterion of *contextual dependency* of the system. It is easy to design *ad hoc* contextually dependent systems, e.g., with dice games. By contrast, physical setups with entangled photons would probably require components of different nature like non reciprocal polarizers because the simple 2-setting EPR experiment is symmetrical with respect to bit reversal.

Conversely, when $d(\theta_0, \theta_1) = d(\theta_1, \theta_0)$, the two distributions $g_0(j)$ and $g_1(j)$ are identical. The concept of gauge distribution degenerates and thus each ignition state $\hat{\lambda}_j$ has a definite probability, $g(j)$, whatever the settings. Therefore, the ignition states can be regarded in this case as random *hidden variables* located at the ignition point.

(x_0, x_1)	(θ_0, θ_0)	(θ_1, θ_1)	(θ_0, θ_1)	(θ_1, θ_0)
(0, 0)	$1 - q_1$	$1 - q_2$	$1 - q_3$	$1 - q_4$
(0, 1)	0	0	$q_3 - q_1$	$q_4 - q_2$
(1, 0)	0	0	$q_3 - q_2$	$q_4 - q_1$
(1, 1)	q_1	q_2	$q_1 + q_2 - q_3$	$q_1 + q_2 - q_4$
$d(u_0, u_1)$	0	0	$2q_3 - q_1 - q_2$	$2q_4 - q_1 - q_2$

Table 2: General *locally consistent totally correlated* Bell-type system with two possible settings θ_0 and θ_1 depending on 4 parameters q_1, q_2, q_3 and q_4 . The table gives the conditional probabilities $P(x_0, x_1 | u_0, u_1)$ as well as the mean Hamming distance $d(u_0, u_1)$ between binary strings.

j	$g_0(j)$	$g_1(j)$
0	$1 - q_3$	$1 - q_4$
1	$q_3 - q_2$	$q_4 - q_2$
2	$q_3 - q_1$	$q_4 - q_1$
3	$q_1 + q_2 - q_3$	$q_1 + q_2 - q_4$

Table 3: Gauge probability distributions corresponding to Table 2.

2.3 Simulation of the EPR spin experiment

We are going to classically simulate conventional EPR-B experiments, i.e., Bell-type setups for a pair of totally correlated spins with particular probabilities computed from quantum mechanics. In 2D-configuration, let Θ be the unit circle $\mathbf{U}(1) = [0, 2\pi]$. From quantum mechanics we know that for any $\theta_a, \theta_b \in \Theta$ the target probabilities are given by the following equations and Table (4).

$$P(0; 1 | \theta_a; \theta_b) = P(1; 0 | \theta_a; \theta_b) = (1/4)[1 - \cos(\theta_a - \theta_b)]. \quad (21)$$

$$P(0; 0|\theta_a; \theta_b) = P(1; 1|\theta_a; \theta_b) = (1/4)[1 + \cos(\theta_a - \theta_b)]. \quad (22)$$

Clearly, the system is locally consistent and totally correlated. In addition the two entities are symmetrical and therefore the two families of solutions are identical.

$(u_0, u_1) \rightarrow$ (x_0, x_1) \downarrow	(θ_a, θ_a) (θ_b, θ_b)	(θ_a, θ_b) (θ_b, θ_a)
(0, 0)	1/2	(1/4)(1 + cos θ)
(0, 1)	0	(1/4)(1 - cos θ)
(1, 0)	0	(1/4)(1 - cos θ)
(1, 1)	1/2	(1/4)(1 + cos θ)
$d(u_0, u_1)$	0	(1/2)(1 - cos θ)

Table 4: Conditional probability $P(x_0, x_1|u_0; u_1)$ of a typical EPR-B system for two different settings θ_a and θ_b with $\theta = \theta_a - \theta_b$.

2.3.1 2-setting EPR-B experiment

Consider first a 2-setting EPR-B experiment: Let θ_0 and θ_1 be the polarization angles and $\theta_{01} = \theta_1 - \theta_0$. This is obtained with $q_1 = q_2 = 1/2$ and $q_3 = q_4 = (1/4)(3 - \cos \theta_{01})$ in Table 2. Then Table 3 reduces to Table 5. The two gauge distributions are identical and the ignition states can be viewed as hidden variables with a definite probability. Such 2-setting EPR-pairs are referred to as ‘Bell’s states’ when $\theta_{01} = \pi/2$.

j	$4 \times g_0(j) = 4 \times g_1(j)$
0	$1 + \cos \theta_{01}$
1	$1 - \cos \theta_{01}$
2	$1 - \cos \theta_{01}$
3	$1 + \cos \theta_{01}$

Table 5: Gauge probability distributions for a symmetric EPRB system with 2 settings, θ_0 and θ_1 . (We have $\theta_{01} = \theta_1 - \theta_0$). The two gauge distributions are identical

2.3.2 3-setting EPR-B experiment

We now consider an EPR-B system with $K = 3$ different settings, leading to a violation of Bell’s inequalities Eq.(1).

The linear system Eq.(16) admits $2^3 = 8$ unknowns and $2K = 6$ independent equations: Each probability distribution depends on 2 arbitrary parameters and therefore the general solution depends on 6 arbitrary parameters. We can take advantage of this degeneracy to cancel the gauge probability of two ignition states. It is possible to use the 6 double-plateau indices \mathbf{D}_3 ,

$$\mathbf{D}_3 = \{j\} = \{3, 7, 6, 4, 0, 1\}.$$

The linear system admits then only one solution provided that all gauge probabilities p_{kj} are non negative.

Let θ_0, θ_1 and θ_2 be the three settings with $0 \leq \theta_0 \leq \theta_1 \leq \theta_2 < \pi$. Let $\theta_{ab} = \theta_b - \theta_a$. The solutions of Eq.(16) define a working set of effective gauge ignition states given by Table 6. The maximum violation of Bell’s inequality, Eq.(1) or (3), is obtained, e.g., for $\theta_0 = 0, \theta_1 = \pi/5$ and $\theta_2 = \pi/2$. This could be checked by a computer simulation using Table 7.

2.3.3 4-setting EPR-B experiment

We now consider an EPR-B system with $K = 4$ different settings, leading to a violation of CHSH inequalities Eq.(4). We proceed similarly as for 3 settings.

Let $\theta_0, \theta_1, \theta_2$ and θ_3 be the four settings. The maximum violation of CHSH inequality is obtained, e.g., for $\theta_0 = 0, \theta_1 = \pi/8, \theta_2 = \pi/4$ and $\theta_3 = 3\pi/8$.

The linear system Eq.(16) admits $2^4 = 16$ unknowns and $2K = 8$ independent equations: Each probability distribution depends on 8 arbitrary parameters and the general solution depends on

j	$4 \times g_0(j)$	$4 \times g_1(j)$	$4 \times g_2(j)$
0	$1 + \cos \theta_{02}$	$\cos \theta_{01} + \cos \theta_{21}$	$1 + \cos \theta_{02}$
1	$1 - \cos \theta_{01}$	$1 - \cos \theta_{01}$	$\cos \theta_{12} - \cos \theta_{02}$
3	$\cos \theta_{01} - \cos \theta_{02}$	$1 - \cos \theta_{12}$	$1 - \cos \theta_{12}$
4	$\cos \theta_{01} - \cos \theta_{02}$	$1 - \cos \theta_{12}$	$1 - \cos \theta_{12}$
6	$1 - \cos \theta_{01}$	$1 - \cos \theta_{01}$	$\cos \theta_{12} - \cos \theta_{02}$
7	$1 + \cos \theta_{02}$	$\cos \theta_{01} + \cos \theta_{21}$	$1 + \cos \theta_{02}$

Table 6: Working set of gauge probability distributions using only 6 ignition states (out of 8), $j \in \mathbf{D}_3 = \{0, 1, 3, 4, 6, 7\}$ for 3 settings. This working set is only valid when all entries are positive.

j	$g_0(j)$	$g_1(j)$	$g_2(j)$
0	0.250	0.349	0.250
1	0.048	0.048	0.147
3	0.202	0.103	0.103
4	0.202	0.103	0.103
6	0.048	0.048	0.147
7	0.250	0.349	0.250

Table 7: Gauge probability distribution with 6 ignition states derived from Table 6, for the 3 settings: $\theta_0 = 0; \theta_1 = \pi/5; \theta_2 = \pi/2$ leading to the maximum violation of Bell's inequality, Eq.(1) or (3).

32 arbitrary parameters. We will take advantage of this degeneracy to force to zero the gauge probability of 8 ignition states. It is possible to keep the double-plateau indices,

$$\mathbf{D}_4 = \{j\} = \{3, 7, 15, 14, 12, 8, 0, 1\}.$$

Let again $\theta_{ab} = \theta_b - \theta_a$. The linear system admits then only one solution given in Table 8, which turns out to be positive in the range of interest (Table 9).

j	$4 \times g_0(j)$	$4 \times g_1(j)$	$4 \times g_2(j)$	$4 \times g_3(j)$
0	$1 + \cos \theta_{30}$	$\cos \theta_{10} + \cos \theta_{13}$	$\cos \theta_{20} + \cos \theta_{23}$	$1 + \cos \theta_{30}$
1	$1 - \cos \theta_{10}$	$1 - \cos \theta_{10}$	$\cos \theta_{21} - \cos \theta_{20}$	$\cos \theta_{31} - \cos \theta_{30}$
3	$\cos \theta_{10} - \cos \theta_{20}$	$1 - \cos \theta_{12}$	$1 - \cos \theta_{12}$	$\cos \theta_{32} - \cos \theta_{31}$
7	$\cos \theta_{20} - \cos \theta_{30}$	$\cos \theta_{21} - \cos \theta_{31}$	$1 - \cos \theta_{23}$	$1 - \cos \theta_{23}$
8	$\cos \theta_{20} - \cos \theta_{30}$	$\cos \theta_{21} - \cos \theta_{31}$	$1 - \cos \theta_{23}$	$1 - \cos \theta_{23}$
12	$\cos \theta_{10} - \cos \theta_{20}$	$1 - \cos \theta_{12}$	$1 - \cos \theta_{12}$	$\cos \theta_{32} - \cos \theta_{31}$
14	$1 - \cos \theta_{10}$	$1 - \cos \theta_{10}$	$\cos \theta_{21} - \cos \theta_{20}$	$\cos \theta_{31} - \cos \theta_{30}$
15	$1 + \cos \theta_{30}$	$\cos \theta_{10} + \cos \theta_{13}$	$\cos \theta_{20} + \cos \theta_{23}$	$1 + \cos \theta_{30}$

Table 8: Working set of gauge probability distributions using only 8 ignition states (out of 16), $j \in \mathbf{D}_4 = \{0, 1, 3, 7, 8, 12, 14, 15\}$, for 4 settings. This working set is only valid when all entries are positive.

2.3.4 Discrete regular K -setting EPR-B experiment

Finally consider a discrete EPR-B system with K regular settings

$$k \in \llbracket 0, K-1 \rrbracket : k \mapsto \theta_k = \frac{k\pi}{K}.$$

We proceed similarly as above. The rank of the system is $2K$ and the relevant ignition states λ_j are defined by the double-plateau function $j = D_K(r)$, Eq.(14), with $r \in \llbracket 0, 2K-1 \rrbracket$. If we use the same projection function $\Pi(k, j)$, Eq.(50), we have,

$$\Pi[k, D_K(r)] = \Pi[0, D_K(r-k)] \quad ; \quad g_k[D_K(r)] = g_0[D_K(r-k)]$$

where $r-k \geq 0$ is computed modulo $2K$.

j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$
0	0.346	0.408	0.408	0.346
1	0.019	0.019	0.054	0.081
3	0.054	0.019	0.019	0.054
7	0.081	0.054	0.019	0.019
8	0.081	0.054	0.019	0.019
12	0.054	0.019	0.019	0.054
14	0.019	0.019	0.054	0.081
15	0.346	0.408	0.408	0.346

Table 9: Gauge probability distributions derived from Table 8, for the four settings: $\theta_0 = 0; \theta_1 = \pi/8; \theta_2 = \pi/4; \theta_3 = 3\pi/8$. This setup leads to the maximum violation of the CHSH inequalities, Eq. (4).

An alternative formulation is to define a number of $2K$ ignition states,

$$\lambda_r^D \stackrel{\text{(def)}}{=} \lambda_{D_K(r)},$$

a new projection function

$$\Pi^D(k, r) \stackrel{\text{(def)}}{=} \Pi[k, D_K(r)]$$

and then

$$g_k^D(r) \stackrel{\text{(def)}}{=} g_k[D_K(r)]$$

The functions,

$$g(r) \stackrel{\text{(def)}}{=} g_0^D(r), \quad \text{and} \quad \Pi(r) \stackrel{\text{(def)}}{=} \Pi^D(0, r)$$

with $r \in \llbracket 0, 2K - 1 \rrbracket$ are sufficient to describe all distributions since

$$g_k^D(r) = g(r - k) \quad \text{and} \quad \Pi^D(k, r) = \Pi(r - k) \quad (23)$$

where $r - k \geq 0$ is computed modulo $2K$. It is a simple exercise to derive the following expression,

$$g(r) = \frac{1}{2} \sin \alpha \times \begin{cases} |\cos(2r + 1)\alpha| & K \equiv 0 \pmod{2} \\ |\cos 2r\alpha| & K \equiv 1 \pmod{2} \end{cases} \quad (24)$$

where $\alpha = \pi/(2K)$, and,

$$\Pi(r) = \begin{cases} 0 & \text{if } 2r \leq K \text{ (computed modulo } 2K) \\ 1 & \text{if } 2r > K \text{ (computed modulo } 2K) \end{cases} \quad (25)$$

(where the residues modulo $2K$ are taken into $\llbracket 0, 2K - 1 \rrbracket$). For instance, Table 10 gives the result for $K = 5$. For large values of K , regular settings approach continuous settings which deserve a special derivation.

r	$g_0^D(r) : M$	$g_1^D(r) : M$	$g_2^D(r) : M$	$g_3^D(r) : M$	$g_4^D(r) : M$
0	1	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$
1	$\cos 2\alpha$	1	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$
2	$\cos 4\alpha$	$\cos 2\alpha$	1	$\cos 2\alpha$	$\cos 4\alpha$
3	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$	1	$\cos 2\alpha$
4	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$	1
5	1	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$
6	$\cos 2\alpha$	1	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$
7	$\cos 4\alpha$	$\cos 2\alpha$	1	$\cos 2\alpha$	$\cos 4\alpha$
8	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$	1	$\cos 2\alpha$
9	$\cos 2\alpha$	$\cos 4\alpha$	$\cos 4\alpha$	$\cos 2\alpha$	1

Table 10: Gauge distributions $g_k^D(r)/M$ for 5 settings $M = (1/2) \sin \alpha$; $\alpha = \pi/10$

2.3.5 Continuous 2D-EPR-B experiment

For a continuous ensemble of settings, we have to construct a continuous ignition set (i.e., of cardinality \aleph_1) and thus to define a convenient projection function. Next we will derive a continuous set of gauge probability distributions.

Ignition set For any setting labelled θ let $(\xi|\theta)$ be the local target $\xi \in \mathbf{X}$, given $\theta \in \Theta$. For any real λ , define the ignition state $\hat{\lambda}$ by the intersection of local targets:

$$\hat{\lambda} = \bigcap_{\theta \in [0, \pi]} \{(\Pi(\theta, \lambda)|\theta)\} \in \mathbf{\Lambda}, \quad (26)$$

where according to Eq.(12), the binary digit $\xi \in \mathbf{X}$ is computed by a projection function $\Pi(\theta, \lambda)$. It is convenient to define the following dyadic square wave function of θ and λ guessed by analogy with Eq.(23):

$$\Pi(\theta, \lambda) = \frac{1}{2}[1 + \text{sgn} \cos(\theta - \lambda)]. \quad (27)$$

Then, we will replace $\hat{\lambda}$ by λ when necessary and we will use a subset of $\mathbf{\Lambda}$ defined by Eq.(26) and corresponding to $0 \leq \lambda \leq 2\pi$. This leads to a single solution. When compared with the regular discrete settings, we find a close similarity. This justifies the name of *trigonometric order* for the double-plateau functions in Sec. 2.2.1.

Gauge probability distributions For each $\theta \in \Theta$ we are going to derive a gauge distribution density, $g_\theta(\lambda)d\lambda$. Define the interval

$$\mathbf{I}(x, \theta) = \{\lambda \mid \Pi(\theta, \lambda) = x\}$$

and let $\mathbf{J}(x_0, x_1, u_0, u_1) = \mathbf{I}(x_0, u_0) \cap \mathbf{I}(x_1, u_1)$. To compute the solution, let $u_0 = \theta$. For each setting $u_1 \in \Theta$ and each outcome pair x_0, x_1 , we have the functional equation:

$$\forall u_1, \forall x_0, \forall x_1 : \int_{\mathbf{J}(x_0, x_1, u_0, u_1)} g_\theta(\hat{\lambda})d\lambda = P(x_0; x_1 | u_0 = \theta; u_1) \quad (28)$$

where the unknown function is $g_\theta(\hat{\lambda})$ or simply, $g_\theta(\lambda)$. The solution has to be computed for each interval. For instance, for $u_1 > \theta$, we have $\mathbf{J}(1, 1, \theta, u_1) = [u_1 - \pi/2, \theta + \pi/2]$ and $P(1; 1 | \theta; u_1) = (1/4)[1 + \cos(\theta - u_1)]$. Finally, the result is the following working set of gauge distributions [24]:

$$g_\theta(\lambda) = (1/4)|\cos(\theta - \lambda)| \quad (29)$$

Eqs. (27) and (29) are closely related with Eq. (23). Now the classical collapse is described by the process of Sec. 2.2.3 with some straightforward changes. Clearly, this model exactly emulates the 2D-conventional EPR-B experiment and leads to a violation of Bell's inequalities.

3 Simulation of multipartite entangled systems

We will now extend the above results beyond the particular case of Bell-type system to general systems with any number of regions. Even if we do not necessarily require that the system mimics a quantum situation, the concept of local consistency remains the key property of classical analogues of quantum systems. This notion is closely related to the nonsignaling quantum correlation of Popescu and Rohrlich [35].

3.1 Classical analogues of quantum systems

Consider an extended object in equilibrium composed of n *correlated* entities $\{\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_{n-1}\}$, located in n space regions $\mathcal{R}_0, \mathcal{R}_1, \dots, \mathcal{R}_{n-1}$. The equilibrium can be broken and the future outcome x_i will depend randomly upon both the symmetry of the object and the free choice u_i of the observer. When $u_i = \theta_k$, it is convenient to give a name to the pair (i, k) of one setting in one region:

Definition (Configuration). *We will call configuration a pair (i, k) of one region \mathcal{R}_i and one setting θ_k . Each configuration is uniquely labelled by an index $\gamma = k + iK$, $\gamma \in \llbracket 0, nK - 1 \rrbracket$.*

Let \mathbf{x} be the outcome vector $(x_0; x_1; \dots; x_{n-1})$ and \mathbf{u} the setting vector $(u_0; u_1, \dots; u_{n-1})$. We will name again *local target* the entry $(x_i|u_i)$ and *global target* or simply *target* the pair $(\mathbf{x}|\mathbf{u})$. Let $\mathfrak{T} = \{(\mathbf{x}|\mathbf{u})\}$ be the set of all targets. Depending upon the symmetry of the object, each target has

a given conditional probability $P(\mathbf{x}|\mathbf{u})$. There are $\text{card}(\mathfrak{T}) = 2^n \times K^n = (2K)^n$ distinct targets, subjected to K^n normalization constraints, namely:

$$\sum_{x_0=0}^1 \cdots \sum_{x_{n-1}=0}^1 P(x_0; x_1; \dots; x_{n-1} | u_0; u_1; \dots; u_{n-1}) = 1$$

Given a particular probability $P(\mathbf{x}|\mathbf{u})$, we aim to describe classically a break of equilibrium of the multipartite system by means of stochastic gauge distributions. Beforehand, we will clarify the concepts of local consistency, ignition states and gauge distributions in this context.

3.1.1 Nonsignaling correlations, complete local consistency and degrees of entanglement

We no more necessarily require total correlation, i.e., equal outcomes for identical settings in different regions, and accept any degree of entanglement. By contrast, we do conserve *local consistency* in order to allow independent partial measurements. This means that a single local probability is defined in each region, irrespective of the settings in the other regions. More generally, we will define the concept of *complete local consistency* when it is possible to derive consistent subsystems in $n - r$ regions simply by ignoring r regions. Again, this condition only allows nonsignaling correlations between any pair of regions. A limit case concerns independent regions, when the probability of the whole system is simply the product of the probabilities of each region. Then, the system is separable and the entities are no more correlated.

Definition (Complete local consistency). *A set of n entangled classical entities $\{\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_{n-1}\}$, is completely local-consistent when the measurement in one or several regions does not affect the marginal probabilities in the other regions.*

For any partition of $\{0, 1, \dots, n-1\}$ into two subsets, say (after reordering the indices if necessary), $\{0, 1, \dots, r-1\}$ and $\{r, r+1, \dots, n-1\}$ with $1 \leq r \leq n-1$, irrespective of $u_r, u_{r+1}, \dots, u_{n-1}$, we have

$$\text{Prob}(x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1}) = \sum_{x_r=0}^1 \cdots \sum_{x_{n-1}=0}^1 P(x_0; \dots; x_{n-1} | u_0; \dots; u_{n-1}). \quad (30)$$

In words, $\text{Prob}(x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1})$ only depends on the local settings, $(u_0; \dots; u_{r-1})$. Thus, the free choices of $u_r, u_{r+1}, \dots, u_{n-1}$ and the subsequent measurements of x_r, \dots, x_{n-1} do not affect the probabilities in regions $\mathcal{R}_0, \mathcal{R}_1, \dots, \mathcal{R}_{r-1}$, as far as these choices and the subsequent outcomes are ignored. It will be convenient to name *partial target* the entry $(x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1})$. Therefore, *when complete local consistency holds, any partial target has a definite probability.*

Conversely, it is possible to increase the number of regions simply by adding successively new regions. This is similar to the process of quantum purification. Any subset of regions is consistent and the global and local viewpoints are compatible. Then, a multipartite system can be decomposed into its subsets without local perturbation. Note that computation of the partial probabilities $\text{Prob}(x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1})$ implies that the regions $\mathcal{R}_r, \mathcal{R}_{r+1}, \dots$ are ignored. As stressed in Sec.(2.1.2), partial probabilities are subjective by definition.

3.1.2 Degrees of freedom

Besides the K^n normalization constraints, complete local consistency provides a number of additional constraints between the target probabilities $P(\mathbf{x}|\mathbf{u})$. Since any subset of regions is consistent, it is possible to enumerate the degrees of freedom in each m -partite subsystem in isolation. In each single region, we have K local independent parameters, e.g. $\text{Prob}(0|u_i)$, because $\text{Prob}(1|u_i)$ is derived by normalization, and thus nK degrees of freedom for n regions and $2K$ for 2 regions. In each pair of regions, we have seen in Sec. 2.2.2, that we have $K^2 + 2K$ degrees of freedom. Since $2K$ local parameters are already counted we have K^2 *joint bipartite* degrees of freedom, e.g., $\text{Prob}(0; 0|u_i; u_j)$. As a result, we have $\binom{n}{2} K^2$ bipartite degrees of freedom for the general n -region system. Similarly, it is easily shown by induction that there are K^m *joint m -partite* degrees of freedom for a general m -region system and hence $\binom{n}{m} K^m$ m -partite degrees of freedom for the general n -region system. Finally, in the general multipartite system, the number of degrees of freedom is

$$\binom{n}{1} K + \binom{n}{2} K^2 + \binom{n}{3} K^3 + \cdots + \binom{n}{n} K^n = (K+1)^n - 1.$$

Because only nK degrees of freedom are local, in general, $(K+1)^n - nK - 1$ degrees of freedom are *shared*. These results are summarized in Table 11. Note that only the total number of constraints is meaningful, and for $n > 1$ one may prefer to consider that there is only 1 constraint of normalization

and $2^n K^n - (K + 1)^n$ constraints of local consistency. One or several degrees of freedom can be suppressed by additional constraints. For instance, in order to force all degrees of freedom to be shared, besides the constraints of normalization and total consistency, we have to add a number of nK new constraints. For $n = 2$, these $2K$ new constraints may be $P(0; 1|\theta_k; \theta_k) = P(1; 0|\theta_k; \theta_k) = 0$ forcing total correlation. Similarly, for $n > 2$, we can suppress both only-local and only-bipartite correlations with a number of $nK + \binom{n}{2}K^2$ additional constraints.

Number of regions	n
Number of settings	K
Number of targets ($\mathbf{x} \mathbf{u}$)	$(2K)^n$
Total number of local targets	$2nK$
Constraints of normalization	K^n
Constraints of local consistency	$K^n(2^n - 1) - (K + 1)^n + 1$
m -partite degrees of freedom	$\binom{n}{m}K^m$
Total degrees of freedom	$(K + 1)^n - 1$
Shared degrees of freedom	$(K + 1)^n - Kn - 1$
Number of gauge distributions $g_\gamma(\lambda_j)$	nK
Rank of the gauge equations	$2(K + 1)^{n-1}$

Table 11: Degrees of freedom in general multipartite locally consistent systems. Gauge parameters refer to one-step collapse.

The degrees of freedom are only relevant for general systems. Indeed, a particular system is completely definite and has thus no free parameter. It is useful to define instead a number of *coefficients of entanglement*, characterizing the behaviour of the system subject to splitting. The main issue, arguably surprising, is the emergence of *randomness*.

3.2 Randomness and entropy

We regard the classical equivalent of a stationary quantum system as a spatially extended object in equilibrium. As long as the equilibrium holds, the system is not probabilistic but completely deterministic. The function $P(\mathbf{x}|\mathbf{u})$ describes only the symmetry of the object, exactly as a spinning roulette can be characterized by the prior probabilities of its wheel pockets, but before any break of equilibrium, by definition, the entropy remains zero. Similarly, in quantum physics the von Neumann entropy of a pure state in unitary evolution is zero.

Consider for a start the case of bipartite systems. Randomness can arise from two causes, measurement and partition of the system.

3.2.1 Entropy of measurement in bipartite systems

Firstly, randomness arises by measurement, because the initial equilibrium is broken. This is a well known situation in classical physics: After a break of symmetry, several final outcomes are possible at random. The randomness is then described by the conventional Shannon entropy [43], accounting for all constraints at the very best. Since the break of equilibrium follows a free choice of the settings, there are as many entropy coefficients as possibilities. For a finite number of K settings in the 2-region context, the entropy is described by a square K -matrix, $S_1(u_0, u_1)$. For a global measurement, we have,

$$S_1(u_0, u_1) = \sum_{x_0=0}^1 \sum_{x_1=0}^1 -P(x_0; x_1|u_0; u_1) \log_2 P(x_0; x_1|u_0; u_1) \quad (31)$$

It is possible also to consider only a partial a measurement, e.g., in region \mathcal{R}_0 , and we have

$$S_1(u_0) = \sum_{x_0=0}^1 -\text{Prob}(x_0|u_0) \log_2 \text{Prob}(x_0|u_0) \quad (32)$$

where $\text{Prob}(x_0|u_0)$ is computed from Eq.(5) or Eq.(19).

3.2.2 Entropy of entanglement in bipartite systems

Less expected, a second source of randomness arises in entangled systems without break of equilibrium. Indeed, it is possible to split the whole system into its two parts and consider each region

separately. Then two cases are possible: either the two subsystems are independent, or the two regions are entangled. In the first case, the two regions are separable, all information is strictly local and no randomness appears. The entropy of the whole system is the sum of the entropy of each subsystem, i.e., *the entropy is extensive* [44].

By contrast, in the second case, local information is insufficient to completely describes the whole system and additional information is needed. By splitting, we decide to ignore all non local information and our description of the system becomes incomplete. In other words, the system is no more completely definite. Therefore, without any break of equilibrium, the entropy of the two separated subsystems is now positive, clearly greater than the zero entropy of the full system. As a result, *entropy is no more extensive*.

In quantum theory, this is generally viewed as an astonishing aspect of entanglement. Actually, this situation is well known in classical thermodynamics in the face of *long range interactions* [30, 31]. In the present case, the only possible paradox lies perhaps in the fact that the splitting is nonsignaling, and thus each region remains locally consistent.

Quantitatively, we aim to evaluate non-local correlations between a pair of local targets, $(x_0|u_0)$ and $(x_1|u_1)$, where x_i is regarded as a random variable while u_i is viewed as a parameter. A similar issue holds in the conventional Shannon transmission theory. The concept of *mutual information* was indeed devised to evaluate the correlation between the data at the two ends of a communication link. This concept was extended to multipartite systems by R. Fano [45]. In this background, the entropy of measurement S_1 can also be called *self information*. For ease of exposition, it is convenient to use a shorthand X_i for the local target, $(x_i|u_i)$. Then $(X_0; X_1)$ describes the global target $(x_0; x_1|u_0; u_1)$. Mutual information is defined as [46],

$$I(X_0; X_1) \stackrel{\text{(def)}}{=} \mathbf{E}[\log_2 \frac{\text{Prob}(X_0; X_1)}{\text{Prob}(X_0)\text{Prob}(X_1)}]$$

where $\mathbf{E}[\cdot]$ stands for the expectation value with respect to $\text{Prob}(X_0; X_1) = P(x_0; x_1|u_0; u_1)$. Mutual information does not depend on the order of the two targets,

$$I(X_0; X_1) = I(X_1; X_0)$$

Define,

$$S_2(u_0, u_1) \stackrel{\text{(def)}}{=} I[(x_0|u_0); (x_1|u_1)] = I[(x_1|u_1); (x_0|u_0)]$$

We compute easily,

$$S_2(u_0, u_1) = \sum_{x_0=0}^1 \sum_{x_1=0}^1 P(x_0; x_1|u_0; u_1) \log_2 \left(\frac{P(x_0; x_1|u_0; u_1)}{\text{Prob}(x_0|u_0)\text{Prob}(x_1|u_1)} \right) \quad (33)$$

In the present context, it is convenient to call this coefficient *entanglement entropy*. Depending upon the background, *mutual information* is also named (apart from the sign) *relative entropy* [47] or *Kullback-Leibler divergence* [48]. Clearly, the entanglement entropy is invariant by interchange of the two regions (but not, in general, by permutation of the settings). It is zero for a pair of independent regions. This classical definition is closely related to the quantum definition of entanglement entropy [49, 50].

For a discrete system with K possible settings, the entanglement entropy is a square matrix of order K , in general non symmetrical. This result is not surprising because we know that we have K^2 bipartite degrees of freedom (Tables 1 and 11). Therefore we can identify these K^2 parameters with the K^2 coefficients of bipartite entanglement. As a result, a bipartite system is separable if and only if its entanglement entropy matrix is zero.

The entanglement entropy measures the amount of information required to account for non-local correlation. For instance, in a previous paper [25], we have described a dice game between two parties, Alice and Bob, to illustrate the concept of classical entanglement. In this example, it is easy to compute an improbable ‘entanglement entropy’ between the two players as a 3-matrix, given in Table 12!

For a continuous system, the entanglement entropy is a function $S_2(u_0, u_1)$. Specially, for a pair of EPR totally entangled entities, the entanglement entropy with respect to a pair of settings θ_a and θ_b is

$$S_2(\theta_a, \theta_b) = (1/2)[(1 + \cos \theta) \log_2(1 + \cos \theta) + (1 - \cos \theta) \log_2(1 - \cos \theta)]$$

where $\theta = \theta_a - \theta_b$. Clearly, $S_2 \leq 1$ and the maximum is equal to 1 bit, obtained for $\theta = 0$. This is just the amount of information required to transmit classically the non-local correlation between the two entities of the pair [51].

$S_2(\Delta_a, \Delta_b)$	Δ_1	Δ_2	Δ_3
Δ_1	1	0.35	0
Δ_2	0.35	1	0.35
Δ_3	0	0.35	1

Table 12: Example of classical entanglement entropy (in bits) between two players, Alice and Bob, in a maximally entangled classical dice game described in Ref. [25]. The game is implemented with three dice labelled Δ_k ($k = 1, 2$ and 3). At each run, each player selects freely one die, Δ_a and Δ_b respectively. The coefficients measure the amount of information required to account for non-local correlation.

3.2.3 Multipartite entanglements

When considering multipartite *measurements*, there is no new topic. By contrast, *splitting* of the system raises different questions because there is a number of different ways to group the regions together. Again, the same issue is encountered in communication networks: In transmission theory, the natural extension of the bipartite mutual information is the multivariate mutual information [45]. Similarly, we will define multivariate mutual information coefficients between local targets, computed recursively as [46],

$$I(X_0; X_1; \dots; X_{n-1}) \stackrel{(\text{def})}{=} I(X_0; X_1; \dots; X_{n-3}; X_{n-2}) - I[(X_0; X_1; \dots; X_{n-3}; X_{n-2})|X_{n-1}] \quad (34)$$

In Eq.(34), X_i is still a local target, shorthand for $(x_i|u_i)$, where x_i is a random variable and u_i a parameter. It can be shown that multivariate mutual information is completely symmetrical with respect to X_0, X_1, \dots, X_{n-1} even if this is not manifest in the definition. Contrary to bivariate mutual information which is always positive, multivariate mutual information can be either positive or negative.

Shannon's information theory has a nice set-theoretic structure [46, 52, 53] recalled in Table 13. Thanks to this structure, computation of entropic expressions is simplified by a graphical representation, named *information diagram* and similar to the conventional Venn diagram. Entropy or mutual information I are associated with a signed measure μ . For simplicity, and without loss of generality, we will use the same symbol I to describe both entropy and information (while entropy is usually symbolized by H). Multivariate mutual information, Eq. (34), is directly associated

Information theory	Set theory
$I(X)$	$\mu(X)$
$(X_1; X_2)$	$X_1 \cap X_2$
(X_1, X_2)	$X_1 \cup X_2$
$(X_1 X_2)$	$X_1 - X_2$

Table 13: Set-theoretic structure of Shannon's information theory. Each variable X is represented by a set X . Entropy or mutual information I are associated with a signed measure μ . Information expressions, like $I(X_1, X_2)$, $I(X_1; X_2)$ or $I[(X_1|X_2)]$ are respectively associated with $\mu(X_1 \cup X_2)$, $\mu(X_1 \cap X_2)$ and $\mu(X_1 - X_2)$ (where $X_1 - X_2 = X_1 \cap X_2^c$).

with

$$I(X_0; \dots; X_{n-1}) = \mu(X_0 \cap \dots \cap X_{n-1}) \quad (35)$$

and thus $I(X_0; \dots; X_{n-1})$ is clearly completely symmetrical with respect to X_0, \dots, X_{n-1} . For instance, when $n = 3$, Eq.(34) is represented by the information diagram shown in Fig.1. Given a setting vector \mathbf{u} , each local distribution X_i is represented by an area and $I(X_i)$ is the partial measurement entropy, or the self-information of this variable. For instance, Eq.(32), describes the self-information $I(X_0)$. The n regions delimit a number of $2^n - 1$ atoms of information, $\mu_1, \dots, \mu_{2^n-1}$. For $n = 3$, we have $I(X_0; X_1; X_2) = \mu_7$.

In practice, it is convenient to derive first the measurement entropy $I(X_{i_0}, \dots, X_{i_{r-1}})$ of each partial subsystem $(\mathcal{R}_{i_0}, \dots, \mathcal{R}_{i_r})$ for $r = 1, \dots, n$, because each probability distribution is easily computed. For instance, if $\{i_0, \dots, i_{r-1}\} = \{0, \dots, r-1\}$, the relevant partial probability in the r -region subsystem is given by Eq.(30). We have $2^n - 1$ such partial subsystems. Next, we express the measure μ_i of each atom of the information diagram (Fig. 1) in terms of these measurement entropies. We have $2^n - 1$ atoms and therefore we obtain the mutual information $I(X_0; X_1; \dots; X_{n-1}) = \mu_{2^n-1}$ by solving a linear system of $2^n - 1$ equations for $2^n - 1$ unknowns.

For example, when $n = 3$, we derive easily the following equation by inspection of Fig. 1,

$$\begin{bmatrix} I(X_0) \\ I(X_1) \\ I(X_1, X_0) \\ I(X_2) \\ I(X_2, X_0) \\ I(X_2, X_1) \\ I(X_2, X_1, X_0) \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \\ \mu_5 \\ \mu_6 \\ \mu_7 \end{bmatrix},$$

and we obtain

$$\begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \\ \mu_5 \\ \mu_6 \\ \mu_7 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 & 1 & -1 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 0 & 0 & 1 & -1 \\ -1 & 0 & 1 & 0 & 1 & 0 & -1 \\ 1 & 1 & -1 & 1 & 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} I(X_0) \\ I(X_1) \\ I(X_1, X_0) \\ I(X_2) \\ I(X_2, X_0) \\ I(X_2, X_1) \\ I(X_2, X_1, X_0) \end{bmatrix}$$

The last row reads

$$\begin{aligned} \mu_7 = I(X_0; X_1; X_2) &= I(X_0) + I(X_1) + I(X_2) \\ &\quad - I(X_0, X_1) - I(X_0, X_2) - I(X_1, X_2) + I(X_0, X_1, X_2) \end{aligned} \quad (36)$$

More generally we have,

$$\begin{aligned} I(X_0; X_1; \dots; X_{n-1}) &= \sum_i I(X_i) - \sum_{i,j} I(X_i, X_j) + \\ &\quad \sum_{i,j,k} I(X_i, X_j, X_k) - \dots + (1)^{n-1} I(X_0, \dots, X_{n-1}) \end{aligned} \quad (37)$$

where the summation is taken over all possible combinations of subscripts and,

$$\begin{aligned} I(X_i) = S_1(u_i) &= \sum_{x_i=0}^1 -\text{Prob}(x_i|u_i) \log_2 \text{Prob}(x_i|u_i) \\ &= \sum_{x_0=0}^1 \dots \sum_{x_{n-1}=0}^1 -P(\mathbf{x}, \mathbf{u}) \log_2 \text{Prob}(x_i|u_i), \end{aligned}$$

$$\begin{aligned} I(X_i, X_j) = S_1(u_i, u_j) &= \sum_{x_j=0}^1 \sum_{x_i=0}^1 -\text{Prob}(x_i; x_j|u_i; u_j) \log_2 \text{Prob}(x_i; x_j|u_i; u_j) \\ &= \sum_{x_0=0}^1 \dots \sum_{x_{n-1}=0}^1 -P(\mathbf{x}, \mathbf{u}) \log_2 \text{Prob}(x_i; x_j|u_i; u_j), \end{aligned}$$

etc. In the present context, we will use the following terminology:

Definition (n -partite entanglement entropy, degree and coefficients of entanglement). *We will name n -partite entanglement entropy the expression*

$$S_n(\mathbf{u}) = I(X_0; X_1; \dots; X_{n-1}). \quad (38)$$

When different from zero, $S_n(\mathbf{u})$ defines one degree of n -partite entanglement, $e_n = 1$. We will call entanglement coefficients its K^n components.

The K^n entanglement coefficients can be identified with the K^n n -partite degrees of freedom of a general n -region system. As a result, if $S_n \neq 0$, we will regard the n -region system as n -partite entangled. In general, $S_n(\mathbf{u})$ is not invariant by permutation of the settings but the object describes the n -partite entanglement as a whole and does not depends on the order of the n regions. For instance, when $n = 3$, we have K^3 coefficients of tripartite entanglement,

$$\begin{aligned} S_3(\mathbf{u}) &= \sum_{x_0=0}^1 \sum_{x_1=0}^1 \sum_{x_2=0}^1 P(\mathbf{x}|\mathbf{u}) \times \\ &\quad \log_2 \left(\frac{\text{Prob}(x_0; x_1|u_0; u_1) \text{Prob}(x_1; x_2|u_1; u_2) \text{Prob}(x_2; x_0|u_2; u_0)}{P(x_0; x_1; x_2|u_0; u_1; u_2) \text{Prob}(x_0|u_0) \text{Prob}(x_1|u_1) \text{Prob}(x_2|u_2)} \right) \end{aligned} \quad (39)$$




Figure 1: Information diagram of three parties, X_0 , X_1 and X_2 . Using Table 13, we have for instance $I(X_0) = \mu_1 + \mu_3 + \mu_5 + \mu_7$, $I(X_0; X_1) = \mu_3 + \mu_7$, $I(X_0; X_1; X_2) = \mu_7$.

3.2.4 Characterizing entanglement

Definition (Degree of entanglement). *When a n -region system is n -partite entangled, we will say that its degree of entanglement is n and set $e_n=1$. Otherwise, its degree of entanglement is the larger number of regions, m , which constitutes a m -partite entangled subsystem.*

Finally, any given system has a particular structure described by the scheme (e_2, e_3, \dots, e_n) of its bipartite, tripartite, \dots , n -partite degrees of entanglement. The number of m -partite coefficients of entanglement non identically vanishing is thus $e_m K^m$, with $e_m \leq \binom{n}{m}$.

Definition (Entanglement scheme). *Given a particular n -partite system with K settings, we will call entanglement scheme the sequence (e_2, e_3, \dots, e_n) of its bipartite, tripartite, \dots , n -partite degrees of entanglements, where $e_m \leq \binom{n}{m}$. The overall number of entanglement coefficients is*

$$N = \sum_{m=2}^n e_m K^m.$$

In addition, we will define later K^n total entanglement coefficients by Eq. (40) in Sec. (3.2.5).

Given the probability distribution $P(\mathbf{x}|\mathbf{u})$ the coefficients of entanglement are easily computed. For instance, Bell-type systems are bipartite, totally correlated systems, characterized by the scheme $e_2 = 1$ (with $\binom{2}{2}K^2 = K^2$ coefficients of entanglement). Among tripartite systems (Sec. 3.4.3 below), GHZ-states (considered with only two settings) are characterized by the scheme $(e_2, e_3) = (0, 1)$ and $K = 2$, with $\binom{3}{3}K^3 = 8$ coefficients of entanglement, while general W-states (with two settings) are characterized by the scheme $(e_2, e_3) = (3, 1)$ and $K = 2$, with $\binom{3}{2}K^2 + K^3 = 20$ coefficients of entanglement, but only 2 tripartite coefficients (up to 8) are non zero. Separable systems are trivially characterized by $e_2 = e_3 = \dots = e_n = 0$ and thus zero coefficient of entanglement.

The current results are rather paradoxical. We have elaborated upon a close analogy with signaling networks except that the present system is \dots nonsignaling! However that may be, information theory has established a number of helpful inequalities between entropy functions, which are of course currently valid.

3.2.5 Information inequalities

In quantum information theory, entanglement in multipartite systems is not yet fully understood. By contrast, in the present model, the information of all atoms is easily computed and we have a number of tools at disposal stemmed from information theory. Let α, β be two subsets of $\llbracket 0, n-1 \rrbracket$. If $\alpha = \{i_1, i_2, \dots, i_r\}$, define

$$I(\alpha) \stackrel{(\text{def})}{=} I(X_{i_1}, \dots, X_{i_r})$$

(for simplicity, we conserve the same symbol I).

The following *polymatroidal axioms* [46] describe the Shannon information inequalities.

1. $I(\emptyset) = 0$

2. If $\alpha \subset \beta$ then $I(\alpha) \leq I(\beta)$
3. $I(\alpha) + I(\beta) \geq I(\alpha \cap \beta) + I(\alpha \cup \beta)$

This tool is powerful to derive important entropic relations. Note that this is not the complete story since there are inequalities not derivable from this set of axioms, called *non-Shannon inequalities*.

We will just give one example, and define the concept of *total entanglement*. In transmission theory, Watanabe [54] has defined a global coefficient C , named *total correlation* as,

$$C(X_0; X_1; \dots; X_{n-1}) = \sum_{i=0}^{n-1} I(X_i) - I(X_0, X_1, \dots, X_{n-1})$$

Total correlation is always positive and is zero if and only if all regions are independent. For instance, in Fig.1, we have

$$C(X_0; X_1; X_2) = 2\mu_7 + \mu_3 + \mu_5 + \mu_6$$

In the present context, it is convenient to use a special term,

Definition (Total entanglement). *Given a particular n -partite system with K settings, we will call total entanglement the function*

$$C(\mathbf{u}) \stackrel{\text{(def)}}{=} \sum_{i=0}^{n-1} I(X_i) - I(X_0, X_1, \dots, X_{n-1}). \quad (40)$$

Total entanglement is defined for each setting vector \mathbf{u} . Since we only consider binary outcomes, we have

$$I(X_i) \leq 1 \text{ bit}$$

As a result, we derive the inequality $0 \leq C(\mathbf{u}) \leq n$ bits. Actually this maximum can be improved to $n-1$ bits. The proof is trivial for $n=1$. Now, thanks to the polymatroidal axioms, the addition of one region X_m can only increase the total entanglement by $I(X_m) \leq 1$ bit. Therefore, we have for a n -region system,

$$0 \leq C(\mathbf{u}) \leq n-1 \text{ bits.} \quad (41)$$

The upper bound can be saturated, e.g., when the measures μ_i of all atoms of information are zero, except for $I(X_0; X_1; \dots; X_{n-1}) = 1$ bit. This correspond to the general n -region GHZ-states, when the same setting, corresponding to the axis of entanglement, is used in each region.

Definition (Maximally entangled system). *We will say that a n -region system is maximally entangled when its total entanglement for at least one setting-vector is equal to $n-1$ bits.*

For instance, EPR-pairs are maximally entangled bipartite systems (Table 24 below) and W-states are maximally entangled tripartite systems (Table 25 below). Surprisingly, with only two settings, GHZ-states are not maximally entangled (Table 28 below). However, when we add a third setting Z , corresponding to the axis of common entanglement, it is readily shown that the setting-vector $(Z; Z; Z)$ leads to a set of atom measures $\mu_i = 0$ for $i \in [0, 6]$ and $\mu_7 = 1$ bit (Fig.1). Therefore, we have $C(Z; Z; Z) = 2\mu_7 + \mu_3 + \mu_5 + \mu_6 = 2$ bits, and the general 3-regions GHZ states are indeed maximally entangled.

This framework is likely to be helpful to discuss a number of issues like ‘entanglement monogamy’ [55] or collapse mechanism (Sec. 3.3.1 below). These discussions are beyond the scope of this paper.

3.3 Classical multipartite collapse

We will now describe the break of equilibrium of a n -region classical system. The equilibrium can be broken as follows: In one or several regions \mathcal{R}_i , an observer, \mathcal{O}_i , (1) selects freely a setting u_i , element of a given set Θ of K different settings, (2) breaks the local equilibrium in a way reliant on u_i , and (3) measures on \mathcal{E}_i an observable x_i , element of a set $\mathbf{X} = \{0, 1\}$.

3.3.1 Collapse mechanism

We aim to discuss first the collapse mechanism, in the sense of ‘reaction mechanism’ in chemical kinetics. Indeed, the simple process defined for Bell-type systems turns out to be insufficient to describe multipartite system collapses.

The EPR paradox was cleared up by a direct gauge implementation of the probability distributions in the two regions. Actually, a gauge selection defined a *leading region*, say \mathcal{R}_0 , but afterward the two regions entered into the model on just the same footing. We have computed a probability distribution, $g_{k_0}(\hat{\lambda}_j)$, located on the boundary of the two regions, accounting for the free choice $u_0 = k_0$ in the leading region and governing *both* $(x_0|k_0)$ and $(x_1|k_0; u_1)$.

However, local realism does not require that the same distribution governs the two processes. One can construct an *indirect collapse* as follows. (1) The settings (k_0, k_1) are sent to an ignition point \mathcal{I} . (2) A gauge selection defines the leading region, e.g., \mathcal{R}_0 . (3) At the boundary of the two regions, the outcome $x_0 = \xi_0$ of the leading region is drawn at once, using $\text{Prob}(x_0|k_0)$. (4) A gauge distribution $\text{Prob}(x_1|\xi_0; k_0; u_1)$ is used in the second region \mathcal{R}_1 . Of course, thanks to the well known product rule [56], accounting for local consistency, the two processes give exactly the same final result,

$$P(\xi_0; x_1|k_0; u_1) = \text{Prob}(\xi_0|k_0)\text{Prob}(x_1|\xi_0; k_0; u_1)$$

Clearly, the indirect process requires two successive trials. As a result, the number of gauge distributions is doubled. On the other hand, each distribution $\text{Prob}(x_1|\xi_0; k_0; u_1)$ is simpler, and even trivial in bipartite system, because we have only to account for compatibility conditions in one region instead of two. Therefore, in the computation of the gauge distributions, the direct route can fail, while the two-step process turns out to be successful. As in chemical kinetics, this may be physically meaningful because the two-step route may require a delay, while the direct process can be viewed as quasi-instantaneous at the ignition point.

In multipartite systems, we will proceed similarly. Thanks again to the product rule and local consistency, it is possible to define multi-step collapses, for instance,

$$P(x_0; x_1; x_2; x_3|u_0; u_1; u_2; u_3) = \text{Prob}(x_0|u_0) \times \text{Prob}(x_1|x_0; u_0; u_1) \\ \times \text{Prob}(x_2|x_0; x_1; u_0; u_1; u_2) \times \text{Prob}(x_3|x_0; x_1; x_2; u_0; u_1; u_2; u_3) \quad (42)$$

At each step, we have to select a new gauge region. For m steps, we have m gauge trials per run.

Suppose, for instance, that several observers decide to proceed to a measurement and select freely a setting θ_{k_i} . They send their setting towards an ignition point, located if possible at the boundary of all regions. A first gauge selection at the ignition point defines a leading configuration, say $\gamma_0 = k_0 + Ki_0$, i.e., a gauge region \mathcal{R}_0 and a gauge setting θ_{k_0} . The first outcome ξ_0 is drawn using the local probability $\text{Prob}(x_0|k_0)$. Now, the probability distribution in the remaining $n - 1$ regions is a gauge function

$$\text{Prob}(x_1; \dots; x_{n-1}|\xi_0; k_0; u_1; \dots; u_{n-1}) = M \times P(\xi_0; x_1; \dots; x_{n-1}|k_0; u_1; \dots; u_{n-1}) \quad (43)$$

where $M = M(\xi_0, k_0, u_1, \dots, u_{n-1})$ is a normalization factor. Note that Eq.(43) only describes the updated probability of (x_1, \dots, x_{n-1}) for the observer \mathcal{O}_0 , because he has got alone a new information, namely $u_0 = k_0$ and $x_0 = \xi_0$. By contrast, the $n - 1$ other observers are not aware of what happens in region \mathcal{R}_0 , and their view of the $n - 1$ regions is still given by Eq.(30),

$$\text{Prob}(x_1; \dots; x_{n-1}|u_1; \dots; u_{n-1}) = \sum_{x_0=0}^1 P(x_0; \dots; x_{n-1}|u_0; u_1; \dots; u_{n-1}). \quad (44)$$

Due to local consistency, the right hand-side of Eq.(44) does not depend on u_0 and we can plug k_0 in u_0 . In general, Eq.(43) and Eq.(44) are different, because they describe different level of knowledge.

Even if the original system mimics a quantum situation, the probability distribution Eq.(43) may be more general. Nevertheless, it can be seen that Eq.(43) is still locally consistent. Therefore we can define a second-step probability distribution P' as

$$P'(x_1; \dots; x_{n-1}|u_1; \dots; u_{n-1}) \stackrel{(\text{def})}{=} \text{Prob}(x_1; \dots; x_{n-1}|\xi_0; k_0; u_1; \dots; u_{n-1}). \quad (45)$$

Owing to Eq.(42), this distribution P' describes now the $n - 1$ remaining regions (instead of $P(\mathbf{x}|\mathbf{u})$). It is possible to iterate when wanted. Let m be the number of iterations. If $m = n$, the overall collapse is completed. Each observer gets his own outcome, $x_i = \xi_i$ but ignores the other outcomes. If $m < n$, the last gauge trial has to be performed in a subsystem of $n - m$ regions. For this purpose, we will construct in the next section a convenient stochastic gauge system on the model of the Bell-type systems. If we fail to obtain a set of positive gauge probability distributions, this means that there is no locally realist collapse in m steps and we have to try with $m + 1$ steps. The last stage in just one region is trivial, and therefore, it is always possible to find a locally realist route whatever the number of regions and the degree of entanglement. In other words, given any locally consistent system, the global collapse can always be implemented by an arbitrary chain of n one-region collapses.

Definition (m-step collapse). *We will call m-step collapse a collapse composed of m-1 cascaded one-region trials and one final trial into a subsystem of $n - m + 1$ regions.*

The feasibility of a m -step collapse with $m < n$ is likely to be characterized by inequalities between the target probabilities. In this respect, Bell inequalities may be viewed as feasibility conditions for a form of ‘zero step’ collapse. Note that a n -step collapse, requiring a minimum of computational resources, can be helpful to simulate quantum algorithms.

In bipartite systems, we have always found that the direct collapse in just one step is feasible. By contrast, we will see that a one-step process fails in some tripartite systems with strong entanglement but in all our simulations, the two-step route was successful.

3.3.2 One-step collapse

In order to describe a classical one-step collapse in multipartite systems, we have to generalize the concepts defined in Bell-type systems, namely, ignition states, gauge probability distributions, and measurement process.

Ignition states The ignition states will define the potential outcomes in the n regions. In Sec (2.2.1) we have constructed a fine-grained description of Bell-type systems. Now, we aim to construct a fine-grained description of multipartite systems by the same token.

We have defined the *local target* $(x_i|u_i) = (\xi|\theta_k)_i$ in a region \mathcal{R}_i by $x_i = \xi$ given $u_i = \theta_k$. When the system is completely local-consistent, the probability $\text{Prob}(\xi|\theta_k)_i$ in a single region is uniquely defined. The number of distinct local targets is thus $2K$ for one region and $2nK$ for the whole. Next, we construct the subset $\mathfrak{T}_{(\xi|\theta_k)_i}$ of \mathfrak{T} composed of all global targets compatible with one particular local target $(\xi|\theta_k)_i$.

$$\mathfrak{T}_{(\xi|\theta_k)_i} = \{(\mathbf{x}|\mathbf{u}) / x_i = \xi ; u_i = \theta_k\}$$

Any local target $(\xi|\theta_k)_i$ in region \mathcal{R}_i can be regarded as an union of global targets

$$(\xi|\theta_k)_i = \bigcup_{(\mathbf{x}|\mathbf{u}) \in \mathfrak{T}_{(\xi|\theta_k)_i}} \{(\mathbf{x}|\mathbf{u})\}.$$

Conversely, any global target $(\mathbf{x}|\mathbf{u})$ can be viewed as the intersection of n local targets in regions $\mathcal{R}_0, \dots, \mathcal{R}_{n-1}$

$$(\mathbf{x}|\mathbf{u}) = (x_0|u_0) \cap (x_1|u_1) \cap \dots \cap (x_{n-1}|u_{n-1})$$

Let $\mathcal{J} = \{j\}$ be a set of integers. For $i \in \llbracket 0, n-1 \rrbracket$, define n applications $\Pi_i(k, j)$ of $\llbracket 0, K-1 \rrbracket \times \mathcal{J} \rightarrow \mathbf{X} = \{0, 1\}$

$$\xi_i = \Pi_i(k, j). \quad (46)$$

We will call *projection function* in region \mathcal{R}_i the function $\Pi_i(k, j)$. More concisely, it is possible to define a global projection function $\Pi(\gamma, j)$ in terms of configuration as

$$\Pi(\gamma, j) \stackrel{(\text{def})}{=} \Pi_i(k, j) \text{ with } \gamma = k + iK \quad (47)$$

Now, we construct a sequence of $\text{card}(\mathcal{J})$ fine grains $\hat{\lambda}_j$ by the following intersection of local targets:

$$\begin{aligned} j \in \mathcal{J} &\mapsto \hat{\lambda}_j \\ \hat{\lambda}_j &\stackrel{(\text{def})}{=} \bigcap_{i=0}^{n-1} \bigcap_{k_i=0}^{K-1} \left(\Pi_i(k_i, j) | \theta_{k_i} \right)_i. \end{aligned} \quad (48)$$

There are K^n different setting vectors, labelled $(k_0, k_1, \dots, k_{n-1})$. Therefore, the maximum number of distinct fine grains is 2^{K^n} . On the other hand, we have to define the outcome x_i for each configuration (i, k) given j . There are only nK configurations $\gamma = k + iK$ and the maximum cardinality of \mathcal{J} is thus 2^{nK} . We need a selection rule to select 2^{nK} ignition states among 2^{nK} potential fine grains. Let

$$j = \sum_{\mu=0}^{K^n-1} j_\mu 2^\mu \quad (49)$$

be the binary expansion of an integer j , $(0 \leq j \leq 2^{K^n} - 1)$. For any $\mu \in \llbracket 0, K^n - 1 \rrbracket$, there is a unique base- K expansion of μ as

$$\mu = \sum_{i=0}^{n-1} k_i K^i \quad (0 \leq k_i \leq K-1)$$

Define a selection rule σ ,

$$\begin{aligned} \sigma : \llbracket 0, K^n - 1 \rrbracket &\mapsto \llbracket 0, K^n - 1 \rrbracket \\ (k, i) &\rightarrow \mu = \sigma(k, i) \end{aligned}$$

A priori, any function σ can be used... provided that the linear system, Eq.(51) below, leads to feasible solutions. The simplest selection function is $\sigma_1(k, i) = k + iK = \gamma$. We have found by simulation that this simple function is sufficient, and more sophisticated selection rules only give

a permutation of the label of the ignition states. Therefore, we will use this selection function throughout the rest of the paper, unless stated otherwise explicitly.

Let j_μ be the coefficient of 2^μ in the binary expansion of j in Eq.(49). A convenient definition of the projection function is

$$\Pi_i(k, j) \stackrel{(\text{def})}{=} j_{\sigma(k, i)} = j_{k+iK} = j_\gamma. \quad (50)$$

For ease of exposition, when no confusion is possible, we will write alternatively

$$\Pi_i(\theta_k, \hat{\lambda}_j) \stackrel{(\text{def})}{=} \Pi_i(k, j).$$

Now, we are going to show that this problem is roughly similar to a bipartite Bell-type system with Kn settings.

Gauge probability distributions In the n -region- K -setting context, we will need nK gauge probability distributions $g_\gamma(\hat{\lambda}_j)$ (or simply, $g_\gamma(j)$) for $\gamma = k + iK \in \llbracket 0, nK - 1 \rrbracket$ with $i \in \llbracket 0, n - 1 \rrbracket$ and $k \in \llbracket 0, K - 1 \rrbracket$. Suppose that the observer in region \mathcal{R}_{i_0} selects the setting k_0 . Let us derive the gauge distribution $g_{\gamma_0}(j)$ for $\gamma_0 = k_0 + i_0K$. Let $p_{\gamma_0 j}$ be the unknown probability of $\hat{\lambda}_j$ for $j \in \mathcal{J}$. Thus, we have to solve a linear system.

For each setting vector \mathbf{u} with $u_{i_0} = \theta_{k_0}$ and each outcome vector \mathbf{x} , we will write one equation:

$$u_{i_0} = \theta_{k_0} : \forall u_i (i \neq i_0), \forall x_i : \sum_{j \in \mathbf{J}(\mathbf{x}, \mathbf{u})} p_{\gamma_0 j} = P(\mathbf{x}|\mathbf{u}) \quad (51)$$

$j \in \mathbf{J}(\mathbf{x}, \mathbf{u})$ means that $\forall i : x_i = \Pi_i(u_i, \hat{\lambda}_j)$, i.e., $\hat{\lambda}_j \in (\mathbf{x}|\mathbf{u})$. For each gauge distribution $g_{\gamma_0}(j)$ we have $\text{card}(\mathcal{J})$ unknowns $p_{\gamma_0 j}$ and $2^n K^{n-1}$ equations among which it can be shown that only $2(K+1)^{n-1}$ are independent. This system is generally degenerate and provides a set of solutions, $g_{\gamma_0}(j) = p_{\gamma_0 j}$. The computation fails if it is impossible to find non-negative solutions. Then, we have to try an indirect collapse, with more than one step.

Consistency conditions The gauge distributions are not independent. As in Bell-type system, Eq. (19), we have a number of consistency conditions, translating complete local consistency, Eq.(30), into gauge formulation. Whatever the gauge distribution g_γ with $\gamma = k + Ki$, provided that $u_i = k$, we have:

$$\text{Prob}(x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1}) = \sum_{\lambda_j \in (x_0; \dots; x_{r-1} | u_0; \dots; u_{r-1})} g_\gamma(j) \quad (52)$$

Measurement We proceed similarly as for totally correlated Bell-type pairs: In each region i , each observer selects his own setting $u_i = \theta_k$. Each configuration $\gamma_i = k + iK$ is transmitted with finite velocity towards the ignition point \mathcal{I} . Only *one configuration*, say γ_{ign_i} , is kept at random. Now, at \mathcal{I} , we perform a trial in the ignition set Λ using the only probability distribution $g_{\gamma_{\text{ign}_i}}(\hat{\lambda}_j)$ to draw a single ignition state $\hat{\lambda}_{j_{\text{ign}_i}}$. Let j_{γ_i} be the coefficients of 2^{γ_i} in the binary expansion of j_{ign_i} . These coefficients, $j_{\gamma_i} = \Pi(\gamma_i, j_{\text{ign}_i})$ are transmitted with finite velocity towards the end regions \mathcal{R}_i , where the final outcomes are thus $x_i = \Pi(\gamma_i, j_{\text{ign}_i})$.

Conversely, let $g_{\gamma_{\text{ign}_i}}(\hat{\lambda}_j)$ be any gauge distribution, with $\gamma_{\text{ign}_i} = k_{\text{ign}_i} + i_{\text{ign}_i}K$. This gauge selection implies that the setting in region i_{ign_i} is $\theta_{k_{\text{ign}_i}}$. It is possible to compute the probability $P(\mathbf{x}|\mathbf{u})$. We have to collect all ignition states $\hat{\lambda}_j$ within the target (\mathbf{x}, \mathbf{u}) , given that $u_{i_{\text{ign}_i}} = \theta_{k_{\text{ign}_i}}$: Then, we have,

$$P(\mathbf{x}|\mathbf{u}) = \sum_{\hat{\lambda}_j \in (\mathbf{x}|\mathbf{u})} g_{\gamma_{\text{ign}_i}}(\hat{\lambda}_j) \quad (53)$$

where $\hat{\lambda}_j \in (\mathbf{x}|\mathbf{u})$ means that $\forall i, x_i = \Pi_i(u_i, \hat{\lambda}_j)$.

3.3.3 Multi-step collapse

If we fail in computing a feasible set of gauge distributions, $g_\gamma(\lambda_j)$, we try a two-step collapse. This means a gauge selection of one region and a local trial to check again a new one-step collapse, but now in only $n - 1$ regions. If we fail again, we iterate for a three-step collapse, etc. Finally, any n -region system admits at least a classical collapse in n steps.

3.4 Typical examples

We are now going to give some examples. In bipartite systems, we will consider the cases of partially entangled regions as well as ‘super-quantum’ states. In tripartite system, we will deal with the GHZ and the W states. In each example, we will derive the multipartite entanglement entropies.

3.4.1 One-region systems

Even if a system composed of a single entity \mathcal{E}_0 is trivially not entangled, it is convenient to have a general gauge formalism, valid in this limit case. Only the direct collapse is relevant. A one-region system with K settings (Table 14) has K configurations and K gauges distributions. It requires a working set of 2 ignition states (Table 11), as described in Table 15.

x_0	θ_0	θ_1	θ_2	θ_3	θ_4
0	p_0	p_1	p_2	p_3	p_4
1	$1 - p_0$	$1 - p_1$	$1 - p_2$	$1 - p_3$	$1 - p_4$

Table 14: One region: Typical probabilities $P(x|u)$ for 5 settings.

j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$	$g_4(j)$
0	p_0	p_1	p_2	p_3	p_4
31	$1 - p_0$	$1 - p_1$	$1 - p_2$	$1 - p_3$	$1 - p_4$

Table 15: One region: Typical gauge distributions $g_k(j)$ with 2 ignition states, $j = 0$ and $j = 2^K - 1 = 31$, for $K = 5$ settings.

3.4.2 Bipartite systems

We have already described Bell-type system within a particular formalism, devoted to totally correlated entities. Now, we will use the general formalism, valid for any degree of entanglement, including separable states.

Separable states Consider the case of two *independent* entities \mathcal{E}_0 and \mathcal{E}_1 . By definition, the entities are not entangled and the joint probability of the pair is defined by the product

$$P(x_0; x_1 | u_0; u_1) = \text{Prob}(x_0 | u_0) \times \text{Prob}(x_1 | u_1)$$

The entangled entropy matrix of separable states is of course zero.

The collapse mechanism involves physically two steps. Nevertheless, we have always found that a one-step collapse is successful. This opens the possibility of ‘purification’: a third region entangled with each of the two separable regions can be added to the system and a one-step overall collapse is likely to be triggered from this third region.

General bipartite 2-setting system Let $K = 2$. When we drop the condition of total correlation, Table 2 is expanded into Table 16, depending now on 8 parameters, q_1, q_2, \dots, q_8 . It is possible to define a one-step collapse as follows: Each gauge distribution is defined by a linear system. We have $2 \times 2 = 4$ gauge distributions. For each gauge distribution, there are 16 unknowns for 6 independent equations when accounting for local consistency. We have then 16 ignition states but the rank of the linear system is 6 and thus each distribution can be computed with only 6 non zero components. It is easy to compute the partial distribution in each region (Tables 17 and 18). The maximum bipartite entanglement entropy is equal to 1 bit. Especially, this maximum is obtained in totally correlated systems like Bell’s states (Table. 24) but also in more exotic objects like the PR-Box (Table. 20) below.

(x_0, x_1)	(θ_0, θ_0)	(θ_1, θ_1)	(θ_0, θ_1)	(θ_1, θ_0)
(0, 0)	$1 - q_1$	$1 - q_2$	$1 - q_3$	$1 - q_4$
(0, 1)	q_5	q_6	$q_3 - q_1 + q_5$	$q_4 - q_2 + q_6$
(1, 0)	q_7	q_8	$q_3 - q_2 + q_8$	$q_4 - q_1 + q_7$
(1, 1)	$q_1 - q_5 - q_7$	$q_2 - q_6 - q_8$	$q_1 + q_2 - q_3 - q_5 - q_8$	$q_1 + q_2 - q_4 - q_6 - q_7$

Table 16: Two regions: Conditional probabilities $P(x_0; x_1 | u_0; u_1)$ of the general *locally consistent* 2-region system with 2 possible settings θ_0 and θ_1 depending on 8 parameters q_1 to q_8 . All entries have to be non negative

\mathcal{R}_0	θ_0	θ_1
0	$1 - q_1 + q_5$	$1 - q_2 + q_6$
1	$q_1 - q_5$	$q_2 - q_6$

Table 17: Partial probabilities $\text{Prob}(x_0|u_0)$ for region \mathcal{R}_0 derived from Table 16.

\mathcal{R}_1	θ_0	θ_1
0	$1 - q_1 + q_7$	$1 - q_2 + q_8$
1	$q_1 - q_7$	$q_2 - q_8$

Table 18: Partial probabilities $\text{Prob}(x_1|u_1)$ for region \mathcal{R}_1 derived from Table 16.

Totally correlated pairs Of course, the general case includes the particular case of totally correlated pairs and provides new gauge distributions for this case. This inflation of distributions raises no real problem because ignition states and gauge distributions are not physical entities but only gauge parameters. For instance, we can recover the previous distributions of the EPR pairs by using a particular working set of $2K$ ignition states, namely $j = (2^K + 1)D_K(r)$ (with $0 \leq r \leq 2K - 1$, $D_K(r)$ being the double-plateau function of order K defined in Sec. 2.2.1). Then the gauge distributions g_k are duplicate and $g_{k+K} = g_k$.

PR-Box Another example of interest is the so-called ‘PR-Box’ proposed by Popescu and Rohrlich [35] and recalled in Table 19. This example is supposed to be hypothetical or ‘super-quantum’ because it exceeds the Tsirelson bound [57]. Now, we see that the box corresponds to an ‘ordinary’ 2-region-2-setting-system, with $q_1 = q_3 = q_4 = q_6 = q_8 = 1/2$, $q_2 = 1$ and $q_5 = q_7 = 0$ in Table 16. Therefore the collapse can be completed in classical physics, using e.g. the gauge distributions given in Table 21. A possible implementation as a classical game using a fair coin is proposed in Table 21 caption. The system is clearly maximally entangled and its entanglement entropy matrix is given in Table 20. This is an example of maximally entangled state which is not totally correlated (as defined in Sec. 2.1.3). The box is known to solve the problem of ‘communication complexity’ [58] in the sense that all distributed computations can be performed with a trivial amount of communication, i.e. with one bit [59]. This seems very surprising because the device can be easily implemented, but on the other hand, this classical realization involves a stage of communication. Finally, we have found that the box also describes the second-step collapse of the GHZ-states (Sec. 3.4.3 below).

x_0x_1	$(\theta_0\theta_0)$	$(\theta_0\theta_1)$	$(\theta_1\theta_0)$	$(\theta_1\theta_1)$
00	1/2	1/2	1/2	0
01	0	0	0	1/2
10	0	0	0	1/2
11	1/2	1/2	1/2	0

Table 19: PR-Box: Probabilities $P(x|u)$ proposed by Popescu and Rohrlich [35]. This system is maximally entangled (Table 20) and the process can be implemented in classical physics, using the gauge distributions given in Table 21.

3.4.3 Tripartite entangled systems

It is well known that there are two irreducible families of tripartite quantum entangled systems, referred to as GHZ states and W-states. We will first analyse the W-states, because the collapse is obtained in one step, without surprise, apart from the fact that these states are compatible with a simple quadruplet of hidden variables located at the ignition point. By contrast, the second family, the emblematic GHZ states with uniform tripartite entanglement, requires a two-step collapse.

W-type states Consider an ensemble of three entities $\{\mathcal{E}_0, \mathcal{E}_1, \mathcal{E}_2\}$, located respectively in three distant space regions $\mathcal{R}_0, \mathcal{R}_1$ and \mathcal{R}_2 located in a plane (X, Y) . The physical system is equivalent

$u_0 \backslash u_1$	0	1
0	1	1
1	1	1

Table 20: Entanglement entropy (in bits) of the PR-Box. This bipartite distribution ($n = 2$) is maximally entangled because $S_2(u_0, u_1) = C(u_0, u_1) = 1 = n - 1$ but not totally correlated.

j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$
0	1/2	0	1/2	0
6	0	1/2	0	1/2
9	0	1/2	0	1/2
15	1/2	0	1/2	0

Table 21: PR-Box: Gauge distribution working set with 4 ignition states (out of 16). This proves that the PR-box, while exceeding the Tsirelson bound [57], can be implemented in classical physics. We note that each distribution has only two equally likely outcomes. Therefore, it is possible to emulate the system as a simple game between two players \mathcal{O}_0 and \mathcal{O}_1 using a fair coin: Each player \mathcal{O}_i chooses her/his setting $k_i = 0$ or 1 and sends to a referee (located at the ignition point) her/his configuration, $\gamma_i = k_i + 2i$. The referee selects at random between γ_0 and γ_1 a gauge configuration, $\gamma_{\text{igni}} \in [0, 3]$, and tosses the coin. The coin is regarded as a gauge distribution $g_{\gamma_{\text{igni}}}(j)$ with an assignment of the relevant ignition states j to ‘head’ and ‘tail’ respectively, e.g., ‘head’ for $j \in \{3, 6\}$ and ‘tail’ for $j \in \{9, 15\}$. Then, the referee draws an ignition state j . For example, $j = 0$ or 15 if $\gamma_{\text{igni}} = 0$. At last, the final outcomes are $x_i = \Pi(\gamma_i, j)$ respectively for $i = 0, 1$. For example, if $\gamma_0 = 0$, $\gamma_1 = 1$, $\gamma_{\text{igni}} = 0$ and $j = 0$, we have $x_0 = x_1 = 0$.

to a set of three spin 1/2 particles, initially entangled in the state

$$|\psi\rangle = \frac{1}{\sqrt{3}}(|001\rangle_Z + |010\rangle_Z + |100\rangle_Z)$$

specifies along an axis Z perpendicular to the plane (X, Y) , where, e.g., $|0\rangle_Z$ stands for spin up and $|1\rangle_Z$ for spin down. Later the spins of the three particles are measured independently in the distant regions along either the axis X referred to as $\theta_0 \in \Theta$ or the axis $Y = \theta_1$. In each region, an observer selects freely a setting u_i ($i = 0, 1$ or 2). The conditional probabilities

$(u_0, u_1, u_2) \rightarrow$ (x_0, x_1, x_2) \downarrow	$(\theta_1, \theta_0, \theta_0)$ $(\theta_0, \theta_1, \theta_1)$	$(\theta_0, \theta_1, \theta_0)$ $(\theta_1, \theta_0, \theta_1)$	$(\theta_0, \theta_0, \theta_1)$ $(\theta_1, \theta_1, \theta_0)$	$(\theta_1, \theta_1, \theta_1)$ $(\theta_0, \theta_0, \theta_0)$
(0, 0, 0)	1/4	1/4	1/4	1/2
(0, 0, 1)	0	0	1/4	0
(0, 1, 0)	0	1/4	0	0
(0, 1, 1)	1/4	0	0	0
(1, 0, 0)	1/4	0	0	0
(1, 0, 1)	0	1/4	0	0
(1, 1, 0)	0	0	1/4	0
(1, 1, 1)	1/4	1/4	1/4	1/2

Table 22: W-states: Example of conditional probability $P(x_0; x_1; x_2 | u_0; u_1; u_2)$ of a tripartite-2-setting system.

$P(x_0; x_1; x_2 | u_0; u_1; u_2) = P(\mathbf{x} | \mathbf{u})$ are given in Table 22 and clearly depends upon the settings in the three regions. Complete consistency holds and, e.g., the partial probability $\text{Prob}(x_0 | u_0)$ in region \mathcal{R}_0 does not depend on the setting in \mathcal{R}_1 and \mathcal{R}_2 . Thus, we have $(\forall u_0, \forall u_1, \forall u_2 \in \Theta)$:

$$\text{Prob}(x_0 | u_0) = \sum_{x_1=0}^1 \sum_{x_2=0}^1 P(x_0; x_1; x_2 | u_0; u_1; u_2) = \frac{1}{2} \quad (54)$$

and similarly for ports \mathcal{R}_1 and \mathcal{R}_2 . If we ignore the last port, the probability $\text{Prob}(x_0; x_1|u_0; u_1)$ in regions $(\mathcal{R}_0, \mathcal{R}_1)$, does not depend on the setting in \mathcal{R}_2 . Therefore, we have $(\forall u_0, \forall u_1, \forall u_2 \in \Theta)$:

$$\text{Prob}(x_0; x_1|u_0; u_1) = \sum_{x_2=0}^1 P(x_0; x_1; x_2|u_0; u_1; u_2) \quad (55)$$

The result is given in Table 23. This is the probability distribution of a Bell's state, i.e., an EPR

(x_0, x_1)	(θ_0, θ_0)	(θ_0, θ_1)	(θ_1, θ_0)	(θ_1, θ_1)
(0, 0)	1/2	1/4	1/4	1/2
(0, 1)	0	1/4	1/4	0
(1, 0)	0	1/4	1/4	0
(1, 1)	1/2	1/4	1/4	1/2

Table 23: W-states: Partial probability $\text{Prob}(x_0; x_1|u_0; u_1)$. This is just the probability distribution of a Bell's state

pair with $\theta_0 = 0$ and $\theta_1 = \pi/2$, obtained from the general 2-region 2-setting system (Table 16) for $q_1 = q_2 = 1/2$, $q_3 = q_4 = 3/4$ and $q_5 = q_6 = q_7 = q_8 = 0$. Thus *total correlation holds between any pair of regions*. As a result the bipartite degree of entanglement is $e_2 = 3$. It is easy to compute the bipartite entanglement entropy matrices. The result for one pair of regions is given in Table 24.

$u_0 \backslash u_1$	0	1
0	1	0
1	0	1

Table 24: Partial W-states: Bipartite entanglement entropy $S_2(u_0, u_1)$ (in bits) of a Bell's state.

(u_0, u_1, u_2)	(000)	(001)	(010)	011	(100)	(101)	(110)	(111)
$S_3(u_0, u_1, u_2)$	1	0	0	0	0	0	0	1
$C(u_0, u_1, u_2)$	2	1	1	1	1	1	1	2

Table 25: W-states: Tripartite entanglement entropy $S_3(u_0, u_1, u_2)$ and total entanglement $C((u_0, u_1, u_2))$ (in bits). Since $C(0, 0, 0) = C(1, 1, 1) = 2$ bits, this state is maximally entangled. For simplicity, (ijk) stands for $(\theta_i, \theta_j, \theta_k)$.

Coming back to the full system, the number of configurations is $nK = 3 \times 2 = 6$. Therefore, the number of fine grains is $2^{K^n} = 2^8 = 256$. However, we can use the standard projection function, Eq. (47), with just $2^{nK} = 2^6 = 64$ ignition states and 6 stochastic gauge distributions, $g_\gamma(\lambda_j)$ with $\gamma = 0, \dots, 5$. Surprisingly, it is possible to compute a particular working set of six *identical* gauge distributions with only 4 ignition states, namely $j \in \{0, 21, 42, 63\}$, as described in Table 26. This means that the system is degenerate and compatible with a set of random *hidden variables* located at the ignition point. For instance, suppose that a global break of equilibrium is triggered from region \mathcal{R}_0 ($i = 0$) with the setting $u_0 = \theta_0$ ($k = 0$, $\gamma_0 = k + 2i = 0$). In principle, we have to select the gauge distribution $g_0(j)$ but here all distributions are identical. We perform a trial and we can only draw one of the 4 equally likely ignition states $j = 0, 21, 42$ or 63 , say e.g., $j = 42 = 2^5 + 2^3 + 2$. If for instance, we have $u_1 = \theta_1$ ($\gamma_1 = 3$) and $u_2 = \theta_0$ ($\gamma_2 = 2$), we obtain $x_0 = 0$, $x_1 = 1$, $x_2 = 0$.

GHZ-states The second family of tripartite quantum entangled system is the GHZ-states, defined by Greenberger et al [60]. The physical system is a set of three spin 1/2 particles, initially entangled in the state

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|000\rangle_Z + |111\rangle_Z)$$

specifies along an axis Z perpendicular to the plane (X, Y) . Again, the spins of the three particles are measured along either the axis $X = \theta_0$ or the axis $Y = \theta_1$.

The target probabilities $P(\mathbf{x}|\mathbf{u})$ are given in Table 27. In non-contextual systems, one would expect that each local outcome only depends on a local trial. This assumption is easily checked

j	$g(j)$
0	1/4
21	1/4
42	1/4
63	1/4

Table 26: W-states: Gauge distribution working set with 4 ignition states (out of 64). The 6 gauge distributions are identical. This means that the W-states can be described by a quadruplet of hidden variables, $\lambda_0, \lambda_{21}, \lambda_{42}$ and λ_{63} located at the ignition point.

by assuming the existence of local random functions, defined as $X(\mathcal{R}_i|\theta_k) = x_i$ when the entity \mathcal{E}_i is measured with the setting θ_k . Let $f(u_0, u_1, u_2) \equiv X(\mathcal{R}_0|u_0) + X(\mathcal{R}_1|u_1) + X(\mathcal{R}_2|u_2) \pmod{2}$. Clearly, $f(\theta_1, \theta_1, \theta_1) \equiv f(\theta_0, \theta_0, \theta_1) + f(\theta_0, \theta_1, \theta_0) + f(\theta_1, \theta_0, \theta_0) \pmod{2}$. The assumption is falsified by Table 27, since $f(\theta_1, \theta_1, \theta_1) \equiv 1$ with certainty, while $f(\theta_0, \theta_0, \theta_1) \equiv 0$, $f(\theta_0, \theta_1, \theta_0) \equiv 0$ and $f(\theta_1, \theta_0, \theta_0) \equiv 0$ with certainty.

$(u_0, u_1, u_2) \rightarrow$ (x_0, x_1, x_2) \downarrow	$(\theta_0, \theta_0, \theta_0)$ $(\theta_1, \theta_1, \theta_1)$	$(\theta_0, \theta_0, \theta_1)$ $(\theta_1, \theta_1, \theta_0)$	$(\theta_0, \theta_1, \theta_0)$ $(\theta_1, \theta_0, \theta_1)$	$(\theta_0, \theta_1, \theta_1)$ $(\theta_1, \theta_0, \theta_0)$
(0, 0, 0)	0	1/4	1/4	1/4
(0, 0, 1)	1/4	0	0	0
(0, 1, 0)	1/4	0	0	0
(0, 1, 1)	0	1/4	1/4	1/4
(1, 0, 0)	1/4	0	0	0
(1, 0, 1)	0	1/4	1/4	1/4
(1, 1, 0)	0	1/4	1/4	1/4
(1, 1, 1)	1/4	0	0	0

Table 27: Strict GHZ states: Conditional probability $P(x_0; x_1; x_2|u_0; u_1; u_2)$ of a strict GHZ system for two different settings θ_0 and θ_1 . Local realism is supposed to be falsified by computing a convenient function, $f(u_0, u_1, u_2) \equiv x_0 + x_1 + x_2 \pmod{2}$. Whatever the outcomes, we see by inspection that $f(\theta_1, \theta_1, \theta_1) \equiv 1$ with certainty, while $f(\theta_0, \theta_0, \theta_1) \equiv 0$, $f(\theta_0, \theta_1, \theta_0) \equiv 0$ and $f(\theta_1, \theta_0, \theta_0) \equiv 0$ with certainty. Furthermore, this strict GHZ states is an example of tripartite system that can only collapse in two steps

Complete consistency holds. For instance, the partial probability $\text{Prob}(x_0|u_0)$ in region \mathcal{R}_0 does not depend on the setting in \mathcal{R}_1 and \mathcal{R}_2 . Thus, we have $(\forall u_0, \forall u_1, \forall u_2 \in \Theta)$:

$$\text{Prob}(x_0|u_0) = \sum_{x_1=0}^1 \sum_{x_2=0}^1 P(x_0; x_1; x_2|u_0; u_1; u_2) = \frac{1}{2} \quad (56)$$

and similarly for ports \mathcal{R}_1 and \mathcal{R}_2 . If we ignore the last port, the probability $\text{Prob}(x_0; x_1|u_0; u_1)$ in regions $(\mathcal{R}_0, \mathcal{R}_1)$, does not depend on the setting in \mathcal{R}_2 . Therefore, we have $(\forall u_0, \forall u_1, \forall u_2 \in \Theta)$:

$$\text{Prob}(x_0; x_1|u_0; u_1) = \sum_{x_2=0}^1 P(x_0; x_1; x_2|u_0; u_1; u_2) = \frac{1}{4} \quad (57)$$

This is obtained from the general 2-region 2-setting system (Table 16) for $q_1 = q_2 = q_3 = q_4 = 3/4$ and $q_6 = q_6 = q_7 = q_8 = 1/4$. Thus, the two regions are separable and the probabilities given by Eq(57) are simply the product of the probability obtained in Eq.(56). Therefore the entanglement scheme is (0, 1). The bipartite entanglement entropy for any pair of region is zero. Whatever the settings, the signed tripartite entanglement entropy is equal to -1 bit and the total entanglement to 1 bit (Table 28).

We are now going to break the global equilibrium. We first check a one-step collapse. The number of configurations is $3 \times 2 = 6$. In order to classically encode Table 27, we construct an ignition set $\Lambda = \{\lambda_j\}$ with $2^6 = 64$ ignition states, labelled j from $j = 0$ to 63, and 6 stochastic gauge distributions, $g_\gamma(\lambda_j)$ with $\gamma = 0, \dots, 5$. However, none of the six linear systems admits non negative solution. To circumvent this problem, we can try a different ignition set. The maximum number of distinct fine grains is $2^{2^3} = 256$. However, it is easily shown that these 256 fine grains

(u_0, u_1, u_2)	(000)	(001)	(010)	(011)	(100)	(101)	(110)	(111)
$S_3(u_0, u_1, u_2)$	-1	-1	-1	-1	-1	-1	-1	-1
$C(u_0, u_1, u_2)$	1	1	1	1	1	1	1	1

Table 28: GHZ-states: Tripartite entanglement entropy $S_3(u_0, u_1, u_2)$ and total entanglement $C(u_0, u_1, u_2)$ (in bits). Bipartite entanglement entropies $S_2(u_i, u_j)$ are identically zero. These coefficients do not depend on the settings.

are all included into at least one target of zero probability. Therefore, whatever the gauge, the probability of all ignition states will always be zero. In conclusion, within the realm of local realism, the strict GHZ state *cannot collapse in one step*.

We have thus to try a two-step collapse. Let \mathcal{I} be the ignition point at the boundary of the three regions. Each observer selects freely his own setting, $u_i = k_i$ and sends his choice towards \mathcal{I} , with a finite velocity. Suppose that the first received configuration is $u_2 = k_2$ from region \mathcal{R}_2 . As a result, we break the equilibrium at \mathcal{I} from region \mathcal{R}_2 . We draw the outcome $x_2 = \xi_2$ from the local probability distribution $\text{Prob}(x_2|k_2)$, Eq.(56). The two possible outcomes $\xi_2 \in \{0, 1\}$ are equally likely. Next, in regions $\mathcal{R}_0, \mathcal{R}_1$ we compute the second-step probability distribution $P'(x_0; x_1|u_0; u_1) = \text{Prob}(x_0; x_1|\xi_2; k_2; u_0; u_1)$. The resulting distribution is given in Table 29. Note that this distribution does not depend on ξ_2 . On the other hand, if the observer \mathcal{O}_2 had chosen $k_2 = 1$ instead of $k_2 = 0$, the distribution $P'(x_0; x_1|u_0; u_1)$, now given in Table 30, would have been different. These probability distributions are identical to the probabilities of the PR-box, Table 19, (apart from the labelling). These second-step distributions are locally consistent and

x_0x_1	$\theta_0\theta_0$	$\theta_0\theta_1$	$\theta_1\theta_0$	$\theta_1\theta_1$
00	0	1/2	1/2	1/2
01	1/2	0	0	0
10	1/2	0	0	0
11	0	1/2	1/2	1/2

Table 29: GHZ-states: Second-step probability $P'(x_0; x_1|u_0; u_1)$ in regions $\mathcal{R}_0, \mathcal{R}_1$, following a gauge selection of region \mathcal{R}_2 , a free choice $u_2 = \theta_0$ and a first trial in \mathcal{R}_2 . This distributions is identical to the probability distribution of the PR-box, Table 19, (apart from the labelling).

x_0x_1	$(\theta_0\theta_0)$	$(\theta_0\theta_1)$	$(\theta_1\theta_0)$	$(\theta_1\theta_1)$
00	1/2	0	0	0
01	0	1/2	1/2	1/2
10	0	1/2	1/2	1/2
11	1/2	0	0	0

Table 30: GHZ-states: Second-step probability $P'(x_0; x_1|u_0; u_1)$ in regions $\mathcal{R}_0, \mathcal{R}_1$, following a gauge selection of region \mathcal{R}_2 , a free choice $u_2 = \theta_1$ (different from the free choice of Table 29) and a first trial in \mathcal{R}_2 . This distributions is identical to the probability distribution of the PR-box, Table 19, (apart from the labelling).

the local probabilities, e.g. in region \mathcal{R}_0 are computed as

$$\text{Prob}(x_0|u_0) = \sum_{x_1=0}^1 P'(x_0; x_1|u_0; u_1) = \frac{1}{2} \quad (58)$$

in accordance with Eq.(56). Furthermore, they are maximally entangled but not totally correlated (see above Sec.2.1.3) and their single bipartite entropy matrix is given in Table 20.

To complete the two-step collapse, we have now to construct an ignition set for the two regions \mathcal{R}_0 and \mathcal{R}_1 , accounting for the relevant distribution $P'(x_0; x_1|u_0; u_1)$. This is easily achieved because the new system is actually a PR-box. Each gauge distribution requires just two ignition states, but we need four ignition states for the whole, as shown in Table 31. This proves that the strict GHZ-states can be classically simulated, but only a two-step collapse is feasible.

We have checked whether a *quasi-GHZ* state can collapse in one step. Indeed, in the strict GHZ state, a number of targets have zero probability. It is possible to replace 0 by a small

j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$
0	0	1/2	0	1/2
6	1/2	0	1/2	0
9	1/2	0	1/2	0
15	0	1/2	0	1/2

Table 31: Gauge distribution working set with 4 ignition states (out of 16) for the second-step probability $P'(x_0; x_1|u_0; u_1)$ given in Table 29. This is a relabelling of Table 21.

$(u_0, u_1, u_2) \rightarrow$ (x_0, x_1, x_2) \downarrow	$(\theta_0, \theta_0, \theta_0)$ $(\theta_1, \theta_1, \theta_1)$	$(\theta_0, \theta_0, \theta_1)$ $(\theta_1, \theta_1, \theta_0)$	$(\theta_0, \theta_1, \theta_0)$ $(\theta_1, \theta_0, \theta_1)$	$(\theta_0, \theta_1, \theta_1)$ $(\theta_1, \theta_0, \theta_0)$
(0, 0, 0)	ϵ	$1/4 - \epsilon$	$1/4 - \epsilon$	$1/4 - \epsilon$
(0, 0, 1)	$1/4 - \epsilon$	ϵ	ϵ	ϵ
(0, 1, 0)	$1/4 - \epsilon$	ϵ	ϵ	ϵ
(0, 1, 1)	ϵ	$1/4 - \epsilon$	$1/4 - \epsilon$	$1/4 - \epsilon$
(1, 0, 0)	$1/4 - \epsilon$	ϵ	ϵ	ϵ
(1, 0, 1)	ϵ	$1/4 - \epsilon$	$1/4 - \epsilon$	$1/4 - \epsilon$
(1, 1, 0)	ϵ	$1/4 - \epsilon$	$1/4 - \epsilon$	$1/4 - \epsilon$
(1, 1, 1)	$1/4 - \epsilon$	ϵ	ϵ	ϵ

Table 32: Quasi-GHZ states: Conditional probability $P(x_0; x_1; x_2|u_0; u_1; u_2)$ of a quasi-GHZ system. Target probabilities are derived from the strict GHZ state by replacing 0 by ϵ in Table 27. A one-step collapse is feasible for $1/16 < \epsilon < 3/16$. Partial probabilities Eq.(56 and 57) are unchanged.

probability ϵ as shown in Table 32. In spite of this change, the partial probabilities, Eq. (56) and (57) are strictly conserved and therefore, any pair of regions remains separable. As a result, the entanglement scheme is still (0, 1), except when the three regions become independent. Thus, the new system may actually be viewed as a *quasi-GHZ* state, provided that ϵ remains small. By computer simulation, we have found that a one-step collapse is feasible only when ϵ is in the range $]1/16, 3/16[$. For $\epsilon = 2/16$, the three regions become independent. For instance, when $\epsilon = 1/16^+$, the tripartite entanglement entropy is given in Table 33. The strength of the tripartite entanglement is then rather low (-0.19 bit instead of -1 bit). A one-step collapse is feasible and we have computed a working set of gauge probability distribution given in Table 34.

For $\epsilon = 1/16^+$, we have checked whether the system passes the test of ‘local realism’ (in fact, non-contextuality). With the same notations as above, the probability that $f(\theta_1, \theta_1, \theta_1) \equiv 1 \pmod{2}$ is now 0.75 (instead of 1), while the probability that $f(\theta_0, \theta_0, \theta_1) \equiv 0$, $f(\theta_0, \theta_1, \theta_0) \equiv 0$ and $f(\theta_1, \theta_0, \theta_0) \equiv 0$ is 0.56 (instead of 1). We can conclude that the behaviour of a quasi-GHZ state is similar to a strict mode, but there is no dramatic all-versus-nothing outcome as in the strict state.

These results have to be compared with the experiment [61]. The GHZ-modes were constructed from bipartite entangled pairs [62] and thus there is no way to distinguish between strict and quasi-mode at this stage. In addition, due to low efficiency of the source, the experimental results of the ‘local realism’ test are smooth and seem compatible with the behaviour of the quasi-mode.

More generally, the current model should open the possibility to *classically* synthesize any locally consistent system, whether quantum or not, whatever its degree of entanglement. One has to define a number of regions, n , a number of settings, K , n local measurement entropies and $\binom{n}{m} K^m$ coefficients of entanglement for $m = 2$ to n . This is similar to the analysis of the

(u_0, u_1, u_2)	(000)	(001)	(010)	(011)	(100)	(101)	(110)	(111)
$S_3(u_0, u_1, u_2)$	-0.19	-0.19	-0.19	-0.19	-0.19	-0.19	-0.19	-0.19
$C(u_0, u_1, u_2)$	0.19	0.19	0.19	0.19	0.19	0.19	0.19	0.19

Table 33: Quasi-GHZ state with $\epsilon = 1/16$: Tripartite entanglement entropy and total entanglement (in bits). Whatever the settings, the tripartite entropy is equal to -0.19 bit.

j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$	$g_4(j)$	$g_5(j)$	j	$g_0(j)$	$g_1(j)$	$g_2(j)$	$g_3(j)$	$g_4(j)$	$g_5(j)$
0	0.062	0.062	0.062	0.062	0.062	0.063	1	0.000	0.000	0.063	0.000	0.063	0.000
2	0.000	0.000	0.000	0.063	0.000	0.062	4	0.063	0.000	0.000	0.000	0.063	0.000
6	0.000	0.000	0.063	0.000	0.062	0.000	7	0.000	0.063	0.000	0.000	0.000	0.000
8	0.000	0.063	0.000	0.000	0.000	0.062	9	0.063	0.000	0.000	0.063	0.063	0.063
11	0.000	0.000	0.000	0.000	0.062	0.000	13	0.000	0.000	0.000	0.000	0.000	0.063
14	0.000	0.000	0.000	0.000	0.063	0.000	15	0.062	0.062	0.062	0.062	0.063	0.000
16	0.063	0.000	0.063	0.000	0.000	0.000	18	0.000	0.062	0.063	0.063	0.063	0.000
19	0.000	0.000	0.000	0.062	0.062	0.000	21	0.063	0.000	0.063	0.000	0.063	0.000
22	0.000	0.063	0.000	0.000	0.000	0.063	23	0.000	0.000	0.063	0.000	0.063	0.063
24	0.063	0.063	0.000	0.062	0.062	0.000	25	0.000	0.000	0.000	0.063	0.000	0.000
26	0.000	0.000	0.000	0.000	0.063	0.000	28	0.000	0.062	0.000	0.000	0.063	0.000
29	0.063	0.000	0.000	0.000	0.062	0.000	31	0.000	0.000	0.000	0.000	0.000	0.062
32	0.000	0.063	0.000	0.063	0.000	0.000	33	0.062	0.000	0.063	0.063	0.000	0.063
35	0.000	0.000	0.062	0.000	0.000	0.062	36	0.063	0.063	0.062	0.000	0.000	0.062
37	0.000	0.000	0.000	0.000	0.000	0.063	38	0.000	0.000	0.063	0.000	0.000	0.000
41	0.063	0.000	0.000	0.000	0.000	0.000	42	0.000	0.063	0.000	0.063	0.000	0.063
43	0.000	0.000	0.000	0.063	0.000	0.063	44	0.062	0.000	0.000	0.000	0.000	0.063
46	0.000	0.063	0.000	0.000	0.000	0.062	49	0.063	0.000	0.000	0.062	0.000	0.000
50	0.000	0.063	0.062	0.000	0.000	0.000	51	0.000	0.000	0.063	0.063	0.000	0.000
52	0.000	0.062	0.063	0.000	0.000	0.000	53	0.062	0.000	0.062	0.000	0.000	0.000
56	0.062	0.000	0.000	0.063	0.000	0.000	58	0.000	0.062	0.000	0.062	0.000	0.000
60	0.063	0.063	0.000	0.000	0.000	0.000							

Table 34: One step collapse of the quasi-GHZ mode with $\epsilon = 1/16$ using a gauge distribution working set with 43 ignition states for the whole (out of 64).

so-called ‘non-signaling polytope’ in quantum information. This discussion is beyond the scope of the present paper.

4 Conclusion

Contextuality is not the privilege of the quantum world. Quite the reverse, we have shown in a previous paper [25] that a strictly classical dice game can be contextually dependent and even non-signaling in the quantum sense. Therefore, the difficult task to reconcile quantum mechanics with non-contextual logic is unnecessary. In the current paper we have constructed a model named ‘stochastic gauge system’ to compute contextually dependent classical systems. We have shown that the theory described quantum randomness as well and can be used to simulate quantum states and even non-local boxes in classical physics, e.g., EPR pairs, GHZ states or PR-boxes. Quantum systems are compared with classical extended objects in equilibrium and quantum collapses are therefore identified with classical breaks of equilibrium. Furthermore, the model gives a straightforward classical interpretation of entanglement and entanglement entropy and provides simple tools for characterizing multipartite entanglements.

In our opinion, the present theory opens the way for a number of important questionings, regarding quantum theory, quantum information and even the general conception of physics, obviously beyond the scope of this paper.

In quantum mechanics, we have shown that classical local realism is compatible with the conventional formulation of the theory, contrary to a general belief. This result is likely to explain a number of difficulties and opens up also new topics. For instance, in the relativistic domain, the conventional interpretation fails to define a consistent probability distribution. In our opinion, the concept of gauge probability should be a key ingredient to settle this problem. The notion of quantum collapse should be also revisited and compared with a classical break of equilibrium. New issues, like collapse kinetics should probably be investigated.

The model proposes also an interpretation of quantum parallelism. A quantum system can be viewed as a complex entangled object in equilibrium. When the equilibrium is broken, a number of new situations become potentially possible. Quantum parallelism describes the set of all these potential situations. In quantum information theory, a final measurement will draw just one outcome. The theory uses entanglement as a resource, but again, this is not a quantum privilege. In our opinion, contextuality should be a resource in classical information as well. Indeed, ‘quantum randomness’ can be viewed as strictly classical, provided that one includes an additional concept in the conventional stochastic theory, namely, the notion of gauge distribution. Classical simulation of quantum algorithms should open up new paths in classical computation, beside deterministic and probabilistic algorithms. More generally, a number of notions like quantum cryptography,

quantum complexity or communication complexity (see above Sec. 3.4.2) should probably deserve to be revisited.

Nowadays, the principle of gauge invariance plays a key role in modern physics. This is not very surprising because every physical measurement is performed within a particular framework while physical reality has to be independent of any referential and any system of units. Therefore, mathematical tools have been devised: Dimensional analysis, covariance principle, and more generally, gauge theories. The gauge invariance principle has proved to be extraordinary powerful in all branches of physics.

Surprisingly, the measurement in quantum theory remains nevertheless thought as *absolute*. The seed of this conception lies probably in the very origin of the theory, i.e., the Copenhagen interpretation of quantum mechanics, according to which the squared amplitude of the wave function is an *absolute probability*. In our opinion, this belief is inconsistent, as highlighted by the EPR paradox or the problematic generalization of this interpretation in special relativity.

The view of an absolute probability space is also the origin of a controversy between ‘Bayesians’ and ‘Frequentists’ in the probability domain. In our opinion, the ‘Frequentist’ position is simply not sustainable. For instance, we have emphasized in the present paper that a number of fundamental concepts like partial measurements are subjective *by definition*.

Actually, physics aims to understand the entirety of the universe, in which a number of long range interactions are recorded. By contrast, each observer is embedded into *his* own limited *causal bubble* or *causal diamond*, bounded by light sheets, black hole and cosmological horizons. Therefore, he can only describe a finite part of the world and, apart from long range interactions, he has to account for the rest of the universe by *his* best evaluation of *his* irreducible ignorance, technically in form of entropy located on *his* causal horizons.

This vision of the universe seems very new in physics: Unpredictability is no more in contradiction with determinism but, quite the reverse, an inevitable consequence. This implies a break between the general Laplace concept of *determinism* and the technical notion of *causality*. In this respect, the dispute concerning the completeness of quantum mechanics appears totally unrealistic.

References

- [1] A. Einstein, B. Podolsky, N. Rosen, Can quantum mechanical description of reality be considered complete?, Phys. Rev. 47 (1935) 777.
- [2] D. Bohm, A suggested interpretation of the quantum theory in terms of ‘hidden variables’, Phys. Rev. 85 (1952) 166–193.
- [3] J. S. Bell, On the Einstein-Podolsky-Rosen paradox, Physics 195 (1964) 1.
- [4] J. S. Bell, Speakable and unspeakable in quantum mechanics, Cambridge University Press, Cambridge, 1987.
- [5] S. Kochen, E. Specker, The problem of hidden variables in quantum mechanics, Journal of Mathematics and Mechanics 17 (1967) 59–87.
- [6] M. Genovese, Research on hidden variable theory: a review of recent progresses, Physics Reports 413 (2005) 319. [arXiv:quant-ph/0701071v1](https://arxiv.org/abs/quant-ph/0701071v1).
- [7] S. P. Gudder, Quantum Probability, Academic Press, San Diego, Ca, USA, 1988.
- [8] J. Evans, A. S. Thorndike, Quantum Mechanics at the Crossroads: New Perspectives from History, Philosophy and Physics, Springer, Cambridge, 2006.
- [9] P. Goyal, Information-geometric reconstruction of quantum theory, Phys. Rev. A 78 (5) (2008) 052120. [arXiv:0805.2761](https://arxiv.org/abs/0805.2761).
- [10] R. B. Griffiths, Consistent Quantum Theory, Cambridge University Press, 2003.
- [11] C. Rovelli, Relational quantum mechanics, International Journal of Theoretical Physics 35 (1996) 1637. [arXiv:quant-ph/9609002](https://arxiv.org/abs/quant-ph/9609002).
- [12] C. A. Fuchs, Quantum mechanics as quantum information (and only a little more) (2002). [arXiv:quant-ph/0205039](https://arxiv.org/abs/quant-ph/0205039).
- [13] E. T. Jaynes, Clearing up mysteries, in: J. Skilling (Ed.), Maximum entropy and Bayesian methods, Kluwer, Dordrecht, 1989, pp. 1–29.
URL <http://bayes.wustl.edu/etj/articles/cmystery.pdf>
- [14] A. F. Kracklauer, The error in Bell’s theorem (1998). [arXiv:quant-ph/9810081](https://arxiv.org/abs/quant-ph/9810081).

- [15] K. Hess, W. Philipp, Breakdown of Bell's theorem for certain objective local parameter spaces, *Proc. Nat. Acad. Sciences USA* 101 (7) (2004) 1799.
- [16] K. Hess, W. Philipp, *Europhys. Lett.* 57 (2002) 775. [arXiv:quant-ph/0103028](#).
- [17] K. Hess, W. Philipp, Bell's theorem: Critique of proofs with and without inequalities (2004). [arXiv:quant-ph/0410015](#).
- [18] A. Y. Khrennikov, *Interpretations of Probability*, VSP Int. Sc. Publishers, Utrecht, 1999.
- [19] A. Khrennikov, I. Volovich, Local realism, contextualism and loopholes in Bell's experiments (2002). [arXiv:quant-ph/0212127](#).
- [20] L. Accardi, *Urne e camaleonti. Dialogo sulla realtà, le leggi del caso e la teoria quantistica*, Il Saggiatore, 1997.
- [21] L. Accardi, M. Regoli, Locality and Bell's inequality (2000). [arXiv:quant-ph/0007005](#).
- [22] M. Clover, Bell's theorem: A critique (2005). [arXiv:quant-ph/0502016](#).
- [23] G. Adenier, A refutation of Bell's theorem (2000). [arXiv:quant-ph/0006014](#).
- [24] M. Feldmann, New loophole for the Einstein-Podolsky-Rosen paradox, *Found. Phys. Lett.* 8 (1) (1995) 41–53. [arXiv:quant-ph/9904051](#), [doi:10.1007/BF02187530](#).
- [25] M. Feldmann, Classical counterexample to Bell's theorem (2008). [arXiv:0812.4506\[physics.gen-ph\]](#).
- [26] M. B. Plenio, V. Vedral, Entanglement in quantum information theory, *Contemporary Physics* 39 (1998) 431. [arXiv:quant-ph/9804075](#).
- [27] M. B. Plenio, S. Virmani, An introduction to entanglement measures, *Quantum information & computation* 7 (2007) 1. [arXiv:quant-ph/0504163](#).
- [28] G. 't Hooft, The holographic principle (2000). [arXiv:hep-th/0003004](#).
- [29] L. Susskind, The world as a hologram, *J.Math.Phys.* 36 (1995) 6377–6396. [arXiv:hep-th/9409089v2](#).
- [30] T. Dauxois, S. Ruffo, E. Arimondo, M. Wilkens, Dynamics and thermodynamics of systems with long range interactions: An introduction, in: *Lecture Notes in Physics*, volume 602, Springer, Berlin, Heidelberg, 2002, pp. 1–19. [arXiv:cond-mat/0208455v1](#).
- [31] M. Antoni, S. Ruffo, A. Torcini, Dynamics and statistics of simple models with infinite-range attractive interaction (1999). [arXiv:cond-mat/9908336](#).
- [32] R. W. Hamming, Error detecting and error correcting codes, *Bell System Technical Journal* 26 (2) (1950) 147–160.
URL <http://guest.engelschall.com/~sb/hamming/>
- [33] E. Santos, Relation of the Bell inequalities with quantum logic, hidden variables and information theory, *Physics Letters A* 115 (1986) 363. [arXiv:quant-ph/0207062](#).
- [34] J. F. Clauser, M. A. Horne, A. Shimony, R. A. Holt, Proposed experiment to test local hidden-variable theories, *Phys. Rev. Lett.* 23 (15) (1969) 880–884. [doi:10.1103/PhysRevLett.23.880](#).
- [35] S. Popescu, D. Rohrlich, Quantum nonlocality as an axiom, *Foundations of Physics* 24 (1994) 379.
- [36] D. Rohrlich, S. Popescu, Nonlocality as an axiom for quantum theory (1995). [arXiv:quant-ph/9508009](#).
- [37] L. Masanes, A. Acin, N. Gisin, General properties of nonsignaling theories, *Physical Review A* 73 (2006) 012112. [arXiv:quant-ph/0508016](#).
- [38] G. Rousseaux, C. Mathis, P. Maissa, T. G. Philbin, U. Leonhardt, Observation of negative phase velocity waves in a water tank: A classical analogue to the Hawking effect? (2007). [arXiv:0711.4767](#).
- [39] T. G. Philbin, C. Kuklewicz, S. Robertson, S. Hill, F. Konig, U. Leonhardt, Fiber-optical analogue of the event horizon (2007). [arXiv:0711.4796](#).

- [40] P. Pearle, How stands collapse (2006). [arXiv:quant-ph/0611211;quant-ph/0611212](#).
- [41] R. Bousso, The holographic principle, *Reviews of Modern Physics* 74 (2002) 825. [arXiv:hep-th/0203101](#).
- [42] T. Jacobson, R. Parentani, Horizon entropy, *Found. Phys.* 33 (2003) 323. [arXiv:gr-qc/0302099](#).
- [43] C. E. Shannon, A mathematical theory of communication, *Bell System Technical Journal* 27 (1948) 379–423 , 656–715.
- [44] E. T. Jaynes, The Gibbs paradox, in: G. Erickson, P. Neudorfer, C. R. Smith (Eds.), *Maximum entropy and Bayesian methods*, Kluwer, Dordrecht, 1992. URL <http://bayes.wustl.edu/etj/articles/gibbs.paradox.pdf>
- [45] R. M. Fano, The transmission of information, Technical Report No. 65. Research Laboratory of Electronics, M.I.T., Cambridge (Mass.), USA. (1949).
- [46] R. W. Yeung, *Information Theory and network coding*, Springer-Verlag, New York, 2008.
- [47] A. Caticha, *Lectures on probability, entropy, and statistical physics* (2008). [arXiv:0808.0012v1\[physics.data-an\]](#).
- [48] S. Kullback, R. A. Leibler, On information and sufficiency, *Annals of Mathematical Statistics* 22 (1951) 79–86.
- [49] R. Horodecki, P. Horodecki, M. Horodecki, K. Horodecki, Quantum entanglement, *Reviews of Modern Physics* 81 (2009) 865. [arXiv:quant-ph/0702225](#).
- [50] V. P. Belavkin, M. Ohya, Entanglement, quantum entropy and mutual information, *PROC. R. SOC. LOND. A* 458 (2002) 209. [arXiv:quant-ph/0208111](#).
- [51] B. F. Toner, D. Bacon, The communication cost of simulating Bell correlations, *Physical Review Letters* 91 (2003) 187904. [arXiv:quant-ph/0304076v3](#).
- [52] G.-D. Hu, On the amount of information, *Teor. Veroyatnost. i Primenen* 4 (1962) 447–455, (in Russian).
- [53] R. W. Yeung, A new outlook on Shannon’s information measures, *IEEE Transactions on Information Theory* 37 (1991) 466–474.
- [54] S. Watanabe, Information theoretical analysis of multivariate correlation, *IBM Journal of Research and Development* 4 (1) (1960) 66–82.
- [55] V. Coffman, J. Kundu, W. K. Wootters, Distributed entanglement, *Physical Review A* 61 (2000) 052306. [arXiv:quant-ph/9907047](#).
- [56] E. T. Jaynes, *Probability Theory: The Logic of Science*, Cambridge University Press, Cambridge, UK, 2003.
- [57] B. S. Cirel’son, Quantum generalization of Bell’s inequalities, *Letters in Mathematical Physics* 4 (1) (1980) 93.
- [58] A. C.-C. Yao, Some complexity questions related to distributive computing (preliminary report), in: *STOC ’79: Proceedings of the eleventh annual ACM symposium on Theory of computing*, ACM, New York, NY, USA, 1979, pp. 209–213.
- [59] W. van Dam, Implausible consequences of superstrong nonlocality (2005). [arXiv:quant-ph/0501159](#).
- [60] D. M. Greenberger, M. A. Horne, A. Zeilinger, Going beyond Bell’s theorem, in: M. Kafatos (Ed.), *Bell’s Theorem, Quantum Theory, and Conceptions of the Universe*, Kluwer, Dordrecht, 1989, pp. 69–72. [arXiv:0712.0921v1\[quant-ph\]](#).
- [61] J.-W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter, A. Zeilinger, Experimental test of quantum nonlocality in three-photon Greenberger-Horne-Zeilinger entanglement, *Nature* 403 (2000) 515 – 519.
- [62] D. Bouwmeester, J.-W. Pan, M. Daniell, H. Weinfurter, A. Zeilinger, Observation of three-photon Greenberger-Horne-Zeilinger entanglement, *Physical Review Letters* 82 (1999) 1345. [arXiv:quant-ph/9810035](#).