

# Harnack Inequalities and Applications for SDEs with Non-Additive Noise and Neumann Semigroup on Non-Convex Manifolds \*

Feng-Yu Wang

School of Mathematical Sci. and Lab. Math. Com. Sys., Beijing Normal University, Beijing 100875, China

and

Department of Mathematics, Swansea University, Singleton Park, SA2 8PP, UK

Email: wangfy@bnu.edu.cn; F.Y.Wang@swansea.ac.uk

January 26, 2023

## Abstract

By constructing a coupling with unbounded time-dependent drift, dimension-free Harnack inequalities are established for the semigroup associated to SDEs with non-additive noise. These inequalities are applied to the study of heat kernel upper bound and contractivity properties of the semigroup. The main results are extended to non-constant diffusions on manifolds with (non-convex) boundary where the dimension-free Harnack inequality has been unknown for a long time.

AMS subject Classification: 60H10, 58G32.

Keywords: Stochastic differential equation, Harnack inequality, Neumann semigroup, Riemannian manifold with boundary.

## 1 Introduction

Consider the following SDE on  $\mathbb{R}^d$ :

---

\*Supported in part by WIMCS, NNSFC(10721091) and the 973-Project.

$$(1.1) \quad dX_t = \sigma(t, X_t)dB_t + b(t, X_t)dt,$$

where  $B_t$  is the  $d$ -dimensional Brownian motion on a complete filtered probability space  $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ , and

$$\sigma : [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d, \quad b : [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d$$

are progressively measurable and continuous in the second variable. Throughout the paper we assume that for any  $X_0 \in \mathbb{R}^d$  the equation (1.1) has a unique strong solution which is non-explosive.

To derive explicit Harnack type inequalities, we shall make use of the following standard dissipative type assumptions. Let  $\|\cdot\|_{HS}$  denote the Hilbert-Schmidt norm.

(A1) There exists an increasing function  $K : [0, \infty) \rightarrow \mathbb{R}$  such that

$$\|\sigma(t, x) - \sigma(t, y)\|_{HS}^2 + 2\langle b(t, x) - b(t, y), x - y \rangle \leq K_t |x - y|^2, \quad x, y \in \mathbb{R}^d, t \geq 0.$$

(A2) There exists a decreasing function  $\lambda : [0, \infty) \rightarrow (0, \infty)$  such that  $\sigma(t, x)\sigma(t, x)^* \geq \lambda_t^2 I$ ,  $x \in \mathbb{R}^d, t \geq 0$ .

(A3) There exists an increasing function  $\delta : [0, \infty) \rightarrow (0, \infty)$  such that

$$|(\sigma(t, x) - \sigma(t, y))(x - y)| \leq \delta_t |x - y|, \quad x, y \in \mathbb{R}^d, t \geq 0.$$

(A4) For  $n \geq 1$  there exists a constant  $c_n > 0$  such that

$$\|\sigma(t, x) - \sigma(t, y)\|_{HS} + |b(t, x) - b(t, y)| \leq c_n |x - y|, \quad |x|, |y|, t \leq n.$$

It is well known that (A1) ensures the uniqueness of the solution to (1.1) while (A4) implies the existence and the uniqueness of the solution (see e.g. [11] and references within for weaker conditions). When  $b$  and  $\sigma$  depend only on the variable  $x \in \mathbb{R}^d$ , the continuity will be enough to imply the existence of weak solution (see [13, Theorem 2.3]), so that by the pathwise uniqueness the Yamada-Watanabe principle [27] ensures the existence and uniqueness of the strong solution.

Let  $X_t^x$  be the solution to (1.1) for  $X_0 = x \in \mathbb{R}^d$ . We aim to establish the Harnack inequality for the operator  $P_t$ :

$$P_t f(x) := \mathbb{E}f(X_t^x), \quad t \geq 0, x \in \mathbb{R}^d, f \in \mathcal{B}_b^+(\mathbb{R}^d),$$

where  $\mathcal{B}_b^+(\mathbb{R}^d)$  is the class of all bounded non-negative measurable functions on  $\mathbb{R}^d$ .

Note that if  $\sigma(t, x)$  and  $b(t, x)$  are deterministic and independent of  $t$ , then the solution is a time-homogeneous Markov process generated by

$$L := \frac{1}{2} \sum_{i,j=1}^d a_{ij} \partial_i \partial_j + \sum_{i=1}^d b_i \partial_i,$$

where  $a := \sigma \sigma^*$ . If further more  $\sigma$  and  $b$  are smooth, we may consider the Bakry-Emery curvature condition [5]:

$$(1.2) \quad \Gamma_2(f, f) \geq -K\Gamma(f, f), \quad f \in C^\infty(\mathbb{R}^d)$$

for some constant  $K \in \mathbb{R}$ , where

$$\begin{aligned} \Gamma(f, g) &:= \frac{1}{2} \sum_{i,j=1}^d a_{ij} (\partial_i f)(\partial_j g), \quad f, g \in C^1(\mathbb{R}^d), \\ \Gamma_2(f, f) &:= \frac{1}{2} L\Gamma(f, f) - \Gamma(f, Lf), \quad f \in C^\infty(\mathbb{R}^d). \end{aligned}$$

According to [21, Lemma 2.2] and [22, Theorem 1.2], the curvature condition (4.1) is equivalent to the dimension-free Harnack inequality

$$(1.3) \quad (P_t f(x))^p \leq (P_t f^p(y)) \exp \left[ \frac{p\rho_a(x, y)^2}{2(p-1)(1 - e^{-2Kt})} \right], \quad f \in \mathcal{B}_b^+(\mathbb{R}^d), x, y \in \mathbb{R}^d$$

for all  $t > 1$  and  $p > 1$ . This type inequality has been extended and applied to the study of heat kernel (or transition probability) and contractivity properties for diffusion semigroups for some finite- and infinite-dimensional models, see e.g. [1, 17, 4] for diffusions on manifolds with possibly unbounded below curvature, see [23, 15] for stochastic generalized porous media and fast diffusion equations [2, 3, 8, 16, 14, 18, 10, 28] for the study of some other SPDEs, with additive noise (i.e. constant diffusion coefficients).

Let us come back to the equation (1.1). When the dimension  $d$  is large and  $\sigma$  is non-constant, it is normally very hard to verify the curvature condition (4.1). Indeed, this condition depends on second order derivatives of  $a^{-1}$ , the inverse matrix of  $a$ . This is the main reason why existed results in this direction only focus on the additive-noise case, except the recent paper [19] where the log-Harnack inequality, a weaker version of (1.3), was established for a class of non-additive noise SDEs by using gradient estimates.

The main purpose of the paper is to develop a general coupling argument to derive Harnack inequalities for SDEs with non-additive noise, and for the Neumann semigroup on manifolds with non-convex boundary. In both cases the validity of dimension-free Harnack inequality of type (1.3) has been open for a long time.

For the additive noise case, i.e.  $\sigma$  is independent of the second component  $x$ , the dimension-free Harnack inequality can be derived by using the coupling argument developed in [4]. This argument has been also applied to the study of SPDEs in the above

mentioned references. To see the difficulty in the study for non-constant diffusions, let us briefly recall the main idea of the coupling method in the study of the Harnack inequality.

For simplicity, let  $\sigma$  and  $b$  be independent of the second variable. For  $x \neq y$  and  $T > 0$ , let  $X_t$  solve (1.1) with  $X_0 = x$  and  $Y_t$  solve

$$dY_t = \sigma(t)dB_t + b(t)dt + \frac{|x-y|(X_t - Y_t)}{T|X_t - Y_t|}dt, \quad Y_0 = y.$$

Then  $Y_t$  is well defined up to the coupling time

$$\tau := \inf\{t \geq 0 : X_t = Y_t\}.$$

Let  $X_t = Y_t$  for  $t \geq \tau$ . We have

$$d|X_t - Y_t| = -\frac{|x-y|}{T}dt, \quad t \leq \tau.$$

This implies  $\tau = T$  and hence,  $X_T = Y_T$ . On the other hand, by the Girsanov theorem we have

$$P_T f(y) = \mathbb{E}[Rf(Y_T)]$$

for

$$R := \exp \left[ -\frac{|x-y|}{T} \int_0^T \frac{\langle \sigma(t)^{-1}(X_t - Y_t), dB_t \rangle}{|X_t - Y_t|} - \frac{|x-y|^2}{2T^2} \int_0^T \frac{|\sigma(t)^{-1}(X_t - Y_t)|^2}{|X_t - Y_t|^2} dt \right].$$

Since  $X_T = Y_T$ , it follows that

$$(P_T f(y))^p = (\mathbb{E}[Rf(X_T)])^p \leq (P_T f^p(x))(\mathbb{E}R^{p/(p-1)})^{p-1}.$$

Since by (A1) and (A2) it is easy to estimate the moments of  $R$ , the desired Harnack inequality follows immediately.

In generally, when  $\sigma(t, x)$  depends on  $x$ , the process  $X_t - Y_t$  contains a non-trivial martingale term, which can not be dominated by any bounded drift. So, in this case, any additional bounded drift put in the equation for  $Y_t$  is not enough to make the coupling successful before a fixed time  $T$ . This is the main difficulty to establish the Harnack inequality for diffusion semigroups with non-constant diffusion coefficient.

In this paper, under assumptions (A1) and (A2), we are able to constructed a coupling with a drift which is infinite at the fixed time  $T$ , such that the coupling is successful before  $T$ . In this case the corresponding exponential martingale has finite entropy such that the log-Harnack inequality holds; if further more (A3) holds then the exponential martingale is  $L^p$ -integrable for some  $p > 1$  such that the Harnack inequality with power hold. More precisely, we have the following result.

**Theorem 1.1.** *Let  $\sigma(t, x)$  and  $b(t, x)$  either be deterministic and independent of  $t$ , or satisfy (A4).*

(1) If (A1) and (A2) hold then

$$(1.4) \quad P_T \log f(y) \leq \log P_T f(y) + \frac{K_T |x - y|^2}{2\lambda_T^2(1 - e^{-K_T T})}, \quad f \geq 1, x, y \in \mathbb{R}^d, T > 0.$$

(2) If (A1), (A2) and (A3) hold, then for  $p > (1 + \frac{\delta_T}{\lambda_T})^2$  and  $\delta_{p,T} := \max\{\delta_T, \frac{\lambda_T}{2}(\sqrt{p} - 1)\}$ ,

$$(1.5) \quad (P_T f(y))^p \leq (P_T f^p(x)) \exp \left[ \frac{K_T \sqrt{p} (\sqrt{p} - 1) |x - y|^2}{4\delta_{p,T}[(\sqrt{p} - 1)\lambda_T - \delta_{p,T}](1 - e^{-K_T T})} \right]$$

holds for all  $T > 0, x, y \in \mathbb{R}^d$  and  $f \in \mathcal{B}_b^+(\mathbb{R}^d)$ .

As standard applications of the dimension-free Harnack inequality, we present below two consequences on the heat kernel inequalities and contractivity properties of the semigroup.

Let  $p_t(x, y)$  be the transition density of  $P_t$  w.r.t. a Radon measure  $\mu$ . Then according to [25, Proposition 2.4], the above log-Harnack inequality (1.4) and Harnack inequality (1.5) are equivalent to the following heat kernel inequalities respectively:

$$(1.6) \quad \int_{\mathbb{R}^d} p_T(x, z) \log \frac{p_T(x, z)}{p_T(y, z)} \mu(dz) \leq \frac{K_T |x - y|^2}{4\lambda_T^2(1 - e^{-K_T T})}, \quad x, y \in \mathbb{R}^d, T > 0$$

and

$$(1.7) \quad \int_{\mathbb{R}^d} p_T(x, z) \left( \frac{p_T(x, z)}{p_T(y, z)} \right)^{1/(p-1)} \mu(dz) \leq \exp \left[ \frac{K_T \sqrt{p} |x - y|^2}{4\delta_{p,T}(\sqrt{p} + 1)[(\sqrt{p} - 1)\lambda_T - \delta_{p,T}](1 - e^{-K_T T})} \right], \quad x, y \in \mathbb{R}^d, T > 0.$$

So, the following is a direct consequence of Theorem 1.1.

**Corollary 1.2.** *Let  $\sigma(t, x)$  and  $b(t, x)$  either be deterministic and independent of  $t$ , or satisfy (A4). Let  $P_t$  has a strictly positive density w.r.t. a Radon measure  $\mu$ . Then (A1) and (A2) imply (1.6), while (A1)-(A3) imply (1.7).*

**Corollary 1.3.** *Let  $\sigma(t, x)$  and  $b(t, x)$  be deterministic and independent of  $t$ , such that (A1)-(A3) hold for constant  $K, \lambda$  and  $\delta$ . Let  $P_t$  have an invariant probability measure  $\mu$ .*

- (1) *If there exists  $r > K^+/\lambda^2$  such that  $\mu(e^{r|\cdot|^2}) < \infty$ , then  $P_t$  is hypercontractive, i.e.  $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} = 1$  holds for some  $t > 0$ .*
- (2) *If  $\mu(e^{r|\cdot|^2}) < \infty$  holds for all  $r > 0$ , then  $P_t$  is supercontractive, i.e.  $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} < \infty$  holds for all  $t > 0$ .*
- (3) *If  $P_t e^{r|\cdot|^2}$  is bounded for any  $t, r > 0$ , then  $P_t$  is ultracontractive, i.e.  $\|P_t\|_{L^2(\mu) \rightarrow L^\infty(\mu)} < \infty$  for any  $t > 0$ .*

**Remark 1.1.** In the situation of Corollary 1.3 with  $\mu$  a probability measure, when  $P_t$  is symmetric w.r.t.  $\mu$ , the hypercontractivity is equivalent to the validity of the log-Sobolev inequality

$$\mu(f^2 \log f^2) \leq C \mu(\Gamma(f, f)), \quad f \in C_b^\infty(\mathbb{R}^d), \mu(f^2) = 1$$

for some constant  $C > 0$ . Moreover, by (A2) and (A3) there exists a constant  $C > 0$  such that

$$C^{-1} |\nabla f|^2 \leq \Gamma(f, f) \leq C |\nabla f|^2$$

holds for all  $f \in C^\infty(\mathbb{R}^d)$ . So, by the concentration of measure for the log-Sobolev inequality, the hypercontractivity of  $P_t$  implies that  $\mu(e^{r|\cdot|^2}) < \infty$  for some  $r > 0$ , while the supercontractivity of  $P_t$  implies that  $\mu(e^{r|\cdot|^2}) < \infty$ . Combining this with Corollary 1.3, we have the following assertions.

- (i) Let  $K \leq 0$ . Then  $P_t$  is hypercontractive if and only if  $\mu(e^{r|\cdot|^2}) < \infty$  holds for some  $r > 0$ .
- (ii)  $P_t$  is supercontractive if and only if  $\mu(e^{r|\cdot|^2}) < \infty$  holds for all  $r > 0$ .
- (iii)  $P_t$  is ultracontractive if and only if  $P_t e^{r|\cdot|^2}$  is bounded for any  $t, r > 0$ .

Therefore, when  $\sigma$  is bounded, Corollary 1.3 (2)-(3) are sharp for the supercontractivity and ultracontractivity of  $P_t$ . Moreover, as shown in [7] that when  $\sigma$  is constant, the sufficient condition  $\mu(e^{r|\cdot|^2}) < \infty$  for some  $r > K^+/\lambda^2$  is optimal for the hypercontractivity of  $P_t$ . So, Corollary 1.3(1) also provides a sharp sufficient condition for the hypercontractivity of  $P_t$ .

We will prove Theorem 1.1 and Corollary 1.3 in the next section. In section 3 we extend these results to SDEs on Riemannian manifolds without boundary or a convex boundary. Combining the results derived in Section 3 with a conformal change of metric introduced in [23], we are able to establish Harnack inequalities in Section 4 for the Neumann semigroup on a class of non-convex manifolds.

## 2 Proofs of Theorem 1.1 and Corollary 1.3

Let  $x, y \in \mathbb{R}^d, T > 0$  and  $p > (1 + \delta_T/\lambda_T)^2$  be fixed such that  $x \neq y$ . We have

$$(2.1) \quad \theta_T := \frac{2\delta_T}{(\sqrt{p} - 1)\lambda_T} \in (0, 2).$$

For  $\theta \in (0, 2)$ , let

$$\xi_t = \frac{2 - \theta}{K_T}(1 - e^{K_T(t-T)}), \quad t \in [0, T].$$

Then  $\xi$  is smooth and strictly positive on  $[0, T)$  such that

$$(2.2) \quad 2 - K_T \xi_t + \xi'_t = \theta, \quad t \in [0, T].$$

Consider the coupling

$$(2.3) \quad \begin{aligned} dX_t &= \sigma(t, X_t)dB_t + b(t, X_t)dt, \quad X_0 = x, \\ dY_t &= \sigma(t, Y_t)dB_t + b(t, Y_t)dt + \frac{1}{\xi_t}\sigma(t, Y_t)\sigma(t, X_t)^{-1}(X_t - Y_t)dt, \quad Y_0 = y. \end{aligned}$$

Since the additional drift term  $\xi_t^{-1}\sigma(t, y)\sigma(t, x)^{-1}(x - y)$  is locally Lipschitzian in  $y$  if (A4) holds, and continuous in  $y$  when  $\sigma$  and  $b$  are deterministic and time independent, the coupling  $(X_t, Y_t)$  is well defined for  $t < T \wedge \zeta$ , where  $\zeta$  is the explosive time of  $Y_t$ ; namely,  $\zeta = \lim_{n \rightarrow \infty} \zeta_n$  for

$$\zeta_n := \inf\{t \in [0, T) : |Y_t| \geq n\}.$$

Here and in what follows we set  $\inf \emptyset = \infty$  by convention. Let

$$d\tilde{B}_t = dB_t + \frac{1}{\xi_t}\sigma(t, X_t)^{-1}(X_t - Y_t)dt, \quad t < T \wedge \zeta.$$

If  $\zeta = \infty$  and the exponential martingale

$$R_s := \exp \left[ - \int_0^s \xi_t^{-1} \langle \sigma(t, X_t)^{-1}(X_t - Y_t), dB_t \rangle - \frac{1}{2} \int_0^s \xi_t^{-2} |\sigma(t, X_t)^{-1}(X_t - Y_t)|^2 dt \right]$$

is uniformly integrable for  $s \in [0, T)$ , i.e.

$$\lim_{n \rightarrow \infty} \sup_{s \in [0, T)} \mathbb{E}[R_s 1_{\{R_s > n\}}] = 0,$$

then by the martingale convergence theorem,  $R_T := \lim_{t \uparrow T} R_t$  exists and  $\{R_t\}_{t \in [0, T]}$  is a martingale. In this case, by the Girsanov theorem  $\{\tilde{B}_t\}_{t \in [0, T)}$  is a  $d$ -dimensional Brownian motion under the probability  $R_T \mathbb{P}$ . Reformulate (2.3) as

$$(2.4) \quad \begin{aligned} dX_t &= \sigma(t, X_t)d\tilde{B}_t + b(t, X_t)dt - \frac{X_t - Y_t}{\xi_t}dt, \quad X_0 = x, \\ dY_t &= \sigma(t, Y_t)d\tilde{B}_t + b(t, Y_t)dt, \quad Y_0 = y. \end{aligned}$$

Since  $\int_0^T \xi_t^{-1} dt = \infty$ , we will see that the additional drift  $-\frac{X_t - Y_t}{\xi_t}dt$  is strong enough to make the coupling successful up to time  $T$ . So, we first study the uniform integrability of  $\{R_s\}_{s \in [0, T)}$ .

**Lemma 2.1.** Assume (A1) and (A2) and let  $\theta \in (0, 2)$ ,  $x, y \in \mathbb{R}^d$  and  $T > 0$  be fixed. Then  $\zeta = \infty$  and

$$\sup_{s \in [0, T)} \mathbb{E} R_s \log R_s \leq \frac{K_T |x - y|^2}{2\lambda_T^2 \theta (2 - \theta) (1 - e^{-K_T T})}.$$

In particular,  $\{R_s\}_{s \in [0, T)}$  is uniformly integrable so that  $R_T := \lim_{s \uparrow T} R_s$  exists and  $\{R_s\}_{s \in [0, T]}$  is a uniformly integrable martingale.

*Proof.* Let  $s \in [0, T)$  be fixed. For any  $n \geq 1$ , let

$$\tau_n = \inf\{t \in [0, T) : |X_t| + |Y_t| \geq n\}.$$

By (2.4), (A1) and the Itô formula,

$$d\|X_t - Y_t\|^2 \leq 2\langle(\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t\rangle + K_T |X_t - Y_t|^2 dt - \frac{2}{\xi_t} |X_t - Y_t|^2 dt$$

holds for  $t \leq s \wedge \tau_n$ . Combining this with (2.2) we obtain

$$\begin{aligned} (2.5) \quad & d \frac{|X_t - Y_t|^2}{\xi_t} \\ & \leq \frac{2}{\xi_t} \langle(\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t\rangle - \frac{|X_t - Y_t|^2}{\xi_t^2} (2 - K_T \xi_t + \xi_t') dt \\ & = \frac{2}{\xi_t} \langle(\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t\rangle - \frac{\theta}{\xi_t^2} |X_t - Y_t|^2 dt, \quad t \leq s \wedge \tau_n. \end{aligned}$$

Since due to the Girsanov theorem  $\{\tilde{B}_t\}_{t \leq \tau_n \wedge s}$  is the  $d$ -dimensional Brownian motion under the probability measure  $R_{s \wedge \tau_n} \mathbb{P}$ , letting  $\mathbb{E}_{s, n}$  denote the corresponding expectation we obtain

$$(2.6) \quad \mathbb{E}_{s, n} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \leq \frac{|x - y|^2}{\theta \xi_0}, \quad s \in [0, T), n \geq 1.$$

Since  $X_t$  is non-explosive, this in particular implies that  $Y_t$  is non-explosive before time  $T$ , i.e.  $\zeta = \infty$ . By (A2) and the definitions of  $R_t$  and  $\tilde{B}_t$ , we have

$$\begin{aligned} \log R_r &= - \int_0^r \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1} (X_t - Y_t), d\tilde{B}_t \rangle + \frac{1}{2} \int_0^r \frac{|\sigma(t, X_t)^{-1} (X_t - Y_t)|^2}{\xi_t^2} dt \\ &\leq - \int_0^r \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1} (X_t - Y_t), d\tilde{B}_t \rangle + \frac{1}{2\lambda_T^2} \int_0^r \frac{|X_t - Y_t|^2}{\xi_t^2} dt, \quad r \leq s \wedge \tau_n. \end{aligned}$$

Since  $\{\tilde{B}_t\}$  is the  $d$ -dimensional Brownian motion under  $R_{s \wedge \tau_n} \mathbb{P}$  up to  $s \wedge \tau_n$ , combining this with (2.6) we obtain



$$\mathbb{E}R_{s \wedge \tau_n} \log R_{s \wedge \tau_n} = \mathbb{E}_{s,n} \log R_{s \wedge \tau_n} \leq \frac{|x-y|^2}{2\lambda_T^2 \theta \xi_0}, \quad s \in [0, T], n \geq 1.$$

By the Fatou lemma and noting that  $\lim_{n \rightarrow \infty} \tau_n \geq T$  since  $X_t$  and  $Y_t$  are non-explosive before  $T$ , we arrive at

$$\mathbb{E}R_s \log R_s \leq \frac{|x-y|^2}{2\lambda_T^2 \theta \xi_0} = \frac{K_T |x-y|^2}{2\lambda_T^2 \theta (2-\theta)(1-e^{-K_T T})}, \quad s \in [0, T].$$

This completes the proof.  $\square$

Lemma 2.1 ensures that the coupling  $(X_t, Y_t)$  is well-constructed for  $t \in [0, T]$ . Since  $\int_0^T \xi_t^{-1} dt = \infty$ , we shall see that the coupling is successful up to time  $T$ , so that  $X_T = Y_T$  holds (see the proof of Theorem 1.1 below). This will provide the desired Harnack inequality for  $P_t$  as explained in Section 1 as soon as  $R_T$  has finite  $p/(p-1)$ -moment. The next lemma provides an explicit upper bound on the moments of  $R_T$ .

**Lemma 2.2.** *Assume (A1)-(A3). Let  $R_t$  and  $\xi_t$  be fixed for  $\theta = \theta_T$ . We have*

$$(2.7) \quad \sup_{s \in [0, T]} \mathbb{E} \left\{ R_s \exp \left[ \frac{\theta_T^2}{8\delta_T^2} \int_0^s \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right\} \leq \exp \left[ \frac{\theta_T K_T |x-y|^2}{4\delta_T^2 (2-\theta_T)(1-e^{-K_T T})} \right].$$

Consequently,

$$(2.8) \quad \sup_{s \in [0, T]} \mathbb{E} R_s^{1+r_T} \leq \exp \left[ \frac{\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x-y|^2}{8\delta_T^2 (2-\theta_T)(\delta_T + \theta_T \lambda_T)(1-e^{-K_T T})} \right]$$

holds for

$$r_T = \frac{\lambda_T^2 \theta_T^2}{4\delta_T^2 + 4\theta_T \lambda_T \delta_T}.$$

*Proof.* Let  $\theta = \theta_T$ . By (2.5), for any  $r > 0$  we have

$$\begin{aligned} & \mathbb{E}_{s,n} \exp \left[ r \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \\ & \leq \exp \left[ \frac{r|x-y|^2}{\theta_T \xi_0} \right] \cdot \mathbb{E}_{s,n} \exp \left[ \frac{2r}{\theta_T} \int_0^{s \wedge \tau_n} \frac{1}{\xi_t} \langle (\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t \rangle \right] \\ & \leq \exp \left[ \frac{r K_T |x-y|^2}{\theta_T (2-\theta_T)(1-e^{-K_T T})} \right] \left( \mathbb{E}_{s,n} \exp \left[ \frac{8r^2 \delta_T^2}{\theta_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right)^{1/2}, \end{aligned}$$

where the last step is due to (A3) and the fact that

$$\mathbb{E}e^{Mt} \leq (\mathbb{E}e^{2\langle M \rangle_t})^{1/2}$$

holds for a continuous exponential integrable martingale  $M_t$ . Taking  $r = \theta_T^2/(8\delta_T^2)$ , we arrive at

$$\mathbb{E}_{s,n} \exp \left[ \frac{\theta_T^2}{8\delta_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \leq \left[ \frac{\theta_T K_T |x - y|^2}{4\delta_T^2(2 - \theta_T)(1 - e^{-K_T T})} \right], \quad n \geq 1.$$

This implies (2.7) by letting  $n \rightarrow \infty$ .

Next, let

$$M_t = - \int_0^{s \wedge \tau_n} \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1} (X_t - Y_t), d\tilde{B}_t \rangle, \quad t \leq s \wedge \tau_n,$$

which is a martingale w.r.t.  $R_{s \wedge \tau_n} \mathbb{P}$ . By the definition of  $R_t$  we have

$$\begin{aligned} \mathbb{E}R_{s \wedge \tau_n}^{1+r_T} &= \mathbb{E}_{s,n} R_{s \wedge \tau_n}^{r_T} = \mathbb{E}_{s,n} \exp \left[ r_T M_s + \frac{r_T}{2} \langle M \rangle_s \right] \\ &= \mathbb{E}_{s,n} \exp \left[ r_T M_s - \frac{r_T^2 q}{2} \langle M \rangle_s + \frac{r_T}{2} (qr_T + 1) \langle M \rangle_s \right] \\ &\leq \left( \mathbb{E}_{s,n} \exp \left[ r_T q M_s - \frac{r_T^2 q^2}{2} \langle M \rangle_s \right] \right)^{1/q} \left( \mathbb{E}_{s,n} \exp \left[ \frac{r_T q (r_T q + 1)}{2(q-1)} \langle M \rangle_s \right] \right)^{(q-1)/q} \\ &= \left( \mathbb{E}_{s,n} \exp \left[ \frac{r_T q (r_T q + 1)}{2(q-1)} \langle M \rangle_s \right] \right)^{(q-1)/q}, \quad q > 1. \end{aligned}$$

Combining this with (A2) we obtain

$$(2.9) \quad \mathbb{E}R_{s \wedge \tau_n}^{1+r_T} \leq \left( \mathbb{E}_{s,n} \exp \left[ \frac{qr_T(qr_T + 1)}{2(q-1)\lambda_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right)^{(q-1)/q}, \quad q > 1.$$

Taking

$$(2.10) \quad q = 1 + \sqrt{1 + r_T^{-1}}$$

which minimizes  $q(qr_T + 1)/(q - 1)$  such that, due to the definition of  $r_T$ ,

$$\begin{aligned} (2.11) \quad \frac{qr_T(qr_T + 1)}{2\lambda_T^2(q-1)} &= \frac{r_T + \sqrt{r_T(r_T + 1)}}{2\lambda_T^2 \sqrt{1 + r_T^{-1}}} (r_T + 1 + \sqrt{r_T(r_T + 1)}) \\ &= \frac{(r_T + \sqrt{r_T^2 + r_T})^2}{2\lambda_T^2} = \frac{\theta_T^2}{8\delta_T^2}. \end{aligned}$$

Combining (2.9) with (2.7) and (2.11), and noting that due to (2.10) and the definition of  $r_T$  one has

$$\frac{q-1}{q} = \frac{\sqrt{1+r_T^{-1}}}{1+\sqrt{1+r_T^{-1}}} = \frac{2\delta_T + \theta_T \lambda_T}{2\delta_T + 2\theta_T \lambda_T},$$

we obtain

$$\mathbb{E}R_{s \wedge \tau_n}^{1+r_T} \leq \exp \left[ \frac{\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x-y|^2}{8\delta_T^2 (2-\theta_T)(\delta_T + \theta_T \lambda_T)(1-e^{-K_T T})} \right].$$

According to the Fatou lemma, the proof is then completed by letting  $n \rightarrow \infty$ .  $\square$

*Proof of Theorem 1.1.* (1) By Lemma 2.1,  $\{R_s\}_{s \in [0, T]}$  is an uniformly integrable martingale. Moreover, by the Girsanov theorem,  $\{\tilde{B}_t\}_{t \leq T}$  is a  $d$ -dimensional Brownian motion under the probability  $R_T \mathbb{P}$ . Thus,  $Y_t$  can be solved up to time  $T$ . Let

$$\tau = \inf\{t \in [0, T] : X_t = Y_t\}.$$

We claim that  $\tau \leq T$  so that  $X_T = Y_T$  a.s. Indeed, if for some  $\omega \in \Omega$  such that  $\tau(\omega) > T$ , by the continuity of the diffusions we have

$$R_T(\omega) \inf_{t \in [0, T]} |X_t - Y_t|^2(\omega) > 0.$$

Noting that  $\int_0^T \xi_t^{-2} dt = \infty$ , this implies that

$$R_T \int_0^T \frac{|X_t - Y_t|^2}{\xi_t^2} dt = \infty$$

holds on the set  $\{\tau > T\}$ . But according to Lemma 2.2 we have

$$\mathbb{E} \left\{ R_T \int_0^T \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right\} < \infty.$$

Therefore,  $\mathbb{P}(\tau > T) = 0$  and hence,  $X_T = Y_T$  a.s.

Now, combining Lemma 2.1 with  $X_T = Y_T$  and using the Young inequality, for  $f \geq 1$  we have

$$\begin{aligned} P_T \log f(y) &= \mathbb{E}[R_T \log f(Y_T)] = \mathbb{E}[R_T \log f(X_T)] \\ &\leq \mathbb{E}[R_T \log R_T] + \log[\mathbb{E}f(X_T)] \leq \log P_T f(x) + \frac{K_T |x-y|^2}{2\lambda_T^2 \theta (2-\theta)(1-e^{-K_T T})}. \end{aligned}$$

This completes the proof of (1) by taking  $\theta = 1$ .

(2) Let  $\theta = \theta_T$ . Since  $X_T = Y_T$  and  $\{\tilde{B}_t\}_{t \in [0, T]}$  is the  $d$ -dimensional Brownian motion under  $R_T \mathbb{P}$ , we have

$$(2.12) \quad (P_T f(y))^p = (\mathbb{E}[R_T f(Y_T)])^p = (\mathbb{E}[R_T f(X_T)])^p \leq (P_T f^p(x))(\mathbb{E} R_T^{p/(p-1)})^{p-1}.$$

Due to (2.1) we have

$$\frac{p}{p-1} = 1 + \frac{\lambda_T^2 \theta_T^2}{4\delta_T(\delta_T + \theta_T \lambda_T)}.$$

So, it follows from Lemma 2.2 and (2.1) that

$$\begin{aligned} (\mathbb{E} R_T^{p/(p-1)})^{p-1} &= (\mathbb{E} R_T^{1+r_T})^{p-1} \leq \exp \left[ \frac{(p-1)\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x-y|^2}{8\delta_T^2 (2-\theta_T)(\delta_T + \theta_T \lambda_T)(1-e^{-K_T T})} \right] \\ &= \exp \left[ \frac{K_T \sqrt{p} (\sqrt{p}-1) |x-y|^2}{4\delta_T [(\sqrt{p}-1)\lambda_T - \delta_T](1-e^{-K_T T})} \right]. \end{aligned}$$

Combining this with (2.12) we obtain

$$(2.13) \quad (P_T f(y))^p \leq (P_T f^p(x)) \exp \left[ \frac{K_T \sqrt{p} (\sqrt{p}-1) |x-y|^2}{4\delta_T [(\sqrt{p}-1)\lambda_T - \delta_T](1-e^{-K_T T})} \right].$$

Finally, since  $\delta_{p,t} \geq \delta_t$  so that (A3) holds for  $\delta_{p,t}$  in place of  $\delta_t$ , we conclude that (2.13) also holds with  $\delta_{p,T}$  in place of  $\delta_T$ .  $\square$

*Proof of Corollary 1.3.* Let  $f \in \mathcal{B}_b^+(\mathbb{R}^d)$  be such that  $\mu(f^p) \leq 1$ . Let  $p > (1 + \delta/\lambda)^2$ . By Theorem 1.1(2), we have

$$(P_t f(y))^p \exp \left[ - \frac{K \sqrt{p} (\sqrt{p}-1) |x-y|^2}{4\tilde{\delta}_p [(\sqrt{p}-1)\lambda - \tilde{\delta}_p](1-e^{-Kt})} \right] \leq P_t f^p(x), \quad x, y \in \mathbb{R}^d,$$

where  $\tilde{\delta}_p = \max\{\delta, \frac{\lambda}{2}(\sqrt{p}-1)\}$ . Integrating w.r.t.  $\mu(dx)$  and noting that  $\mu$  is  $P_t$ -invariant, we obtain

$$(2.14) \quad (P_t f(y))^p \int_{\mathbb{R}^d} \exp \left[ - \frac{K \sqrt{p} (\sqrt{p}-1) |x-y|^2}{4\tilde{\delta}_p [(\sqrt{p}-1)\lambda - \tilde{\delta}_p](1-e^{-Kt})} \right] \mu(dx) \leq 1.$$

Now, let  $B(0, 1) = \{x \in \mathbb{R}^d : |x| \leq 1\}$ . Since  $\mu$  is an invariant measure, it has a strictly positive density w.r.t. the Lebesgue measure (cf. [6]). In particular,  $\mu(B(0, 1)) > 0$ . Let  $p \geq (1 + 2\delta/\lambda)^2$ . We have  $\tilde{\delta}_p = (\sqrt{p}-1)\lambda/2$  and thus,

$$\frac{\sqrt{p}(\sqrt{p}-1)}{4\tilde{\delta}_p[(\sqrt{p}-1)\lambda-\tilde{\delta}_p]} = \frac{\sqrt{p}}{\lambda^2(\sqrt{p}-1)}.$$

Combining this with (2.14) and noting that

$$\begin{aligned} & \int_{\mathbb{R}^d} \exp \left[ -\frac{K\sqrt{p}(\sqrt{p}-1)|x-y|^2}{4\tilde{\delta}_p[(\sqrt{p}-1)\lambda-\tilde{\delta}_p](1-e^{-Kt})} \right] \mu(dx) \\ & \geq \mu(B(0,1)) \exp \left[ -\frac{K\sqrt{p}(\sqrt{p}-1)(1+|y|)^2}{4\tilde{\delta}_p[(\sqrt{p}-1)\lambda-\tilde{\delta}_p](1-e^{-Kt})} \right], \end{aligned}$$

we obtain

$$(2.15) \quad (P_t f(y))^p \leq C_1 \exp \left[ \frac{K\sqrt{p}(1+|y|)^2}{\lambda_T^2(\sqrt{p}-1)(1-e^{-Kt})} \right], \quad t > 0, y \in \mathbb{R}^d$$

for some constant  $C_1 > 0$  and all  $f \in \mathcal{B}_b^+(\mathbb{R}^d)$  with  $\mu(f^p) \leq 1$ . Since

$$\lim_{p \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{K\sqrt{p}}{\lambda^2(\sqrt{p}-1)(1-e^{-Kt})} = \frac{K^+}{\lambda^2},$$

for any  $r > K^+/\lambda^2$  there exist  $p > (1+2\delta/\lambda)^2$ ,  $\beta > 1$  and  $t_1 > 0$  such that

$$(P_{t_1} f(y))^{\beta p} \leq C_2 e^{r|y|^2}, \quad y \in \mathbb{R}^d, f \in \mathcal{B}_b^+(\mathbb{R}^d), \mu(f^p) \leq 1$$

holds for some constant  $C_2 > 0$ . Thus,  $\mu(e^{r|\cdot|^2}) < \infty$  implies that  $\|P_{t_1}\|_{L^p(\mu) \rightarrow L^{p\beta}(\mu)} < \infty$ . Since  $\|P_s\|_{L^q(\mu)} = 1$  holds for any  $q \in [1, \infty]$ , by the interpolation theorem and the semigroup property one may find  $t_2 > t_1$  such that

$$(2.16) \quad \|P_{t_2}\|_{L^2(\mu) \rightarrow L^4(\mu)} < \infty.$$

Moreover, by [12, Theorem 3.6(ii)], there exist some constants  $\eta, C_3 > 0$  such that

$$\|P_t - \mu\|_{L^2(\mu)} \leq C_3 e^{-\eta t}, \quad t \geq 0.$$

Combining this with (2.16) we conclude that  $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} \leq 1$  holds for sufficiently large  $t > 0$ , i.e. (2) holds.

Finally, (3) and (4) follow immediately from (2.15) and the interpolation theorem.  $\square$

### 3 Extension to manifolds with convex boundary

Let  $M$  be a  $d$ -dimensional complete, connected Riemannian manifold, possibly with a convex boundary  $\partial M$ . Let  $N$  be the inward unit normal vector field of  $\partial M$  when  $\partial M \neq \emptyset$ . Let  $P_t$  be the (Neumann) semigroup generated by

$$L := \psi^2(\Delta + Z)$$

on  $M$ , where  $\psi \in C^1(M)$  and  $Z$  is a  $C^1$  vector field on  $M$ . Assume that  $\psi$  is bounded and

$$(3.1) \quad \text{Ric} - \nabla Z \geq -K_0$$

holds for some constant  $K_0 \geq 0$ . Then the (reflecting) diffusion process generated by  $L$  is non-explosive.

To formulae  $P_t$  as the semigroup associated as SDE like (1.1), we set

$$(3.2) \quad \sigma = \sqrt{2}\psi, \quad b = \psi^2 Z.$$

Then  $P_t$  is the semigroup for the solution to the following Itô SDE (see e.g. [9] or [4] for the definition of  $d_I$  on Riemannian manifolds):

$$(3.3) \quad d_I X_t = \sigma(X_t) \Phi_t dB_t + b(X_t) dt + N(X_t) dl_t,$$

where  $B_t$  is the  $d$ -dimensional Brownian motion on a complete filtered probability space  $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ ,  $\Phi_t$  is the horizontal lift of  $X_t$  onto the frame bundle  $O(M)$ , and  $l_t$  is the local time of  $X_t$  on  $\partial M$ . When  $\partial M = \emptyset$ , we simply set  $l_t = 0$ .

To derive the Harnack inequality as in Section 2, we assume that

$$(3.4) \quad \lambda := \inf \sigma > 0, \quad \delta := \sup \sigma - \inf \sigma < \infty.$$

Now, let  $x, y \in M$  and  $T > 0$  be fixed. Let  $\rho$  be the Riemannian distance on  $M$ , i.e.  $\rho(x, y)$  is the length of the minimal geodesic on  $M$  linking  $x$  and  $y$ , which exists if  $\partial M$  is either convex or empty.

Let  $X_t$  solve (3.3) with  $X_0 = x$ . Next any strictly positive function  $\xi \in C([0, T])$ , let  $Y_t$  solve

$$d_I Y_t = \sigma(Y_t) P_{X_t, Y_t} \Phi_t dB_t + b(X_t) dt - \frac{\sigma(Y_t) \rho(X_t, Y_t)}{\sigma(X_t) \xi_t} \nabla \rho(X_t, \cdot)(Y_t) dt + N(Y_t) d\tilde{l}_t$$

for  $Y_0 = y$ , where  $\tilde{l}_t$  is the local time of  $Y_t$  on  $\partial M$ , and  $P_{X_t, Y_t} : T_{X_t}M \rightarrow T_{Y_t}M$  is the parallel displacement along the minimal geodesic from  $X_t$  to  $Y_t$ , which exists since  $\partial M$  is convex or empty. As explained in [4, Section 3], we may and do assume that the cut-locus of  $M$  is empty such that the parallel displacement is smooth. Let

$$d\tilde{B}_t = dB_t + \frac{\rho(X_t, Y_t)}{\xi_t \sigma(X_t)} \Phi_t^{-1} \nabla \rho(\cdot, Y_t)(X_t) dt, \quad t < T.$$

By the Girsanov theorem, for any  $s \in (0, T)$  the process  $\{\tilde{B}_t\}_{t \in [0, s]}$  is the  $d$ -dimensional Brownian motion under the weighted probability measure  $R_s \mathbb{P}$ , where

$$(3.5) \quad R_s := \exp \left[ - \int_0^s \frac{\rho(X_t, Y_t)}{\xi_t \sigma(X_t)} \langle \nabla \rho(\cdot, Y_t)(X_t), \phi_t dB_t \rangle - \frac{1}{2} \int_0^s \frac{\rho(X_t, Y_t)^2}{\xi_t^2 \sigma(X_t)^2} dt \right].$$

Thus, by (3.2) we have

$$\begin{aligned} d_I X_t &= \sqrt{2} \psi(X_t) \Phi_t d\tilde{B}_t + (\psi^2 Z)(X_t) dt - \frac{\rho(X_t, Y_t)}{\xi_t} \nabla \rho(\cdot, Y_t)(X_t) dt + N(X_t) dl_t, \\ d_I Y_t &= \sqrt{2} \psi(Y_t) \Phi_t d\tilde{B}_t + (\psi^2 Z)(Y_t) dt + N(Y_t) d\tilde{l}_t. \end{aligned}$$

Let  $\xi \in C^1([0, T])$  be strictly positive and take

$$\beta_t = - \frac{\rho(X_t, Y_t)}{\sqrt{2} \xi_t \psi(X_t)} \Phi_t^{-1} \nabla(\cdot, Y_t)(X_t).$$

Repeating the proof of (4.10) in [26] we obtain

$$d\rho(X_t, Y_t) \leq (\sigma(X_t) - \sigma(Y_t)) \langle \nabla \rho(\cdot, Y_t)(X_t), \Phi_t d\tilde{B}_t \rangle + K_1 \rho(X_t, Y_t) dt - \frac{\rho(X_t, Y_t)}{\xi_t} dt, \quad t < T,$$

where

$$K_1 = K_0 \|\psi\|_\infty^2 + 2\|Z\|_\infty \|\nabla \psi\|_\infty \|\psi\|_\infty.$$

This implies that

$$d \frac{\rho(X_t, Y_t)^2}{\xi_t} \leq \frac{2}{\xi_t} \rho(X_t, Y_t) (\sigma(X_t) - \sigma(Y_t)) \langle \nabla \rho(\cdot, Y_t)(X_t), \Phi_t d\tilde{B}_t \rangle - \frac{\rho(X_t, Y_t)^2}{\xi_t^2} (2 - K\xi_t + \xi_t') dt$$

holds for

$$(3.6) \quad \begin{aligned} K &:= 2K_1 + \|\nabla \sigma\|_\infty^2 \\ &= 2K_0 \|\psi\|_\infty^2 + 4\|Z\|_\infty \|\nabla \psi\|_\infty \|\psi\|_\infty + 2\|\nabla \psi\|_\infty^2. \end{aligned}$$

In particular, letting

$$\xi_t = \frac{2-\theta}{K}(1 - e^{K(t-T)}), \quad t \in [0, T], \theta \in (0, 2),$$

we have

$$2 - K\xi_t + \xi'_t = \theta.$$

Therefore, the following result follows immediately by repeating calculations in Section 2.

**Theorem 3.1.** *Assume that  $\partial M$  is either empty or convex. Let (4.1) and  $Z, \phi$  be bounded such that*

$$K := 2K_0\|\psi\|_\infty^2 + 4\|Z\|_\infty\|\nabla\psi\|_\infty\|\psi\|_\infty + 2\|\nabla\psi\|_\infty^2 < \infty.$$

*Then all assertions in Theorem 1.1 and Corollaries 1.2, 1.3 hold for  $P_t$  the (Neumann) semigroup generated on  $M$  by  $L = \psi^2(\Delta + Z)$  on  $M$  for constant functions  $K, \delta := \sqrt{2}(\sup \psi - \inf \psi)$  and  $\lambda := \sqrt{2} \inf |\psi|$ .*

## 4 Neumann semigroup on non-convex manifolds

Following the line of [23], we are able to make the boundary from non-convex to convex by using a conformal change of metric. This will enable us to extend our results to the Neumann semigroup on a class of non-convex manifolds.

Let  $\partial M \neq \emptyset$  with  $N$  the inward normal unit vector field. Then the second fundamental form of  $\partial M$  is a two-tensor on the tangent space of  $\partial M$  defined by

$$\mathbb{I}(X, Y) := -\langle \nabla_X N, Y \rangle, \quad X, Y \in T\partial M.$$

Assume that there exists  $\kappa > 0$  and  $K_0 \in \mathbb{R}$  such that

$$(4.1) \quad \text{Ric} - \nabla Z \geq -K_0, \quad \mathbb{I} \geq -\kappa$$

holds for  $M$  and a  $C^1$  vector field  $Z$ . We shall consider the Harnack inequality for the Neumann semigroup  $P_t$  generated by

$$L = \Delta + Z.$$

To make the boundary convex, let  $f \in C_b^\infty(M)$  such that  $f \geq 1$  and  $N \log f|_{\partial M} \geq \kappa$ . By [23, Lemma 2.1],  $\partial M$  is convex under the metric

$$\langle \cdot, \cdot \rangle' = f^{-2} \langle \cdot, \cdot \rangle.$$

Let  $\Delta'$  and  $\nabla'$  be the Laplacian and gradient induced by the new metric. We have (see (2.2) in [20])



$$L = f^{-2}(\Delta' + Z'), \quad Z' = f^2 Z + \frac{d-2}{2} \nabla f^2.$$

So, Theorem 3.1 applies for the convex manifold  $(M, \langle \cdot, \cdot \rangle')$ . Let  $\text{Ric}'$  be the Ricci curvature induced by the metric  $\langle \cdot, \cdot \rangle'$ . We have (see the proof of [26, Theorem 5.1])

$$\text{Ric}' - \nabla' Z' \geq -K_f \langle \cdot, \cdot \rangle'$$

for

$$(4.2) \quad K_f = \sup \left\{ K f^2 - d \Delta f + (d-3) |\nabla f|^2 + 3 |Z| f |\nabla f| \right\}.$$

Thus, Theorem 3.1 holds for  $\psi = f^{-1}$  and

$$\begin{aligned} K &= 2K_f^+ \|f^{-1}\|_\infty + 4\|Z'\|'_\infty \|\nabla' f^{-1}\|'_\infty \|f^{-1}\|_\infty + 2\|\nabla' f^{-1}\|'^2_\infty \\ &\leq 2K_f^+ + 4\|fZ + (d-2)\nabla\|_\infty \|\nabla f\|_\infty + 2\|\nabla f\|_\infty^2, \end{aligned}$$

where  $\|\cdot\|'$  is the norm induced by  $\langle \cdot, \cdot \rangle'$  and we have used that  $f \geq 1$ .

**Theorem 4.1.** *Let (4.1) hold for some  $\kappa > 0$  and  $K_0 \in \mathbb{R}$ , and let  $P_t$  be the Neumann semigroup generated by  $L = \Delta + Z$  on  $M$ . Then for any  $f \in C_b^\infty(M)$  such that  $\inf f = 1$ ,  $N \log f|_{\partial M} \geq \kappa$  and  $K < \infty$ , where  $K$  is fixed by (4.2) and (4), all assertions in Theorem 1.1 and Corollaries 1.2 and 1.3 hold for constant functions  $K, \delta := \sqrt{2}(\sup f^{-1} - \inf f^{-1})$ , and  $\lambda := \sqrt{2} \inf f^{-1}$ .*

**Remark 4.1** A simple choice of  $f$  in Theorem 4.1 is  $f = \phi \circ \rho_\partial$ , where  $\rho_p$  is the Riemannian distance to the boundary which is smooth on  $\{\rho_\partial \leq r_0\}$  for some  $r_0 > 0$  provided the injectivity radius of the boundary is positive, and  $f \in C_b^\infty([0, \infty))$  is such that  $f(0) = 1, f'(0) = \kappa$  and  $f(r) = f(r_0)$  for  $r \geq r_0$ . In general,  $f$  is taken account to  $r_T$  and bounds of the second fundamental form and sectional curvatures, see e.g. [23, 26] for details. With specific choices of  $f$  Theorem 4.1 provides explicit Harnack type inequalities, heat kernels estimates and criteria on contractivity properties for the Neumann semigroup on manifolds with non-convex boundary.

## References

- [1] S. Aida, *Uniform positivity improving property, Sobolev inequalities, and spectral gaps*, J. Funct. Anal. 158(1998), 152–185.
- [2] S. Aida and H. Kawabi, *Short time asymptotics of a certain infinite dimensional diffusion process*, Stochastic analysis and related topics, VII (Kusadasi, 1998), Progr. Probab., vol. 48, Birkhäuser Boston, Boston, 2001, pp. 77–124.

- [3] S. Aida and T. Zhang, *On the small time asymptotics of diffusion processes on path groups*, Potential Anal. 16(2002), 67–78.
- [4] M. Arnaudon, A. Thalmaier, F.-Y. Wang, *Harnack inequality and heat kernel estimates on manifolds with curvature unbounded below*, Bull. Sci. Math. 130(2006), 223–233.
- [5] D. Bakry, M. Emery, *Hypercontractivité de semi-groupes de diffusion*, C. R. Acad. Sci. Paris. Sér. I Math. 299(1984), 775–778.
- [6] V. I. Bogachev, N. V. Krylov, M. Röckner, *On regularity of transition probabilities and invariant measures of singular diffusions under minimal conditions*, Comm. Partial Differential Equations 26 (2001), 2037–2080.
- [7] X. Chen, F.-Y. Wang, *Optimal integrability condition for the log-Sobolev inequality*, Quart. J. Math. (Oxford) 58(2007), 17–22.
- [8] G. Da Prato, M. Röckner, F.-Y. Wang, *Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups*, J. Funct. Anal. 257(2009), 992–1017.
- [9] M. Emery, *Stochastic calculus in manifolds*, Springer-Verlag, Berlin, 1989.
- [10] A. Es-Sarhir, M.-K. von Renesse, M. Scheutzew, *Harnack inequality for functional SDEs with bounded memory*, arXiv:0910.4536.
- [11] S. Fang, T.-S. Zhang, *A study of a class of stochastic differential equations with non-Lipschitzian coefficients*, Probab. Theory Related Fields 132 (2005), 356–390.
- [12] M. Hino, *Exponential decay of positivity preserving semigroups on  $L^p$* , Osaka J. Math. 37 (2000), 603–624.
- [13] N. Ikeda, S. Watanabe, *Stochastic Differential Equations and Diffusion Processes (Second Edition)*, North-Holland, Amsterdam, 1989.
- [14] W. Liu, *Fine properties of stochastic evolution equations and their applications*, Doctor-Thesis, Bielefeld University, 2009.
- [15] W. Liu and F.-Y. Wang, *Harnack inequality and strong Feller property for stochastic fast-diffusion equations*, J. Math. Anal. Appl. 342(2008), 651–662.
- [16] S.-X. Ouyang, *Harnack inequalities and applications for stochastic equations*, Ph.D. thesis, Bielefeld University, 2009.
- [17] M. Röckner and F.-Y. Wang, *Supercontractivity and ultracontractivity for (non-symmetric) diffusion semigroups on manifolds*, Forum Math. 15(2003), no. 6, 893–921.

- [18] M. Röckner, F.-Y. Wang, *Harnack and functional inequalities for generalized Mehler semigroups*, J. Funct. Anal. 203(2003), 237–261.
- [19] M. Röckner and F.-Y. Wang, *Log-Harnack inequality for stochastic differential equations in Hilbert spaces and its consequences*, arXiv:0911.0290.
- [20] A. Thalmaier, F.-Y. Wang, *Gradient estimates for harmonic functions on regular domains in Riemannian manifolds*, J. Funct. Anal. 155:1(1998), 109–124.
- [21] F.-Y. Wang, *Logarithmic Sobolev inequalities on noncompact Riemannian manifolds*, Probab. Theory Related Fields 109(1997), 417–424.
- [22] F.-Y. Wang, *Equivalence of dimension-free Harnack inequality and curvature condition*, Integral Equations Operator Theory 48(2004), no. 4, 547–552.
- [23] F.-Y. Wang, *Harnack inequality and applications for stochastic generalized porous media equations*, Ann. Probab. 35(2007), 1333–1350.
- [24] F.-Y. Wang, *Estimates of the first Neumann eigenvalue and the log-Sobolev constant on nonconvex manifolds*, Math. Nachr. 280(2007), 1431–1439.
- [25] F.-Y. Wang *Harnack inequalities on manifolds with boundary and applications*, arXiv:0908.2888.
- [26] F.-Y. Wang *Transportation-cost inequalities on path space over manifolds with boundary*, arXiv:0908.2891.
- [27] T. Yamada, S. Watanabe, *On the uniqueness of solutions of stochastic differential equations*, J. Math. Kyoto Univ. 11(1971), 155–167.
- [28] T.-S. Zhang, *White noise driven SPDEs with reflection: strong Feller properties and Harnack inequalities*, 2008 preprint.