

Harnack Inequalities for Semigroups with Non-Constant Diffusion Coefficients*

Feng-Yu Wang

School of Mathematical Sci. and Lab. Math. Com. Sys., Beijing Normal University, Beijing 100875, China
and

Department of Mathematics, Swansea University, Singleton Park, SA2 8PP, UK

Email: wangfy@bnu.edu.cn; F.Y.Wang@swansea.ac.uk

March 25, 2019

Abstract

By constructing a coupling with unbounded time-dependent drift, dimension-free Harnack inequalities are established for a large class of diffusion semigroups with non-constant diffusion coefficients. These inequalities are applied to the study of heat kernel upper bound and contractivity properties of the semigroup. The main results are extended to non-constant diffusions on manifolds possibly with (non-convex) boundary.

1 Introduction

Consider the following SDE on \mathbb{R}^d :

$$(1.1) \quad dX_t = \sigma(t, X_t)dB_t + b(t, X_t)dt,$$

where B_t is the d -dimensional Brownian motion on a complete filtered probability space $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$, and

$$\sigma : [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d, \quad b : [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d$$

are progressively measurable and continuous in the second variable. Throughout the paper we assume that for any $X_0 \in \mathbb{R}^d$ the equation (1.1) has a unique strong solution which is non-explosive.

*Supported in part by WIMCS, NNSFC(10721091) and the 973-Project.

To derive explicit Harnack type inequalities, we shall make use of the following assumptions.

(A1) There exists an increasing function $K : [0, \infty) \rightarrow \mathbb{R}$ such that

$$\|\sigma(t, x) - \sigma(t, y)\|_{HS}^2 + 2\langle b(t, x) - b(t, y), x - y \rangle \leq K_t |x - y|^2, \quad x, y \in \mathbb{R}^d, t \geq 0.$$

(A2) There exists a decreasing function $\lambda : [0, \infty) \rightarrow (0, \infty)$ such that $\sigma(t, x)^* \sigma(t, x) \geq \lambda_t^2 I$, $x \in \mathbb{R}^d, t \geq 0$.

(A3) There exists an increasing function $\delta : [0, \infty) \rightarrow (0, \infty)$ such that

$$|(\sigma(t, x) - \sigma(t, y))(x - y)| \leq \delta_t |x - y|, \quad x, y \in \mathbb{R}^d, t \geq 0.$$

(A4) For $n \geq 1$ there exists a constant $c_n > 0$ such that

$$\|\sigma(t, x) - \sigma(t, y)\|_{HS} + |b(t, x) - b(t, y)| \leq c_n |x - y|, \quad |x|, |y|, t \leq n.$$

It is well known that (A1) ensures the uniqueness of the solution to (1.1) while (A4) implies the existence and the uniqueness of the solution (see e.g. [12] and references within for weaker conditions), and when b and σ depend only on the variable $x \in \mathbb{R}^d$, the continuity will be enough to imply the existence of weak solution (see [14, Theorem 2.3]). By the Yamada-Watanabe principle [28], this implies the existence and uniqueness of the strong solution.

Let X_t^x be the solution to (1.1) for $X_0 = x$. We aim to establish the Harnack inequality for the operator P_t :

$$P_t f(x) := \mathbb{E} f(X_t^x), \quad t \geq 0, x \in \mathbb{R}^d, f \in \mathbf{B}_b^+(\mathbb{R}^d),$$

where $\mathbf{B}_b^+(\mathbb{R}^d)$ is the class of all bounded non-negative measurable functions on \mathbb{R}^d .

Note that if $\sigma(t, x)$ and $b(t, x)$ are deterministic and independent of t , then the solution is a time-homogeneous Markov process generated by

$$L := \frac{1}{2} \sum_{i,j=1}^d a_{ij} \partial_i \partial_j + \sum_{i=1}^d b_i \partial_i,$$

where $a := \sigma \sigma^*$. If further more σ and b are smooth, we may consider the Bakry-Emery curvature condition [5]:

$$(1.2) \quad \Gamma_2(f, f) \geq -K \Gamma(f, f), \quad f \in C^\infty(\mathbb{R}^d)$$

for some constant $K \in \mathbb{R}$, where

$$\Gamma(f, g) := \frac{1}{2} \sum_{i,j=1}^d a_{ij}(\partial_i f)(\partial_j g), \quad f, g \in C^1(\mathbb{R}^d),$$

$$\Gamma_2(f, f) := \frac{1}{2} L\Gamma(f, f) - \Gamma(f, Lf), \quad f \in C^\infty(\mathbb{R}^d).$$

According to [22, Lemma 2.2] and [23, Theorem 1.2], the curvature condition (4.1) is equivalent to the dimension-free Harnack inequality

$$(P_t f(x))^p \leq (P_t f^p(y)) \exp \left[\frac{p\rho_a(x, y)^2}{2(p-1)(1 - e^{-2Kt})} \right], \quad t \geq 0, p > 1, f \in \mathbf{B}_b^+(\mathbb{R}^d), x, y \in \mathbb{R}^d.$$

This type of inequality has been extended and applied to the study of heat kernel (or transition probability) and contractivity properties for diffusion semigroups for some finite- and infinite-dimensional models, see e.g. [1, 18, 4] for diffusions on manifolds with possibly unbounded below curvature, see [24, 16] for stochastic generalized porous media and fast diffusion equations [2, 3, 8, 17, 15, 19, 11, 29] for the study of some other SPDEs, with additive noise (i.e. constant diffusion coefficients).

Let us come back to the equation (1.1). When the dimension d is large and σ is non-constant, it is normally very hard to verify the curvature condition (4.1). Indeed, this condition depends on second order derivatives of a^{-1} , the inverse matrix of a . This is the main reason why existed results on the dimension-free Harnack inequality for SPDEs are only proved for the additive noise case.

For the additive noise case, i.e. σ is independent of the second component x , the dimension-free Harnack inequality can be derived by following the coupling argument developed in [4]. This argument has been also applied to the study of SPDEs in the above mentioned references. To see the difficulty in the study for non-constant diffusions, let us briefly recall the main idea of this argument.

To explain the key idea of the coupling, let e.g. σ and b be independent of the second variable. For $x \neq y$ and $T > 0$, let X_t solve (1.1) with $X_0 = x$ and Y_t solve

$$dY_t = \sigma(t)dB_t + b(t)dt + \frac{|x - y|(X_t - Y_t)}{T|X_t - Y_t|}dt, \quad Y_0 = y.$$

Then Y_t is well defined up to the coupling time

$$\tau := \inf\{t \geq 0 : X_t = Y_t\}.$$

Let $X_t = Y_t$ for $t \geq \tau$. We have

$$d|X_t - Y_t| = -\frac{|x - y|}{T}dt, \quad t \leq \tau.$$

This implies $\tau = T$ and hence, $X_T = Y_T$. On the other hand, by the Girsanov theorem we have

$$P_T f(y) = \mathbb{E}[Rf(Y_T)]$$

for

$$R := \exp \left[-\frac{|x-y|}{T} \int_0^T \frac{\langle \sigma(t)^{-1}(X_t - Y_t), dB_t \rangle}{|X_t - Y_t|} - \frac{|x-y|^2}{2T^2} \int_0^T \frac{|\sigma(t)^{-1}(X_t - Y_t)|^2}{|X_t - Y_t|^2} dt \right].$$

Therefore, since $X_T = Y_T$, it follows that

$$(P_T f(y))^p = (\mathbb{E}[Rf(X_T)])^p \leq (P_T f^p(x))(\mathbb{E}R^{p/(p-1)})^{p-1}.$$

Since by (A1) and (A2) it is easy to estimate moments of R , the desired Harnack inequality follows immediately.

In generally, if $\sigma(t, x)$ depends on x , then the process $X_t - Y_t$ contains a non-trivial martingale term, which can not be dominated by and bounded drift. So, in this case, any additional bounded drift put in the equation for Y_t is not enough to make the coupling successful before a fixed time T . This is the main difficulty to establish the Harnack inequality for diffusion semigroups with non-constant diffusion coefficient.

In this paper, under assumptions (A1) and (A2), we are able to constructed a coupling with a drift which is unbounded around a fixed time T , such that the coupling is successful before T . In this case the corresponding exponential martingale has finite entropy such that the log-Harnack inequality holds; if further more (A3) holds then the exponential martingale is L^p -integrable for some $p > 1$ such that the Harnack inequality with power hold. More precisely, we have the following result.

Theorem 1.1. *Let $\sigma(t, x)$ and $b(t, x)$ either be deterministic and independent of t , or satisfy (A4).*

(1) *If (A1) and (A2) hold then the log-Harnack inequality*

$$P_T \log f(y) \leq \log P_T f(y) + \frac{K_T |x - y|^2}{2\lambda_T^2(1 - e^{-K_T T})}, \quad f \geq 1, x, y \in \mathbb{R}^d, T > 0.$$

(2) *If (A1), (A2) and (A3) hold, then for $p > (1 + \frac{\delta_T}{\lambda_T})^2$ and $\delta_{p,T} := \max\{\delta_T, \frac{\lambda_T}{2}(\sqrt{p}-1)\}$, the Harnack inequality*

$$(P_T f(y))^p \leq (P_T f^p(x)) \exp \left[\frac{K_T \sqrt{p}(\sqrt{p}-1)|x-y|^2}{4\delta_{p,T}[(\sqrt{p}-1)\lambda_T - \delta_{p,T}](1 - e^{-K_T T})} \right]$$

holds for all $T > 0, x, y \in \mathbb{R}^d$ and $f \in \mathbf{B}_b^+(\mathbb{R}^d)$.

Theorem 1.1(1) generalizes a recent result in [20] on the log-Harnack inequality by using the gradient estimate on the diffusion semigroup.

Let $p_t(x, y)$ be the transition density of P_t w.r.t. a Radon measure μ . Then according to [26, Proposition 2.4], the above log-Harnack inequality and Harnack inequality are equivalent to the following heat kernel inequalities respectively:

$$(1.3) \quad \int_{\mathbb{R}^d} p_T(x, z) \log \frac{p_T(x, z)}{p_T(y, z)} \mu(dz) \leq \frac{K|x-y|^2}{2\lambda_T^2(1-e^{-K_T T})}, \quad x, y \in \mathbb{R}^d, T > 0$$

and

$$(1.4) \quad \int_{\mathbb{R}^d} p_T(x, z) \left(\frac{p_t(x, z)}{p_t(y, z)} \right)^{1/(p-1)} \mu(dz) \leq \exp \left[\frac{K_T \sqrt{p} |x-y|^2}{4\delta_{p,T}(\sqrt{p}+1)[(\sqrt{p}-1)\lambda_T - \delta_{p,T}](1-e^{-K_T T})} \right], \quad x, y \in \mathbb{R}^d, T > 0.$$

So, the following is a direct consequence of Theorem 1.1.

Corollary 1.2. *Let $\sigma(t, x)$ and $b(t, x)$ either be deterministic and independent of t , or satisfy (A4). Let P_t has a strictly positive density w.r.t. a Radon measure μ . Then (A1) and (A2) imply (1.3), while (A1)-(A3) imply (1.4).*

Next, by standard applications of the Harnack inequality with power, we have the following consequence of Theorem 1.1 on contractivity properties of P_t .

Corollary 1.3. *Let $\sigma(t, x)$ and $b(t, x)$ be deterministic and independent of t , such that (A1)-(A3) hold for constant K, λ and δ . Let P_t have an invariant probability measure μ .*

- (1) *If there exists $r > K^+/\lambda^2$ such that $\mu(e^{r|\cdot|^2}) < \infty$, then P_t is hypercontractive, i.e. $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} = 1$ holds for some $t > 0$.*
- (2) *If $\mu(e^{r|\cdot|^2}) < \infty$ holds for all $r > 0$, then P_t is supercontractive, i.e. $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} < \infty$ holds for all $t > 0$.*
- (3) *If $P_t e^{r|\cdot|^2}$ is bounded for any $t, r > 0$, then P_t is ultracontractive, i.e. $\|P_t\|_{L^2(\mu) \rightarrow L^\infty(\mu)} < \infty$ for any $t > 0$.*

Remark 1.1. In the situation of Corollary 1.3 with μ a probability measure, when P_t is symmetric w.r.t. μ , the hypercontractivity is equivalent to the validity of the log-Sobolev inequality

$$\mu(f^2 \log f^2) \leq C \mu(\Gamma(f, f)), \quad f \in C_b^\infty(\mathbb{R}^d), \mu(f^2) = 1$$

for some constant $C > 0$. Moreover, by (A2) and (A3) there exists a constant $R > 0$ such that

$$R^{-1}|\nabla f|^2 \leq \Gamma(f, f) \leq R|\nabla f|^2$$

holds for all $f \in C^\infty(\mathbb{R}^d)$. So, by the concentration of measure for the log-Sobolev inequality, the hypercontractivity implies that $\mu(e^{r|\cdot|^2}) < \infty$ for some $r > 0$, while the supercontractivity implies that $\mu(e^{r|\cdot|^2}) < \infty$. Combining this with Corollary 1.3, we have the following assertions.

- (i) Let $K \leq 0$. Then P_t is hypercontractive if and only if $\mu(e^{r|\cdot|^2}) < \infty$ holds for some $r > 0$.
- (ii) P_t is supercontractive if and only if $\mu(e^{r|\cdot|^2}) < \infty$ holds for all $r > 0$.
- (iii) P_t is ultracontractive if and only if $P_t e^{r|\cdot|^2}$ is bounded for any $t, r > 0$.

Therefore, conditions in Corollary 1.3(2) and (3) are sharp for the supercontractivity and ultracontractivity of P_t . Moreover, as shown in [7] that when σ is constant, the sufficient condition $\mu(e^{r|\cdot|^2}) < \infty$ for some $r > K^+/\lambda^2$ is optimal for the hypercontractivity of P_t . So, Corollary 1.3(1) also provides a sharp sufficient condition for the hypercontractivity of P_t .

We will prove Theorem 1.1 and Corollary 1.3 in the next section. In section 3 we extend these results to SDEs on Riemannian manifolds possibly with a convex boundary. Combining the results in Section 3 with a conformal change method introduced in [24], we are able to establish Harnack inequalities in Section 4 for the Neumann semigroup on a class of non-convex manifolds. Finally, we extend Theorem 1.1 and its consequences to semi-linear stochastic equations on Hilbert spaces.

2 Proofs of Theorem 1.1 and Corollary 1.3

Let $x, y \in \mathbb{R}^d, T > 0$ and $p > (1 + \delta_T/\lambda_T)^2$ be fixed such that $x \neq y$. We have

$$(2.1) \quad \theta_T := \frac{2\delta_T}{(\sqrt{p}-1)\lambda_T} \in (0, 2).$$

For $\theta \in (0, 2)$, let

$$\xi_t = \frac{2-\theta}{K_T}(1 - e^{K_T(t-T)}), \quad t \in [0, T].$$

Then ξ is smooth and strictly positive on $[0, T)$ such that

$$(2.2) \quad 2 - K_T \xi_t + \xi'_t = \theta, \quad t \in [0, T].$$

Consider the coupling

$$(2.3) \quad \begin{aligned} dX_t &= \sigma(t, X_t)dB_t + b(t, X_t)dt, \quad X_0 = x, \\ dY_t &= \sigma(t, Y_t)dB_t + b(t, Y_t)dt + \frac{1}{\xi_t}\sigma(t, Y_t)\sigma(t, X_t)^{-1}(X_t - Y_t)dt, \quad Y_0 = y. \end{aligned}$$

Since the additional drift term $\xi_t^{-1}\sigma(t, y)\sigma(t, x)^{-1}(x - y)$ is locally Lipschitzian in y if (A4) holds, and continuous in y when σ and b are deterministic and time independent, the coupling (X_t, Y_t) is well defined for $t < T \wedge \zeta$, where ζ is the explosive time of Y_t ; namely, $\zeta = \lim_{n \rightarrow \infty} \zeta_n$ for

$$\zeta_n := \inf\{t \in [0, T) : |Y_t| \geq n\},$$

here and in what follows we set $\inf \emptyset = \infty$ by convention. Let

$$d\tilde{B}_t = dB_t + \frac{1}{\xi_t}\sigma(t, X_t)^{-1}(X_t - Y_t)dt, \quad t < T \wedge \zeta.$$

If $\zeta = \infty$ and the exponential martingale

$$R_s := \exp \left[- \int_0^s \xi_t^{-1} \langle \sigma(t, X_t)^{-1}(X_t - Y_t), dB_t \rangle - \frac{1}{2} \int_0^s \xi_t^{-2} |\sigma(t, X_t)^{-1}(X_t - Y_t)|^2 dt \right]$$

is uniformly integrable for $s \in [0, T)$, i.e.

$$\lim_{n \rightarrow \infty} \sup_{s \in [0, T)} \mathbb{E}[R_s 1_{\{R_s > n\}}] = 0,$$

then by the martingale convergence theorem, $R_T := \lim_{t \uparrow T} R_t$ exists and $\{R_t\}_{t \in [0, T]}$ is a martingale. In this case, by the Girsanov theorem $\{\tilde{B}_t\}_{t \in [0, T)}$ is a d -dimensional Brownian motion under the probability $R_T \mathbb{P}$. Rewrite (2.3) as

$$(2.4) \quad \begin{aligned} dX_t &= \sigma(t, X_t)d\tilde{B}_t + b(t, X_t)dt - \frac{X_t - Y_t}{\xi_t}dt, \quad X_0 = x, \\ dY_t &= \sigma(t, Y_t)d\tilde{B}_t + b(t, Y_t)dt, \quad Y_0 = y. \end{aligned}$$

Since $\int_0^T \xi_t^{-1} dt = \infty$, we will see that the additional drift $-\frac{X_t - Y_t}{\xi_t}dt$ is strong enough to force the coupling successful up to time T . So, we first study the uniform integrability of $\{R_s\}_{s \in [0, T)}$.

In general where ζ might be finite under \mathbb{P} , we shall construct a weighted probability measure \mathbb{Q} such that $\zeta = \infty$ \mathbb{Q} -a.s. Let

$$\tau_n = \inf\{t \in [0, T) : |X_t| + |Y_t| \geq n\}.$$

Since X_t is non-explosive as assumed, we have $\tau_n \uparrow \zeta$ as $n \uparrow \infty$.

Lemma 2.1. Assume (A1) and (A2) and let $\theta \in (0, 2)$, $x, y \in \mathbb{R}^d$ and $T > 0$ be fixed.

(1) There holds

$$\sup_{s \in [0, T], n \geq 1} \mathbb{E} R_{s \wedge \tau_n} \log R_{s \wedge \tau_n} \leq \frac{K_T |x - y|^2}{2\lambda_T^2 \theta (2 - \theta) (1 - e^{-K_T T})}.$$

Consequently,

$$R_{s \wedge \zeta} := \lim_{n \uparrow \infty} R_{s \wedge \tau_n}, \quad s \in [0, T), \quad R_{T \wedge \zeta} := \lim_{s \uparrow T} R_{s \wedge \zeta}$$

exist such that $\{R_{s \wedge \zeta}\}_{s \in [0, T]}$ is a uniformly integrable martingale.

(2) Let $\mathbb{Q} = R_{T \wedge \zeta} \mathbb{P}$. Then $\mathbb{Q}(\zeta = \infty) = 1$.

Proof. (1) Let $s \in [0, T]$ be fixed. By (2.4), (A1) and the Itô formula,

$$d\|X_t - Y_t\|^2 \leq 2\langle (\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t \rangle + K_T |X_t - Y_t|^2 dt - \frac{2}{\xi_t} |X_t - Y_t|^2 dt$$

holds for $t \leq s \wedge \tau_n$. Combining this with (2.2) we obtain

$$\begin{aligned} & d \frac{|X_t - Y_t|^2}{\xi_t} \\ (2.5) \quad & \leq \frac{2}{\xi_t} \langle (\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t \rangle - \frac{|X_t - Y_t|^2}{\xi_t^2} (2 - K_T \xi_t + \xi_t') dt \\ & = \frac{2}{\xi_t} \langle (\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t \rangle - \frac{\theta}{\xi_t^2} |X_t - Y_t|^2 dt, \quad t \leq s \wedge \tau_n. \end{aligned}$$

Since due to the Girsanov theorem $\{\tilde{B}_t\}_{t \leq \tau_n \wedge s}$ is the d -dimensional Brownian motion under the probability measure $R_{s \wedge \tau_n} \mathbb{P}$, letting $\mathbb{E}_{s, n}$ denote the corresponding expectation we obtain

$$(2.6) \quad \mathbb{E}_{s, n} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \leq \frac{|x - y|^2}{\theta \xi_0}, \quad s \in [0, T], n \geq 1.$$

By (A2) and the definitions of R_t and \tilde{B}_t , we have

$$\begin{aligned} \log R_r &= - \int_0^r \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1} (X_t - Y_t), d\tilde{B}_t \rangle + \frac{1}{2} \int_0^r \frac{|\sigma(t, X_t)^{-1} (X_t - Y_t)|^2}{\xi_t^2} dt \\ &\leq - \int_0^r \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1} (X_t - Y_t), d\tilde{B}_t \rangle + \frac{1}{2\lambda_T^2} \int_0^r \frac{|X_t - Y_t|^2}{\xi_t^2} dt, \quad r \leq s \wedge \tau_n. \end{aligned}$$

Since $\{\tilde{B}_t\}$ is the d -dimensional Brownian motion under $R_{s \wedge \tau_n} \mathbb{P}$ up to $s \wedge \tau_n$, combining this with (2.6) we obtain

$$\mathbb{E} R_{s \wedge \tau_n} \log R_{s \wedge \tau_n} = \mathbb{E}_{s,n} \log R_{s \wedge \tau_n} \leq \frac{|x - y|^2}{2\lambda_T^2 \theta \xi_0}, \quad s \in [0, T], n \geq 1.$$

By the martingale convergence theorem and the Fatou lemma, $\{R_{s \wedge \zeta} : s \in [0, T]\}$ is a well-defined martingale with

$$\mathbb{E} R_{s \wedge \zeta} \log R_{s \wedge \zeta} \leq \frac{|x - y|^2}{2\lambda_T^2 \theta \xi_0} = \frac{K_T |x - y|^2}{2\lambda_T^2 \theta (2 - \theta)(1 - e^{-K_T T})}, \quad s \in [0, T].$$

(2) Let $\sigma_n = \inf\{t \geq 0 : |X_t| \geq n\}$. We have $\sigma_n \uparrow \infty$ \mathbb{P} -a.s and hence, also \mathbb{Q} -a.s. Since $\{\tilde{B}_t\}$ is a \mathbb{Q} -Brownian motion up to $T \wedge \zeta$, it follows from (2.5) that

$$\frac{(n - m)^2}{\xi_0} \mathbb{Q}(\sigma_m > t, \zeta_n \leq t) \leq \mathbb{E}_{\mathbb{Q}} \frac{|X_{t \wedge \sigma_m \wedge \zeta_n} - X_{t \wedge \sigma_m \wedge \zeta_n}|^2}{\xi_{t \wedge \sigma_m \wedge \zeta_n}} \leq \frac{|x - y|^2}{\xi_0}$$

holds for all $n > m > 0$ and $t \in [0, T]$. By letting first $n \uparrow \infty$ then $m \uparrow \infty$, we obtain $\mathbb{Q}(\zeta \leq t) = 0$ for all $t \in [0, T]$. This is equivalent to $\mathbb{Q}(\zeta = \infty) = 1$ according to the definition of ζ . \square

Lemma 2.1 ensures that under $\mathbb{Q} := R_{T \wedge \zeta} \mathbb{P}$, $\{\tilde{B}_t\}_{t \in [0, T]}$ is a Brownian motion, so that the coupling (X_t, Y_t) is well-constructed for $t \in [0, T]$ by (2.4). Since $\int_0^T \xi_t^{-1} dt = \infty$, we shall see that the coupling is successful up to time T , so that $X_T = Y_T$ holds (see the proof of Theorem 1.1 below). This will provide the desired Harnack inequality for P_t as explained in Section 1 as soon as $R_{T \wedge \zeta}$ has finite $p/(p - 1)$ -moment. The next lemma provides an explicit upper bound on moments of $R_{T \wedge \zeta}$.

Lemma 2.2. *Assume (A1)-(A3). Let R_t and ξ_t be fixed for $\theta = \theta_T$. We have*

$$(2.7) \quad \sup_{s \in [0, T]} \mathbb{E} \left\{ R_{s \wedge \zeta} \exp \left[\frac{\theta_T^2}{8\delta_T^2} \int_0^{s \wedge \zeta} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right\} \leq \exp \left[\frac{\theta_T K_T |x - y|^2}{4\delta_T^2 (2 - \theta_T)(1 - e^{-K_T T})} \right].$$

Consequently,

$$(2.8) \quad \sup_{s \in [0, T]} \mathbb{E} R_{s \wedge \zeta}^{1+r_T} \leq \exp \left[\frac{\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x - y|^2}{8\delta_T^2 (2 - \theta_T)(\delta_T + \theta_T \lambda_T)(1 - e^{-K_T T})} \right]$$

holds for

$$r_T = \frac{\lambda_T^2 \theta_T^2}{4\delta_T^2 + 4\theta_T \lambda_T \delta_T}.$$

Proof. Let $\theta = \theta_T$. By (2.5), for an $r > 0$ we have

$$\begin{aligned} & \mathbb{E}_{s,n} \exp \left[r \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \\ & \leq \exp \left[\frac{r|x-y|^2}{\theta_T \xi_0} \right] \cdot \mathbb{E}_{s,n} \exp \left[\frac{2r}{\theta_T} \int_0^{s \wedge \tau_n} \frac{1}{\xi_t} \langle (\sigma(t, X_t) - \sigma(t, Y_t))(X_t - Y_t), d\tilde{B}_t \rangle \right] \\ & \leq \exp \left[\frac{rK_T|x-y|^2}{\theta_T(2-\theta_T)(1-e^{-K_T T})} \right] \left(\mathbb{E}_{s,n} \exp \left[\frac{8r^2\delta_T^2}{\theta_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right)^{1/2}, \end{aligned}$$

where the last step is due to (A3) and the fact that

$$\mathbb{E}e^{M_t} \leq (\mathbb{E}e^{2\langle M \rangle_t})^{1/2}$$

for a continuous exponential integrable martingale M_t . Taking $r = \theta_T^2/(8\delta_T^2)$, we arrive at

$$\mathbb{E}_{s,n} \exp \left[\frac{\theta_T^2}{8\delta_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \leq \left[\frac{\theta_T K_T |x-y|^2}{4\delta_T^2(2-\theta_T)(1-e^{-K_T T})} \right], \quad n \geq 1.$$

This implies (2.7) by letting $n \rightarrow \infty$.

Next, by (A2) and the definition of R_s we have

$$\begin{aligned} (2.9) \quad & \mathbb{E}R_{s \wedge \tau_n}^{1+r_T} = \mathbb{E}_{s,n} R_{s \wedge \tau_n}^{r_T} \\ & = \mathbb{E}_{s,n} \exp \left[-r_T \int_0^{s \wedge \tau_n} \frac{1}{\xi_t} \langle \sigma(t, X_t)^{-1}(X_t - Y_t), d\tilde{B}_t \rangle \right. \\ & \quad \left. + \frac{r_T}{2} \int_0^{s \wedge \tau_n} \frac{|\sigma(t, X_t)^{-1}(X_t - Y_t)|^2}{\xi_t^2} dt \right]. \end{aligned}$$

Noting that for any exponential integrable martingale M_t w.r.t. $R_{s \wedge \tau_n} \mathbb{P}$, one has

$$\begin{aligned} & \mathbb{E}_{s,n} \exp[r_T M_t + r_T \langle M \rangle_t / 2] = \mathbb{E}_{s,n} \exp[r_T M_t - r_T^2 q \langle M \rangle_t / 2 + r_T(qr_T + 1) \langle M \rangle_t / 2] \\ & \leq (\mathbb{E}_{s,n} \exp[r_T q M_t - r_T^2 q^2 \langle M \rangle_t / 2])^{1/q} \left(\mathbb{E}_{s,n} \exp \left[\frac{r_T q(r_T q + 1)}{2(q-1)} \langle M \rangle_t \right] \right)^{(q-1)/q} \\ & = \left(\mathbb{E}_{s,n} \exp \left[\frac{r_T q(r_T q + 1)}{2(q-1)} \langle M \rangle_t \right] \right)^{(q-1)/q}, \quad q > 1, \end{aligned}$$

it follows from (2.9) that

$$(2.10) \quad \mathbb{E}R_{s \wedge \tau_n}^{1+r_T} \leq \left(\mathbb{E}_{s,n} \exp \left[\frac{qr_T(qr_T + 1)}{2(q-1)\lambda_T^2} \int_0^{s \wedge \tau_n} \frac{|X_t - Y_t|^2}{\xi_t^2} dt \right] \right)^{(q-1)/q}.$$

Take

$$(2.11) \quad q = 1 + \sqrt{1 + r_T^{-1}},$$

which minimizes $q(qr_T + 1)/(q - 1)$ such that

$$(2.12) \quad \begin{aligned} \frac{qr_T(qr_T + 1)}{2\lambda_T^2(q - 1)} &= \frac{r_T + \sqrt{r_T(r_T + 1)}}{2\lambda_T^2\sqrt{1 + r_T^{-1}}} (r_T + 1 + \sqrt{r_T(r_T + 1)}) \\ &= \frac{(r_T + \sqrt{r_T^2 + r_T})^2}{2\lambda_T^2} = \frac{\theta_T^2}{8\delta_T^2}. \end{aligned}$$

Combining (2.10) with (2.7) and (2.12), and noting that due to (2.11) and the definition of r_T

$$\frac{q - 1}{q} = \frac{\sqrt{1 + r_T^{-1}}}{1 + \sqrt{1 + r_T^{-1}}} = \frac{2\delta_T + \theta_T\lambda_T}{2\delta_T + 2\theta_T\lambda_T},$$

we obtain

$$\mathbb{E}R_{s \wedge \tau_n}^{1+r_T} \leq \exp \left[\frac{\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x - y|^2}{8\delta_T^2 (2 - \theta_T) (\delta_T + \theta_T \lambda_T) (1 - e^{-K_T T})} \right].$$

According to the Fatou lemma, the proof is then completed by letting $n \rightarrow \infty$. \square

Proof of Theorem 1.1. Since (A3) also holds for $\delta_{p,T}$ in place of δ_T , it suffices to prove the desired Harnack inequality for δ_T in place of $\delta_{p,T}$.

(1) By Lemma 2.1, $\{R_{s \wedge \zeta}\}_{s \in [0, T]}$ is a uniformly integrable martingale and $\{\tilde{B}_t\}_{t \leq T}$ is a d -dimensional Brownian motion under the probability \mathbb{Q} . Thus, Y_t can be solved up to time T . Let

$$\tau = \inf\{t \in [0, T] : X_t = Y_t\}.$$

We claim that under \mathbb{Q} , $\tau \leq T$ a.s. so that $X_T = Y_T$ a.s. Indeed, if for some $\omega \in \Omega$ such that $\tau(\omega) > T$, by the continuity of the diffusions we have

$$\inf_{t \in [0, T]} |X_t - Y_t|^2(\omega) > 0.$$

So,

$$\int_0^T \frac{|X_t - Y_t|^2}{\xi_t^2} dt = \infty$$

holds on the set $\{\tau > T\}$. But according to Lemma 2.2 we have

$$\mathbb{E}_{\mathbb{Q}} \int_0^T \frac{|X_t - Y_t|^2}{\xi_t^2} dt < \infty,$$

we conclude that $\mathbb{Q}(\tau > T) = 0$. Therefore, $X_T = Y_T$ \mathbb{Q} -a.s.

Now, combining Lemma 2.1 with $X_T = Y_T$ and using the Young inequality, for $f \geq 1$ we have

$$\begin{aligned} P_T \log f(y) &= \mathbb{E}_{\mathbb{Q}}[\log f(Y_T)] = \mathbb{E}[R_{T \wedge \zeta} \log f(X_T)] \\ &\leq \mathbb{E} R_{T \wedge \zeta} \log R_{T \wedge \zeta} + \log \mathbb{R} f(X_T) \leq \log P_T f(x) + \frac{K_T |x - y|^2}{2\lambda_T^2 \theta (2 - \theta) (1 - e^{-K_T T})}. \end{aligned}$$

This completes the proof of (1) by taking $\theta = 1$.

(2) Let $\theta = \theta_T$. Since $X_T = Y_T$ and $\{\tilde{B}_t\}_{t \in [0, T]}$ is the d -dimensional Brownian motion under \mathbb{Q} , we have

$$(2.13) \quad (P_T f(y))^p = (\mathbb{E}_{\mathbb{Q}}[f(Y_T)])^p = (\mathbb{E}[R_{T \wedge \zeta} f(X_T)])^p \leq (P_T f^p(x)) (\mathbb{E} R_{T \wedge \zeta}^{p/(p-1)})^{p-1}.$$

Due to (2.1) we have

$$\frac{p}{p-1} = 1 + \frac{\lambda_T^2 \theta_T^2}{4\delta_T(\delta_T + \theta_T \lambda_T)}.$$

So, it follows from Lemma 2.2 and (2.1) that

$$\begin{aligned} (\mathbb{E} R_{T \wedge \zeta}^{p/(p-1)})^{p-1} &= (\mathbb{E} R_{T \wedge \zeta}^{1+r_T})^{p-1} \leq \exp \left[\frac{(p-1)\theta_T K_T (2\delta_T + \theta_T \lambda_T) |x - y|^2}{8\delta_T^2 (2 - \theta_T) (\delta_T + \theta_T \lambda_T) (1 - e^{-K_T T})} \right] \\ &= \exp \left[\frac{K_T \sqrt{p} (\sqrt{p} - 1) |x - y|^2}{4\delta_T [(\sqrt{p} - 1)\lambda_T - \delta_T] (1 - e^{-K_T T})} \right]. \end{aligned}$$

The proof is then completed by combining this with (2.13). \square

Proof of Corollary 1.3. Let $f \in \mathbf{B}_b^+(\mathbb{R}^d)$ be such that $\mu(f^p) \leq 1$. Let $p > (1 + \delta/\lambda)^2$. By Theorem 1.1(2), we have

$$(P_t f(y))^p \exp \left[- \frac{K \sqrt{p} (\sqrt{p} - 1) |x - y|^2}{4\delta_p [(\sqrt{p} - 1)\lambda - \delta_p] (1 - e^{-Kt})} \right] \leq P_t f^p(x), \quad x, y \in \mathbb{R}^d,$$

where $\delta_p = \max\{\delta, \frac{\lambda}{2}(\sqrt{p} - 1)\}$. Integrating w.r.t. $\mu(dx)$ and noting that μ is P_t -invariant, we obtain

$$(2.14) \quad (P_t f(y))^p \int_{\mathbb{R}^d} \exp \left[- \frac{K\sqrt{p}(\sqrt{p}-1)|x-y|^2}{4\delta[(\sqrt{p}-1)\lambda-\delta](1-e^{-Kt})} \right] \mu(dx) \leq 1.$$

Taking $f = n \wedge (p_t(y, \cdot))^{1/p}$ and letting $n \uparrow \infty$, we prove the first assertion.

Next, let $B(0, 1) = \{x \in \mathbb{R}^d : |x| \leq 1\}$. Since μ is an invariant measure, it has a strictly positive density w.r.t. the Lebesgue measure so that $\mu(B(0, 1)) > 0$ (cf. [6]). Let $p \geq (1 + 2\delta/\lambda)^2$. We have $\delta_p = (\sqrt{p}-1)\lambda/2$ and thus,

$$\frac{\sqrt{p}(\sqrt{p}-1)}{4\delta_p[(\sqrt{p}-1)\lambda-\delta_p]} = \frac{\sqrt{p}}{\lambda^2(\sqrt{p}-1)}.$$

Combining this with (2.14) and noting that

$$\begin{aligned} & \int_{\mathbb{R}^d} \exp \left[- \frac{K\sqrt{p}(\sqrt{p}-1)|x-y|^2}{4\delta[(\sqrt{p}-1)\lambda-\delta](1-e^{-Kt})} \right] \mu(dx) \\ & \geq \mu(B(0, 1)) \exp \left[- \frac{K\sqrt{p}(\sqrt{p}-1)(1+|y|)^2}{4\delta[(\sqrt{p}-1)\lambda-\delta](1-e^{-Kt})} \right], \end{aligned}$$

we obtain

$$(2.15) \quad (P_t f(y))^p \leq C_1 \exp \left[\frac{K\sqrt{p}(1+|y|)^2}{\lambda_T^2(\sqrt{p}-1)(1-e^{-Kt})} \right], \quad t > 0, y \in \mathbb{R}^d$$

for some constant $C_1 > 0$ and all $f \in \mathbf{B}_b^+(\mathbb{R}^d)$ with $\mu(f^p) \leq 1$. Since

$$\lim_{p \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{K\sqrt{p}}{\lambda^2(\sqrt{p}-1)(1-e^{-Kt})} = \frac{K^+}{\lambda^2},$$

for any $r > K^+/\lambda^2$ there exist $p > (1 + 2\delta_T/\lambda)^2, \beta > 1$ and $t_1 > 0$ such that

$$(P_{t_1} f(y))^{\beta p} \leq C_2 e^{r|y|^2}, \quad y \in \mathbb{R}^d, f \in \mathbf{B}_b^+(\mathbb{R}^d), \mu(f^p) \leq 1$$

holds for some constant $C_2 > 0$. Thus, $\mu(e^{r|\cdot|^2}) < \infty$ implies that $\|P_{t_1}\|_{L^p(\mu) \rightarrow L^{p\beta}(\mu)} < \infty$. Since $\|P_s\|_{L^q(\mu)} = 1$ holds for any $q \in [1, \infty]$, by the interpolation theorem and the semigroup property one may find $t_2 > t_1$ such that

$$(2.16) \quad \|P_{t_2}\|_{L^2(\mu) \rightarrow L^4(\mu)} < \infty.$$

Moreover, by [13, Theorem 3.6(ii)], there exist some constants $\eta, C_3 > 0$ such that

$$\|P_t - \mu\|_{L^2(\mu)} \leq C_3 e^{-\eta t}, \quad t \geq 0.$$

Combining this with (2.16) we conclude that $\|P_t\|_{L^2(\mu) \rightarrow L^4(\mu)} \leq 1$ holds for sufficiently large $t > 0$, i.e. (2) holds.

Finally, (3) and (4) follow immediately from (2.15) and the interpolation theorem. \square

3 Extension to manifolds with convex boundary

Let M be a d -dimensional complete, connected Riemannian manifold, possibly with a convex boundary ∂M . Let N be the inward unit normal vector field of ∂M when $\partial M \neq \emptyset$. Let P_t be the (Neumann) semigroup generated by

$$L := \psi^2(\Delta + Z)$$

on M , where $\psi \in C^1(M)$ and Z is a C^1 vector field on M . Assume that ψ is bounded and

$$(3.1) \quad \text{Ric} - \nabla Z \geq -K_0$$

holds for some constant $K_0 \geq 0$. Then the (reflecting) diffusion process generated by L is non-explosive.

To formulae P_t as the semigroup associated as SDE like (1.1), we set

$$(3.2) \quad \sigma = \sqrt{2}\psi, \quad b = \psi^2 Z.$$

Then P_t is the semigroup for the solution to the following Itô SDE (see e.g. [10] or [4] for the definition of d_I on Riemannian manifolds):

$$(3.3) \quad d_I X_t = \sigma(X_t) \Phi_t dB_t + b(X_t) dt + N(X_t) dl_t,$$

where B_t is the d -dimensional Brownian motion on a complete filtered probability space $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$, Φ_t is the horizontal lift of X_t onto the frame bundle $O(M)$, and l_t is the local time of X_t on ∂M . When $\partial M = \emptyset$, we simply set $l_t = 0$.

To derive the Harnack inequality as in Section 2, we assume that

$$(3.4) \quad \lambda := \inf \sigma > 0, \quad \delta := \sup \sigma - \inf \sigma < \infty.$$

Now, let $x, y \in M$ and $T > 0$ be fixed. Let ρ be the Riemannian distance on M , i.e. $\rho(x, y)$ is the length of the minimal geodesic on M linking x and y , which exists if ∂M is either convex or empty.

Let X_t solve (3.3) with $X_0 = x$. Next any strictly positive function $\xi \in C([0, T])$, let Y_t solve

$$d_I Y_t = \sigma(Y_t) P_{X_t, Y_t} \Phi_t dB_t + b(X_t) dt - \frac{\sigma(Y_t) \rho(X_t, Y_t)}{\sigma(X_t) \xi_t} \nabla \rho(X_t, \cdot)(Y_t) dt + N(Y_t) d\tilde{l}_t$$

for $Y_0 = y$, where \tilde{l}_t is the local time of Y_t on ∂M , and $P_{X_t, Y_t} : T_{X_t}M \rightarrow T_{Y_t}M$ is the parallel displacement along the minimal geodesic from X_t to Y_t , which exists since ∂M is convex or empty. As explained in [4, Section 3], we may and do assume that the cut-locus of M is empty such that the parallel displacement is smooth. Let

$$d\tilde{B}_t = dB_t + \frac{\rho(X_t, Y_t)}{\xi_t \sigma(X_t)} \Phi_t^{-1} \nabla \rho(\cdot, Y_t)(X_t) dt, \quad t < T.$$

By the Girsanov theorem, for any $s \in (0, T)$ the process $\{\tilde{B}_t\}_{t \in [0, s]}$ is the d -dimensional Brownian motion under the weighted probability measure $R_s \mathbb{P}$, where

$$(3.5) \quad R_s := \exp \left[- \int_0^s \frac{\rho(X_t, Y_t)}{\xi_t \sigma(X_t)} \langle \nabla \rho(\cdot, Y_t)(X_t), \phi_t dB_t \rangle - \frac{1}{2} \int_0^s \frac{\rho(X_t, Y_t)^2}{\xi_t^2 \sigma(X_t)^2} dt \right].$$

Thus, by (3.2) we have

$$\begin{aligned} d_I X_t &= \sqrt{2} \psi(X_t) \Phi_t d\tilde{B}_t + (\psi^2 Z)(X_t) dt - \frac{\rho(X_t, Y_t)}{\xi_t} \nabla \rho(\cdot, Y_t)(X_t) dt + N(X_t) dl_t, \\ d_I Y_t &= \sqrt{2} \psi(Y_t) \Phi_t d\tilde{B}_t + (\psi^2 Z)(Y_t) dt + N(Y_t) d\tilde{l}_t. \end{aligned}$$

Let $\xi \in C^1([0, T])$ be strictly positive and take

$$\beta_t = - \frac{\rho(X_t, Y_t)}{\sqrt{2} \xi_t \psi(X_t)} \Phi_t^{-1} \nabla(\cdot, Y_t)(X_t).$$

Repeating the proof of (4.10) in [27] we obtain

$$d\rho(X_t, Y_t) \leq (\sigma(X_t) - \sigma(Y_t)) \langle \nabla \rho(\cdot, Y_t)(X_t), \Phi_t d\tilde{B}_t \rangle + K_1 \rho(X_t, Y_t) dt - \frac{\rho(X_t, Y_t)}{\xi_t} dt, \quad t < T,$$

where

$$K_1 = K_0 \|\psi\|_\infty^2 + 2\|Z\|_\infty \|\nabla \psi\|_\infty \|\psi\|_\infty.$$

This implies that

$$d \frac{\rho(X_t, Y_t)^2}{\xi_t} \leq \frac{2}{\xi_t} \rho(X_t, Y_t) (\sigma(X_t) - \sigma(Y_t)) \langle \nabla \rho(\cdot, Y_t)(X_t), \Phi_t d\tilde{B}_t \rangle - \frac{\rho(X_t, Y_t)^2}{\xi_t^2} (2 - K\xi_t + \xi_t') dt$$

holds for

$$(3.6) \quad \begin{aligned} K &:= 2K_1 + \|\nabla \sigma\|_\infty^2 \\ &= 2K_0 \|\psi\|_\infty^2 + 4\|Z\|_\infty \|\nabla \psi\|_\infty \|\psi\|_\infty + 2\|\nabla \psi\|_\infty^2. \end{aligned}$$

In particular, letting

$$\xi_t = \frac{2-\theta}{K}(1 - e^{K(t-T)}), \quad t \in [0, T], \theta \in (0, 2),$$

we have

$$2 - K\xi_t + \xi'_t = \theta.$$

Therefore, the following result follows immediately by repeating calculations in Section 2.

Theorem 3.1. *Assume that ∂M is either empty or convex. Let (4.1) and Z, ϕ be bounded such that*

$$K := 2K_0\|\psi\|_\infty^2 + 4\|Z\|_\infty\|\nabla\psi\|_\infty\|\psi\|_\infty + 2\|\nabla\psi\|_\infty^2 < \infty.$$

Then all assertions in Theorem 1.1 and Corollaries 1.2, 1.3 hold for P_t the (Neumann) semigroup generated on M in place of \mathbb{R}^d by $L = \psi^2(\Delta + Z)$ on M for constant functions $K, \delta := \sup \psi - \inf \psi$ and $\lambda := \inf |\psi|$.

4 Neumann semigroup on non-convex manifolds

Following the line of [24], we are able to make the boundary from non-convex to convex by using a conformal change of metric. This will enable us to extend our results to the Neumann semigroup on a class of non-convex manifolds.

Let $\partial M \neq \emptyset$ with N the inward normal unit vector field. Then the second fundamental form of ∂M is a two-tensor on the tangent space of ∂M defined by

$$\mathbb{I}(X, Y) := -\langle \nabla_X N, Y \rangle, \quad X, Y \in T\partial M.$$

Assume that there exists $\kappa > 0$ and $K_0 \in \mathbb{R}$ such that

$$(4.1) \quad \text{Ric} - \nabla Z \geq -K_0, \quad \mathbb{I} \geq -\kappa$$

holds for M and a C^1 vector field Z . We shall consider the Harnack inequality for the Neumann semigroup P_t generated by

$$L = \Delta + Z.$$

To make the boundary convex, let $f \in C_b^\infty(M)$ such that $f \geq 1$ and $N \log f|_{\partial M} \geq \kappa$. By [24, Lemma 2.1], ∂M is convex under the metric

$$\langle \cdot, \cdot \rangle' = f^{-2} \langle \cdot, \cdot \rangle.$$

Let Δ' and ∇' be the Laplacian and gradient induced by the new metric. We have (see (2.2) in [21])

$$L = f^{-2}(\Delta' + Z'), \quad Z' = f^2 Z + \frac{d-2}{2} \nabla f^2.$$

So, Theorem 3.1 applies for the convex manifold $(M, \langle \cdot, \cdot \rangle')$. Let Ric' be the Ricci curvature induced by the metric $\langle \cdot, \cdot \rangle'$. We have (see the proof of [27, Theorem 5.1])

$$\text{Ric}' - \nabla' Z' \geq -K_f \langle \cdot, \cdot \rangle'$$

for

$$(4.2) \quad K_f = \sup \left\{ K f^2 - d \Delta f + (d-3) |\nabla f|^2 + 3 |Z| f |\nabla f| \right\}.$$

Thus, Theorem 3.1 holds for $\psi = f^{-1}$ and

$$\begin{aligned} K &= 2K_f^+ \|f^{-1}\|_\infty + 4\|Z'\|'_\infty \|\nabla' f^{-1}\|'_\infty \|f^{-1}\|_\infty + 2\|\nabla' f^{-1}\|'^2_\infty \\ &\leq 2K_f^+ + 4\|fZ + (d-2)\nabla\|_\infty \|\nabla f\|_\infty + 2\|\nabla f\|_\infty^2, \end{aligned}$$

where $\|\cdot\|'$ is the norm induced by $\langle \cdot, \cdot \rangle'$ and we have used that $f \geq 1$.

Theorem 4.1. *Let (4.1) hold for some $\kappa > 0$ and $K_0 \in \mathbb{R}$, and let P_t be the Neumann semigroup generated by $L = \Delta + Z$ on M . Then for any $f \in C_b^\infty(M)$ such that $\inf f = 1$, $N \log f|_{\partial M} \geq \kappa$ and $K < \infty$, where K is fixed by (4.2) and (4), all assertions in Theorem 1.1 and Corollaries 1.2 and 1.3 hold for constant functions $K, \delta := \sup f^{-1} - \inf f^{-1}$, and $\lambda := \inf f^{-1}$.*

Remark 4.1 A simple choice of f in Theorem 4.1 is $f = \phi \circ \rho_\partial$, where ρ_p is the Riemannian distance to the boundary which is smooth on $\{\rho_\partial \leq r_T\}$ for some $r_T > 0$ provided the injectivity radius of the boundary is positive, and $f \in C_b^\infty([0, \infty))$ is such that $f(0) = 1$, $f'(0) = \kappa$ and $f(r) = f(r_T)$ for $r \geq r_T$. In general, f is taken account to r_T and bounds of the second fundamental form and sectional curvatures, see e.g. [24, 27] for details. With specific choices of f Theorem 4.1 provides explicit Harnack type inequalities, heat kernels estimates and criteria on contractivity properties for the Neumann semigroup on manifolds with non-convex boundary.

References

- [1] S. Aida, *Uniform positivity improving property, Sobolev inequalities, and spectral gaps*, J. Funct. Anal. 158(1998), 152–185.
- [2] S. Aida and H. Kawabi, *Short time asymptotics of a certain infinite dimensional diffusion process*, Stochastic analysis and related topics, VII (Kusadasi, 1998), Progr. Probab., vol. 48, Birkhäuser Boston, Boston, 2001, pp. 77–124.

- [3] S. Aida and T. Zhang, *On the small time asymptotics of diffusion processes on path groups*, Potential Anal. 16(2002), 67–78.
- [4] M. Arnaudon, A. Thalmaier, F.-Y. Wang, *Harnack inequality and heat kernel estimates on manifolds with curvature unbounded below*, Bull. Sci. Math. 130(2006), 223–233.
- [5] D. Bakry, M. Emery, *Hypercontractivité de semi-groupes de diffusion*, C. R. Acad. Sci. Paris. Sér. I Math. 299(1984), 775–778.
- [6] V. I. Bogachev, N. V. Krylov, M. Röckner, *On regularity of transition probabilities and invariant measures of singular diffusions under minimal conditions*, Comm. Partial Differential Equations 26 (2001), 2037–2080.
- [7] X. Chen, F.-Y. Wang, *Optimal integrability condition for the log-Sobolev inequality*, Quart. J. Math. (Oxford) 58(2007), 17–22.
- [8] G. Da Prato, M. Röckner, F.-Y. Wang, *Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups*, J. Funct. Anal. 257(2009), 992–1017.
- [9] G. Da Prato and J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, Cambridge Univ. Press, 1992.
- [10] M. Emery, *Stochastic calculus in manifolds*, Springer-Verlag, Berlin, 1989.
- [11] A. Es-Sarhir, M.-K. von Renesse, M. Scheutzew, *Harnack inequality for functional SDEs with bounded memory*, arXiv:0910.4536.
- [12] S. Fang, T.-S. Zhang, *A study of a class of stochastic differential equations with non-Lipschitzian coefficients*, Probab. Theory Related Fields 132 (2005), 356–390.
- [13] M. Hino, *Exponential decay of positivity preserving semigroups on L^p* , Osaka J. Math. 37 (2000), 603–624.
- [14] N. Ikeda, S. Watanabe, *Stochastic Differential Equations and Diffusion Processes (Second Edition)*, North-Holland, Amsterdam, 1989.
- [15] W. Liu, *Fine properties of stochastic evolution equations and their applications*, Doctor-Thesis, Bielefeld University, 2009.
- [16] W. Liu and F.-Y. Wang, *Harnack inequality and strong Feller property for stochastic fast-diffusion equations*, J. Math. Anal. Appl. 342(2008), 651–662.
- [17] S.-X. Ouyang, *Harnack inequalities and applications for stochastic equations*, Ph.D. thesis, Bielefeld University, 2009.

- [18] M. Röckner and F.-Y. Wang, *Supercontractivity and ultracontractivity for (non-symmetric) diffusion semigroups on manifolds*, Forum Math. 15(2003), no. 6, 893–921.
- [19] M. Röckner, F.-Y. Wang, *Harnack and functional inequalities for generalized Mehler semigroups*, J. Funct. Anal. 203(2003), 237–261.
- [20] M. Röckner and F.-Y. Wang, *Log-Harnack inequality for stochastic differential equations in Hilbert spaces and its consequences*, to appear in Inf. Dim. Anal. Quant. Proba. Relat. Top.
- [21] A. Thalmaier, F.-Y. Wang, *Gradient estimates for harmonic functions on regular domains in Riemannian manifolds*, J. Funct. Anal. 155:1(1998), 109–124.
- [22] F.-Y. Wang, *Logarithmic Sobolev inequalities on noncompact Riemannian manifolds*, Probab. Theory Related Fields 109(1997), 417–424.
- [23] F.-Y. Wang, *Equivalence of dimension-free Harnack inequality and curvature condition*, Integral Equations Operator Theory 48(2004), no. 4, 547–552.
- [24] F.-Y. Wang, *Harnack inequality and applications for stochastic generalized porous media equations*, Ann. Probab. 35(2007), 1333–1350.
- [25] F.-Y. Wang, *Estimates of the first Neumann eigenvalue and the log-Sobolev constant on nonconvex manifolds*, Math. Nachr. 280(2007), 1431–1439.
- [26] F.-Y. Wang *Harnack inequalities on manifolds with boundary and applications*, to appear in JMPA. arXiv:0908.2888
- [27] F.-Y. Wang *Transportation-cost inequalities on path space over manifolds with boundary*, arXiv:0908.2891
- [28] T. Yamada, S. Watanabe, *On the uniqueness of solutions of stochastic differential equations*, J. Math. Kyoto Univ. 11(1971), 155–167.
- [29] T.-S. Zhang, *White noise driven SPDEs with reflection: strong Feller properties and Harnack inequalities*, 2008 preprint.