

Angular momentum-mass inequality for axisymmetric black holes

Sergio Dain

Facultad de Matemática, Astronomía y Física,
Universidad Nacional de Córdoba,
Ciudad Universitaria, (5000) Córdoba,
Argentina

January 21, 2021

Abstract

In these notes we describe recent results concerning the inequality $m \geq \sqrt{|J|}$ for axially symmetric black holes.

1 Introduction

The following conjectures constitute the essence of the current standard picture of the gravitational collapse: i) Gravitational collapse results in a black hole (weak cosmic censorship) ii) The spacetime settles down to a stationary final state. If we further assume that at some finite time all the matter fields have fallen into the black hole and hence the exterior region is pure vacuum (for simplicity we discard electromagnetic fields in the exterior), then the black hole uniqueness theorem implies that the final state should be the Kerr black hole. The Kerr black hole is uniquely characterized by its mass m_0 and angular momentum J_0 . These quantities satisfy the following remarkable inequality

$$\sqrt{|J_0|} \leq m_0. \quad (1)$$

From Newtonian considerations, we can interpret this inequality as follows[14]: in a collapse the gravitational attraction ($\approx m_0^2/r^2$) at the horizon ($r \approx m_0$) dominates over the centrifugal repulsive forces ($\approx J_0^2/m_0r^3$).

If the initial conditions for a collapse violate (1) then the extra angular momentum should be radiated away in gravitational waves. However, in an

axially symmetric spacetime the angular momentum is a conserved quantity (the Komar integral of the Killing vector, see, for example, [16]). In this case angular momentum cannot be radiated: the angular momentum J of the initial conditions must be equal to the final one J_0 . On the other hand, the mass of the initial conditions m satisfies $m \geq m_0$ because gravitational radiation carries positive energy. Then, from inequality (1) we obtain

$$\sqrt{|J|} \leq m. \tag{2}$$

More precisely, i)-ii) imply that a complete, vacuum, axisymmetric, asymptotically flat data should satisfy inequality (2), where m and J are the mass and angular momentum of the data. Moreover, the equality in (2) should imply that the data are a slice of extreme Kerr. This is a similar argument to the one used by Penrose [13] to obtain the inequality between mass and the area of the horizon on the initial data. As in the case of Penrose inequality, a counter example of (2) will imply that either i) or ii) is not true. Conversely a proof of (2) gives indirect evidence of the validity of i)-ii), since it is very hard to understand why this highly nontrivial inequality should hold unless i)-ii) can be thought of as providing the underlying physical reason behind it (see the discussion in [15]).

Inequality (2) is a property of the spacetime and not only of the data, since both quantities m and J are independent of the slicing. It is in fact a property of axisymmetric, vacuum, black holes space-times, because a non zero J (in vacuum) implies a non trivial topology on the data and this is expected to signal the presence of a black hole. The physical interpretation of (2) is the following: if we have a stationary vacuum black hole (i.e. Kerr) and add to it axisymmetric gravitational waves, then the spacetime will still have a (non-stationary) black hole, these waves will only increase the mass and not the angular momentum of the spacetime because they are axially symmetric. Since inequality (1) is satisfied for Kerr we get (2).

In this note, we review some recent results (see [10], [5], [9], [6], [7], [4]) in which inequality (2) is proved for one black hole and describe the open problems for the other cases.

2 Variational principle for the mass

Inequality (2) suggests the following variational principle:

The extreme Kerr initial data are the absolute minimum of the mass among all axisymmetric, vacuum, asymptotically flat and complete initial data with fixed angular momentum.

However, it is important to note that for two related inequalities, the positive mass theorem and the Penrose inequality, a variational formulation was not successful. In the case of the positive mass theorem only a local version was proved using a variational principle [2].

The key difference in the present case is axial symmetry. As we will see, in that case it is possible to write the mass (in an appropriate gauge) as a positive definite integral on a spacelike hypersurface. The reason for this particular behavior of the mass is the following. In the presence of a symmetry, vacuum Einstein equations can be reduced à la Kaluza-Klein to a system on a 3-dimensional manifold where it takes the form of 3-dimensional Einstein equations coupled to a matter source. Since in 3-dimension there is no radiation (the Weyl tensor is zero), this source represents the true gravitational degree of freedom that have descended from 4-dimensions to appear as “matter” in 3-dimension. Since all the energy is produced by these effective matter sources, one would expect in that, as in other field theories, the total energy of the system can be expressed as a positive definite integral over them. This was in fact proved by Brill [1] in some restricted cases and then generalized in [12] [10][3]. Using this formula and with the extra assumption that the data are maximal, the variational principle can be formulated in a very simple form [7].

To write the mass formula for axially symmetric spacetimes we follow [8]. Consider a vacuum solution of Einstein’s equations, i.e., a four dimensional manifold \mathcal{V} with metric g_{ab} for which the Ricci tensor ${}^{(4)}\mathcal{R}_{ab}$ vanishes. Suppose, in addition, that there exists a spacetime Killing vector η^a . We define the norm and the twist of η^a , respectively, by

$$\lambda^2 = \eta^a \eta^b g_{ab}, \quad \omega_a = \epsilon_{abcd} \eta^b \bar{\nabla}^c \eta^d, \quad (3)$$

where $\bar{\nabla}_a$ is the connection and ϵ_{abcd} the volume element with respect to g_{ab} . Assuming that the manifold is simply connected and using ${}^{(4)}\mathcal{R}_{ab} = 0$ it is possible to prove that ω_a is the gradient of a scalar field ω

$$\omega_a = \bar{\nabla}_a \omega. \quad (4)$$

In our case the Killing field will be spacelike, i.e. $\lambda \geq 0$.

As we mention above, in the presence of a Killing field, there exists a well known procedure to reduce the field equations [11]. Let \mathcal{N} denote the collection of all trajectories of η^a , and assume that it is a differential 3-manifold. We define the Lorentzian metric h_{ab} on \mathcal{N} by

$$g_{ab} = h_{ab} + \frac{\eta_a \eta_b}{\lambda^2}. \quad (5)$$

Four dimensional Einstein vacuum equation are equivalent to Einstein equations in three dimension on \mathcal{N} coupled to effective matter fields determined by λ and ω . We make a $2 + 1$ decomposition of these equations. Let n^a be the unit normal vector orthogonal to a spacelike, 2-dimensional slice S . The intrinsic metric on S is denoted by q_{AB} and the trace free part of the second fundamental form of the slice is denoted by k_{AB} . On (\mathcal{N}, h) we fix a gauge: the maximal-isothermal gauge (see [8] for details) and the corresponding coordinates system (t, ρ, z) . It is convenient to define the function σ by

$$\lambda = \rho e^{\sigma/2}. \quad (6)$$

In this gauge the mass can be written in the following form

$$m = \frac{1}{16} \int_S \left(2k^{AB}k_{AB} + 3\frac{\lambda'^2}{\lambda^2} + \frac{\omega'^2}{\lambda^4} + |D\sigma|^2 + \frac{|D\omega|^2}{\lambda^4} \right) \rho dV_q. \quad (7)$$

where $dV_q = e^{2u}d\rho dz$ denote the volume element with respect to q_{ab} , D is the covariant derivative with respect to q_{AB} with $|D\sigma|^2 = D^A\sigma D_A\sigma$, and the prime denotes directional derivative with respect to n^a , that is

$$\lambda' = n^a \bar{\nabla}_a \lambda. \quad (8)$$

This is essentially a derivative with respect to t . Note that all the terms in the integrand of (7) are positive definite. The first three terms contain the dynamical part of the data, they vanish for stationary solutions, in particular for the Kerr solution. The last two terms, contain the stationary part of the fields. It is important to note that the integral of these terms does not depends on the metric q_{AB} . In effect, the integral of these terms can be written as

$$\mathcal{M}(\sigma, \omega) = \frac{1}{16} \int_{-\infty}^{\infty} dz \int_0^{\infty} d\rho (|\partial\sigma|^2 + \rho^{-4} e^{-2\sigma} |\partial\omega|^2) \rho, \quad (9)$$

where ∂ denotes partial derivatives with respect to (ρ, z) . The integral (9) depends only on σ and ω . Since we have

$$m \geq \mathcal{M}, \quad (10)$$

to find the minimum of m is equivalent as to find the minimums of \mathcal{M} .

In order to write the variational principle, it only remains to discuss the boundary conditions. Physically, we want to prescribe boundary conditions such that the total angular momentum is fixed. The information of the angular momentum is determined by the value of the twist potential ω at




Figure 1: N asymptotic ends

the axis $\rho = 0$ (see [7]). To include more than one black hole, we prescribe the following topology. Let i_k be a finite collection of points located at the axis $\rho = 0$. Define the intervals I_k , $0 \leq k \leq N - 1$, to be the open sets in the axis between i_k and i_{k-1} , we also define I_0 and I_N as $z < i_0$ and $z > i_N$ respectively. See figure 2. Each point i_k will correspond to an asymptotic end of the data, and hence we will say that the data contain N black holes.

To fix the total angular momentum J (where J is an arbitrary constant) of the data is equivalent as to prescribe the following boundary condition for ω (see [10])

$$\omega|_{I_0} = 4J, \quad \omega|_{I_N} = -4J. \quad (11)$$

We want to study the minimums of the functional \mathcal{M} with these boundary conditions.

We are now in position to write the precise form of the variational principle.

Conjecture 2.1. *Let σ, ω be arbitrary functions such that ω satisfies the boundary condition (11). Then we have*

$$\mathcal{M}(\sigma, \omega) \geq \sqrt{|J|}. \quad (12)$$

Moreover, the equality implies that σ, ω are given by the extreme Kerr solution.

This conjecture was proved for the case $N = 1$ in [10]. This result was extended in [4] to include more generic data.

The conjecture is open for the case $N \geq 2$. For this case, the variational problem is fixed if we impose the boundary condition

$$\omega|_{I_i} = 4J_i, \tag{13}$$

with $0 < i < N$, for arbitrary constants J_i . Note however, that conjecture 2.1 is independent of the values J_i .

Remarkably, in [4] it is proved that the variational problem has a solution (i.e. a minimum) for arbitrary N , but the value of \mathcal{M} for this solution is not known. In order to prove the conjecture for $N \geq 2$, one need to compute a lower bound for this quantity. This problem is related with the uniqueness of the Kerr black hole with degenerate and disconnected horizons.

References

- [1] D. Brill. On the positive definite mass of the Bondi-Weber-Wheeler time-symmetric gravitational waves. *Ann. Phys.*, 7:466–483, 1959.
- [2] Y. Choquet-Bruhat and J. E. Marsden. Solution of the local mass problem in general relativity. *Commun. Math. Phys.*, 51(3):283–296, 1976.
- [3] P. T. Chruściel. Mass and angular-momentum inequalities for axisymmetric initial data sets I. Positivity of mass. 2007, arXiv:0710.3680 [gr-qc].
- [4] P. T. Chruściel, Y. Li, and G. Weinstein. Mass and angular-momentum inequalities for axisymmetric initial data sets. II. Angular-momentum. 2007, arXiv:0712.4064 [gr-qc].
- [5] S. Dain. Angular momentum-mass inequality for axisymmetric black holes. *Phys. Rev. Lett.*, 96:101101, 2006, gr-qc/0511101.
- [6] S. Dain. Proof of the (local) angular momentum-mass inequality for axisymmetric black holes. *Class. Quantum. Grav.*, 23:6845–6855, 2006, gr-qc/0511087.
- [7] S. Dain. A variational principle for stationary, axisymmetric solutions of einstein’s equations. *Class. Quantum. Grav.*, 23:6857–6871, 2006, gr-qc/0508061.
- [8] S. Dain. Axisymmetric evolution of Einstein equations and mass conservation. *Class. Quantum. Grav.*, 25:145021, 2008, 0804.2679.

- [9] S. Dain. The inequality between mass and angular momentum for axially symmetric black holes. *International Journal of Modern Physics D*, 17(3-4):519–523, 2008, arXiv:0707.3118 [gr-qc].
- [10] S. Dain. Proof of the angular momentum-mass inequality for axisymmetric black holes. *J. Differential Geometry*, 79(1):33–67, 2008, gr-qc/0606105.
- [11] R. Geroch. A method for generating solutions of Einstein’s equations. *J. Math. Phys.*, 12(6):918–924, 1971.
- [12] G. W. Gibbons and G. Holzegel. The positive mass and isoperimetric inequalities for axisymmetric black holes in four and five dimensions. *Class. Quant. Grav.*, 23:6459–6478, 2006, gr-qc/0606116.
- [13] R. Penrose. Gravitational collapse: The role of general relativity. *Riv. Nuovo Cimento*, 1:252–276, 1969.
- [14] R. Wald. Final states of gravitational collapse. *Phys. Rev. Lett.*, 26(26):1653–1655, 1971.
- [15] R. Wald. Gravitational collapse and cosmic censorship. In B. R. Iyer and B. Bhawal, editors, *Black Holes, Gravitational Radiation and the Universe*, volume 100 of *Fundamental Theories of Physics*, pages 69–85. Kluwer Academic, Dordrecht, 1999, gr-qc/9710068.
- [16] R. M. Wald. *General Relativity*. The University of Chicago Press, Chicago, 1984.