

MONOTONICITY OF TOPOLOGICAL ENTROPY UNDER NORMALISED RICCI FLOW

DANIEL J. THOMPSON

ABSTRACT. We prove that the topological entropy of the geodesic flow for a compact Riemannian manifold (M, g) decreases as the metric g evolves under the normalised Ricci flow provided that M admits a metric of constant negative sectional curvature, and g is in a sufficiently small C^2 neighbourhood of the constant curvature metric. More generally, the same phenomenon occurs if g satisfies a certain negative curvature pinching condition, where the pinching constant depends on both the dimension and the diameter of (M, g) . This provides an affirmative answer to an open question posed in Manning's paper 'The volume entropy of a surface decreases along the Ricci flow' [*Ergodic Theory Dynam. Systems*, 24:171-76, 2004].

1. INTRODUCTION

We consider the normalised Ricci flow (NRF) $\{g_t\}$ for an initial metric g_0 on an n dimensional compact manifold M which admits a metric of constant negative sectional curvature g_∞ . We let $h_{top}(g)$ denote the topological entropy of the geodesic flow for a metric g and we consider the map

$$t \mapsto h_{top}(g_t).$$

If the initial metric g_0 is close to g_∞ in the C^2 topology, or close to g_∞ in the sense that g_0 satisfies a suitable curvature pinching assumption, we show that the map $t \mapsto h_{top}(g_t)$ is decreasing. This result is an affirmative answer to a question of Manning [13], who proved that the map $t \mapsto h_{top}(g_t)$ is decreasing for a surface of negative Gauss curvature, and asked five natural questions about the relationship between various dynamical quantities and Ricci flow. To date, the other questions remain open, although some related progress was made by Jane [8].

The conclusion of our results, ' $t \mapsto h_{top}(g_t)$ is decreasing for $t \in [0, \infty)$ ' makes sense because the Besson-Courtois-Gallot theorem [1] states that topological entropy is (globally) minimised at g_∞ , and a result of Ye [18] states that if g_0 is in a neighbourhood of g_∞ , or satisfies certain curvature pinching conditions, then the NRF of g_0 exists for all time and converges to g_∞ . Thus, one can hope that topological entropy is decreasing for initial metrics for which Ye's convergence result applies, and we show that this is the case. Our proof relies on both of the results mentioned above, the Katok-Knieper-Weiss formula [11] for the derivative of topological entropy along a path of metrics, and a theorem of Freire-Mañé [5] about solutions of the Riccati equation for manifolds without conjugate points.

We give some dynamical and geometric interpretations of our result. The quantity $h_{top}(g_t)$ can be interpreted as a measure of the exponential complexity of the orbit structure of the geodesic flow on (M, g_t) . For a precise description of this interpretation in the case of a discrete time dynamical system see [17, page 172].

In light of this, we can interpret the monotonicity of topological entropy along the normalised Ricci flow as a decrease in the complexity of the orbit structure of the geodesic flow. This matches our intuition that the normalised Ricci flow is ‘smoothing out’ the manifold. Furthermore, Bowen [2] showed that for g with negative sectional curvatures,

$$(1.1) \quad h_{top}(g) = \lim_{t \rightarrow \infty} \frac{1}{t} \log P(t),$$

where $P(t)$ is the number of closed geodesics of period at most t . Thus, when our result applies, it says that the NRF decreases the exponential growth rate of the number of closed geodesics. For metrics with negative sectional curvature (or more generally, metrics with no conjugate points), $h_{top}(g)$ can be formulated as

$$(1.2) \quad h_{top}(g) = \lim_{R \rightarrow \infty} \frac{1}{R} \log \text{Vol}(B(x, R)),$$

where $B(x, R)$ is the ball of radius R in the universal cover of (M, g) , and the choice of x is arbitrary [12], [5]. This quantity is often called the volume entropy. It is interesting to compare this definition with that of the asymptotic volume ratio, where one looks for a polynomial growth rate for $\text{Vol}(B(x, R))$. Results on the evolution of asymptotic volume ratio under Ricci flow for non-negative curvature manifolds are well known in the Ricci flow literature [12], [14]. The volume entropy can be thought of as a negative curvature analogue of the asymptotic volume ratio, so this provides further evidence that studying the evolution of $h_{top}(g)$ under a geometric flow is a natural thing to do.

In §2, we state our results formally and prove them. In §3, we make some remarks on the results and the proofs.

2. RESULTS

We consider the family of metrics $\{g_t\}$ defined by an initial metric g_0 and the normalised Ricci flow (NRF) equation

$$(2.1) \quad \frac{\partial g}{\partial t} = -2\text{Ric} + \frac{2}{n} \frac{\int S d\text{Vol}}{\text{Vol}(M)} g,$$

where Ric is the Ricci curvature and S is the scalar curvature [16], [7], [3]. Of course, Ric , S and $d\text{Vol}$ depend on the metric, and we sometimes write, for example, S_g when we want to highlight this dependence. We call $\{g_t\}$ the NRF for g_0 . The second term in the NRF equation (2.1) serves to fix the volume of the metric as it evolves. We write M_t for the manifold (M, g_t) and SM_t for the unit tangent bundle of (M, g_t) . We consider the map $t \mapsto h_{top}(g_t)$, where $h_{top}(g)$ is defined by (1.1), or equivalently (1.2).

We first formulate our theorem for initial metrics in a C^2 neighbourhood of a constant negative sectional curvature metric g_∞ . To make the results easier to state, we make a standing assumption that the initial metric has the same volume as g_∞ and is not isometric to g_∞ . A theorem of Ye [18] says that for these initial metrics, the normalised Ricci flow exists for all time and converges to g_∞ .

Theorem 2.1. *Suppose $n \geq 3$ and (M^n, g_∞) is a compact Riemannian manifold with constant negative sectional curvature. There is a C^2 neighbourhood U of g_∞ where Ye’s convergence result applies and such that if $g_0 \in U$, then the map $t \mapsto h_{top}(g_t)$ is decreasing for all $t \in [0, \infty)$.*

Our proof follows the same strategy that Manning used in the case of surfaces, and is based on the Katok-Kneiper-Weiss formula [11] for the derivative of topological entropy along a path of metrics of negative sectional curvature (the formula applies in all dimensions). Like Manning's result, the proof is not long but every line is an application of a major theorem. We require that the topological entropy is minimised at g_∞ . For $n \geq 3$, this is the celebrated theorem of Besson-Courtois-Gallot [1]. The result was originally conjectured by Katok, who proved it for surfaces [9]. Another ingredient is a theorem of Freire-Mané [5] which relates $\int_{SM} \text{Ric}(v, v) d\mu(v)$ and h_μ for any probability measure μ on SM which is invariant under the geodesic flow (this is a corollary of the main result of [5] which relates the ergodic average of certain solutions of the matrix Riccati equation to Lyapunov exponents of invariant measures). The only result that Manning uses which does not have a higher dimensional analogue that we can use is the Gauss-Bonnet theorem. We get round the fact that the Gauss-Bonnet theorem is not available by making sure that the total scalar curvature stays controlled along the flow. This follows from estimates of Ye [18]. First, we establish a result which is more general but less natural than theorem 2.1. Let $\mathcal{S}(g_t) = \int_{M_t} S d\text{Vol}$ denote the total scalar curvature functional. In the following theorem, unlike the situation in theorem 2.1, there is no reason that the NRF should converge to g_∞ .

Theorem 2.2. *Let $\{g_t\}$ be a NRF on a compact Riemannian manifold which admits a metric of constant negative sectional curvature g_∞ . For a fixed time t , suppose g_t has everywhere negative sectional curvature and that $-\mathcal{S}(g_t) \leq -(n-1)\mathcal{S}(g_\infty)$, then*

$$\frac{d}{ds}(h_{top}(g_s))|_{s=t} < 0.$$

Proof. The Katok-Kneiper-Weiss formula [11] says that

$$\frac{d}{ds}(h_{top}(g_s))|_{s=t} = -\frac{h_{top}(g_t)}{2} \int_{SM_t} (\partial/\partial s)|_{s=t} g_s(v, v) d\mu_t(v),$$

where $t \mapsto g_t$ is a C^2 path of Riemannian metrics with everywhere negative sectional curvature. Here, μ_t is the Bowen-Margulis measure for M_t , which is the unique probability measure that satisfies $h_{\mu_t} = h_{top}(g_t)$, where h_{μ_t} is the measure theoretic entropy of μ_t with respect to the geodesic flow on M_t . On substituting the NRF equation into the Katok-Kneiper-Weiss formula, the problem reduces to showing that

$$\int_{SM_t} -\text{Ric} d\mu_t - \frac{1}{n\text{Vol}(M)} \int_{M_t} -S d\text{Vol} > 0.$$

Corollary II.1 of Freire and Mané [5] states that for a metric without conjugate points on M and any probability measure ν on SM that is invariant under the geodesic flow, then

$$\int_{SM} -\text{Ric} d\nu \geq h_\nu^2,$$

where h_ν is the measure-theoretic entropy of ν . (In [5], an additional factor of $(n-1)^{1/2}$ is present, but as pointed out by Osserman and Sarnak [15], this is a mistake.) Applying this to the Bowen-Margulis measure μ_t , we have

$$\int_{SM_t} -\text{Ric} d\mu_t \geq h_{\mu_t}^2 > h_{\mu_\infty}^2,$$

where the second inequality is an application of the Besson-Courtois-Gallot theorem [1]. Thus,

$$\int_{SM} -\text{Ric}d\mu_t - h_{\mu_\infty}^2 > 0.$$

At g_∞ , suppose the constant curvature is $-C$. Then $h_{\mu_\infty}^2 = (n-1)^2C$ and $S_{g_\infty} = -n(n-1)C$. Therefore,

$$h_{\mu_\infty}^2 = \frac{(n-1)}{n\text{Vol}(M)} \int_{M_\infty} -S_{g_\infty} d\text{Vol}.$$

We have assumed that $-\mathcal{S}(g_t) \leq -(n-1)\mathcal{S}(g_\infty)$ and thus

$$\frac{1}{n\text{Vol}(M)} \int_{M_t} -S_{g_t} d\text{Vol} \leq \frac{(n-1)}{n\text{Vol}(M)} \int_{M_\infty} -S_{g_\infty} d\text{Vol} = h_{\mu_\infty}^2.$$

Therefore,

$$\int_{SM_t} -\text{Ric}d\mu_t - \frac{1}{n\text{Vol}(M)} \int_{M_t} -S d\text{Vol} \geq \int_{SM_t} -\text{Ric}d\mu_t - h_{\mu_\infty}^2 > 0,$$

and thus $\frac{d}{ds}(h_{top}(g_s))|_{s=t} < 0$. \square

Now to prove theorem 2.1, it suffices to show that for an initial metric g_0 which is sufficiently close to g_∞ that we have enough control on the total scalar curvature along the NRF to ensure that theorem 2.2 applies for all time. Since g_t converges to g_∞ and the total scalar curvature functional is continuous, we already have control over $S(g_t)$ for sufficiently small t and sufficiently large t . The required hard analysis to establish control over $S(g_t)$ for all time is present in Ye's paper [18].

Proof of theorem 2.1. Ye states explicitly that along the NRF for an initial metric in a C^2 neighbourhood of g_∞ , the sectional curvatures remain negative and converge exponentially. The proof that the total scalar curvature stays controlled along the flow follows from the same estimates. We sketch the argument. Let $\overline{\mathcal{S}}(g) = \text{Vol}_g(M)^{-1}\mathcal{S}(g)$. The evolution for $\mathcal{S}(g)$ along the NRF is given by

$$\frac{d}{dt}\overline{\mathcal{S}}(g) = \frac{1}{\text{Vol}(M)} \left(2 \int |T|^2 + \left(\frac{2}{n} - 1\right) \int (\delta S)^2 \right),$$

where $T_g = \text{Ric}_g - S_g g/n$, $|T|^2$ is the function $\langle T, T \rangle$ and $\delta S_g = S_g - \overline{\mathcal{S}}(g)$, and thus

$$\frac{d}{dt}\overline{\mathcal{S}}(g) \geq \text{Vol}(M)^{-1} \left(\frac{2}{n} - 1\right) \int (\delta S)^2.$$

Estimate (1.33) of [18] gives for $t \in [1, \infty)$

$$\|(\delta S)^2\|_{C^0(M_t)} \leq C_1 \exp(-at)$$

for a suitable $a > 0$, where C_1 depends on $\int_{M_0} |T|^2$, and $\|(\delta S)^2\|_{C^0(M_t)}$ is the largest absolute value of $(\delta S_{g_t})^2$. We can make C_1 arbitrarily small by making $\int_{M_0} |T|^2$ arbitrarily small, and this follows from choosing the neighbourhood U small enough. Estimate (2.8) of [18] shows that for $t \in [0, 1]$,

$$\|(\delta S)^2\|_{C^0(M_t)} \leq \|(\delta S)^2\|_{C^0(M_0)} + \int_0^t \|T^2\|_{C^0(M_t)}.$$

We can control the integral term using formula (2.5) of Ye, which gives C^0 pinching of the concircular curvature tensor, and from which smallness of the integral follows.

Again, by choosing U sufficiently small, we can make the integral as small as we like. A computation involving the above inequalities gives for all $t \in [0, \infty)$,

$$\bar{S}(g_t) \geq C_2 \exp(-at) - C_3,$$

for certain $C_2 > 0$, $C_3 > -S(g_\infty)$ and by making the neighbourhood U sufficiently small, we can choose C_2 as small as we like and C_3 as close to $-S(g_\infty)$ as we like. It follows that if the neighbourhood U is sufficiently small, and g_t is the NRF starting at $g_0 \in U$, then $\int_{M_t} -S_{g_t} d\text{Vol} \leq (n-1) \int_{M_t} -S_{g_\infty} d\text{Vol}$, and thus theorem 2.2 applies to every metric along the flow. \square

We extend the result of theorem 2.1 to initial metrics which satisfy certain curvature pinching conditions. The (rather stringent) curvature pinching assumptions are sufficient for Ye's convergence results to apply. We obtain the following two theorems.

Theorem 2.3. *For given $n \geq 4$ and $V > 0$, consider an n -dimensional compact Riemannian manifold (M, g_0) with $\text{Vol}_{g_0}(M) \leq V$. There exists $\epsilon > 0$, which depends only on n and V such that if all sectional curvatures for g_0 are between -1 and $-1 - \epsilon$, then the NRF $\{g_t\}$ with initial metric g_0 exists for all time, converges to a metric of constant sectional curvature and the map $t \mapsto h_{\text{top}}(g_t)$ is decreasing for all $t \in [0, \infty)$.*

Theorem 2.4. *For given $n \geq 3$ and $d > 0$, consider the space of n -dimensional compact Riemannian manifolds (M, g_0) with diameter at most d . There exists $\epsilon > 0$, which depends only on n and d such that if all sectional curvatures for g_0 are between -1 and $-1 - \epsilon$, then the NRF $\{g_t\}$ with initial metric g_0 exists for all time, converges to a metric of constant sectional curvatures and the map $t \mapsto h_{\text{top}}(g_t)$ is decreasing for all $t \in [0, \infty)$.*

Proof of theorems 2.3 and 2.4. Theorem 2 of [18] gives the existence and convergence part of these theorems, and also guarantees that the sectional curvatures remain negative along the flow. We need to ensure that if ϵ is chosen small enough, then we have enough control on $S(g_t)$ along the flow so that theorem 2.2 applies at all times. The argument sketched in the proof of theorem 2.1 goes through by replacing ' U small enough' with ' ϵ small enough'. \square

3. SOME REMARKS

3.1. On theorem 2.1. The result still holds under the weaker hypothesis that g_∞ is a locally symmetric metric of negative curvature (see [15] for the classification of such metrics). Such metrics are Einstein, and Ye's convergence result still apply. As before, the entropy is minimised at g_∞ by the Besson-Courtois-Gallot theorem, and the proof goes through unchanged.

3.2. On theorem 2.2. We did not state theorem 2.2 under the most general hypotheses possible. We state a different version here. Let $\{g_t\}$ be a NRF on a compact Riemannian manifold which admits a locally symmetric metric of negative curvature g_∞ . For a fixed time t , suppose g_t is Anosov (i.e. the geodesic flow has the Anosov property [10]) and that $-\mathcal{S}(g_t) < -(n-1)\mathcal{S}(g_\infty)$, then the map $s \mapsto h_{\text{top}}(g_s)$ is decreasing for a small time interval containing t . The proof is like that of theorem 2.2, combined with the fact that, by structural stability [10, page 572], the hypotheses are an open condition.

3.3. On theorems 2.3 and 2.4. Theorem 2.3 follows from theorem 2.4 by work of Gromov [6], who shows that for $n \geq 4$, bounds on diameter imply bounds on volume. We note that as $d \rightarrow \infty$, $\epsilon \rightarrow 0$. In [4], the authors offer a nice clarification of when the hypotheses of Ye's convergence result are satisfied, and when they do not apply.

3.4. Evolution of topological entropy under the unnormalised Ricci flow.

We first point out that the question of how $h_{top}(g_t)$ behaves when g_t evolves by the Ricci flow (RF) equation $\frac{\partial g}{\partial t} = -2\text{Ric}$ is not a good one. This is because $h_{top}(g_t)$ is not scale invariant. For example, it decreases under a homothetic expansion of the manifold. Since the Ricci flow increases volume when the curvature is everywhere negative, we would expect that it is easier to show that $h_{top}(g_t)$ is decreasing under RF than under NRF. This is the case, because as long as Ric is negative, it follows directly from the Katok-Kneiper-Weiss formula that

$$\frac{d}{ds}(h_{top}(g_s))|_{s=t} = h_{top}(g_t) \int_{SM_t} \text{Ric}(v, v) d\mu_t(v) \leq 0.$$

As discussed in the introduction, we can interpret the monotonicity of topological entropy along the normalised Ricci flow as a decrease in the complexity of the orbit structure of the geodesic flow. A priori, monotonicity of topological entropy along the unnormalised Ricci flow could just be due to the expansion of the volume along the flow, and would not necessarily reveal anything about the complexity of the orbit structure.

REFERENCES

- [1] G. Besson, G. Courtois, and S. Gallot. Minimal entropy and Mostow's rigidity theorem. *Ergodic Theory Dynam. Systems*, 16:623–649, 1996.
- [2] R. Bowen. Periodic orbits for hyperbolic flows. *American J. Math.*, 94:1–30, 1972.
- [3] B. Chow and D. Knopf. *The Ricci Flow: An Introduction*, volume 110 of *Mathematical surveys and monographs*. AMS, 2004.
- [4] F.T. Farrell and P. Ontaneda. A caveat on the convergence of the Ricci flow for pinched negatively curved manifolds. *Asian J. Math.*, 9(3):401–406, 2005.
- [5] A. Freire and R. Mané. On the entropy of geodesic flow in manifolds without conjugate points. *Invent. Math.*, 69(3):375–392, 1982.
- [6] M. Gromov. Manifolds of negative curvature. *J. Differential Geom.*, 3:223–230, 1978.
- [7] R. Hamilton. The formation of singularities in the Ricci flow. *Surveys in Differential Geometry*, 2:7–136, 1995.
- [8] D. Jane. An example of how the Ricci flow can increase topological entropy. *Ergodic Theory Dynam. Systems*, 27(6):1919–1932, 2007.
- [9] A. Katok. Entropy and closed geodesics. *Ergodic Theory Dynam. Systems*, 2:339–365, 1982.
- [10] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1995.
- [11] A. Katok, G. Knieper, and H. Weiss. Formulas for the derivative and critical points of topological entropy for Anosov and geodesic flows. *Comm. Math. Phys.*, 138:19–31, 1991.
- [12] A. Manning. Topological entropy for geodesic flows. *Ann. Math.*, 110:567–573, 1979.
- [13] A. Manning. The volume entropy of a surface decreases along the Ricci flow. *Ergodic Theory Dynam. Systems*, 24:171–176, 2004.
- [14] J. Morgan and G. Tian. *Ricci flow and the Poincaré conjecture*, volume 3 of *Clay mathematics monographs*. AMS, 2007.
- [15] R. Osserman and P. Sarnak. A new curvature invariant and entropy of geodesic flows. *Invent. Math.*, 77(3):455–462, 1984.
- [16] P. Topping. *Lectures on the Ricci flow*, volume 325 of *LMS Lecture note Series*. LMS, 2006.

- [17] P. Walters. *An Introduction to Ergodic Theory*, volume 79 of *Graduate Texts in Mathematics*. Springer, New York, 1982.
- [18] R. Ye. Ricci flow, Einstein metrics and space forms. *Trans. Amer. Math. Soc.*, 338(2):871–896, 1993.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, STATE
COLLEGE, PA 16802, USA

E-mail address: `thompson@math.psu.edu`