

Discrete fractional Calculus and Inequalities

George A. Anastassiou
 Department of Mathematical Sciences
 University of Memphis
 Memphis, TN 38152, U.S.A.
 ganastss@memphis.edu

Abstract

Here we define a Caputo like discrete fractional difference and we compare it to the earlier defined Riemann-Liouville fractional discrete analog. Then we produce discrete fractional Taylor formulae for the first time, and we estimate their remainders. Finally, we derive related discrete fractional Ostrowski, Poincare and Sobolev type inequalities.

2000 Mathematics Subject Classification : Primary: 39A12, 34A25, 26A33, Secondary: 26D15, 26D20;

Key words and phrases: Discrete fractional Calculus, discrete inequalities.

1 Preliminaries

We make

Definition 1 We follow [2], [3], [4].

Let $\nu > 0$. The ν -th fractional sum of f is defined by

$$\Delta^{-\nu} f(t, a) = \frac{1}{\Gamma(\nu)} \sum_{s=a}^{t-\nu} (t-s-1)^{(\nu-1)} f(s).$$

Here f is defined for $s = a \bmod (1)$ and $\Delta^{-\nu} f$ is defined for $t = (a + \nu) \bmod (1)$; in particular $\Delta^{-\nu}$ maps functions defined on \mathbb{N}_a to functions defined on $\mathbb{N}_{a+\nu}$, where $\mathbb{N}_t = \{t, t+1, t+2, \dots\}$.

$$\text{Here } t^{(\nu)} = \frac{\Gamma(t+1)}{\Gamma(t-\nu+1)}.$$

From now in this context for convenience we set $\Delta^{-\nu} f(t, a) = \Delta^{-\nu} f(t)$.

We mention

Theorem 2 ([2]) Let f be a real-valued function defined on \mathbb{N}_a and let $\mu, \nu > 0$. Then

$$\Delta^{-\nu} (\Delta^{-\mu} f(t)) = \Delta^{-(\mu+\nu)} f(t) = \Delta^{-\mu} (\Delta^{-\nu} f(t)), \quad \forall t \in \mathbb{N}_{a+\mu+\nu}.$$

We make

Definition 3 Let $\mu > 0$ and $m - 1 < \mu < m$, where m denotes a positive integer, $m = \lceil \mu \rceil$, $\lceil \cdot \rceil$ ceiling of number. Set $\nu = m - \mu$.

The μ -th fractional Caputo like difference is defined as

$$\Delta_*^\mu f(t) = \Delta^{-\nu} (\Delta^m f(t)) = \frac{1}{\Gamma(\nu)} \sum_{s=a}^{t-\nu} (t-s-1)^{(\nu-1)} (\Delta^m f)(s), \quad \forall t \in \mathbb{N}_{a+\nu}.$$

Here Δ^m is the m -th order forward difference operator

$$(\Delta^m f)(s) = \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} f(s+k).$$

We mention

Theorem 4 ([3]) For $\nu > 0$ and p a positive integer we have

$$\Delta^{-\nu} \Delta^p f(t) = \Delta^p \Delta^{-\nu} f(t) - \sum_{k=0}^{\nu-1} \frac{(t-a)^{(\nu-p+k)}}{\Gamma(\nu+k-p+1)} \Delta^k f(a),$$

where f is defined on \mathbb{N}_a .

Remark 5 Let $\mu > 0$ and $m - 1 < \mu < m$, $m = \lceil \mu \rceil$, where m is a positive integer, $\nu = m - \mu > 0$. Then by Theorem 4 we get

$$\Delta^{-\nu} \Delta^m f(t) = \Delta^m \Delta^{-\nu} f(t) - \sum_{k=0}^{m-1} \frac{(t-a)^{(\nu-m+k)}}{\Gamma(\nu+k-m+1)} \Delta^k f(a),$$

where f is defined on \mathbb{N}_a .

So we have proved

$$\Delta_*^\mu f(t) = \Delta^m \Delta^{-\nu} f(t) - \sum_{k=0}^{m-1} \frac{(t-a)^{(\nu-m+k)}}{\Gamma(\nu+k-m+1)} \Delta^k f(a),$$

that is

$$\Delta^m \Delta^{-\nu} f(t) = \Delta_*^\mu f(t) + \sum_{k=0}^{m-1} \frac{(t-a)^{(\nu-m+k)}}{\Gamma(\nu+k-m+1)} \Delta^k f(a), \quad (1)$$

where f is defined on \mathbb{N}_a .

Definition 6 ([3]) The μ -th fractional Riemann-Liouville type difference is defined by

$$\Delta^\mu f(t) := \Delta^{m-\nu} f(t) := \Delta^m (\Delta^{-\nu} f(t)),$$

where $\mu > 0$, $m-1 < \mu < m$, $\nu = m - \mu > 0$.

Remark 7 Consequently from (1) we get

$$\Delta^\mu f(t) = \Delta_*^\mu f(t) + \sum_{k=0}^{m-1} \frac{(t-a)^{(\nu-m+k)}}{\Gamma(\nu+k-m+1)} \Delta^k f(a), \quad (2)$$

where f is defined on \mathbb{N}_a .

2 Results

We give the following Caputo type fractional Taylor's difference formula.

Theorem 8 For $\mu > 0$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$, it holds:

$$f(t) = \sum_{k=0}^{m-1} \frac{(t-a)^{(k)}}{k!} \Delta^k f(a) + \frac{1}{\Gamma(\mu)} \sum_{s=a+\nu}^{t-\mu} (t-s-1)^{(\mu-1)} \Delta_*^\mu f(s), \quad \forall t \in \mathbb{N}_{a+m}, \quad (3)$$

where f is defined on \mathbb{N}_a with $a \in \mathbb{Z}^+$, $\mathbb{Z}^+ := \{0, 1, 2, \dots\}$.

Proof. Notice that by Definition 3,

$$\Delta_*^\mu f(t) = \Delta^{-\nu} (\Delta^m f(t)) = \Delta^{-(m-\mu)} (\Delta^m f(t)), \quad \forall t \in \mathbb{N}_{a+\nu}.$$

Consequently we get $\Delta^{-\mu} \Delta_*^\mu f(t) = \Delta^{-\mu} \Delta^{-(m-\mu)} (\Delta^m f(t))$ (by Theorem 2) $= \Delta^{-(\mu+(m-\mu))} (\Delta^m f(t)) = \Delta^{-m} (\Delta^m f(t))$, $\forall t \in \mathbb{N}_{a+\nu+\mu}$.

So that

$$\Delta^{-\mu} \Delta_*^\mu f(t) = \Delta^{-m} (\Delta^m f(t)), \quad \forall t \in \mathbb{N}_{a+m}. \quad (4)$$

We notice that

$$(t-s-1)^{(m-1)} = \frac{\Gamma(t-s)}{\Gamma(t-s-m+1)} = (t-s-1)(t-s-2)\dots(t-s-m+1), \quad (5)$$

the falling factorial, here we have $t-s-m+1 > 0$.

Therefore we obtain

$$\Delta^{-m} (\Delta^m f(t)) = \frac{1}{(m-1)!} \sum_{s=a}^{t-m} (t-s-1)^{(m-1)} \Delta^m f(s). \quad (6)$$

By ([1], p. 28, Theorem 1.8.5) the discrete Taylor's formula we get

$$f(t) = \sum_{k=0}^{m-1} \frac{(t-a)^{(k)}}{k!} \Delta^k f(a) + \frac{1}{(m-1)!} \sum_{s=a}^{t-m} (t-s-1)^{(m-1)} \Delta^m f(s), \quad (7)$$

where $t^{(k)} = t(t-1)\dots(t-k+1)$.

From the last we derive

$$f(t) = \sum_{k=0}^{m-1} \frac{(t-a)^{(k)}}{k!} \Delta^k f(a) + \Delta^{-\mu} \Delta_*^\mu f(t), \quad (8)$$

where f is defined on \mathbb{N}_a , $\forall t \in \mathbb{N}_{a+m}$, proving the claim. ■

We make

Remark 9 Here $[a, b]$ denotes the discrete interval $[a, b] = [a, a+1, a+2, \dots, b]$, where $a < b$ and $a, b \in \{0, 1, \dots\}$.

Let $\mu > 0$ be non integer such that $m-1 < \mu < m$, i.e. $m = \lceil \mu \rceil$. Consider a function f defined on $[a, b]$. Then clearly the fractional discrete Taylor's formula (3) is valid only for $t \in [a+m, b]$, $a+m < b$.

We need

Theorem 10 ([3]) Let p be a positive integer and let $\nu > p$. Then

$$\Delta^p (\Delta^{-\nu} f(t)) = \Delta^{-(\nu-p)} f(t). \quad (9)$$

We make

Remark 11 Let $\mu > p$, where $p \in \mathbb{N}$. Then

$$\Delta^p (\Delta^{-\mu} \Delta_*^\mu f(t)) \stackrel{(9)}{=} \Delta^{-(\mu-p)} (\Delta_*^\mu f(t)), \quad \forall t \in \mathbb{N}_{a+m-p}. \quad (10)$$

Also notice that

$$\Delta^p \left(\frac{(t-a)^{(k)}}{k!} \right) = \frac{(t-a)^{(k-p)}}{(k-p)!}, \quad \text{for } k \geq p. \quad (11)$$

By the last we obtain the following discrete Caputo type fractional extended Taylor's formula.

Theorem 12 Let $\mu > p$, $p \in \mathbb{N}$, μ not integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$. Then

$$\Delta^p f(t) = \sum_{k=p}^{m-1} \frac{(t-a)^{(k-p)}}{(k-p)!} \Delta^k f(a) + \frac{1}{\Gamma(\mu-p)} \sum_{s=a+\nu}^{t-\mu+p} (t-s-1)^{(\mu-p-1)} \Delta_*^\mu f(s), \quad (12)$$

$\forall t \in \mathbb{N}_{a+m-p}$, f is defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$.

Note 13 Assuming that f is defined on $[a, b]$, then (12) is valid only for $[a+m-p, b]$, with $a+m-p < b$.

Notice for $p = 0$ applied on (12) we get (3).

We give

Proposition 14 For $\mu > 0$, μ not an integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$, f is defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$; and $\Delta^k f(a) = 0$, for $k = 0, \dots, m - 1$, we get

$$f(t) = \frac{1}{\Gamma(\mu)} \sum_{s=a+\nu}^{t-\mu} (t-s-1)^{(\mu-1)} \Delta_*^\mu f(s), \quad \forall t \in \mathbb{N}_{a+m}. \quad (13)$$

Proof. By (3). ■

Also we present

Proposition 15 Let $\mu > p$, $p \in \mathbb{N}$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$; f is defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Assume that $\Delta^k f(a) = 0$, $k = p, \dots, m - 1$. Then

$$\Delta^p f(t) = \frac{1}{\Gamma(\mu-p)} \sum_{s=a+\nu}^{t-\mu+p} (t-s-1)^{(\mu-p-1)} \Delta_*^\mu f(s), \quad \forall t \in \mathbb{N}_{a+m-p}. \quad (14)$$

Proof. By (12). ■

We make

Remark 16 We want to find

$$\sum_{s=a+\nu}^{t-\mu} (t-s-1)^{(\mu-1)} = \sum_{s=a+\nu}^{t-\mu} \frac{\Gamma(t-s)}{\Gamma(t-s-\mu+1)} = \sum_{s=a+\nu}^{t-\mu-1} \frac{\Gamma(t-s)}{\Gamma(t-s-\mu+1)} + \Gamma(\mu). \quad (15)$$

We notice that

$$\frac{\Gamma(x+1)}{\Gamma(k+1)\Gamma(x-k+1)} = \frac{\Gamma(x+2)}{\Gamma(k+2)\Gamma(x-k+1)} - \frac{\Gamma(x+1)}{\Gamma(k+2)\Gamma(x-k)} \quad (16)$$

with $x > k$, $x, k \in \mathbb{R}$; $k > -1$, $x > -1$.

That is

$$\frac{\Gamma(x+1)}{\Gamma(x-k+1)} = \frac{1}{(k+1)} \left(\frac{\Gamma(x+2)}{\Gamma(x-k+1)} - \frac{\Gamma(x+1)}{\Gamma(x-k)} \right). \quad (17)$$

We find $A := \sum_{s=a+\nu}^{t-\mu-1} \frac{\Gamma(t-s)}{\Gamma(t-s-\mu+1)} =$

(by (17) for $x := t-s-1 \geq \mu > 0$, $k := \mu-1 > -1$, and $x > k$)

$$\begin{aligned} & \frac{1}{\mu} \sum_{s=a+\nu}^{t-\mu-1} \left[\frac{\Gamma(t-s+1)}{\Gamma(t-s+1-\mu)} - \frac{\Gamma(t-s)}{\Gamma(t-s-\mu)} \right] = \\ & \frac{1}{\mu} \left[\left(\frac{\Gamma(t-a-\nu+1)}{\Gamma(t-a-\nu+1-\mu)} - \frac{\Gamma(t-a-\nu)}{\Gamma(t-a-\nu-\mu)} \right) + \left(\frac{\Gamma(t-a-\nu)}{\Gamma(t-a-\nu-\mu)} - \frac{\Gamma(t-a-\nu-1)}{\Gamma(t-a-\nu-1-\mu)} \right) + \right. \\ & \left. \left(\frac{\Gamma(t-a-\nu-1)}{\Gamma(t-a-\nu-1-\mu)} - \frac{\Gamma(t-a-\nu-2)}{\Gamma(t-a-\nu-2-\mu)} \right) + \dots \left(\dots - \frac{\Gamma(\mu+1)}{\Gamma(1)} \right) \right] = \\ & \left[\frac{\Gamma(t-a-\nu+1)}{\mu\Gamma(t-a-\nu+1-\mu)} - \Gamma(\mu) \right]. \end{aligned}$$

That is

$$A = \frac{\Gamma(t-a-\nu+1)}{\mu\Gamma(t-a-\nu+1-\mu)} - \Gamma(\mu). \quad (18)$$

Consequently we found

$$\sum_{s=a+\nu}^{t-\mu} (t-s-1)^{(\mu-1)} = \frac{\Gamma(t-a-\nu+1)}{\mu\Gamma(t-a+1-m)} = \frac{(t-a-\nu)^{(\mu)}}{\mu}. \quad (19)$$

Using (19) we give

Corollary 17 (to Theorem 8) Let $\mu > 0$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$, $t \in \mathbb{N}_{a+m}$, f defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Then

$$\left| f(t) - \sum_{k=0}^{m-1} \frac{(t-a)^{(k)}}{k!} \Delta^k f(a) \right| \leq \frac{(t-a-\nu)^{(\mu)}}{\Gamma(\mu+1)} \cdot \max_{s \in \{a+\nu, a+\nu+1, \dots, t-\mu\}} |\Delta_*^\mu f(s)|. \quad (20)$$

Similarly we get

Corollary 18 (to Theorem 12) Let $\mu > p$, $p \in \mathbb{N}$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$, $t \in \mathbb{N}_{a+m-p}$, f defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Then

$$\left| \Delta^p f(t) - \sum_{k=p}^{m-1} \frac{(t-a)^{(k-p)}}{(k-p)!} \Delta^k f(a) \right| \leq \frac{(t-a-\nu)^{(\mu-p)}}{\Gamma(\mu-p+1)} \cdot \max_{s \in \{a+\nu, \dots, t-\mu+p\}} |\Delta_*^\mu f(s)|. \quad (21)$$

We need

Lemma 19 Let $a > \nu$, $a, \nu > -1$, $a, \nu \in \mathbb{R}$, $a \leq b$. Then

$$\sum_{r=a}^b r^{(\nu)} = \frac{1}{(\nu+1)} \left(\frac{\Gamma(b+2)}{\Gamma(b-\nu+1)} - \frac{\Gamma(a+1)}{\Gamma(a-\nu)} \right) = \left(\frac{(b+1)^{(\nu+1)} - a^{(\nu+1)}}{\nu+1} \right). \quad (22)$$

Proof. We have

$$\begin{aligned} \sum_{r=a}^b r^{(\nu)} &= \sum_{r=a}^b \frac{\Gamma(r+1)}{\Gamma(r-\nu+1)} \stackrel{\text{(by (17))}}{=} \frac{1}{(\nu+1)} \sum_{r=a}^b \left(\frac{\Gamma(r+2)}{\Gamma(r-\nu+1)} - \frac{\Gamma(r+1)}{\Gamma(r-\nu)} \right) = \\ &= \frac{1}{(\nu+1)} \left(\sum_{r=a}^b \left\{ \left(\frac{\Gamma(a+2)}{\Gamma(a-\nu+1)} - \frac{\Gamma(a+1)}{\Gamma(a-\nu)} \right) + \left(\frac{\Gamma(a+3)}{\Gamma(a+2-\nu)} - \frac{\Gamma(a+2)}{\Gamma(a+1-\nu)} \right) + \right. \right. \\ &\quad \left. \left(\frac{\Gamma(a+4)}{\Gamma(a+3-\nu)} - \frac{\Gamma(a+3)}{\Gamma(a+2-\nu)} \right) + \dots + \left(\frac{\Gamma(b+1)}{\Gamma(b-\nu)} - \frac{\Gamma(b)}{\Gamma(b-1-\nu)} \right) + \left(\frac{\Gamma(b+2)}{\Gamma(b-\nu+1)} - \frac{\Gamma(b+1)}{\Gamma(b-\nu)} \right) \right\} \right) = \\ &= \frac{1}{(\nu+1)} \left(\frac{\Gamma(b+2)}{\Gamma(b-\nu+1)} - \frac{\Gamma(a+1)}{\Gamma(a-\nu)} \right), \end{aligned}$$

proving the claim. ■

Next we present a discrete fractional Ostrowski type inequality.

Theorem 20 Let $\mu > p$, $p \in \mathbb{Z}^+$, μ not an integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$. Here f is defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$ and $j \in [a+m-p+1, b]$, with $a+m-p < b \in \mathbb{N}$. Assume that $\Delta^k f(a) = 0$, for $k \in [p+1, \dots, m-1]$.

Then

$$\begin{aligned} & \left| \left(\frac{1}{(b-a-m+p)} \sum_{j=a+m-p+1}^b \Delta^p f(j) \right) - \Delta^p f(a) \right| \leq \\ & \frac{1}{(b-a-m+p)\Gamma(\mu-p+2)} \left[(b-a-\nu+1)^{(\mu-p+1)} - \Gamma(\mu-p+2) \right] \\ & \cdot \left(\max_{t \in \{a+\nu, \dots, b-\mu+p\}} |\Delta_*^\mu f(t)| \right). \end{aligned} \quad (23)$$

Proof. By (12) we have

$$\Delta^p f(j) - \Delta^p f(a) = \frac{1}{\Gamma(\mu-p)} \sum_{s=a+\nu}^{j-\mu+p} (j-s-1)^{(\mu-p-1)} \Delta_*^\mu f(s), \quad (24)$$

for all $j \in [a+m-p+1, b]$.

We get that

$$\begin{aligned} & \frac{1}{b-(a+m-p)} \sum_{j=a+m-p+1}^b \Delta^p f(j) - \Delta^p f(a) = \\ & \frac{1}{b-(a+m-p)} \sum_{j=a+m-p+1}^b (\Delta^p f(j) - \Delta^p f(a)) = \\ & \frac{1}{(b-(a+m-p))\Gamma(\mu-p)} \sum_{j=a+m-p+1}^b \left(\sum_{s=a+\nu}^{j-\mu+p} (j-s-1)^{(\mu-p-1)} \Delta_*^\mu f(s) \right). \end{aligned} \quad (25)$$

Therefore we obtain

$$\begin{aligned} & \left| \frac{1}{b-(a+m-p)} \sum_{j=a+m-p+1}^b \Delta^p f(j) - \Delta^p f(a) \right| = \\ & \frac{1}{(b-a-m+p)} \left| \sum_{j=a+m-p+1}^b (\Delta^p f(j) - \Delta^p f(a)) \right| \leq \\ & \frac{1}{(b-a-m+p)} \sum_{j=a+m-p+1}^b |\Delta^p f(j) - \Delta^p f(a)| \leq \\ & \frac{1}{(b-a-m+p)\Gamma(\mu-p)} \sum_{j=a+m-p+1}^b \left(\sum_{s=a+\nu}^{j-\mu+p} (j-s-1)^{(\mu-p-1)} \cdot |\Delta_*^\mu f(s)| \right) \stackrel{\text{(by (19))}}{\leq} \\ & \frac{1}{(b-a-m+p)\Gamma(\mu-p+1)} \sum_{j=a+m-p+1}^b (j-a-\nu)^{(\mu-p)} \cdot \max_{s \in \{a+\nu, \dots, j-\mu+p\}} |\Delta_*^\mu f(s)| \leq \\ & \frac{1}{(b-a-m+p)\Gamma(\mu-p+1)} \left(\sum_{j=a+m-p+1}^b (j-a-\nu)^{(\mu-p)} \right) \cdot \max_{s \in \{a+\nu, \dots, b-\mu+p\}} |\Delta_*^\mu f(s)| =: (*) \end{aligned} \quad (26)$$

Next we use (22). We notice that

$$\sum_{j=a+m-p+1}^b (j-a-\nu)^{(\mu-p)} = \sum_{r=\mu-p+1}^{b-a-\nu} r^{(\mu-p)} =$$

$$\frac{1}{(\mu - p + 1)} \left(\frac{\Gamma(b - a - \nu + 2)}{\Gamma(b - a - m + p + 1)} - \Gamma(\mu - p + 2) \right). \quad (27)$$

Therefore

$$\begin{aligned} (*) &= \frac{1}{(b - a - m + p) \Gamma(\mu - p + 2)} \left(\frac{\Gamma(b - a - \nu + 2)}{\Gamma(b - a - m + p + 1)} - \Gamma(\mu - p + 2) \right) \\ &\quad \cdot \left(\max_{t \in \{a + \nu, \dots, b - \mu + p\}} |\Delta_*^\mu f(t)| \right). \end{aligned} \quad (28)$$

The last completes the proof. ■

Next we give a discrete fractional Poincaré inequality.

Theorem 21 *Let $\mu > p$, $p \in \mathbb{Z}^+$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$. Assume that $\Delta^k f(a) = 0$, $k = p, \dots, m - 1$, f defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Let $\gamma, \delta > 1$: $\frac{1}{\gamma} + \frac{1}{\delta} = 1$. Then*

$$\begin{aligned} \sum_{j=a+m-p}^b |\Delta^p f(j)|^\delta &\leq \frac{1}{(\Gamma(\mu - p))^\delta} \left[\sum_{j=a+m-p}^b \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{\delta}{\gamma}} \right] \\ &\quad \cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right). \end{aligned} \quad (29)$$

Proof. We have

$$\Delta^p f(j) = \frac{1}{\Gamma(\mu - p)} \sum_{s=a+\nu}^{j-\mu+p} (j-s-1)^{(\mu-p-1)} \Delta_*^\mu f(s), \quad \forall j \in [a+m-p, b]. \quad (30)$$

Let $\gamma, \delta > 1$ such that $\frac{1}{\gamma} + \frac{1}{\delta} = 1$.

We observe that

$$|\Delta^p f(j)| \leq \frac{1}{\Gamma(\mu - p)} \sum_{s=a+\nu}^{j-\mu+p} (j-s-1)^{(\mu-p-1)} |\Delta_*^\mu f(s)|$$

(by discrete Hölder's inequality)

$$\leq \frac{1}{\Gamma(\mu - p)} \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{1}{\gamma}} \cdot \left(\sum_{s=a+\nu}^{j-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)^{\frac{1}{\delta}}. \quad (31)$$

I.e. it holds

$$|\Delta^p f(j)|^\delta \leq \frac{1}{(\Gamma(\mu - p))^\delta} \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{\delta}{\gamma}} \cdot \left(\sum_{s=a+\nu}^{j-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)$$

$$\begin{aligned} &\leq \frac{1}{(\Gamma(\mu-p))^\delta} \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{\delta}{\gamma}} \\ &\quad \cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right), \quad \forall j \in [a+m-p, b]. \end{aligned} \quad (32)$$

Applying $\sum_{j=a+m-p}^b$ on (32) we establish (29). ■

It follows a discrete Sobolev type fractional inequality.

Theorem 22 *Let $\mu > p$, $p \in \mathbb{Z}^+$, μ non-integer, $m = \lceil \mu \rceil$, $\nu = m - \mu$. Assume that $\Delta^k f(a) = 0$, $k = p, \dots, m-1$; f defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Let $\gamma, \delta > 1$: $\frac{1}{\gamma} + \frac{1}{\delta} = 1$, and $r \geq 1$. Then*

$$\begin{aligned} &\left(\sum_{j=a+m-p}^b |\Delta^p f(j)|^r \right)^{\frac{1}{r}} \leq \\ &\frac{1}{\Gamma(\mu-p)} \left[\sum_{j=a+m-p}^b \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{r}{\gamma}} \right]^{\frac{1}{r}} \cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)^{\frac{1}{\delta}}. \end{aligned} \quad (33)$$

Proof. By (31) and discrete Hölder's inequality, we have

$$\begin{aligned} |\Delta^p f(j)| &\leq \frac{1}{\Gamma(\mu-p)} \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{1}{\gamma}} \\ &\quad \cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)^{\frac{1}{\delta}}, \quad \forall j \in [a+m-p, b], \end{aligned} \quad (34)$$

where $\gamma, \delta > 1$: $\frac{1}{\gamma} + \frac{1}{\delta} = 1$.

Hence, by $r \geq 1$ we obtain

$$\begin{aligned} |\Delta^p f(j)|^r &\leq \frac{1}{(\Gamma(\mu-p))^r} \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{r}{\gamma}} \\ &\quad \cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)^{\frac{r}{\delta}}, \quad \forall j \in [a+m-p, b]. \end{aligned} \quad (35)$$

Consequently we get

$$\sum_{j=a+m-p}^b |\Delta^p f(j)|^r \leq \frac{1}{(\Gamma(\mu-p))^r} \left[\sum_{j=a+m-p}^b \left(\sum_{s=a+\nu}^{j-\mu+p} ((j-s-1)^{(\mu-p-1)})^\gamma \right)^{\frac{r}{\gamma}} \right]$$

$$\cdot \left(\sum_{s=a+\nu}^{b-\mu+p} |\Delta_*^\mu f(s)|^\delta \right)^{\frac{r}{\delta}}. \quad (36)$$

The last proves the claim. ■

We finish with the following discrete fractional average Sobolev type inequality.

Theorem 23 *Let $0 < \mu_1 < \mu_2 < \dots < \mu_k$; $m_l = \lceil \mu_l \rceil$, $\nu_l = m_l - \mu_l$, $l = 1, \dots, k$, $k \in \mathbb{N}$. Assume that $\Delta^\tau f(a) = 0$, for $\tau = 0, 1, \dots, m_k - 1$: f is defined on \mathbb{N}_a , $a \in \mathbb{Z}^+$. Let $r \geq 1$; $C_l(s) > 0$ defined on $[a + \nu_l, b - \mu_l]$, $l = 1, \dots, k$. Call $B_l := \sum_{s=a+\nu_l}^{b-\mu_l} C_l(s) (\Delta_*^{\mu_l} f(s))^2$,*

$$\delta^* := \max_{1 \leq l \leq k} \left\{ \frac{1}{(\Gamma(\mu_l))^2} \left[\sum_{j=a+m_l}^b \left(\sum_{s=a+\nu_l}^{j-\mu_l} ((j-s-1)^{(\mu_l-1)})^2 \right)^{\frac{r}{2}} \right]^{\frac{2}{r}} \right\},$$

$$\varrho^* := \max_{1 \leq l \leq k} \left| \left(\frac{1}{C_l(s)} \right) \right|_{\infty, [a+\nu_l, b-\mu_l]}.$$

Then

$$\|f\|_{r, [a+m_k, b]} \leq \sqrt{\delta^* \varrho^*} \left(\frac{\sum_{l=1}^k B_l}{k} \right)^{\frac{1}{2}}. \quad (37)$$

Proof. We see that also $\Delta^\tau f(a) = 0$, $\tau = 0, 1, \dots, m_l - 1$, $l = 1, \dots, k - 1$. So the assumptions of Theorem 22 are fulfilled for f and fractional orders μ_l , $l = 1, \dots, k$. Thus by choosing $p = 0$ and $\gamma = \delta = 2$ we apply (33), for $l = 1, \dots, k$, to obtain

$$\begin{aligned} \left(\sum_{j=a+m_l}^b |f(j)|^r \right)^{\frac{1}{r}} &\leq \frac{1}{\Gamma(\mu_l)} \left[\sum_{j=a+m_l}^b \left(\sum_{s=a+\nu_l}^{j-\mu_l} ((j-s-1)^{(\mu_l-1)})^2 \right)^{\frac{r}{2}} \right]^{\frac{1}{r}} \\ &\cdot \left(\sum_{s=a+\nu_l}^{b-\mu_l} (\Delta_*^{\mu_l} f(s))^2 \right)^{\frac{1}{2}}. \end{aligned} \quad (38)$$

Hence it holds

$$\begin{aligned} \left(\sum_{j=a+m_l}^b |f(j)|^r \right)^{\frac{2}{r}} &\leq \\ \frac{1}{(\Gamma(\mu_l))^2} \left[\sum_{j=a+m_l}^b \left(\sum_{s=a+\nu_l}^{j-\mu_l} ((j-s-1)^{(\mu_l-1)})^2 \right)^{\frac{r}{2}} \right]^{\frac{2}{r}} &\cdot \left(\sum_{s=a+\nu_l}^{b-\mu_l} (\Delta_*^{\mu_l} f(s))^2 \right) \\ \leq \delta^* \left(\sum_{s=a+\nu_l}^{b-\mu_l} (\Delta_*^{\mu_l} f(s))^2 \right) &= \delta^* \left(\sum_{s=a+\nu_l}^{b-\mu_l} (C_l(s))^{-1} (C_l(s)) (\Delta_*^{\mu_l} f(s))^2 \right) \end{aligned}$$

$$\leq \delta^* \rho^* \left(\sum_{s=a+\nu_l}^{b-\mu_l} C_l(s) (\Delta_*^{\mu_l} f(s))^2 \right). \quad (39)$$

That is

$$\left(\sum_{j=a+m_k}^b |f(j)|^r \right)^{\frac{2}{r}} \leq \left(\sum_{j=a+m_l}^b |f(j)|^r \right)^{\frac{2}{r}} \leq \delta^* \rho^* \left(\sum_{s=a+\nu_l}^{b-\mu_l} C_l(s) (\Delta_*^{\mu_l} f(s))^2 \right) = \delta^* \rho^* B_l, \quad \text{for } l = 1, \dots, k. \quad (40)$$

Hence

$$\|f\|_{r,[a+m_k,b]}^2 \leq \delta^* \rho^* \left(\frac{\sum_{l=1}^k B_l}{k} \right), \quad (41)$$

proving the claim. ■

References

- [1] R.P. Agarwal, *Difference equations and inequalities*, Marcel Dekker, NY, 2000.
- [2] Ferhan Atici and Paul W. Eloe, *A Transform method in discrete fractional calculus*, Intern.J. of Difference Equations, Vol. 2, no. 2(2007), 165-176.
- [3] Ferhan Atici and Paul W. Eloe, *Initial value problems in discrete fractional calculus*, Proc. AMS, 137, no. 3 (2009), 981-989.
- [4] K.S. Miller and B. Ross, *Fractional difference calculus*, Proc. Intern. Sympos. on Univalent Functions, Fractional calculus and their Appls. Nihon University, Koriyama, Japan, May 1988; Ellis Horwood, Chichester, West Susses, England, 1989, pp. 139-152.