Dickson's conjecture on Z^n

—An equivalent form of Green-Tao's conjecture

Shaohua Zhang

School of Mathematics, Shandong University, Jinan, Shandong, 250100, PRC

Abstract

In [1], we give Dickson's conjecture on N^n . In this paper, we further give Dickson's conjecture on Z^n and obtain an equivalent form of Green-Tao's conjecture [2]. Based on our work, it is possible to establish a general theory that several multivariable integral polynomials on Z^n represent simultaneously prime numbers for infinitely many integral points and generalize the analogy of Chinese Remainder Theorem in [3].

Résumé

Conjecture de Dickson sur Z^n — Une forme équivalente de conjecture de Green-Tao. Dans [1], nous donnons la conjecture de Dickson sur N^n . Dans ce document, en outre nous accordons une conjecture de Dickson sur Z^n et obtenons une forme équivalent de conjecture de Green-Tao [2]. Sur la base de nos travaux, il est possible d'établir une théorie générale que plusieurs polynômes intégraux multivariables sur Z^n représentent simultanément les nombres premiers pour un nombre infini de points entiers et de généraliser les l'analogie de Théorème des Restes Chinois dans [3].

Keywords: Dickson's conjecture, affine-linear form, Generalized Chinese Remainder Theorem, prime, multivariable integral polynomials, admissible map, strongly admissible map, polynomial map on Z^n

2000 MR Subject Classification: 11A41, 11A99

 $^{^1\}mathrm{E}\text{-}\mathrm{mail}$ address: shaohuazhang@mail.sdu.edu.cn

²This work was partially supported by the National Basic Research Program (973) of China (No. 2007CB807902) and the Natural Science Foundation of Shandong Province (No. Y2008G23).

1 Some basic notations

Let Z be the set of integers. Denote the set of all positive prime numbers by P (we do not consider negative primes.) and denote the set of all natural numbers or positive integers by N. We define affine n-space over Z (resp. N), denoted Z^n (resp. N^n), to be the set of all n-tuples of elements of Z (resp. N). An element $x = (x_1, ..., x_n)$ in Z^n is called an integral point and x_i is called the coordinates of x. Let $Z[x_1, ..., x_n]$ be the polynomial ring in n variables over Z. Let $Z_n^* = \{x \in N | 1 \le x \le n, \gcd(x, n) = 1\}$ be the set of positive integers less than or equal to n that are coprime to n. Let $Z_n = \{x \in Z | 0 \le x \le n - 1\}.$

2 Some basic definitions

Let's consider the map $F: \mathbb{Z}^n \to \mathbb{Z}^m$ for all integral points $x = (x_1, ..., x_n) \in \mathbb{Z}^n$, $F(x) = (f_1(x), ..., f_m(x))$ for distinct polynomials $f_1, ..., f_m \in \mathbb{Z}[x_1, ..., x_n]$, where $m, n \in \mathbb{N}$. In this case, we call F a polynomial map on \mathbb{Z}^n . Call F a polynomial map on \mathbb{N}^n if $F: \mathbb{N}^n \to \mathbb{Z}^m$ for all integral points $x \in \mathbb{N}^n$, $F(x) = (f_1(x), ..., f_m(x))$. We call F on \mathbb{Z}^n (resp. \mathbb{N}^n) linear or a system of non-constant affine-linear forms if for any $1 \leq i \leq m$, $f_i(x_1, ..., x_n)$ has degree 1.

We call the polynomial map F on Z^n (resp. N^n) admissible if for every positive integer r there exists an integral point $x = (x_1, ..., x_n) \in Z^n$ (resp. N^n) such that r is coprime to $f_1(x) \times ... \times f_m(x) = \prod_{i=1}^{i=m} f_i(x)$, moreover, $f_i(x) > 1$ for $1 \le i \le m$ (namely, $F(x) \in (N \setminus \{1\})^m$). Generally speaking, when people talk about the definition of "admissible", they do not consider the condition " $f_i(x) > 1$ for $1 \le i \le m$ ". Nevertheless, from this paper, one will see that this is very necessary. Sometimes, in the definition of "admissible", people regard r as a prime. But, this is not always true. For example, let's consider one-variable function $f(x) = -x^2 + 6$. It is easy to test that for any prime p, there is an x such that gcd(f(x), p) = 1 and f(x) > 1. But f(x) is not "admissible".

We call the polynomial map F on Z^n (resp. N^n) strongly admissible if there is a positive integer C such that for every positive integer $k \ge C$, there exists an integral point $x = (x_1, ..., x_n)$ such that $f_1(x) > 1, ..., f_m(x) > 1$ are all in Z_k^* . We call the least positive integer C such that F is strongly admissible a strongly admissible constant. Let S be the set of all solutions of the simultaneous equations:

$$f_1(x_1, ..., x_n) \in P, ..., f_m(x_1, ..., x_n) \in P$$

Let $H = F(S) \in P^m$ be the image of S under F. An element of H is called a prime point. An important question is the following: Let F be a given polynomial map on Z^n (resp. N^n), how to determine whether H is an infinite set or not?

We say that several multivariable integral polynomials $f_1(x), ..., f_m(x)$ on Z^n (resp. N^n) represent simultaneously prime numbers for infinitely many integral points x, if for any $1 \leq i \leq m$, $f_i(x)$ itself can represent primes for infinitely many integral points x, moreover, there is an infinite sequence of integral points $(x_{11}, ..., x_{n1}), ..., (x_{1i}, ..., x_{ni}), ...$ such that for any positive integer r, $f_1(x_{1r}, ..., x_{nr}), ..., f_m(x_{1r}, ..., x_{ni})$ represent simultaneously primes, and for any $i \neq j$, $f_1(x_{1i}, ..., x_{ni}) \neq f_1(x_{1j}, ..., x_{nj}), ...,$ $f_m(x_{1i}, ..., x_{ni}) \neq f_m(x_{1j}, ..., x_{nj})$ hold simultaneously. In this case, we also say that the polynomial map F on Z^n (resp. N^n) represents infinitely many prime points. One of our aims is to study the conditions that the polynomial map F on Z^n represents infinitely many prime points. In the case that F is on N^n , we have initially considered this problem, for the details, see [3].

3 Introductions

In [1], we have considered the case that F on N is linear and obtained an equivalent form of Dickson's conjecture which gives the necessary and sufficient condition that a system of non-constant affine-linear forms on Nrepresents infinitely many prime points. Moreover, we generalized Dickson's conjecture to the case that the linear polynomial map F is on N^n . In this paper, we will further consider the case that the linear polynomial map Fis on Z^n and obtain an equivalent form of Green-Tao's conjecture [2].

Now, let's start with the work of Green and Tao. In 2006 [2], they are the first to generalize Dickson's conjecture and consider the case that F is a system of non-constant affine-linear forms on Z^n . They noticed that if the linear polynomial map $F = (f_1(x), ..., f_m(x))$ on Z^n satisfies the conditions that for any $1 \leq i \neq j \leq m$, $f_i(x)$ and $f_j(x)$ are not rational multiples of each other, then F represents infinitely many prime points. They further gave asymptotic formulae and generalized Hardy-Littlewood Conjecture. But, the conditions that for any $1 \leq i \neq j \leq m$, $f_i(x)$ and $f_j(x)$ are not rational multiples of each other is not sufficient and it is only necessary. For example, $F = (f_1(x), f_2(x), f_3(x))$ never represents prime points, where $f_1(x) = -2x_1 + 3x_2 - 6$, $f_2(x) = 3x_1 - 4x_2 - 6$, $f_3(x) = -9x_1 + 10x_2 - 6$. Based on the idea in [3], we fix this condition as follows:

Condition A: For every positive integer r, there exists an integral point $x = (x_1, x_2, ..., x_n)$ such that for any $1 \le i \le m$, $f_i(x) > 1$ and $gcd(f_i(x), r) = 1$.

Clearly, Condition A is necessary. We hope that it also is sufficient. Thus, we have the following conjecture:

Conjecture 1: If the linear polynomial map F on Z^n is admissible, then F represents infinitely many prime points.

In the case that the linear polynomial map F is on N^n , in 2009, the author [1] obtained the equivalent form of Dickson's conjecture and conjectured that if F on N^n is strongly admissible, then F represents infinitely many prime points. Naturally, one might further conjecture the following:

Conjecture 2: If the linear polynomial map F on Z^n is strongly admissible, then F represents infinitely many prime points.

In order to explain the rationality of Conjecture 2, we go back to Dickson's conjecture. In 1904, Dickson conjectured: Let $m \ge 1$, $f_i(x) = a_i + b_i x$ with a_i and b_i integers, $b_i \ge 1$ (for i = 1, ..., m). If there does not exist any integer n > 1 dividing all the products $\prod_{i=1}^{i=s} f_i(k)$, for every integer k, then there exist infinitely many natural numbers m such that all numbers $f_1(m), ..., f_s(m)$ are primes.

Obviously, from the condition " $b_i \ge 1$ ", we deduce that Dickson only considered the linear polynomial map F on N. Therefore, one might call this conjecture Dickson's conjecture on N. Then, What does Dickson's conjecture on Z look like? One might naively guess: Let $m \ge 1$, $f_i(x) = a_i + b_i x$ with a_i and b_i integers for i = 1, ..., m. If there does not exist any integer n > 1 dividing all the products $\prod_{i=1}^{i=s} f_i(k)$, for every integer k, then there exist infinitely many natural numbers m such that all numbers $f_1(m), ..., f_s(m)$ are primes. Unfortunately, it is always not true. For instance, $f_1(x) = 4x-5$ and $f_2(x) = -3x+4$ satisfy the claimed condition, but they do not represent simultaneously primes since we do not consider negative primes.

It is not hard to see that Dickson's conjecture on Z should be the fol-

lowing: Let $m \ge 1$, $f_i(x) = a_i + b_i x$ with a_i and b_i integers, either $b_i \ge 1$ or $b_i \leq -1$ (for all i = 1, ..., m). If there does not exist any integer n > 1dividing all the products $\prod_{i=1}^{i=s} f_i(k)$, for every integer k, then there exist infinitely many natural numbers m such that all numbers $f_1(m), ..., f_s(m)$ are primes. This can be equivalently stated that if the linear polynomial map F on Z is admissible, then F represents infinitely many prime points. By the method in [1], one can further prove that if the linear polynomial map F on Z is admissible then F also is strongly admissible. Thus we get Conjecture 2. Note that in the multivariable case, the signs of coefficients might be disordered.

As a toy example, we believe that $f_1(x) = 2x_1 - 3x_2$ and $f_2(x) = -3x_1 + 3x_2$ $4x_2$ represent simultaneously primes for infinitely many integral points x = $(x_1, x_2) \in Z^2.$

Theorem 1: Conjecture 1 and Conjecture 2 are equivalent.

For the proof of theorem 1, see next section. Similarly, one can prove that if the linear polynomial map F on N^n is admissible then F also is strongly admissible. Thus, we complement the work in [1]. Based on given equivalent forms, more precisely, by finding "strongly admissible", it is possible to establish a general theory that several multivariable integral polynomials on Z^n represent simultaneously prime numbers for infinitely many integral points and generalize the analogy of Chinese Remainder Theorem in [3].

Proof of Theorem 1 4

In order to prove that Conjecture 1 and Conjecture 2 are equivalent, it is enough to prove that the following conditions B, C are equivalent. Let the linear polynomial map $F(x) = (f_1(x), ..., f_m(x))$ be on Z^n .

Condition B: *F* is admissible.

Condition C: *F* is strongly admissible.

Lemma 1: Let $F = \begin{cases} f_1(x_1, \dots, x_n) = a_{11}x_1 + \dots + a_{1n}x_n + b_1 \\ \dots & be the \\ f_m(x_1, \dots, x_n) = a_{m1}x_1 + \dots + a_{mn}x_n + b_m \end{cases}$ be the linear polynomial map. If F is admissible, then for any positive integer c,

there is an integral point x such that any prime divisor of $\prod_{i=1}^{i=m} f_i(x)$ is

greater than c, moreover

$$\begin{cases} a_{11}x_1 + \dots + a_{1n}x_n > e = \max_{1 \le j \le m} \{|a_{j1}| + \dots + |a_{jn}|\} > 0 \\ \dots \\ a_{m1}x_1 + \dots + a_{mn}x_n > e = \max_{1 \le j \le m} \{|a_{j1}| + \dots + |a_{jn}|\} > 0 \end{cases}$$

Proof: The case $b_i = 0$ for each $1 \le i \le m$ is trivial. So, let's assume that $B = \{b_1, ..., b_m\} \ne \{0\}$ and consider the number $\alpha = \prod_{x \in B, x \ne 0} x$. By the known condition, there is an integral point x such that

$$\gcd(\prod_{i=1}^{i=m} f_i(x), \prod_{p \le 2|\alpha|ce} p) = 1,$$

moreover $f_i(x) > 1$. Notice that for each $1 \le i \le m$, any prime divisor of $f_i(x)$ is greater than $2|\alpha|ce$. Therefore, $a_{i1}x_1 + \ldots + a_{in}x_n > e$ for each $1 \le i \le m$. Since F is admissible, hence e > 0 and Lemma 1 holds.

Proof that Conditions B, C are equivalent: Clearly, Condition $C \Rightarrow$ Condition B. Next, we prove that Condition $B \Rightarrow$ Condition C. We write

$$F = \begin{cases} f_1(x_1, \dots, x_n) = a_{11}x_1 + \dots + a_{1n}x_n + b_1 \\ \dots \\ f_m(x_1, \dots, x_n) = a_{m1}x_1 + \dots + a_{mn}x_n + b_m \end{cases}$$

By lemma 1 and Condition B, we choose an integer point $y = (y_1, ..., y_n)$ such that, for $1 \le i \le m$, $f_i(y) > 1$ and $\begin{cases} a_{11}y_1 + ... + a_{1n}y_n > e > 0\\a_{m1}y_1 + ... + a_{mn}y_n > e > 0 \end{cases}$

By lemma 1 and Condition B again, we can choose an integer point $z = (z_1, ..., z_n)$ such that any prime divisor of $\prod_{i=1}^{i=m} f_i(z)$ is greater than $\prod_{i=1}^{i=m} f_i(y)$. We also can choose an integer point $w = (w_1, ..., w_n)$ such that any prime divisor of $\prod_{i=1}^{i=m} f_i(w)$ is greater than $1 + e + m \times \prod_{i=1}^{i=m} f_i(z) \times \max_{1 \le j \le m} \{|a_{j1}y_1| + ... + |a_{jn}y_n|\}.$

We claim that there is a positive integer $C \leq \prod_{i=1}^{i=m} f_i(z)f_i(w)$ such that for every positive integer $k \geq C$, there exists an integral point $x = (x_1, ..., x_n)$ such that $f_1(x) > 1, ..., f_m(x) > 1$ are all in Z_k^* . Namely, Condition B \Rightarrow Condition C.

Obviously, if $gcd(k, \prod_{i=1}^{i=m} f_i(z)) = 1$, then we can choose x = z. And if $gcd(k, \prod_{i=1}^{i=m} f_i(w)) = 1$, then we can choose x = w. Therefore, it is enough

to consider the case of k = pqt, where p is a prime divisor of $\prod_{i=1}^{i=m} f_i(z)$ and q is a prime divisor of $\prod_{i=1}^{i=m} f_i(w)$, t is any positive integer.

By our assumption that F is admissible, one can choose an integral point $v = (v_1, ..., v_n)$ such that $gcd(t, \prod_{i=1}^{i=m} f_i(v)) = 1$. Write $t = p^e d$ with gcd(p,d) = 1. Clearly, if d = 1, then we can choose x = y such that Condition B \Rightarrow Condition C since $gcd(pqt, \prod_{i=1}^{i=m} f_i(y)) = 1$ and $f_i(y) > 1$. Hence, we might assume that d > 1.

Now, we consider the coordinates. If for each $1 \leq i \leq n$, $y_i \equiv v_i \pmod{d}$, then, we can similarly choose x = y. So, now, we only consider the case that for some $1 \leq i \leq n$, $y_i \neq v_i \pmod{d}$. Without loss of generality, we assume that $y_1 \neq v_1 \pmod{d}$.

Since gcd(p, d) = 1 and $y_1 \neq v_1 \pmod{d}$, hence there must be a positive integer r_1 with $r_1 < d$ such that $y_1 + pr_1 \equiv v_1 \pmod{d}$. Also from gcd(p, d) = 1, we deduce that there is an integer r_i with $0 \leq r_i < d$ such that $y_i + pr_i \equiv v_i \pmod{d}$ for each $2 \leq i \leq n$.

Let $x^{(h)} = (y_1 + pr_1 + pdhy_1, y_2 + pr_2 + pdhy_2, ..., y_n + pr_n + pdhy_n)$ with $h \in N$. It is easy to see that $gcd(pt, \prod_{i=1}^{i=m} f_i(x^{(h)})) = 1$. We also have that for each $1 \le i \le m$, $f_i(x^{(h)}) = f_i(y) + p(a_{i1}r_1 + ... + a_{in}r_n) + pdh(a_{i1}y_1 + ... + a_{in}y_n) > 1 + p(a_{i1}r_1 + ... + a_{in}r_n) + pdhe > 1 + pdhe - pd(|a_{i1}| + ... + |a_{in}|) > 1$.

Finally, let h range over $\{1, 2, ..., m + 1\}$ and let's consider the following matrix:

$$M = (m_{i,j}) = \begin{pmatrix} f_1(x^{(1)}), & \cdots, & f_1(x^{(m+1)}) \\ \cdots, & \cdots, & \cdots \\ f_m(x^{(1)}), & \cdots, & f_m(x^{(m+1)}) \end{pmatrix}$$

Since $q > 1 + e + m \times \prod_{i=1}^{i=m} f_i(z) \times \max_{1 \le j \le m} \{|a_{j1}y_1| + ... + |a_{jn}y_n|\}$, hence there is at most a number which can be divided q in each row of the matrix M. But there are m + 1 columns in M. So, there must be some jwith $1 \le j \le m + 1$ such that $(\prod_{i=1}^{i=m} f_i(x^{(j)}), q) = 1$. Also notice that for each $1 \le i \le m, 1 < f_i(x^{(j)}) < pqt$. This shows immediately that Theorem 1 holds.

5 Further considerations

In this paper, we mainly research the problem that the linear polynomial map F on Z^n represents infinitely many prime points and try to give a necessary and sufficient condition that F represents infinitely many prime

points. Based on this work and also based on the idea in [3], it is possible to further generalize the famous H Hypothesis as follows:

Conjecture 3 (Generalized H Hypothesis): Let F be a polynomial map on Z^n . If F is strongly admissible, and there exists an integral point $y = (y_1, ..., y_n)$ such that $f_1(y) \ge C, ..., f_m(y) \ge C$ are all primes, then F represents infinitely many prime points, where C is the strongly admissible constant.

Lemma 2 (Generalized Chinese Remainder Theorem): If $gcd(a_i, a_j) = 1$ for $1 \le i \ne j \le m$, then the Cartesian product $(Z_{a_1}^*)^n \times \ldots \times (Z_{a_m}^*)^n$ (resp. $(Z_{a_1})^n \times \ldots \times (Z_{a_m})^n$) is isomorphic to $(Z_{a_1 \ldots a_m}^*)^n$ (resp. $(Z_{a_1 \ldots a_m})^n$) for any positive integers a_i, m, n .

By Lemma 2 and the idea in [3], we further generalize the analogy of Chinese Remainder Theorem in [3].

Generalized analogy of Chinese Remainder Theorem: Let $F = (f_1, ..., f_m)$ be a polynomial map on Z^n . If F represents infinitely many prime points, and if $gcd(a_i, a_j) = 1$ for $1 \le i \ne j \le k$, and there exist integral point $x^{(j)} \in Z^n$ such that $F(x^{(j)})$ is in $(Z^*_{a_j} \setminus \{1\})^m$ for $1 \le j \le k$, then there exists an integral point z such that F(z) is in $(Z^*_{a_1...a_k} \setminus \{1\})^m$.

6 Acknowledgements

I am very thankful to the referee for reading the paper, and also to my supervisor Professor Xiaoyun Wang for her suggestions. Thank the key lab of cryptography technology and information security in Shandong University and the Institute for Advanced Study in Tsinghua University, for providing me with excellent conditions.

7 References

[1] Shaohua Zhang, Notes on Dickson's Conjecture, available at: http://arxiv.org/abs/0906.3850

[2] Ben Green, Terence Tao, Linear equations in primes, preprint, available at: http://arxiv.org/abs/math/0606088, Ann. of Math. (2), in press

[3] Shaohua Zhang, On the infinitude of some special kinds of primes, available at: http://arxiv.org/abs/0905.1655