

A NEW APPROXIMATION OF RELAXED ENERGIES FOR HARMONIC MAPS AND THE FADDEEV MODEL

MARIANO GIAQUINTA, MIN-CHUN HONG AND HAO YIN

ABSTRACT. We propose a new approximation for the relaxed energy E of the Dirichlet energy and prove that the minimizers of the approximating functionals converge to a minimizer u of the relaxed energy, and that u is partially regular without using the concept of Cartesian currents. We also use the same approximation method to study the variational problem of the relaxed energy for the Faddeev model and prove the existence of minimizers for the relaxed energy \tilde{E}_F in the class of maps with Hopf degree ± 1 .

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^3$ be an open bounded domain with smooth boundary $\partial\Omega$. For each $p > 0$, set

$$W^{1,p}(\Omega; S^2) = \{u \in W^{1,p}(\Omega; \mathbb{R}^3) : |u| = 1 \text{ a.e. on } \Omega.\}$$

For a map $u \in W^{1,2}(\Omega; S^2)$, we consider

$$(1.1) \quad E(u, \Omega) = \int_{\Omega} |\nabla u|^2 dx.$$

A map $u \in W^{1,2}(\Omega; S^2)$ is said to be a harmonic map if u satisfies

$$\Delta u + |\nabla u|^2 u = 0$$

in the weak sense.

In [14], Hardt-Lin discovered a gap phenomenon for the Dirichlet energy; i.e., there is a given smooth boundary value $\gamma : \partial\Omega \rightarrow S^2$ with $\deg \gamma = 0$ such that

$$\min_{u \in W_{\gamma}^{1,2}(\Omega, S^2)} \int_{\Omega} |\nabla u|^2 dx < \inf_{v \in W_{\gamma}^{1,2} \cap C^{\infty}(\bar{\Omega}, S^2)} \int_{\Omega} |\nabla v|^2 dx.$$

It is a very interesting problem whether the above right-hand term

$$\inf_{v \in H_{\gamma}^1 \cap C^{\infty}(\bar{\Omega}, S^2)} \int_{\Omega} |\nabla v|^2 dx$$

can be attained by a map $u \in W_{\gamma}^{1,2} \cap C^{\infty}(\bar{\Omega}, S^2)$ or not. For $u \in W^{1,2}(\Omega; S^2)$ we consider the vector field $D(u)$ given by

$$D(u) := (u \cdot u_{x_2} \times u_{x_3}, u \cdot u_{x_3} \times u_{x_1}, u \cdot u_{x_1} \times u_{x_2}).$$

Given $u_0 \in C_{\gamma}^{\infty}(\bar{\Omega}; S^2)$ for a map $u \in W_{\gamma}^{1,2}(\bar{\Omega}; S^2)$, we set

$$L(u) := L(u, u_0) = \frac{1}{4\pi} \sup_{\xi: \Omega \rightarrow \mathbb{R}, \|\nabla \xi\|_{L^{\infty}} \leq 1} \int_{\Omega} [D(u) - D(u_0)] \cdot \nabla \xi dx.$$

1991 *Mathematics Subject Classification.* AMS 58E20, 58E50.

Key words and phrases. Relaxed energy, Harmonic maps, Faddeev model.

The relaxed energy functional for the Dirichlet energy $E(u)$ is then defined by

$$F(u) = \int_{\Omega} |\nabla u|^2 dx + 8\pi L(u).$$

Bethuel-Brezis-Coron in [4] proved that F is sequentially lower semi-continuous and satisfies

$$(1.2) \quad \inf_{u \in W_{\gamma}^{1,2} \cap C^{\infty}(\bar{\Omega}, S^2)} \int_{\Omega} |\nabla u|^2 dx = \min_{u \in W_{\gamma}^{1,2}(\Omega, S^2)} F(u).$$

Moreover, each minimizer of F in $W_{\gamma}^{1,2}(\Omega; S^2)$ is also a weak harmonic map.

There is an interesting problem whether a minimizer of the relaxed energy F in (1.2) is regular. Giaquinta, Modica and Soucek in [10] proved that a minimizer u of the relaxed energy F in $W_{\gamma}^{1,2}(\Omega; S^2)$ is smooth in a set $\Omega_0 \subset \Omega$ and $\mathcal{H}^1(\Omega \setminus \Omega_0) < \infty$, where \mathcal{H}^1 is the Hausdorff measure. Bethuel and Brezis in [3] proved that minimizers for a modified relax energy $F_{\lambda}(u) = \int_{\Omega} |\nabla u|^2 dx + \lambda 8\pi L(u)$ with $0 \leq \lambda < 1$ in $W_{\gamma}^{1,2}(\Omega; S^2)$ have at most isolated singularities.

In the first part of this paper, we propose a new approximation for the relaxed energy F of the Dirichlet energy. More precisely, for a parameter ε , we consider the family of functionals

$$(1.3) \quad E_{\varepsilon}(u; \Omega) = \int_{\Omega} (|\nabla u|^2 + \varepsilon^2 |\nabla u|^4) dx$$

for maps $u \in W^{1,4}$. Of course, there is a minimizer u_{ε} of E_{ε} in $W_{\gamma}^{1,4}(\Omega; S^2)$; i.e.,

$$E_{\varepsilon}(u_{\varepsilon}; \Omega) \leq E_{\varepsilon}(v; \Omega); \quad \forall v \in W_{\gamma}^{1,4}(\Omega; S^2)$$

and it can be proved that each u_{ε} is smooth. We shall prove the following.

Theorem 1.1. *For $\varepsilon > 0$, let u_{ε} be a minimizer of E_{ε} in $W_{\gamma}^{1,4}(\Omega; S^2)$. As $\varepsilon_k \rightarrow 0$, minimizers u_{ε_k} weakly converge (possibly passing subsequence) to a minimizer u of the relaxed energy F in $W_{\gamma}^{1,2}(\Omega; S^2)$. Moreover, u is harmonic and smooth in an open subset $\Omega_0 \subset \Omega$ with $\mathcal{H}_{loc}^1(\Omega \setminus \Omega_0) < +\infty$.*

During the proof of Theorem 1.1, we show that the limit map u is partial regular by a new approach, which is different from the one in [10], and does not use the concept of Cartesian currents.

Giaquinta, Modica and Soucek ([10],[11]) viewed the relaxed energy functional in terms of Cartesian currents. Let γ be a smooth function defined in a neighborhood $\tilde{\Omega}$ of $\bar{\Omega}$ with values in S^2 . Set

$$\text{cart}_{\gamma}^{2,1}(\tilde{\Omega} \times S^2) := \{T \in \text{cart}^{2,1}(\tilde{\Omega} \times S^2) \mid (T - G_{\gamma})_{\llcorner}(\tilde{\Omega} \setminus \bar{\Omega}) = 0\}.$$

It was shown in [10] that for any $u \in W_{\gamma}^{1,2}(\tilde{\Omega}, S^2)$, there is a 1-dimensional integer rectifiable current L_u of minimal mass among the integer rectifiable 1-dimensional currents L with $\text{spt} L_u \subset \tilde{\Omega}$ such that $-\partial L = P(u)$ and

$$\mathbf{M}(L_u) = L(u),$$

where $\mathbf{M}(L_u)$ is the mass of the current L_u . Then

$$\mathcal{D}(T_u; \tilde{\Omega}) = \int_{\tilde{\Omega}} |\nabla u|^2 dx + 8\pi \mathbf{M}(L_u) = F(u; \tilde{\Omega}).$$

For each $L = \tau(\mathcal{L}, \theta, \vec{L})$, we denote by $e(T)$ the energy density of the current $T := G_u + L \times [[S^2]]$; i.e.,

$$e(T) := |\nabla u|^2 dx + 8\pi\theta\mathcal{H}^1 \llcorner \mathcal{L}.$$

Then, in the spirit of [11] we improve Theorem 1.1 as:

Theorem 1.2. *For a parameter $\varepsilon > 0$, let u_ε be a minimizer of E_ε in $H_\gamma^{1,4}(\Omega, S^2)$ as in Theorem 1.1. Then, there is a sequence $\varepsilon_k \rightarrow 0$ such that*

$$G_{u_{\varepsilon_k}} \rightharpoonup T_u = G_u + L_u \times [[S^2]]$$

weakly in $\text{cart}_\gamma^{2,1}(\tilde{\Omega} \times S^2)$ with $L_u = \tau(\mathcal{L}, \theta, \vec{L})$, G_u denotes the graph current of u in Ω ,

$$\mathcal{D}(u_{\varepsilon_k}, \tilde{\Omega}) \rightarrow D(T_u, \tilde{\Omega} \times S^2)$$

and

$$|\nabla u_{\varepsilon_k}|^2 dx \rightharpoonup |\nabla u|^2 dx + 8\pi\theta\mathcal{H}^1 \llcorner \mathcal{L}$$

in the sense of Radon measures. Moreover, $u_{\varepsilon_k} \rightarrow u$ strongly in $W^{1,2}(\Omega_0, S^2)$, where Ω_0 is the open set in Theorem 1.1.

Furthermore, Theorem 1.1 can be generalized to the case of maps between two Riemannian manifolds. Let \mathcal{M} be a Riemannian manifold with boundary $\partial\mathcal{M}$ and \mathcal{N} another compact Riemannian manifold without boundary. Let γ be a smooth map from $\partial\mathcal{M}$ to \mathcal{N} which can be extended to a smooth map u_0 from \mathcal{M} to \mathcal{N} . For a map $u : \mathcal{M} \rightarrow \mathcal{N}$, we consider the Dirichlet energy $E_{\mathcal{M}}(u) = \int_{\mathcal{M}} |\nabla u|^2 d\mathcal{M}$. Then, we can define the relaxed energy for the Dirichlet energy $E_{\mathcal{M}}$ by

$$\tilde{E}_{\mathcal{M}}(u) := \inf \left\{ \liminf_{k \rightarrow \infty} E_{\mathcal{M}}(u_k) \mid \{u_k\} \subset C_\gamma^\infty(\mathcal{M}, \mathcal{N}), u_k \rightharpoonup u \text{ weakly in } W^{1,2}(\mathcal{M}, \mathcal{N}) \right\}$$

Under certain topological conditions on the manifold \mathcal{N} , Giaquinta-Mucci [12] found a representation formula of $\tilde{E}_{\mathcal{M}}(u)$. Without using the explicit formula $\tilde{E}_{\mathcal{M}}(u)$ or any topological assumption on \mathcal{N} , we have

Theorem 1.3. *For a parameter $\varepsilon > 0$, let u_ε be a minimizer of*

$$E_\varepsilon(u; \mathcal{M}) = \int_{\mathcal{M}} |\nabla u|^2 + \varepsilon^2 |\nabla u|^{n+1} d\mathcal{M}$$

in $W_\gamma^{1,n+1}(\mathcal{M}, \mathcal{N})$. As $\varepsilon_k \rightarrow 0$, minimizers u_{ε_k} weakly converge (possibly passing subsequence) to a minimizer u of the relaxed energy $\tilde{E}_{\mathcal{M}}$ in $W_\gamma^{1,2}(\mathcal{M}, \mathcal{N})$. Moreover, assuming that \mathcal{N} is a homogenous manifold, u is harmonic and smooth in an open subset $\mathcal{M}_0 \subset \mathcal{M}$ with $\mathcal{H}_{loc}^{n-2}(\mathcal{M} \setminus \mathcal{M}_0) < +\infty$.

After we finished an early version of our paper, Fanghua Lin informed us that a result related to Theorem 1.3 was studied by him in [17], but his proof is different from ours.

In the second part of this paper, we apply the same approximation method to the study of the relaxed energy for the Faddeev model. The Faddeev theory has created a lot of interest in physics as a successful effective field theory ([8]). The underlying idea is very old for, in the 19th century, Lord Kelvin proposed that atoms could be described as knotted vortex tubes in the ether. Modern physics replaces the ether by ‘fields’ of which the classical gravitational and electromagnetic fields are the most studied. It is known from work in condensed matter theory that field theories (such as sigma models) are useful in understanding the behaviour of composite

particles. Thus Faddeev [7] in 1979 proposed that closed, knotted vortices could be constructed in a definite dynamical system; namely the Faddeev model. The study of the dynamics of knots as the solution configurations of suitable Lagrangian field-theory equations has only been brought to light by Faddeev and Niemi in [9]. They employed powerful numerical algorithms to show that a ring-shaped charge one soliton exists. In mathematics, the Faddeev energy for a map u from \mathbb{R}^3 to S^2 is given by

$$E_F(u) = \int_{\mathbb{R}^3} [|\nabla u|^2 + \frac{1}{2} \sum_{1 \leq k < l \leq 3} |\partial_k u \times \partial_l u|^2] dx.$$

For any interesting configuration in physics, it is required that u converges to a constant sufficiently fast near the infinity. Under this assumption, we may compactify \mathbb{R}^3 by adding a point representing the infinity and view u to be a map from S^3 to S^2 . Hence any smooth field configuration u can be characterized by the topological charge $Q(u)$ given by the Hopf degree of u from \mathbb{R}^3 to S^2 . It is well known that the Hopf degree can be expressed analytically as

$$Q(u) = \frac{1}{16\pi^2} \int_{\mathbb{R}^3} \eta \wedge u^*(\omega_{S^2}),$$

where ω_{S^2} is a volume form on S^2 and $d\eta = u^*\omega_{S^2}$. Such an η exists because $u^*\omega_{S^2}$ is closed. Moreover, one can see that $Q(u)$ is independent of the choice of η . In this paper, unless stated otherwise, we will always take

$$(1.4) \quad \eta = \delta\left(-\frac{1}{4\pi|x|} \star u^*\omega_{S^2}\right).$$

Here \star means convolution and δ is the formal adjoint of d .

Vakulenko and Kapitanski in [23] found a lower-bound for the Faddeev energy; i.e. there is a constant $C > 0$ such that

$$E_F(u) \geq CQ(u)^{3/4}.$$

A minimizer of E_F among all maps with the same Hopf degree is called a hopfion, or a Hopf soliton. For each $d \in \mathbb{Z}$, Faddeev [8] suggested that there is a knotted minimizer of E_F in H_d , where H_d is the class of smooth maps $u : \mathbb{R}^3 \rightarrow S^2$ which have bounded energy and approach a constant value sufficiently fast as required by physics at the infinity (for a more precise definition see Section 3). Many contributions have been made by physicists ([21]). A natural functional space for the minimization of Faddeev energy is

$$X = \{u : \mathbb{R}^3 \rightarrow S^2 \mid E_F(u) < +\infty\}.$$

Lin and Yang ([19], [20]) showed that for infinitely many integers d 's, including the case $d = \pm 1$, there is a minimizer of E_F in X_d , where

$$X_d = \{u \in X \mid Q(u) = d\}.$$

(See [13] for further development). However, it is still unknown whether the minimizers obtained by Lin-Yang [19] are smooth or not. Therefore, there might occur a gap phenomenon for the Faddeev energy similarly to one for harmonic maps; i.e.

$$(1.5) \quad \inf_{u \in X_d} E_F(u) < \inf_{u \in H_d} E_F(u).$$

A similar situation occurs in the Skyrme model as pointed out by Esteban and Müller [5].

It is a very challenging problem whether the infimum of Faddeev energy on the righthand of (1.5) can be achieved or not. We consider the relaxed energy for $u \in X_d$,

$$\tilde{E}_F(u) = \inf \left\{ \liminf_{k \rightarrow \infty} E_F(u_k) \mid \{u_k\} \subset H_d, u_k \rightharpoonup u \text{ weakly} \right\},$$

where ‘weakly converging’ means in the sense of bounded Faddeev energy. For a map $u \in X_d$ without any sequence u_k converging weakly to u , we take the value $\tilde{E}_F(u)$ to be $+\infty$.

However, due to the complexity of the Faddeev energy, this time we do not have an explicit formula as, for instance, in [4] and [11]. By using the same approximation method as previously, we will find a minimizer for the relaxed energy \tilde{E}_F in some cases. More precisely, we have

Theorem 1.4. *For $d = \pm 1$, there is a minimizer of the relaxed energy \tilde{E}_F in X_d . Moreover,*

$$(1.6) \quad \inf_{u \in H_{\pm 1}} E_F(u) = \min_{u \in X_{\pm 1}} \tilde{E}_F(u).$$

Our proofs are based on the ideas of Lin-Yang in [19]. We will introduce a perturbed energy $E_{F,\varepsilon}$ with a parameter ε . It turns out that the analysis involved in the perturbed variational problem for $\varepsilon > 0$ is much easier to understand. Modifying an idea of Ward in [24] and Lin-Yang in [20], we show that the new minimizing problem has a solution for $d = \pm 1$. Then, letting ε go to zero, we prove that the minimizers of the perturbed energy $E_{F,\varepsilon}$ in $X_{\pm 1}$ will converge to a minimizer of the relax energy \tilde{E}_F in $X_{\pm 1}$.

2. RELAXED ENERGY FOR HARMONIC MAPS

Lemma 2.1. *Let u_ε be a minimizer of the functional E_ε . Then, for all $0 < \rho \leq R$ with $B_R(x_0) \subset \Omega$, we have*

$$\begin{aligned} & R^{-1} \int_{B_R(x_0)} [|\nabla u_\varepsilon|^2 + \varepsilon^2 |\nabla u_\varepsilon|^4] dx - \rho^{-1} \int_{B_\rho(x_0)} [|\nabla u_\varepsilon|^2 + \varepsilon^2 |\nabla u_\varepsilon|^4] dx \\ &= \int_{B_R \setminus B_\rho(x_0)} [1 + 2\varepsilon^2 |\nabla u_\varepsilon|^2] |\partial_r u_\varepsilon|^2 r^{-1} dx - \int_\rho^R \int_{B_r(x_0)} 2\varepsilon^2 |\nabla u_\varepsilon|^4(y) dy r^{-2} dr. \end{aligned}$$

Proof. Without loss of generality, we assume $x_0 = 0$. Let $\phi(x) = (\phi^1(x), \dots, \phi^n(x)) \in C^1(\Omega; \mathbb{R}^n)$ be a vector field having compact support in Ω . For a function $\phi_t(x) = x + t\phi(x)$ with $x = (x^1, \dots, x^n)$ and for a function $u(x)$, set $u^{t,\phi}(x) := u_\varepsilon(x + t\phi(x))$.

We see $\nabla_{x^i} u^{t,\phi}(x) = u_{x^k}(x + t\phi)[\delta_{ik} + t\phi_{x^i}^k]$. Since u_ε is a minimizer of E_ε , it follows from $\frac{d}{dt} E_\varepsilon(u^{t,\phi}, \Omega)|_{t=0} = 0$ that

$$\int_\Omega (|\nabla u_\varepsilon|^2 + \varepsilon^2 |\nabla u_\varepsilon|^4) \operatorname{div} \phi - 2u_{\varepsilon x^i} u_{\varepsilon x^k} \phi_{x^i}^k (1 + 2\varepsilon^2 |\nabla u_\varepsilon|^2) dx = 0.$$

For a given ball $B_r(0) \subset \Omega$, we choose $\phi(x) = \xi(|x|)x$ with

$$\xi(s) = \begin{cases} 1 & \text{for } s \leq r \\ 1 + \frac{r-s}{h} & \text{for } r \leq s \leq r+h \\ 0 & \text{for } s \geq r+h. \end{cases}$$

Letting $h \rightarrow 0$, we obtain that for almost every r

$$\begin{aligned} & \int_{B_r} [|\nabla u_\varepsilon|^2 + \varepsilon^2 |\nabla u_\varepsilon|^4] dx - r \int_{\partial B_r} [|\nabla u_\varepsilon|^2 + \varepsilon^2 |\nabla u_\varepsilon|^4] dH^{n-1} \\ &= -\frac{2}{r} \int_{\partial B_r} (1 + 2\varepsilon^2 |\nabla u_\varepsilon|^2) |x^i \partial_{x^i} u_\varepsilon|^2 dH^{n-1} + \int_{B_r} 2\varepsilon^2 |\nabla u_\varepsilon|^4 dx. \end{aligned}$$

Multiplying by r^{-2} both sides of the above identity and integrating with respect to r from ρ to R yields the result. \square

We now complete the proof of Theorem 1.1 and Theorem 1.2.

Proof of Theorem 1.1. Let u_ε be a minimizer of the functional E_ε in $W_\gamma^{1,4}(\Omega; S^2)$ and u the weak limit of the sequence in $W^{1,2}$. Since F is lower semi-continuous, we have

$$(2.1) \quad F(u; \Omega) \leq \liminf_{\varepsilon_k \rightarrow 0} \int_{\Omega} |\nabla u_{\varepsilon_k}|^2 dx \leq \liminf_{\varepsilon_k \rightarrow 0} \int_{\Omega} |\nabla u_{\varepsilon_k}|^2 + \varepsilon_k^2 |\nabla u_{\varepsilon_k}|^4 dx$$

$$(2.2) \quad \leq \inf_{v \in C_\gamma^\infty(\Omega; S^2)} \liminf_{\varepsilon_k \rightarrow 0} \int_{\Omega} |\nabla v|^2 + \varepsilon_k^2 |\nabla v|^4 dx = \inf_{v \in C_\gamma^\infty(\Omega; S^2)} \int_{\Omega} |\nabla v|^2 dx.$$

Using (1.2), we note

$$\inf_{v \in C_\gamma^\infty(\Omega; S^2)} \int_{\Omega} |\nabla v|^2 dx = \min_{v \in W_\gamma^{1,2}(\Omega; S^2)} F(v).$$

Then it follows from (2.1) and (2.2) that u is a minimizer of the relaxed energy F in $W^{1,2}(\Omega; S^2)$. Moreover, we have

$$(2.3) \quad \lim_{\varepsilon_k \rightarrow 0} \int_{\Omega} \varepsilon_k^2 |\nabla u_{\varepsilon_k}|^4 dx = 0.$$

We define

$$\Sigma = \bigcap_{R>0} \left\{ x_0 \in \Omega : B_R(x_0) \subset \Omega, \quad \liminf_{\varepsilon_k \rightarrow 0} \frac{1}{R} \int_{B_R(x_0)} |\nabla u_\varepsilon|^2 dx \geq \varepsilon_0 \right\}$$

for a sufficiently small constant ε_0 to be fixed later.

As in [22], we can show that Σ is relatively closed inside Ω . In fact, let x_j be a sequence in Σ and x_j goes to $x \in \Omega$. To see that $x \in \Sigma$, we need to show for all $R > 0$,

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{R} \int_{B_R(x)} |\nabla u_\varepsilon|^2 dx \geq \varepsilon_0.$$

Let $r < R$ (but very close). Since $x_j \in \Sigma$,

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{r} \int_{B_r(x_j)} |\nabla u_\varepsilon|^2 dx \geq \varepsilon_0.$$

Choose some large j so that $B_r(x_j) \subset B_R(x)$, then

$$\frac{1}{r} \int_{B_r(x_j)} |\nabla u_\varepsilon|^2 dx \leq \frac{R}{r} \frac{1}{R} \int_{B_R(x)} |\nabla u_\varepsilon|^2 dx.$$

Hence,

$$\frac{R}{r} \liminf_{\varepsilon_k \rightarrow 0} \frac{1}{R} \int_{B_R(x)} |\nabla u_\varepsilon|^2 dx \geq \varepsilon_0.$$

Since r can be chosen very close to R ,

$$\liminf_{\varepsilon_k \rightarrow 0} \frac{1}{R} \int_{B_R(x)} |\nabla u_\varepsilon|^2 dx \geq \varepsilon_0.$$

This is true for all $R > 0$. Therefore, $x \in \Sigma$ and Σ is closed. The same argument of Schoen [22] yields $\mathcal{H}_{\text{loc}}^1(\Sigma) < +\infty$.

A consequence is that for each $x_0 \notin \Sigma$, there are a sequence $\varepsilon_k \rightarrow 0$ and a ball $B_{R_0}(x_0) \subset \Omega \setminus \Sigma$ such that

$$\lim_{\varepsilon_k \rightarrow 0} R_0^{-1} \int_{B_{R_0}(x_0)} |\nabla u_{\varepsilon_k}|^2 dx < \varepsilon_0.$$

Now, we prove that u is smooth around such points x_0 . In fact, for each $y \in B_{R_0/2}(x_0)$, trivially

$$\lim_{\varepsilon_k \rightarrow 0} \frac{2}{R_0} \int_{B_{R_0/2}(y)} |\nabla u_{\varepsilon_k}|^2 dx \leq \lim_{\varepsilon_k \rightarrow 0} 2 \frac{1}{R} \int_{B_{R_0}(x_0)} |\nabla u_{\varepsilon_k}|^2 dx < 2\varepsilon_0,$$

while, applying (2.3) to Lemma 2.1, for any $0 < \rho < R_0/2$, we have

$$\begin{aligned} & \lim_{\varepsilon_k \rightarrow 0} \frac{2}{R_0} \int_{B_{R_0/2}(y)} |\nabla u_{\varepsilon_k}|^2 dx - \lim_{\varepsilon_k \rightarrow 0} \rho^{-1} \int_{B_\rho(y)} |\nabla u_{\varepsilon_k}|^2 dx \\ &= \lim_{\varepsilon_k \rightarrow 0} \int_{B_{R_0/2} \setminus B_\rho(y)} [|\partial_r u_{\varepsilon_k}|^2 + \varepsilon_k^2 |\nabla u_{\varepsilon_k}|^2 |\partial_r u_{\varepsilon_k}|^2] r^{-1} dx \geq 0. \end{aligned}$$

Therefore, for each $y \in B_{R_0/2}(x_0)$ and for each $\rho \in (0, R_0/2)$, we have

$$(2.4) \quad \frac{1}{\rho} \int_{B_\rho(y)} |\nabla u|^2 dx \leq \lim_{\varepsilon_k \rightarrow 0} \rho^{-1} \int_{B_\rho(y)} |\nabla u_{\varepsilon_k}|^2 dx < 2\varepsilon_0$$

for a sufficiently small constant $\varepsilon_0 > 0$.

Since u is a weak harmonic map with the property (2.4), it follows, similarly to the proof in [6] (see also Lemma 3.3.13 of [18]) for stationary harmonic maps, that u is smooth inside $B_{R_0/2}(x_0)$. This proves Theorem 1.1. \square

Proof of Theorem 1.2. Let T belong to $\text{cart}^{2,1}(\Omega \times S^2)$, i.e.

$$T = G_{u_T} + L_T \times [[S^2]].$$

The Dirichlet integral of T is given by

$$D(T; \Omega \times S^2) = \int_{\Omega} |\nabla u_T|^2 dx + 8\pi M(L_T).$$

As $\varepsilon_k \rightarrow 0$, passing to a subsequence we have

$$e(G_{u_{\varepsilon_k}}) \rightharpoonup \mu_0$$

for a measure μ_0 . For any $\psi \in C_c^0(\tilde{\Omega})$ with $\psi \geq 0$, we consider the functional in $\text{cart}_\gamma^{2,1}(\tilde{\Omega} \times S^2)$

$$\mathcal{E}(T) := \int \psi(x) e(T).$$

We know that \mathcal{E} is lower semi-continuous with respect to the weak convergence in $\text{cart}^{2,1}$ which implies

$$e(T)(\psi) \leq \liminf_{k \rightarrow \infty} e(G_{u_{\varepsilon_k}})(\psi) \leq \mu_0(\psi).$$

This means $e(T) \leq \mu_0$. By Theorem 1.1, we have $\mathcal{D}(u_k, \Omega) \rightarrow \mathcal{D}(T; \Omega \times S^2)$. Then, we have

$$\mu_0(\Omega) = e(T)(\Omega).$$

This shows $\mu_0 = e(T)$ as we claimed.

On the other hand, if U be a subset inside $\Omega \setminus \Omega_0$, according to [11] p. 436, we can prove $L_T \lrcorner U = 0$. Therefore, we have

$$T_u|_U = G_u|_U.$$

Therefore

$$\int_U |\nabla u_{\varepsilon_k}|^2 dx \rightarrow \int_U |\nabla u_{\varepsilon_k}|^2 dx,$$

and this implies $u_k \rightarrow u$ strongly in $W_{\text{loc}}^{1,2}(\Omega \setminus \bar{\Omega}_0; S^2)$. \square

Proof of Theorem 1.3. It is well-known that $C_\gamma^\infty(\mathcal{M}, \mathcal{N})$ is dense in $W_\gamma^{1,n+1}(\mathcal{M}, \mathcal{N})$. Then

$$\begin{aligned} \tilde{E}_\mathcal{M}(u) &= \inf \left\{ \liminf_{k \rightarrow \infty} E_\mathcal{M}(u_k) \mid \{u_k\} \subset C_\gamma^\infty(\mathcal{M}, \mathcal{N}), u_k \rightharpoonup u \text{ weakly in } W^{1,2}(\mathcal{M}, \mathcal{N}) \right\} \\ &= \inf \left\{ \liminf_{k \rightarrow \infty} E_\mathcal{M}(u_k) \mid \{u_k\} \subset W_\gamma^{1,n+1}(\mathcal{M}, \mathcal{N}), u_k \rightharpoonup u \text{ weakly in } W^{1,2}(\mathcal{M}, \mathcal{N}) \right\} \end{aligned}$$

Let u_ε be a minimizer of the functional $E_\varepsilon(\cdot; \mathcal{M})$ in $W_\gamma^{1,n+1}(\mathcal{M}, \mathcal{N})$. Passing to a subsequence, $\nabla_k u_\varepsilon$ converges to $\nabla_k u$ weakly in L^2 . Then, we have

$$(2.5) \quad \tilde{E}_\mathcal{M}(u) \leq \liminf_{\varepsilon_k \rightarrow 0} E_\mathcal{M}(u_{\varepsilon_k}) \leq \liminf_{\varepsilon_k \rightarrow 0} E_{\varepsilon_k}(u_{\varepsilon_k}; \mathcal{M})$$

$$(2.6) \quad \leq \inf_{v \in W_\gamma^{1,n+1}(\mathcal{M}, \mathcal{N})} E_\mathcal{M}(v) = \inf_{v \in C_\gamma^\infty(\mathcal{M}, \mathcal{N})} E_\mathcal{M}(v).$$

From (2.5)-(2.6), u is a minimizer of the relaxed energy $\tilde{E}_\mathcal{M}$ in $H^{1,2}$ and moreover, we have

$$(2.7) \quad \lim_{\varepsilon_k \rightarrow 0} \int_\mathcal{M} \varepsilon_k^2 |\nabla u_{\varepsilon_k}|^{1+n} d\mathcal{M} = 0.$$

If \mathcal{N} is a homogeneous manifold, we claim that u is a weak harmonic map.

Let X_i be the Killing field on \mathcal{N} as in Hélein [15]. Consider the vector field

$$\xi_i = \langle X_i, \nabla u_\varepsilon \rangle + 2\varepsilon^2 |\nabla u_\varepsilon|^{n-1} \langle X_i, \nabla u_\varepsilon \rangle.$$

We claim that $\text{div} \xi_i = 0$.

To see this, let φ be some cut-off function compactly supported in Ω . Since u_ε is the minimizer of E_ε , we can use $\varphi X_i(u)$ as a testing vector field in the Euler-Lagrange equation to get

$$\int_\mathcal{M} \langle \nabla_k(\varphi X_i(u_\varepsilon)), \nabla_k u_\varepsilon + 2\varepsilon^2 |\nabla u_\varepsilon|^{n-1} \nabla_k u_\varepsilon \rangle d\mu = 0.$$

Since X_i is a Killing vector field,

$$(2.8) \quad \int_\mathcal{M} \nabla_k \varphi \langle X_i(u_\varepsilon), \nabla_k u_\varepsilon + 2\varepsilon^2 |\nabla u_\varepsilon|^{n-1} \nabla_k u_\varepsilon \rangle d\mu = 0.$$

Therefore, $\text{div} \xi = 0$ in distribution sense.

Since u_{ε_k} converges to u strongly in L^2 and X_i are smooth vector fields on \mathcal{N} , $X_i(u_{\varepsilon_k})$ converges to $X_i(u)$ strongly in L^2 . Letting ε_k go to zero in equation (2.8) and noting (2.7), we have

$$\int_{\mathcal{M}} \nabla_k \varphi \langle X_i(u), \nabla_k u \rangle d\mu = 0.$$

Since X_i is Killing field,

$$\int_{\mathcal{M}} \langle \nabla_k(\varphi X_i(u)), \nabla_k u \rangle d\mu = 0.$$

Since N is a homogeneous space, due to the construction of X_i by Helein, we can choose φ_i so that

$$\sum_i \varphi_i X_i(u)$$

is any compactly supported vector field (along u). This implies that u is a weak harmonic map.

We define

$$\Sigma = \bigcap_{R>0} \left\{ x_0 \in \Omega : B_R(x_0) \subset \mathcal{M}, \liminf_{\varepsilon_k \rightarrow 0} \frac{1}{R^{n-2}} \int_{B_R(x_0)} |\nabla u_{\varepsilon_k}|^2 dx \geq \varepsilon_0 \right\}$$

for a sufficiently small constant ε_0 . Then, $\mathcal{H}^{n-2}(\Sigma) < +\infty$. As in the proof of Theorem 1.1, for any $x_0 \notin \Sigma$ with $B_{R_0}(x_0) \subset \mathcal{M} \setminus \Sigma$, for each $y \in B_{R_0/2}(x_0)$ and for each $\rho \in (0, R_0/2)$, we have

$$(2.9) \quad \frac{1}{\rho^{n-2}} \int_{B_\rho(y)} |\nabla u|^2 d\mathcal{M} \leq \lim_{\varepsilon_k \rightarrow 0} \rho^{2-n} \int_{B_\rho(y)} |\nabla u_{\varepsilon_k}|^2 d\mathcal{M} < 2\varepsilon_0$$

for a sufficiently small constant $\varepsilon_0 > 0$. It follows from the proof of Bethuel in [2] (see Lemma 3.3.14 of [18]) that u is smooth inside $\mathcal{M} \setminus \Sigma$. \square

3. BASIC SET UP FOR THE RELAXED ENERGY OF FADDEEV MODEL

The space H_d mentioned in the introduction is not rigorously defined since we have not made clear what ‘ u approaches a constant value sufficiently fast at infinity’ means. So the first task of this section is to propose and justify a replacement for H_d . We define

$$M = \{u : \mathbb{R}^3 \rightarrow S^2 \mid \int_{\mathbb{R}^3} |\nabla u|^2 + |\nabla u|^4 dx < +\infty\}.$$

Since $|\partial_k u \times \partial_l u|^2$ involves the ∇u up to the fourth power, M is only a little smaller than the set of finite energy maps. On the other hand, Theorem 3.1 below implies that the set of smooth maps which are constant outside some large compact set is dense in M . Remember whether this set is dense in the set of finite energy maps is an open question and this is the main motivation of our discussion of relaxed energy.

Theorem 3.1. *Let $u : \mathbb{R}^3 \rightarrow S^2$ be a map such that $E_F(u)$ and $\int_{\mathbb{R}^3} |\nabla u|^4 d\mu$ are finite. Then, there exists a sequence of C^1 maps $u_i : \mathbb{R}^3 \rightarrow S^2$ with u_i constant outside some large ball (depending on i) such that ∇u_i converges to ∇u strongly in $L^4(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$.*

Before proving Theorem 3.1, we discuss some immediate consequences of it.

Corollary 3.2. *For each $u \in M$, $Q(u)$ is an integer.*

Proof. Let u_i be the sequence in Theorem 3.1. $Q(u_i)$ is an integer since u_i is of class C^1 and constant at infinity.

$$Q(u_i) = \frac{1}{16\pi^2} \int_{\mathbb{R}^3} \eta_i \wedge u_i^* \omega_{S^2}.$$

By the definition of η_i (see equation (1.4)) and the fact that $\nabla u_i \rightarrow \nabla u$ strongly in $L^4(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ (which implies $u_i^* \omega_{S^2} \rightarrow u^* \omega_{S^2}$ strongly in $L^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$), η_i converges to η in $W^{1,p}$ for $1 < p \leq 2$. In particular, η_i converges strongly to η in $L^2(\mathbb{R}^3)$. Hence

$$\lim_{i \rightarrow \infty} Q(u_i) = Q(u),$$

which implies $Q(u) \in \mathbb{Z}$. □

For any integer d , we define

$$M_d = \{u \in M \mid Q(u) = d\}.$$

As remarked earlier in this section, M is only a little smaller than the set of finite energy maps. We *may assume* $H_d \subset M_d$ and H_d contains all smooth maps which are constant outside some compact set. Another consequence of Theorem 3.1 is

$$(3.1) \quad \inf_{u \in H_d} E_F(u) = \inf_{u \in M_d} E_F(u).$$

In fact, since $H_d \subset M_d$, we know

$$\inf_{u \in H_d} E_F(u) \geq \inf_{u \in M_d} E_F(u).$$

On the other hand, for any $u \in M_d$, due to Theorem 3.1, there exists a sequence of $u_i \in H_d$ such that

$$E_F(u_i) \rightarrow E_F(u).$$

Hence, the reverse inequality is also true. This proves our claim.

Moreover, according to the above, the definition of relaxed energy given in the introduction can be rewritten as

$$\tilde{E}_F(u) = \inf \left\{ \liminf_{k \rightarrow \infty} E_F(u_k) \mid \{u_k\} \subset M_d, u_k \rightharpoonup u \text{ weakly} \right\}.$$

The proof of Theorem 3.1 depends on the following elliptic estimates.

Let B_r be the ball with center at 0 and radius r in \mathbb{R}^3 , let u be a map from B_1 to S^2 such that

$$\int_{\partial B_1} |\nabla u|^2 + R^{-2} |\nabla u|^4 d\sigma = \epsilon,$$

for some positive R (later R will be large and ϵ small), and let ξ be the average of u over ∂B_1 , i.e.

$$\xi = \frac{1}{4\pi} \int_{\partial B_1} u d\sigma.$$

By the Poincaré inequality and using that $|u| = 1$, we have

$$(3.2) \quad 1 - |\xi| \leq 1 - |\xi|^2 = \frac{1}{4\pi} \int_{\partial B_1} |u - \xi|^2 d\sigma \leq C\epsilon.$$

As in [19], consider the harmonic function $V : B_2 \setminus B_1 \rightarrow \mathbb{R}^3$ such that

$$\Delta V = 0, \quad \text{in } B_2 \setminus B_1,$$

and

$$V|_{\partial B_2} = \frac{\xi}{|\xi|}, \quad V|_{\partial B_1} = u.$$

It follows from (3.2) that

$$\|V - \xi\|_{W^{1,4}(\partial B_2)} \leq C(1 - |\xi|) \leq C\epsilon.$$

The average of $V - \xi$ on ∂B_1 is zero. By the Poincaré inequality,

$$\|u - \xi\|_{W^{1,4}(\partial B_1)} \leq C \|\nabla u\|_{L^4(\partial B_1)}.$$

By the trace theorem of the Sobolev space, we find $w : B_2 \setminus B_1 \rightarrow \mathbb{R}^3$ with the same boundary value as $V - \xi$ on $\partial B_2 \cup \partial B_1$ such that

$$\|w\|_{W^{5/4,4}(B_2 \setminus B_1)} \leq C \|V - \xi\|_{W^{1,4}(\partial B_1 \cup \partial B_2)} \leq C(\|\nabla u\|_{L^4(\partial B_1)} + \epsilon).$$

Then, $V - \xi - w$ has zero boundary value and

$$\Delta(V - \xi - w) = -\Delta w.$$

By the elliptic estimate, we have

$$\|V - \xi - w\|_{W^{5/4,4}(B_2 \setminus B_1)} \leq C(\|\nabla u\|_{L^4(\partial B_1)} + \epsilon).$$

Hence,

$$\|V - \xi\|_{W^{5/4,4}(B_2 \setminus B_1)} \leq C(\|\nabla u\|_{L^4(\partial B_1)} + \epsilon).$$

The following is a Sobolev's embedding theorem, which is a special case of Theorem 7.58 in Adams' book [1]. For reader's convenience, we quote it here.

Theorem 3.3. *Let $s > 0$, $1 < p < q < +\infty$, and $\chi = s - \frac{n}{p} + \frac{n}{q} > 0$. Then*

$$W^{s,p}(\mathbb{R}^n) \rightarrow W^{\chi,q}(\mathbb{R}^n).$$

The result holds true for domains with smooth boundary.

Using the above theorem in the case $s = 5/4$, $p = 4$, $\chi = 1$ and $q = 6$,

$$\|V - \xi\|_{W^{1,6}(B_2 \setminus B_1)} \leq C(\|\nabla u\|_{L^4(\partial B_1)} + \epsilon).$$

In particular,

$$(3.3) \quad \int_{B_2 \setminus B_1} |\nabla V|^6 dx \leq C \left(\int_{\partial B(1)} |\nabla u|^4 d\sigma \right)^{3/2} + C\epsilon^6 < C\epsilon^{3/2}R^3 + C\epsilon^6 < C\epsilon^{3/2}R^3.$$

The last inequality holds because, in the form we need it, ϵ is small and R large. Applying the Hölder inequality, we have

$$(3.4) \quad \int_{B_2 \setminus B_1} |\nabla V|^4 dx \leq C\epsilon R^2.$$

A similar argument works for the L^2 norm of ∇u . Therefore, we conclude

$$(3.5) \quad \int_{B_2 \setminus B_1} |\nabla V|^2 + R^{-2} |\nabla V|^4 dx \leq C \int_{\partial B_1} |\nabla u|^2 + R^{-2} |\nabla u|^4 d\sigma.$$

With this estimate, we can now prove Theorem 3.1.

Proof of Theorem 3.1. Let u be a map from $\mathbb{R}^3 \rightarrow S^2$ such that

$$\int_{\mathbb{R}^3} |\nabla u|^2 + |\nabla u|^4 dx < C.$$

It suffices to show that for any $\varepsilon > 0$ there is a C^1 map w from \mathbb{R}^3 to S^2 such that

- (1) w is constant outside a large ball;
- (2)

$$\int_{\mathbb{R}^3} |\nabla(u-w)|^2 + |\nabla(u-w)|^4 dx < \varepsilon.$$

Since

$$\int_0^\infty \int_{\partial B_r} |\nabla u|^2 + |\nabla u|^4 d\sigma dr < C,$$

for any $C_1 > 0$ small, there exists a sequence of R_i going to infinity such that

$$\int_{\partial B_{R_i}} |\nabla u|^2 + |\nabla u|^4 d\sigma < \frac{C_1}{R_i}.$$

Let ξ_i be the average of u over ∂B_{R_i} ,

$$\xi_i = \frac{1}{|\partial B_{R_i}|} \int u d\sigma.$$

Then

$$\frac{1}{|B_{R_i}|} \int_{\partial B_{R_i}} |u - \xi_i|^2 \leq \int_{\partial B_{R_i}} |\nabla u|^2 d\sigma \rightarrow 0.$$

Define V to be the harmonic function defined on $B_{2R_i} \setminus B_{R_i}$ with boundary value

$$V|_{\partial B_{R_i}} = u, \quad V|_{\partial B_{2R_i}} = \frac{\xi_i}{|\xi_i|}.$$

Due to the estimate (3.5) and a scaling, we know

$$\int_{B_{2R_i} \setminus B_{R_i}} |\nabla V|^2 + |\nabla V|^4 dx \leq CR_i \int_{\partial B_{R_i}} |\nabla u|^2 + |\nabla u|^4 d\sigma < C_1.$$

For fixed ε , choose $C_1 < \varepsilon/96$ and R_i so large such that

$$\int_{\mathbb{R}^3 \setminus B_{R_i}} |\nabla u|^2 + |\nabla u|^4 dx < \varepsilon/6.$$

We then define \tilde{w} to be

- (1) u in B_{R_i} ;
- (2) $\frac{V}{|V|}$ in $B_{2R_i} \setminus B_{R_i}$;
- (3) $\frac{\xi_i}{|\xi_i|}$ outside B_{2R_i} .

Since, as it is proved in page 292 of [19] (see also Lemma 9.7 of [13]), the image of V lies in $B_{3/2} \setminus B_{1/2}$ for sufficiently large i , and the nearest-point-projection map restricted to $B_{3/2} \setminus B_{1/2}$ is Lipschitz and the Lipschitz constant is 2, by replacing \tilde{w} with its projection onto S^2 , we conclude that \tilde{w} has values in S^2 ,

$$\int_{B_{2R_i} \setminus B_{R_i}} |\nabla \tilde{w}|^2 + |\nabla \tilde{w}|^4 dx < \frac{\varepsilon}{6}$$

and

$$\int_{\mathbb{R}^3} |\nabla(\tilde{w} - u)|^2 + |\nabla(\tilde{w} - u)|^4 dx < \varepsilon/3.$$

Finally, since we can modify \tilde{w} to a smooth map w with the same properties of \tilde{w} in such a way that

$$\int_{\mathbb{R}^3} |\nabla(w - u)|^2 + |\nabla(w - u)|^4 dx < \varepsilon,$$

we conclude the proof. \square

Remark 3.4. *Theorem 3.1 holds true for $u : \mathbb{R}^3 \rightarrow S^3$ with finite energies $\int_{\mathbb{R}^3} |\nabla u|^4 dx$ and $\int_{\mathbb{R}^3} |\nabla u|^2 dx$, the proof being the same.*

4. HOPF LIFT AND DECOMPOSITION LEMMA

In this section, we first prove a theorem about the Hopf lift. Our proof depends on Theorem 3.1 in the previous section. Then, we use the Hopf lift to prove a key lemma, the cubic decomposition lemma, which will be essential in the proof of our main result.

Let us start from recalling some basic definitions of the Hopf lift (see [16] for details). Let \mathcal{M} be any complete Riemannian manifold whose second cohomology group $H^2(\mathcal{M}, \mathbb{R})$ is trivial. Let Π be the Hopf map from $S^3 \subset \mathbb{R}^4$ to $S^2 \subset \mathbb{R}^3$ given by

$$\Pi(x_1, x_2, x_3, x_4) = \begin{pmatrix} 2(x_1x_3 + x_2x_4) \\ 2(x_2x_3 - x_1x_4) \\ x_1^2 + x_2^2 - x_3^2 - x_4^2 \end{pmatrix}.$$

Pulling back the volume form ω_{S^2} via Π gives

$$\Pi^* \omega_{S^2} = 4(dx_1 dx_2 + dx_3 dx_4) = 2d\alpha,$$

where $\alpha = x_1 dx_2 - x_2 dx_1 + x_3 dx_4 - x_4 dx_3$. One can check by using Stokes theorem that

$$(4.1) \quad \frac{1}{16\pi^2} \int_{S^3} 2\alpha \wedge \Pi^* \omega_{S^2} = 1.$$

For any map $u : \mathcal{M} \rightarrow S^2$, a map \tilde{u} from \mathcal{M} to S^3 is called a Hopf lift of u if

$$\Pi \circ \tilde{u} = u.$$

Any smooth 1 form η on \mathcal{M} satisfying

$$d\eta = u^* \omega_{S^2}$$

is called a gauge of u . For a fixed u , the gauge η is not unique. However, if one fixes a Hopf lift \tilde{u} of u , then \tilde{u} determines a gauge for u by

$$\eta = 2u^* \alpha.$$

On the other hand, any gauge η of u determines a Hopf lift \tilde{u} such that the above equation is true and such a map \tilde{u} is unique up to multiplication by $e^{i\theta}$ for some constant $\theta \in \mathbb{R}$ and

$$(4.2) \quad |\nabla \tilde{u}|^2 = \frac{1}{4} |\eta|^2 + |\nabla u|^2.$$

See Lemma 2.1 of [16] for a proof.

We will generalize the above definition of Hopf lift and equation (4.2) to Sobolev mapping u with bounded Faddeev energy and finite $\int_{\mathbb{R}^3} |\nabla u|^4 dx$. The proof relies on Theorem 3.1. For such u , due to Theorem 3.1, there exists a sequence of smooth

maps $\{u_i\}$ such that u_i is constant near infinity and ∇u_i converges to ∇u strongly in both $L^2(\mathbb{R}^3)$ and $L^4(\mathbb{R}^3)$.

For each u_i , we define the Coulomb gauge η_i by requiring $\eta_i = \delta(-\frac{1}{4\pi|x|} \star u_i^* \omega_{S^2})$. Here \star means convolution and δ is the formal adjoint of d . One can check that $d\eta_i = u_i^* \omega_{S^2}$ and $\delta\eta_i = 0$. Given u_i and η_i , there exists a lift \bar{u}_i (called the Coulomb lift) such that

$$(4.3) \quad |\nabla \bar{u}_i|^2 = \frac{1}{4} |\eta_i|^2 + |\nabla u_i|^2.$$

Moreover,

$$(4.4) \quad \eta_i = \bar{u}_i^*(2\alpha)$$

(see [16] for a proof).

Since ∇u_i strongly converge to ∇u in $L^4(\mathbb{R}^3)$, we know that η_i converge to the Coulomb gauge η of u strongly in $W^{1,2}(\mathbb{R}^3)$. Due to the Sobolev embedding, η_i converge strongly in $L^2(\mathbb{R}^3)$ and $L^4(\mathbb{R}^3)$. By (4.3), we have

$$\int_{\mathbb{R}^3} |\nabla \bar{u}_i|^2 + |\nabla \bar{u}_i|^4 dx < C.$$

Since the image of \bar{u}_i is in S^3 , $u_i \in W_{loc}^{1,4}(\mathbb{R}^3)$. Hence, for each bounded domain Ω , we can take subsequence of $\nabla \bar{u}_i$ such that it converges weakly in L^2 and L^4 on Ω . Moreover, due to equation (4.3) and its square, we know the $L^2(\Omega)$ and $L^4(\Omega)$ norms of $\nabla \bar{u}_i$ also converge. Denote the limit by \bar{u} , then \bar{u}_i converges to \bar{u} strongly in $W_{loc}^{1,4}(\mathbb{R}^3)$. In particular,

$$\int_{\mathbb{R}^3} |\nabla \bar{u}|^4 dx < C.$$

By the definition of Hopf lift and (4.4), we have for each i

$$\bar{u}_i^*(2\alpha \wedge \Pi^* \omega_{S^2}) = \eta_i \wedge u_i^* \omega_{S^2}.$$

For any bounded domain Ω , we integrate the above identity over Ω and take the limit $i \rightarrow \infty$ to get

$$\int_{\Omega} \bar{u}^*(2\alpha \wedge \Pi^* \omega_{S^2}) = \int_{\Omega} \eta \wedge u^* \omega_{S^2}.$$

Since Ω is arbitrary, we have proved that

Theorem 4.1. *For each u with finite Faddeev energy and finite $\int_{\mathbb{R}^3} |\nabla u|^4 dx$, there exists a Hopf lift \bar{u} of u such that*

$$\frac{1}{16\pi^2} \int_{\mathbb{R}^3} \bar{u}^*(2\alpha \wedge \Pi^* \omega_{S^2}) = \frac{1}{16\pi^2} \int_{\mathbb{R}^3} \eta \wedge u^* \omega_{S^2} \in \mathbb{Z}.$$

Moreover,

$$\int_{\mathbb{R}^3} |\nabla \bar{u}|^4 dx \leq C.$$

Remark 4.2. *The lift given in the above theorem is a special one. In our later proof, we will need the fact that for any map $\bar{u} : \mathbb{R}^3 \rightarrow S^3$ with finite $\int_{\mathbb{R}^3} |\nabla \bar{u}|^4 dx$,*

$$\frac{1}{16\pi^2} \int_{\mathbb{R}^3} \bar{u}^*(2\alpha \wedge \Pi^* \omega_{S^2})$$

is an integer. One can give a proof of this by noting that the fact is true for a smooth map which is constant near infinity and Remark 3.4.

We can now use Hopf lift to prove the cubic decomposition lemma. It is an adapted version of Lemma 6.1 in Lin and Yang [19] (see also [13]). One can see from the statement and the proof of the following lemma the advantage of introducing the space M . Notice that for $u \in M$ we only use the fact that $\int_{\mathbb{R}^3} |\nabla u|^4 dx$ is finite, while estimates will depend only on Faddeev energy.

For the statement of the lemma we need some notations used in Section 6 of [19]. Let $Q(R)$ be a cube of side length $R > 0$. Let $\{Q_i(R)\}_{i=1}^{\infty}$ be a cubic decomposition of \mathbb{R}^3 . That is,

$$\mathbb{R}^3 = \cup_{i=1}^{\infty} Q_i(R),$$

where the interiors of the cubes $Q_i(R)$ are mutually disjoint. For all $a \in \mathbb{R}^3$, let

$$Q_i(R, a) = a + Q_i(R)$$

denote a translation of $Q_i(R)$ by a . We write

$$\Sigma_R = \cup_{i=1}^{\infty} \partial Q_i(R)$$

and

$$\Sigma_R(a) = a + \Sigma_R = \cup_{i=1}^{\infty} \partial Q_i(R, a)$$

for the unions of 2-dimensional faces of $Q_i(R)$'s and $Q_i(R, a)$'s.

Lemma 4.3. *Suppose that $u \in M_d$ and $E_F(u) = C_1$. Let ε be a small constant such that*

$$\int_{\mathbb{R}^3} \varepsilon |\nabla u|^4 dx \leq 1.$$

For any $\delta_0 > 0$, there are some constant C depending only on C_1 , a cubic decomposition $\{Q_i(R_0, a_0)\}$ for some large R_0 and a point $a_0 \in \mathbb{R}^3$ with $|a_0| \leq R_0/4$ such that

$$\int_{\Sigma_{R_0}(a_0)} (|\nabla u|^2 + |\eta|^2 + |\eta|^4) + \varepsilon |\nabla u|^4 dx \leq \frac{C}{R_0} \leq \delta_0^2 \ll 1.$$

where η is the Coulomb gauge of u . Moreover, for each Q_i , there is an integer k_i such that

$$\begin{aligned} \left| d - \sum_{i=1}^{\infty} k_i \right| &\leq \sum_{i=1}^{\infty} \left| \frac{1}{16\pi^2} \int_{Q_i} \eta(u) \wedge u^* \omega_{S^2} - k_i \right| \\ &\leq C \int_{\Sigma_{R_0}(a_0)} |\nabla u|^2 + |\eta|^4 + \varepsilon |\nabla u|^4 d\sigma \\ &\leq \frac{C}{R_0} \leq \delta_0^2 \ll 1. \end{aligned}$$

Proof. Since $E_F(u) < C_1$,

$$\int_{\mathbb{R}^3} |\nabla u|^2 + |\eta|^2 + |\eta|^4 + \varepsilon |\nabla u|^4 dx < C.$$

By Fubini's theorem,

$$\int_0^{R/12} da_1 \int_0^{R/12} da_2 \int_0^{R/12} da_3 \int_{\Sigma_R(a)} (|\nabla u|^2 + |\eta|^2 + |\eta|^4 + \varepsilon |\nabla u|^4) d\sigma \leq CR^2.$$

The mean value theorem then implies that there is a point $a_0 \in \mathbb{R}^3$ with $|a_0| \leq R/4$ such that

$$\int_{\Sigma_R(a_0)} (|\nabla u|^2 + |\eta|^2 + |\eta|^4 + \varepsilon |\nabla u|^4) d\sigma \leq \frac{C}{R}.$$

Here C depends only on C_1 . Therefore, we can choose R sufficiently large so that the first statement of the lemma is true. Please note that R depends only on C_1 and δ_0 .

Next, we need to estimate the error from $\int_{Q_i} \eta(u) \wedge u^* \omega$ to an integer. For simplicity, we suppress lower index i and write Q' for the cube whose center is the same as Q and whose side length is twice as big as Q . Set

$$\int_{\partial Q} |\nabla u|^2 + |\eta|^2 + |\eta|^4 + \varepsilon |\nabla u|^4 d\sigma = \varepsilon_0.$$

Since u has finite Faddeev energy and L^4 norm, we can use Theorem 4.1. There is a lift \bar{u} such that

$$(4.5) \quad \int_Q \eta \wedge u^* \omega_{S^2} = \int_Q \bar{u}^* (2\alpha \wedge \Pi^* \omega_{S^2}).$$

We will now modify \bar{u} outside Q . Let ξ be the average of \bar{u} on ∂Q . Since

$$(4.6) \quad |\nabla \bar{u}|^2 = |\nabla u|^2 + \frac{1}{4} |\eta|^2,$$

we know that

$$\int_{\partial Q} |\nabla \bar{u}|^2 \leq \varepsilon_0$$

and

$$\int_{\partial Q} |\nabla \bar{u}|^4 \leq C(\varepsilon).$$

We can now define $V : Q' \setminus Q \rightarrow \mathbb{R}^4$ to be the harmonic functions with boundary value

$$V|_{\partial Q} = \bar{u}, \quad V|_{\partial Q'} = \frac{\xi}{|\xi|}.$$

Set w to a map from \mathbb{R}^3 to S^3 by

- (1) $w = \bar{u}$ in Q ;
- (2) $w = \frac{V}{|V|}$ in $Q' \setminus Q$;
- (3) $w = \frac{\xi}{|\xi|}$ outside Q' .

Similar to the proof of estimate (3.3), we can show that

$$(4.7) \quad \int_{Q' \setminus Q} |\nabla w|^3 dx < C\varepsilon_0$$

and (w is constant outside Q')

$$(4.8) \quad \int_{\mathbb{R}^3} |\nabla w|^2 + |\nabla w|^4 dx < +\infty.$$

Due to equation (4.8) and Remark 4.2, we know

$$(4.9) \quad \frac{1}{16\pi^2} \int_{\mathbb{R}^3} w^* (2\alpha \wedge \Pi^* \omega_{S^2}) = k \in \mathbb{Z}.$$

By (4.7) and (4.9),

$$\begin{aligned} \left| \frac{1}{16\pi^2} \int_Q \eta \wedge u^* \omega_{S^2} - k \right| &= \left| \frac{1}{16\pi^2} \int_{\mathbb{R}^3 \setminus Q} w^* (2\alpha \wedge \Pi^* \omega_{S^2}) \right| \\ &\leq C\varepsilon_0. \end{aligned}$$

□

As an application of Lemma 4.3, we show that the weak limit of a sequence of maps in M_d has also an integer degree. Notice that such a weak limit may not be in M .

Theorem 4.4. *Let u_j be a sequence in M_d with bounded $E_F(u_j)$. Assume that u_j converges weakly to u in the sense that u_j converges strongly in L^2 to u and ∇u_j converges weakly in L^2 to ∇u and $\partial_k u_j \times \partial_l u_j$ converges weakly in L^2 to $\partial_k u \times \partial_l u$. Then*

$$\frac{1}{16\pi^2} \int_{\mathbb{R}^3} \eta \wedge u^* \omega_{S^2} \in \mathbb{Z}$$

where $\eta = \delta(-\frac{1}{4\pi|x|} \star u^* \omega_{S^2})$.

Proof. For any small positive number ε_0 , we will prove that the difference between

$$\frac{1}{16\pi^2} \int_{\mathbb{R}^3} \eta \wedge u^* \omega_{S^2}$$

with some integer is smaller than ε_0 .

For this ε_0 , we can use Lemma 4.3 for each u_j to obtain a cubic decomposition for each u_j . Precisely, there are $a_j \in \mathbb{R}^3$ and $|a_j| \leq \frac{R_0}{4}$ such that $\Sigma_{R_0}(a_j)$ decompose the space into cubes for which we write $Q_{j,i}$. By choosing a subsequence, we conclude that a_j converges to a and each $Q_{j,i}$ converges to Q_i . Let $\mathcal{E}_{j,i}$ denote the difference between

$$\int_{Q_{j,i}} \frac{1}{16\pi^2} \eta_j \wedge u_j^* \omega_{S^2}$$

and the nearest integer. We know from Lemma 4.3

$$\sum_{i=1}^{\infty} \mathcal{E}_{j,i} < \varepsilon_0.$$

Now we claim:

$$\int_{Q_{j,i}} \eta_j \wedge u_j^* \omega_{S^2} \rightarrow \int_{Q_i} \eta \wedge u^* \omega_{S^2}.$$

We know that for each fixed compact domain Ω

$$\int_{\Omega} \eta_j \wedge u_j^* \omega_{S^2} \rightarrow \int_{\Omega} \eta \wedge u^* \omega_{S^2}.$$

Therefore, to prove the claim, it suffices to show that if $|\Omega| \rightarrow 0$, then

$$\int_{\Omega} \eta_j \wedge u_j^* \omega_{S^2} \rightarrow 0$$

uniformly. This is true due to Hölder inequality and the fact that the L^2 norm of $u_j^* \omega$ and L^4 norm of η_j are bounded uniformly.

Given the claim, we know that the difference between

$$\frac{1}{16\pi^2} \int_{Q_i} \eta \wedge u^* \omega_{S^2}$$

and the nearest integer is no bigger than

$$\liminf_{j \rightarrow \infty} \mathcal{E}_{j,i}.$$

Since $\mathcal{E}_{j,i}$ are all positive, it is elementary to prove

$$\sum_{i=1}^{\infty} \liminf_{j \rightarrow \infty} \mathcal{E}_{j,i} \leq \varepsilon_0.$$

This justifies the statement at the beginning of this proof. \square

5. MINIMIZING RELAXED ENERGY VIA PERTURBED FUNCTIONAL

For each $\varepsilon > 0$, we define perturbed energy as follows

$$E_{F,\varepsilon}(u) = \int_{\mathbb{R}^3} |\nabla u|^2 + \frac{1}{2} \sum_{1 \leq k < j \leq 3} |\partial_k u \times \partial_l u|^2 + \varepsilon |\nabla u|^4 dx.$$

Theorem 5.1. *There is an infinite subset S of \mathbb{Z} such that for each $d \in S$ the minimizing problem of $E_{F,\varepsilon}$ in X_d has a solution.*

The proof follows the same argument as in Lin and Yang [19]. The first ingredient is an energy growth lemma.

Lemma 5.2. *There is a universal constant C such that*

$$E_{F,\varepsilon,d} = \inf\{E_{F,\varepsilon}(u) | u \in M_d\} \leq C |d|^{3/4}.$$

The lemma follows from the same proof as Lemma 5.1 in [19], since the test map constructed in their proof lies in M_d .

The second ingredient is an energy splitting inequality.

Lemma 5.3. *If $d \notin S$, then we can find $l(> 1)$ integers $d_1, \dots, d_l \in S$, whose absolute values are bounded by $|d|$, such that*

$$d = d_1 + \dots + d_l$$

and

$$E_{F,\varepsilon,d} \geq E_{F,\varepsilon,d_1} + \dots + E_{F,\varepsilon,d_l}.$$

For any $d \notin S$, we consider a minimizing sequence u_i of $E_{F,\varepsilon}$ in M_d . Then

$$E_F(u_i) \leq E_{F,\varepsilon}(u_i) \leq C.$$

By Theorem 3.1, there are integers $k_{i,j}$ and cubes $Q_j(R_0, a_i)$ such that

$$\begin{aligned} (5.1) \quad & \sum_{j=1}^{\infty} \left| \frac{1}{16\pi^2} \int_{Q_j(R_0, a_i)} \eta_i \wedge u_i^* \omega_{S^2} \right| \\ & \geq \sum_{j=1}^{\infty} |k_{i,j}| - \sum_{j=1}^{\infty} \left| \frac{1}{16\pi^2} \int_{Q_j(R_0, a_i)} \eta_i \wedge u_i^* \omega_{S^2} - k_{i,j} \right| \\ & \geq m_i - \delta_0^2. \end{aligned}$$

Here m_i is the number of nonzero $k_{i,j}$'s and δ_0 is some very small positive number.

The left hand side of equation (5.1) is bounded above by the energy upper bound C . Therefore, by taking subsequence, we may assume $m_i = m$ for all i . For convenience, we write Q_i^1, \dots, Q_i^m for $Q_j(R_0, a_i)$ (with nonzero $k_{i,j}$).

The proof of Lemma 5.3 is the same as that of Theorem 6.2 in [19]. For the completeness, we repeat it here. We introduce the subsets, S_1, S_2, \dots, S_l of the set

$S = \{1 \cdots m\}$ inductively. Let $s_1 = 1$. We choose a subsequence of u_i , still denoted by u_i , such that there is a subset S_1 of S with $s_1 \in S_1$ and

$$\lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_1}, Q_i^k)$$

exists (finite) for each $k \in S_1$. Moreover, for any $k \in S \setminus S_1$, we have

$$\lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_1} - Q_i^k) = \infty.$$

The existence of such a subsequence of u_i can be checked by induction.

If $S \setminus S_1$ is nonempty, we set $s_2 = \min k \in S \setminus S_1$ and repeat the above procedure to find a subsequence of u_i such that there is a subset S_2 of $S \setminus S_1$ so that

$$\lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_2}, Q_i^k)$$

exists (finite) for each $k \in S_2$ and for any $k \in S \setminus (S_1 \cup S_2)$, we have

$$\lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_2}, Q_i^k) = \infty.$$

Continuing this way we can find l numbers $1 = s_1 < s_2 < \cdots < s_l$, the corresponding subsets S_1, \cdots, S_l of S and a subsequence of u_i (denoted again u_i for simplicity) with the following properties:

$$\begin{aligned} S &= \{1, 2, \cdots, m\} = S_1 \cup S_2 \cup \cdots \cup S_l, \\ S_s \cap S_t &= \emptyset \text{ for } s \neq t, s, t = 1, 2, \cdots, l; \\ s_1 &= 1, \\ S_1 &= \{k \in S \mid \lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_1}, Q_i^k) < \infty\}; \\ s_2 &\in S \setminus S_1, \\ S_2 &= \{k \in S \mid \lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_2}, Q_i^k) < \infty\}; \\ \cdots &\cdots \cdots \\ s_l &\in S \setminus \bigcup_{s=1}^{l-1} S_s, \\ S_l &= \{k \in S \mid \lim_{i \rightarrow \infty} \text{dist}(Q_i^{s_l}, Q_i^k) < \infty\}. \end{aligned}$$

Let $x_{i,1}, x_{i,2}, \cdots, x_{i,l}$ be the centers of cubes $Q_i^{s_1}, \cdots, Q_i^{s_l}$, denoted thereafter as $Q_{i,1}, Q_{i,2}, \cdots, Q_{i,l}$. Let $v_{i,s}(x) = u_i(x - x_{i,s})$, $s = 1, 2, \cdots, l$ and set

$$R_s = \max\{\lim_{i \rightarrow \infty} \text{dist}(Q_{i,s}, Q_i^t) \mid t \in S_s\}, \quad s = 1, 2, \cdots, l.$$

Then $|x_{i,s} - x_{i,t}| \rightarrow \infty$ as $i \rightarrow \infty$ for $s \neq t$ and $s, t = 1, 2, \cdots, l$. By further taking subsequences if necessary, we see that for $s = 1, 2, \cdots, l$

$$d_s \equiv \lim_{i \rightarrow \infty} \sum_{t \in S_s} k(Q_i^t)$$

exists and that $v_{i,s}$ converges to v_s weakly as $i \rightarrow \infty$.

By Lemma 4.3, we obtain

$$\sum_{s=1}^l \sum_{t \in S_s} k(Q_i^t) = d$$

for each i . Thus $d = \sum_{s=1}^l d_s$ follows.

Due to the lower semicontinuity of $E_{F,\varepsilon}$, we know

$$E_{F,\varepsilon}(v_s) < C, \quad s = 1, 2, \cdots, l.$$

By Theorem 4.4, the Hopf degree of v_s is an integer. Following Lemma 6.3 in Lin and Yang [19], we conclude that $Q(v_s) = d_s$.

For any large but fixed R , when i is sufficiently large, $B_R(x_{i,s})$ are disjoint for different s . Therefore

$$E_{F,\varepsilon}(u_i) \geq \sum_{s=1}^l E_{F,\varepsilon}(u_i, B_R(x_{i,s})),$$

where $E_{F,\varepsilon}(u, \Omega)$ means the perturbed energy of u integrating over Ω . Taking $i \rightarrow \infty$,

$$E_{F,\varepsilon;d} \geq \sum_{s=1}^l E_{F,\varepsilon}(v_l, B_R(0)).$$

Since R is an arbitrary positive number, we have

$$E_{F,\varepsilon;d} \geq \sum_{s=1}^l E_{F,\varepsilon;d_s}.$$

If d_i is not in S , we can repeat the above argument until we arrive at a decomposition in which each d_i lies in S .

Combining Lemma 5.2, Lemma 5.3 and the fact that for all $d \neq 0$, $E_{F,\varepsilon;d}$ have a uniform lower bound, one can see that S must be an infinite set.

We have another existence result.

Theorem 5.4. *For $d = \pm 1$, the minimizing problem of $E_{F,\varepsilon}$ on M_d has a solution when ε is very small.*

The proof relies on Lemma 5.3, an upper bound estimate of $E_{F,\varepsilon;1}$ and a lower bound estimate of $E_{F,\varepsilon;d}$ for all d .

As for the lower bound, it was proved by Vakulenko and Kapitanski [23] and improved by Lin and Yang [20] that

$$(5.2) \quad E_{F,\varepsilon}(u) \geq E_F(u) \geq 3^{3/8} 8\sqrt{2}\pi^2 |\deg u|^{3/4}.$$

The upper bound was proved by Ward [24] and Lin and Yang [20] for Faddeev energy

$$(5.3) \quad E_{F,0;1} \leq 32\sqrt{2}\pi^2.$$

Remark 5.5. *The proof in [24] and [20] was indirect. In fact, as pointed out by the authors, the test map Φ is nothing but a Hopf map from ball of radius $\frac{1}{\sqrt{2}}$ composed with a stereographic projection. We may calculate directly that the Faddeev energy of Φ is $32\sqrt{2}\pi^2$, which gives the upper bound. Moreover, by direct computation, the L^4 norm of gradient of this map is bounded. We list below some intermediate results of this calculation.*

The test map $\Phi : \mathbb{R}^3 \rightarrow S^2 \subset \mathbb{R}^3$ is given by (see (4.13) in [20])

$$\Phi(x, y, z) = \begin{pmatrix} \frac{4a}{(r^2+a^2)^2} (2axz + (r^2 - a^2)y) \\ \frac{4a}{(r^2+a^2)^2} (2ayz - (r^2 - a^2)x) \\ 1 - \frac{8a^2}{(r^2+a^2)^2} (z^2 - r^2) \end{pmatrix},$$

where $a = 1/\sqrt{2}$ and $r^2 = x^2 + y^2 + z^2$.

One can calculate

$$|\nabla\Phi|^2 = \frac{64}{(1 + 4r^2)^2}.$$

Integrating over \mathbb{R}^3 ,

$$\int_{\mathbb{R}^3} |\nabla\Phi|^2 dx = 16\sqrt{2}\pi^2.$$

Moreover, we can find out that

$$\int_{\mathbb{R}^3} |\nabla\Phi|^4 dx = 128\sqrt{2}\pi^2.$$

It is important for us that this number is finite. For the other part of the energy,

$$|\partial_x\Phi \times \partial_y\Phi|^2 = \frac{1024(1 - 2x^2 - 2y^2 + 2z^2)^2}{(1 + 2r^2)^6}$$

and

$$\sum_{1 \leq k < l \leq 3} |\partial_k\Phi \times \partial_l\Phi|^2 = \frac{1024}{(1 + 2r^2)^4}.$$

Integrating over \mathbb{R}^3 ,

$$\int_{\mathbb{R}^3} \sum_{1 \leq k < l \leq 3} |\partial_k\Phi \times \partial_l\Phi|^2 dx = 32\sqrt{2}\pi^2.$$

In summary,

$$E_F(\Phi) = 16\sqrt{2}\pi^2 + \frac{1}{2}32\sqrt{2}\pi^2 = 32\sqrt{2}\pi^2.$$

Therefore, when ε is very small, we may assume

$$E_{F,\varepsilon,\pm 1} \leq 32\sqrt{2}\pi^2 + 0.01.$$

For such ε , we can show that $\pm 1 \in S$. In fact, if otherwise, we would have a decomposition of $d = 1$. If it is $1 = 1 + (-2)$, then the energy inequality in Lemma 5.3 says

$$(5.4) \quad 32\sqrt{2}\pi^2 + 0.01 \geq 3^{3/8}8\sqrt{2}\pi^2(1 + 2^{3/4}),$$

which is not possible by direct computation. There are other decompositions, such as $1 = 1 + 1 + (-1)$, $1 = 3 + (-2)$ and so on. It is easy to see that the corresponding right hand sides of (5.4) are even larger so that the inequality is not possible for all decomposition. Therefore, $1 \in S$ and so is -1 .

We now prove Theorem 1.4.

Proof of Theorem 1.4. Let ε_j be a sequence of positive number going to zero. Let u_j be the minimizer of E_{F,ε_j} in M_1 .

We can now apply Lemma 4.3 to u_j . The same argument as in Lemma 5.3 works also for u_j . If $l = 1$ which means there is no splitting at all, we get a limiting map u . $E_F(u)$ is finite due to the weakly lower semicontinuity of E_F . By Theorem 4.4, we know that the Hopf degree of u is an integer. Since there is no splitting, we infer from the construction of u that $Q(u)$ is δ_0 close to 1. Therefore, $u \in X_1$ and by the definition of relaxed energy

$$(5.5) \quad \lim_{j \rightarrow \infty} E_F(u_j) \geq \tilde{E}_F(u) \geq \inf_{w \in X_1} \tilde{E}_F(w).$$

On the other hand, for any positive $\delta > 0$, we can find $w \in X_1$ such that

$$\tilde{E}_F(w) \leq \inf_{w \in X_1} \tilde{E}_F(w) + \delta.$$

By the definition of relaxed energy, we can find $w' \in M_1$ such that

$$E_F(w') \leq \inf_{w \in X_1} \tilde{E}_F(w) + \delta.$$

For this fixed w' , $\int_{\mathbb{R}^3} |\nabla w'|^4 dx$ is finite. therefore, when ε_j is sufficiently small,

$$(5.6) \quad E_F(u_j) \leq E_F(w') + \delta \leq \inf_{w \in X_1} \tilde{E}_F(w) + 3\delta.$$

By equation (5.5) and (5.6), u achieves the minimum of \tilde{E}_F in X_1 .

Next, we will prove that the case $l > 1$ is not possible. If otherwise, there exists $d_1, \dots, d_l \in \mathbb{Z}$ and $x_{j,1}, \dots, x_{j,l} \in \mathbb{R}^3$ such that

$$1 = d_1 + \dots + d_l$$

and $u_j(x - x_{j,s})$ converges weakly to v_s . These v_s has finite energy and by Theorem 4.4 the Hopf degree of v_s is the integer d_s . For any fixed $R > 0$, when j is sufficiently large so that $B(R, x_{j,s})$ are disjoint for different s ,

$$E_F(u_j) \geq \sum_{s=1}^l E_F(u_j, B_R(x_{j,s})).$$

Let $j \rightarrow \infty$,

$$\inf_{w \in X_1} \tilde{E}_F(w) \geq \sum_{s=1}^l E_F(v_s, B_R(0)).$$

Since R is arbitrary,

$$\inf_{w \in X_1} \tilde{E}_F(w) \geq \sum_{s=1}^l E_F(v_s).$$

Since v_s has bounded energy and $Q(v_s) = d_s$, the energy lower bound (5.2) is valid for $E_F(v_s)$. Moreover, because Φ in Remark 5.5 is in M_1 , the same test function implies upper bound for relaxed energy

$$\inf_{w \in X_1} \tilde{E}_F(w) \leq 32\sqrt{2}\pi^2.$$

For the same reason as in Theorem 5.4, the splitting is not possible.

Due to (3.1) and (5.5), we have

$$\inf_{u \in H_1} E_F(u) \geq \min_{u \in X_1} \tilde{E}_F(u).$$

(1.6) follows from the definition of the relaxed energy \tilde{E}_F . □

Acknowledgement 1. *The research of the second author and the third author was supported by the Australian Research Council grant DP0985624. Partial work was done when the first author visited the University of Queensland in August 2009.*

REFERENCES

- [1] R. Adams, Sobolev Spaces, Academic Press, New York, San Francisco, London, 1975.
- [2] F. Bethuel, On the singular set of stationary harmonic maps, *Manus. Math.* **78**(1993), 417–443.
- [3] F. Bethuel and H. Brezis, Regularity of minimizers of relaxed problems for harmonic maps, *J. Funct. Anal.* **101** (1991), 145–161.
- [4] F. Bethuel, H. Brezis and J. M. Coron, Relaxed energies for harmonic maps, In *variational methods*, edited by Berestycki, Coron, Ekeland, Birkhäuser, Basel (1990), 37–52.
- [5] M. Esteban and S. Müller, Sobolev maps with integer degree and applications to Skyrme’s problem, *Proceedings: Mathematical and Physical Sciences*, **436** (1992) 197-201.

- [6] L. C. Evans, Partial regularity for stationary harmonic maps into spheres, Arch. Rational Mech. Anal., **116** (1991) 101–113.
- [7] L. D. Faddeev, Einstein and several contemporary tendencies in the theory of elementary particles in Relativity, quanta, and cosmology, Vol. 1, M. Pantaleo and F. De Finis (Eds.), pp. 247–266 (1979).
- [8] L. D. Faddeev, Knotted Solitons Plenary Address, In: ICM 2002, Beijing, Vol. 1, pp. 235–244 (2002).
- [9] L. D. Faddeev and A. J. Niemi, Stable knot-like structures in classical field theory, Nature, **387** (1997) 58–61.
- [10] M. Giaquinta, G. Modica and J. Soucek, The Dirichlet energy of mappings with values into the sphere, Manus. Math. **65** (1989), 489–507.
- [11] M. Giaquinta, G. Modica and J. Soucek, Cartesian currents in the calculus of variations, Part II, Variational integrals, A series of modern surveys in mathematics, **38**, Springer-Verlag, 1998.
- [12] M. Giaquinta and D. Mucci, The relaxed Dirichlet energy of mappings into a manifold, Calc. Var. Partial Differential Equations **24** (2005), 155–166.
- [13] F. Hang, F. Lin and Y. Yang, Existence of Faddeev Knots in General Hopf Dimensions, Surveys in Differential Geometry 13, in press.
- [14] R. Hardt and F.-H. Lin, A remark on H^1 mappings, Manus. Math. **56** (1986)1–10.
- [15] F. Hélein, Regularity of weakly harmonic maps from a surface into a manifold with symmetries, Manus. Math. **70** (1991) 203–218.
- [16] R. Hardt and T. Riviere, Connecting topological Hopf singularities, Ann. Sc. Norm. Super. Pisa Cl. Sci, 2 (2003), 287–344.
- [17] F.-H. Lin, Mapping Problems, Fundamental Groups and Defect Measures, Acta Mathematica Sinica, English Series, **15** (1999) 25–52.
- [18] F.-H. Lin and C.-Y. Wang, The analysis of harmonic maps and their heat flows, World Scientific Publishing Co. Pte. Ltd. (2008).
- [19] F.-H. Lin and Y. Yang, Existence of energy minimizers as stable knotted solitons in the Faddeev model, Commun. Math. Phys., **249** (2004) 273–303.
- [20] F.-H. Lin and Y. Yang, Energy splitting, substantial inequality, and minimization for the Faddeev and Skyrme models, Commun. Math. Phys., **269** (2007) 137–152.
- [21] E. Radu and M. S. Volkov, Stationary ring solitons in field theory - Knots and vortons, Physics Reports, **468** (2008) 101–151.
- [22] R. Schoen, Analytic aspects of the harmonic map problem, Seminar on nonlinear partial differential equations (Berkeley, Calif., 1983), 321–358, Math. Sci. Res. Inst. Publ., 2, Springer, New York, 1984.
- [23] A. Vakulenko and L. Kapitanski, Stability of solitons in S^2 nonlinear σ -model, Sov. Phys. Dokl., **24** (1979) 433–434.
- [24] R. Ward, Hopf solitons on S^3 and R^3 , Nonlinearity, **12** (1999) 241–246.

MARIANO GIAQUINTA, SCUOLA NORMALE SUPERIORE, 56126 PISA, ITALY
E-mail address: m.giaquinta@sns.it

MIN-CHUN HONG, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF QUEENSLAND, BRISBANE, QLD 4072, AUSTRALIA
E-mail address: hong@maths.uq.edu.au

HAO YIN, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF QUEENSLAND, BRISBANE, QLD 4072, AUSTRALIA AND DEPARTMENT OF MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY, SHANGHAI, CHINA
E-mail address: yinhao78@gmail.com