

# An elementary proof of a Rodriguez-Villegas supercongruence

ROBERTO TAURASO  
Dipartimento di Matematica  
Università di Roma "Tor Vergata", Italy  
tauraso@mat.uniroma2.it  
<http://www.mat.uniroma2.it/~tauraso>

## Abstract

We give a short proof of the following known congruence: for every odd prime  $p$

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 16^{-k} \equiv (-1)^{\frac{p-1}{2}} \pmod{p^2}.$$

Moreover, we provide some new results connected with the above congruence.

## 1 Introduction

The congruence mentioned in the abstract had been conjectured by Rodriguez-Villegas in [4] and it is connected with some properties of the hypergeometric families of Calabi-Yau manifolds. In [3], Mortenson proved the congruence by using the theory of Gaussian hypergeometric series, the properties of the  $p$ -adic  $\Gamma$ -function, and a strange combinatorial identity. In this short note we would like to present a new proof of the congruence which is elementary and uses a different combinatorial identity. The following theorem establishes a more general result, and the conjecture follows immediately by taking  $\alpha = 0$  and  $\beta = -1$  and by noting that  $p^2$  divides  $\binom{2k}{k}^2$  for  $\frac{p-1}{2} < k < p-1$ .

**Theorem** *Let  $p$  be an odd prime and let  $\alpha, \beta \in \mathbb{Z}_p$ , then*

$$\sum_{k=0}^{\frac{p-1}{2}} \binom{2k}{k}^2 \frac{(\alpha - \beta)^{\frac{p-1}{2}-k} (-\beta)^k}{16^k} \equiv \sum_{k=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{k}^2 \alpha^{\frac{p-1}{2}-k} \beta^k \pmod{p^2}.$$

**Remarks.** If we denote by  $T(\alpha, \beta)$  the LHS of the above congruence then it is easy to realize (see the RHS) that this function has some interesting symmetries

$$T(\alpha, \beta) \equiv T(\beta, \alpha) \equiv (-1)^{\frac{p-1}{2}} T(-\alpha, -\beta) \pmod{p^2}.$$

Moreover, by setting in a proper way the parameters  $\alpha$  and  $\beta$  we can show several results:

---

2000 *Mathematics Subject Classification*: 11A07, 11B65 (Primary) 05A10, 05A19 (Secondary)

(i) By letting  $\alpha = 1 - t$  and  $\beta = -t$  then by the above symmetry we have that

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \left(\frac{t}{16}\right)^k \equiv (-1)^{\frac{p-1}{2}} \sum_{k=0}^{p-1} \binom{2k}{k}^2 \left(\frac{1-t}{16}\right)^k \pmod{p^2}.$$

Hence, since the Fibonacci and the Lucas sequences are respectively given by  $F_k = (t^k - (1-t)^k)/\sqrt{5}$  and  $L_k = (t^k + (1-t)^k)$  for  $t = (1 + \sqrt{5})/2$  then for any odd prime  $p$ :

$$\begin{aligned} \text{if } p \equiv 1 \pmod{4} \text{ then } & \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{F_k}{16^k} \equiv 0 \pmod{p^2} \\ \text{if } p \equiv 3 \pmod{4} \text{ then } & \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{L_k}{16^k} \equiv 0 \pmod{p^2}. \end{aligned}$$

(ii) The following result had been conjectured by Zhi-Wei Sun in [5].

If  $p \equiv 3 \pmod{4}$ , by (ii) for  $t = 1/2$  we have

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{32^k} \equiv 0 \pmod{p^2}$$

On the other hand, if  $p \equiv 1 \pmod{4}$  then let  $f = (p-1)/4$  and let  $x \equiv 1 \pmod{4}$  such that  $p = x^2 + y^2$ : Thus by our theorem for  $\alpha = 1/2$  and  $\beta = -1/2$  we obtain

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{32^k} \equiv 4^{-f} \sum_{k=0}^{2f} \binom{2f}{k}^2 (-1)^k = (-4)^{-f} \binom{2f}{f} \equiv 2x - \frac{p}{2x} \pmod{p^2}$$

where in the last step we used Theorem 9.4.3 in [1].

## 2 Proof of the Theorem

The following identity is well-known (see for example 3.18 in [2]): for any non negative integer  $n$ , and any real numbers  $\alpha, \beta$

$$\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} (\alpha - \beta)^{n-k} \beta^k = \sum_{k=0}^n \binom{n}{k}^2 \alpha^{n-k} \beta^k.$$

We give here a straightforward proof via generating functions. Let  $F(z) = \sum_{k \geq 0} \binom{n+k}{k} z^k = (1-z)^{-(n+1)}$  then the LHS is a binomial transform and therefore it is given by

$$[z^n] \frac{1}{1 - (\alpha - \beta)z} F\left(\frac{\beta z}{1 - (\alpha - \beta)z}\right) = \frac{(1 - (\alpha - \beta)z)^n}{(1 - \alpha z)^{n+1}}.$$

Let  $G(z) = \sum_{k \geq 0} \binom{n}{k} z^k = (1+z)^n$  then the RHS is another binomial transform and it is equal to

$$[z^n] \frac{1}{1 - \alpha z} G\left(\frac{\beta z}{1 - \alpha z}\right) = \frac{(1 - (\alpha - \beta)z)^n}{(1 - \alpha z)^{n+1}}.$$

Thus the identity holds and since

$$\binom{n}{k} \binom{n+k}{k} = \binom{n+k}{2k} \binom{2k}{k}$$

it can be restated as follows

$$\sum_{k=0}^n \binom{n+k}{2k} \binom{2k}{k} (\alpha - \beta)^{n-k} \beta^k = \sum_{k=0}^n \binom{n}{k}^2 \alpha^{n-k} \beta^k.$$

In order to complete the proof it suffices to note that for  $n = (p-1)/2$  we have that

$$\binom{n+k}{2k} = \frac{\prod_{j=1}^k (p^2 - (2j-1)^2)}{4^k (2k)!} \equiv \frac{\prod_{j=1}^k (2j-1)^2}{(-4)^k (2k)!} = \binom{2k}{k} (-16)^{-k} \pmod{p^2}.$$

This key-congruence appears both in [5] (as an observation of Z. H. Sun) and in [3] (see Proposition 2.5).  $\square$

## References

- [1] B. C. Berndt, R. J. Evans and K. S. Williams, *Gauss and Jacobi Sums*, Wiley, 1998.
- [2] H. W. Gould, *Combinatorial Identities*, Morgantown, 1972.
- [3] E. Mortenson, *A supercongruence conjecture of Rodriguez-Villegas for a certain truncated hypergeometric function*, J. Number Theory **99** (2003), 139–147.
- [4] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds*, in: Calabi-Yau Varieties and Mirror Symmetry (Toronto, ON, 2001), pp. 223–231, Fields Inst. Commun., **38**, Amer. Math. Soc., Providence, RI, 2003.
- [5] Z. W. Sun, *On congruences related to central binomial coefficients*, preprint arXiv:math.NT/0911.2415v5 (2009).