

# BEST APPROXIMATION WITH WAVELETS IN WEIGHTED ORLICZ SPACES

MARIA DE NATIVIDADE

ABSTRACT. Democracy functions of wavelet admissible bases are computed for weighted Orlicz Spaces  $L^\Phi(w)$  in terms of the fundamental function of  $L^\Phi(w)$ . In particular, we prove that these bases are greedy in  $L^\Phi(w)$  if and only if  $L^\Phi(w) = L^p(w)$ ,  $1 < p < \infty$ . Also, sharp embeddings for the approximation spaces are given in terms of weighted discrete Lorentz spaces. For  $L^p(w)$  the approximation spaces are identified with weighted Besov spaces.

## 1. INTRODUCTION

Let  $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$  be a quasi-Banach space with a countable unconditional basis  $\mathcal{B} = \{e_j : j \in \mathbb{N}\}$ ; that is, every  $x \in \mathbb{B}$  can be uniquely represented as an unconditionally convergent series  $x = \sum_{j \in \mathbb{N}} s_j e_j$ , for some sequences of scalars  $\{s_j : j \in \mathbb{N}\}$ . Let  $\Sigma_N$  denote the set of all elements  $y \in \mathbb{B}$  with at most  $N$  non-null coefficients in the basis representation  $y = \sum_{j \in \mathbb{N}} s_j e_j$ . For  $x \in \mathbb{B}$ , the  $N$ -term error of approximation (with respect to  $\mathcal{B}$ ) is defined by

$$\sigma_N(x)_{\mathbb{B}} \equiv \inf_{y \in \Sigma_N} \|x - y\|_{\mathbb{B}}. \quad (1.1)$$

Two main questions in approximation theory concern the construction of efficient algorithms for  $N$ -term approximation and the characterization of the **approximation spaces**  $\mathcal{A}_q^\alpha(\mathcal{B}, \mathbb{B})$ , which consists of all  $x \in \mathbb{B}$  such that the quantity

$$\|x\|_{\mathcal{A}_q^\alpha(\mathcal{B}, \mathbb{B})} = \begin{cases} \left( \sum_{N \geq 1} (N^\alpha \sigma_N(x)_{\mathbb{B}})^q \frac{1}{N} \right)^{\frac{1}{q}}, & \text{if } 0 < q < \infty; \\ \sup_{N \geq 1} [N^\alpha \sigma_N(x)_{\mathbb{B}}], & \text{if } q = \infty, \end{cases} \quad (1.2)$$

is finite. A computational efficient method to produce  $N$ -term approximations, which has been widely investigate in recent years, is the so called **greedy algorithm** (see e.g [21]). If  $x = \sum_{j \in \mathbb{N}} s_j e_j$  and we order the basis elements in such a way that

$$\|s_{j_1} e_{j_1}\|_{\mathbb{B}} \geq \|s_{j_2} e_{j_2}\|_{\mathbb{B}} \geq \dots \quad (1.3)$$

---

*Date:* May 28, 2018.

*2000 Mathematics Subject Classification.* 41A17, 42C40.

*Key words and phrases.* Greedy algorithm, non-linear approximation, weighted Lebesgue spaces, wavelets, approximation spaces, weighted Besov spaces.

*Acknowledgements and Notes:* This research was supported by Instituto Nacional de Bolsas de Estudo de Angola, INABE and by Grant MTM2007-60952 of Spain. The author wishes to thank J. M. Martell and G. Garrigós for useful discussions on this topic.

(handling ties arbitrarily), the **greedy algorithm of step  $N$**  is defined by the correspondence

$$x = \sum_{j \in \mathbb{N}} s_j e_j \in \mathbb{B} \longrightarrow G_N(x) = \sum_{k=1}^N s_{j_k} e_{j_k} \in \Sigma_N. \quad (1.4)$$

S.V. Konyagin and V. N. Temlyakov ([21]) defined the basis  $\mathcal{B}$  to be **greedy** in  $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$  if the greedy algorithm is optimal in the sense that  $G_N(x)$  is essentially the best  $N$ -term approximation to  $x$  using the basis vectors, i.e, there exists a constant  $C$  such that for all  $x \in \mathbb{B}$  we have

$$\|x - G_N(x)\|_{\mathbb{B}} \leq C \sigma_N(x)_{\mathbb{B}}, \quad N = 1, 2, \dots$$

Thus, for such bases the greedy algorithm produces an almost optimal  $N$ -term approximation, which leads often to a precise identification of the approximation spaces  $\mathcal{A}_q^{\alpha}(\mathcal{B}, \mathbb{B})$ . In [21] greedy basis in a quasi-Banach space  $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$  are characterized as those which are unconditional and **democratic**, the latter meaning that there exists some constant  $\Delta > 0$  such that

$$\left\| \sum_{j \in \Gamma} \frac{e_j}{\|e_j\|_{\mathbb{B}}} \right\|_{\mathbb{B}} \leq \Delta \left\| \sum_{j \in \Gamma'} \frac{e_j}{\|e_j\|_{\mathbb{B}}} \right\|_{\mathbb{B}},$$

holds for all finite sets of indices  $\Gamma, \Gamma' \subset \mathbb{N}$  with the same cardinality. Wavelet systems are well known examples of greedy bases for many function and distribution spaces. Indeed, V.N. Temlyakov showed in [31] that the Haar basis (and any wavelet system  $L^p$ -equivalent to it) is greedy in the Lebesgue space  $L^p([0, 1])$  for  $1 < p < \infty$ . When wavelet have sufficient smoothness and decay, they are also greedy bases for the more general Sobolev and Triebel-Lizorkin classes (see e.g [17, 13]).

The purpose of this paper is to study the efficiency of wavelet greedy algorithms in the weighted Orlicz spaces  $L^{\Phi}(w)$  defined for functions on  $\mathbb{R}^d$ . In Theorem 2.2 (see section 2 ) we show that wavelet bases are unconditional in weighted Orlicz spaces  $L^{\Phi}(w)$  with nontrivial Boyd indices for all  $w \in A_{p, \Phi}(\mathbb{R}^d)$ . We give in section 3 a simple proof of the fact that admissible wavelet bases (see definition below) are not democratic in weighted Orlicz spaces  $L^{\Phi}(w)$  if  $L^{\Phi}(w) \neq L^p(w)$ .

In view of this result it have interest to ask how far wavelet bases are from being democratic in  $L^{\Phi}(w) \neq L^p(w)$ . To quantify democracy of a basis  $\mathcal{B} = \{e_j\}_{j \in \mathbb{N}}$  we shall study the following functions:

$$h_r(N; \mathcal{B}, \mathbb{B}) = \sup_{\text{Card}(\Gamma)=N} \left\| \sum_{\gamma \in \Gamma} \frac{e_{\gamma}}{\|e_{\gamma}\|_{\mathbb{B}}} \right\|_{\mathbb{B}} \quad \text{and} \quad h_l(N; \mathcal{B}, \mathbb{B}) = \inf_{\text{Card}(\Gamma)=N} \left\| \sum_{\gamma \in \Gamma} \frac{e_{\gamma}}{\|e_{\gamma}\|_{\mathbb{B}}} \right\|_{\mathbb{B}}$$

which we call right and left democracy functions of  $\mathcal{B}$  (see also [8, 18, 14]). Observe that a basis is democratic if and only if these two quantities are comparable for all  $N \geq 1$ . Our main result gives a precise value (except for multiplicative constants) of these functions in terms of intrinsic properties of the space  $L^{\Phi}(w)$ . Namely, let  $h_{\varphi}^{+}(t) = \sup_{s>0} \frac{\varphi(st)}{\varphi(s)}$  denote the dilation function associated with the *fundamental function*  $\varphi$  of  $L^{\Phi}(w)$ , and let  $h_{\varphi}^{-}(t)$  be the same quantity with “sup” replaced by “inf” (see Section 2 for precise definitions).

**Theorem 1.1.** *Let  $L^\Phi(w)$  be a weighted Orlicz space with non trivial Boyd indices,  $w \in A_{p^\Phi}$ , a weight on  $\mathbb{R}^d$ , and  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  be an admissible wavelet basis. Then for all  $\Gamma \subset \mathcal{D}$*

$$h_r(N; \mathcal{B}, L^\Phi(w)) \approx h_\varphi^-(N), \quad h_l(N; \mathcal{B}, L^\Phi(w)) \approx h_\varphi^+(N). \quad (1.5)$$

(Here  $p^\Phi = \frac{1}{I_\varphi}$ , where  $I_\varphi$  is the upper Boyd index of  $L^\Phi(w)$ . See definition of Boyd indices in subsection 2.1.)

This result will have applications in the study of approximation spaces (defined using admissible wavelet basis) in weighted Orlicz spaces. We take up this task in the section 4, where we investigate Jackson and Bernstein type estimates and corresponding inclusions for  $N$ -term approximation spaces. In the  $L^p$  case, these estimates are naturally given in terms of the class of discrete Lorentz spaces  $\ell^{\tau,q}$  (see [6, 13, 15, 17, 19]). In the case of weighted Orlicz spaces we shall need *weighted Lorentz sequence spaces*  $\Lambda_\eta^q$ , defined by

$$\Lambda_\eta^q = \left\{ \mathbf{s} : \|\mathbf{s}\|_{\Lambda_\eta^q} = \left[ \sum_{k \geq 1} (\eta_k |s_k^*|)^q \frac{1}{k} \right]^{\frac{1}{q}} < \infty \right\}. \quad (1.6)$$

where  $\{s_k^*\}$  is the non-increasing rearrangement of  $\mathbf{s}$  and the weight  $\eta = \{\eta_k\}$  is a fixed increasing and doubling sequence (see [14]). In particular,  $\Lambda_\eta^q = \ell^{\tau,q}$  when  $\eta_k = k^{1/\tau}$ .

For  $f \in L^\Phi(w)$ , and  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  a wavelet basis in  $L^\Phi(w)$ , write  $f = \sum_{Q \in \mathcal{D}} \langle f, \psi_Q \rangle \psi_Q$ . Then we define  $\Lambda_\eta^q(\mathcal{B}, L^\Phi(w))$  as the set of all  $f \in L^\Phi(w)$  such that the sequence  $\{\|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} : k \geq 1\} \in \Lambda_\eta^q$  and

$$\|f\|_{\Lambda_\eta^q(\mathcal{B}, L^\Phi(w))} = \left\| \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} \right\|_{\Lambda_\eta^q}$$

where  $\|\langle f, \psi_{Q_1} \rangle \psi_{Q_1}\|_{L^\Phi(w)} \geq \|\langle f, \psi_{Q_2} \rangle \psi_{Q_2}\|_{L^\Phi(w)} \geq \dots$  (handling ties arbitrarily).

**Theorem 1.2.** *Let  $L^\Phi(w)$  be a weighted Orlicz space with Boyd indices  $0 < i_\varphi \leq I_\varphi < 1$ , and  $w \in A_{p^\Phi}$  a weight on  $\mathbb{R}^d$ , where  $p^\Phi = \frac{1}{I_\varphi}$ . Then*

$$\Lambda_{k^\alpha h_r(k)}^q(\mathcal{B}, L^\Phi(w)) \hookrightarrow \mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) \hookrightarrow \Lambda_{k^\alpha h_l(k)}^q(\mathcal{B}, L^\Phi(w)), \quad (1.7)$$

These embeddings are optimal, in the sense that the largest and smallest weighted Lorentz spaces  $\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))$  that one can place on the left- and right-hand side of (1.7) are respectively  $\Lambda_{k^\alpha h_r(k)}^q(\mathcal{B}, L^\Phi(w))$  and  $\Lambda_{k^\alpha h_l(k)}^q(\mathcal{B}, L^\Phi(w))$  (see section 4). We point out that a sufficient condition for these two spaces to be equal is that  $h_r(N) \approx h_l(N)$ , in which case the basis is necessarily democratic and  $L^\Phi(w) = L^p(w)$  (see Lemma 5.2 in [14]). Then Theorem 1.2 leads the following identification of Approximation spaces for  $L^p(w)$  in terms of classical Lorentz spaces.

**Corollary 1.3.** *Let  $\alpha > 0$ ,  $1 < p < \infty$ ,  $0 < q \leq \infty$ , and  $w \in A_p$  a weight on  $\mathbb{R}^d$ . Then, for a wavelet basis  $\mathcal{B}$ , we have*

$$\mathcal{A}_q^\alpha(\mathcal{B}, L^p(w)) = \ell^{\tau,q}(\mathcal{B}, L^p(w)), \quad \frac{1}{\tau} = \alpha + \frac{1}{p}. \quad (1.8)$$

Finally we point out that the inclusions in Corollary 1.3 can be described in terms of weighted Besov spaces ([29, 30]), namely

$$\dot{B}_{p,q}^\alpha(w) = \{f \in \mathcal{S}' / \mathcal{P} : (2^{kd} \|\varphi_k * f\|_{L^p(w)})_k \in \ell_q(\mathbb{Z})\}. \quad (1.9)$$

(See the definition of  $\varphi_k$  in section 5).

**Theorem 1.4.** *Let  $\gamma > 0$ ,  $1 < p < \infty$ , and  $w \in A_p$  a weight on  $\mathbb{R}^d$ . Suppose that  $\mathcal{B} = \Psi$  is a family of  $d$ -dimensional Lemarié-Meyer wavelets or a family of  $d$ -dimensional compactly supported Daubechies  $D_N$  wavelets with  $N$  sufficiently large. Then*

$$\mathcal{A}_\tau^\gamma(\Psi, L^p(w)) = \dot{B}_{\tau, \tau}^\gamma(w^{\frac{\tau}{p}}) \quad \text{whenever} \quad \frac{1}{\tau} = \frac{\gamma}{d} + \frac{1}{p}. \quad (1.10)$$

The organization of this article is as follows. Basic facts concerning weights, wavelet bases and greedy bases are given in section 2. Section 3 is devoted to prove Theorem 1.1. Jackson and Bernstein type estimates, as well as the inclusions described in Theorem 1.2 are proved in section 4. Corollary 1.3 and Theorem 1.4 are proved in section 5.

## 2. PRELIMINARIES

**2.1. Basics in weighted Orlicz spaces.** In this subsection we recall some basic facts about weights, weighted Orlicz spaces and wavelet bases on weighted Orlicz spaces, referring to [2, 11] for a complete account on these topics. By a weight on a given measure space, we shall always mean a non-negative locally integrable function  $w$  with values in  $[0, \infty)$  a.e. Let  $w(x)$  be a weight on  $\mathbb{R}^d$ , and for a measurable  $Q \subset \mathbb{R}^d$  write  $w(Q) = \int_Q w(x)dx$ . We say that  $w \in A_p = A_p(\mathbb{R}^d)$ , ( $1 < p < \infty$ ) if there exists a constant  $C_w$  such that

$$\left( \frac{1}{|Q|} \int_Q w(x)dx \right) \left( \frac{1}{|Q|} \int_Q w(x)^{-\frac{1}{p-1}} dx \right)^{p-1} \leq C_w, \quad (2.1)$$

for all  $Q \subset \mathbb{R}^d$ , where  $|Q|$  denote the usual Lebesgue measure of  $Q$ . The condition  $A_1$  can be viewed as limiting case of the condition  $A_p$  for  $p \downarrow 1$ , i.e., (2.1) is viewed as

$$\left( \frac{1}{|Q|} \int_Q w(x)dx \right) \text{ess}_Q \sup(w^{-1}) \leq C_w. \quad (2.2)$$

If  $w \in A_p$  for some  $p \in [1, \infty)$ , then there exist  $C_w^1, C_w^2 > 0$  and  $\delta > 0$  such that

$$C_w^1 \left( \frac{|A|}{|Q|} \right)^p \leq \frac{w(A)}{w(Q)} \leq C_w^2 \left( \frac{|A|}{|Q|} \right)^\delta \quad (2.3)$$

for all subsets  $A \subset Q$ . (For the left hand inequality take  $f = \chi_A$  in part b) of Theorem 2.1, Chapter IV, of [11]; for the right hand inequality see Theorem 2.9, Chapter IV, of [11]).

A Young function is a convex non-decreasing function  $\Phi : [0, \infty) \rightarrow [0, \infty]$  so that  $\lim_{t \rightarrow +\infty} \Phi(t) = \infty$ . Throughout this paper we shall assume that  $\Phi(0) = 0$ ,  $\Phi$  is strictly increasing and everywhere finite, so that it is a continuous bijection of  $[0, \infty)$ . Given such  $\Phi$  and  $w \in A_\infty = \cup_{p \geq 1} A_p$ , the weighted Orlicz space  $L^\Phi(w)$  is the class of all measurable functions  $f : \mathbb{R}^d \rightarrow \mathbb{C}$  so that  $\Phi\left(\frac{|f(x)|}{\lambda}\right) \in L^1(w)$  for some  $\lambda > 0$ . The space  $L^\Phi(w)$  becomes a weighted rearrangement invariant Banach function space when endowed with the corresponding Luxemburg norm

$$\|f\|_{L^\Phi(w)} = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^d} \Phi\left(\frac{|f(x)|}{\lambda}\right) w(x)dx \leq 1 \right\} \quad (2.4)$$

It is not difficult to prove that if  $E \subset \mathbb{R}^d$  is any measurable set

$$\|\chi_E\|_{L^\Phi(w)} = \frac{1}{\Phi^{-1}\left(\frac{1}{w(E)}\right)}. \quad (2.5)$$

The function  $\varphi(t) = \frac{1}{\Phi^{-1}\left(\frac{1}{t}\right)}$ ,  $0 < t < \infty$ , satisfies  $\varphi(t) = \|\chi_E\|_{L^\Phi(w)}$  for any measurable set  $E \subset \mathbb{R}^d$  such that  $w(E) = t$ , and it is called the *fundamental function of  $L^\Phi(w)$* .

The Boyd indices of the weighted Orlicz space  $L^\Phi(w)$  can be computed directly from the Young function  $\Phi$  or from the fundamental function  $\varphi$ . Set

$$h_\varphi^+(t) = \sup_{s>0} \frac{\varphi(st)}{\varphi(s)}, \quad 0 < t < \infty. \quad (2.6)$$

Then, the lower and upper Boyd indices  $i_\varphi$  and  $I_\varphi$  of  $L^\Phi(w)$  are given by

$$i_\varphi = \lim_{t \rightarrow 0} \frac{\log h_\varphi^+(t)}{\log t} = \sup_{0 < t \leq 1} \frac{\log h_\varphi^+(t)}{\log t} \quad (2.7)$$

and

$$I_\varphi = \lim_{t \rightarrow \infty} \frac{\log h_\varphi^+(t)}{\log t} = \inf_{1 < t < \infty} \frac{\log h_\varphi^+(t)}{\log t} \quad (2.8)$$

respectively (see [2], p. 277 or [22], p. 54). It is known that  $0 \leq i_\varphi \leq I_\varphi \leq 1$  (see Proposition 5.15 of [2], p. 149). Assuming further that  $i_\varphi > 0$  it follows that

$$\varphi(st) \leq C_\epsilon \max\{s^{i_\varphi - \epsilon}, s^{I_\varphi + \epsilon}\} \varphi(t), \quad s, t > 0 \quad (2.9)$$

and

$$\varphi(st) \geq C_\epsilon \min\{s^{i_\varphi - \epsilon}, s^{I_\varphi + \epsilon}\} \varphi(t), \quad s, t > 0 \quad (2.10)$$

for every  $\epsilon > 0$  and some constant  $C_\epsilon > 0$  (see [20], p. 3). In this paper we shall only consider weighted Orlicz spaces with non trivial Boyd indices, that is  $0 < i_\varphi \leq I_\varphi < 1$ .

**Example 2.1.** When  $\Phi(t) = t^p$ ,  $1 \leq p < \infty$ , then  $L^\Phi(w) = L^p(w)$  and  $\varphi(t) = t^{\frac{1}{p}}$ . Hence,  $h_\varphi^+(t) = t^{\frac{1}{p}}$ , which implies  $i_\varphi = I_\varphi = \frac{1}{p}$ .

**2.2. Wavelet bases and weighted Orlicz spaces.** Let  $\mathcal{D} = \{Q_{j,k} = 2^{-j}([0, 1]^d + k) : j \in \mathbb{Z}, k \in \mathbb{Z}^d\}$  denote the set of all dyadic cubes in  $\mathbb{R}^d$ . We say that a finite collection of functions  $\{\psi^1, \dots, \psi^L\} \subset L^2(\mathbb{R}^d)$  is an *orthonormal wavelet family* if the system

$$\left\{ \psi_{Q_{j,k}}^l(x) = 2^{\frac{jd}{2}} \psi^l(2^j x - k) : j \in \mathbb{Z}, k \in \mathbb{R}^d, l = 1, \dots, L \right\}, \quad (2.11)$$

forms an orthonormal basis of  $L^2(\mathbb{R}^d)$ . We will say that the wavelet family is *admissible* if for all  $1 < p < \infty$ ,

$$\|S_\psi f(\cdot)\|_{L^p(\mathbb{R}^d)} \approx \|f(\cdot)\|_{L^p(\mathbb{R}^d)}, \quad (2.12)$$

where

$$S_\psi f(\cdot) = \left( \sum_{l=1}^L \sum_{I \in \mathcal{D}} |\langle f(\cdot), \psi_I(\cdot) \rangle|^2 \chi_I(\cdot) |I|^{-1} \right)^{\frac{1}{2}}. \quad (2.13)$$

This implies that wavelet admissible bases are unconditional in  $L^p(\mathbb{R}^d)$ ,  $1 < p < \infty$ . The reader can consult [5, 16, 23, 25], for constructions, examples, and properties of

orthonormal wavelets. Admissible wavelets include the  $d$ -dimensional Haar system, wavelet arising from multiresolution analysis (see [25], p. 22), wavelets belonging to the regularity class  $\mathcal{R}^0$  (as defined in [16], p. 64 for  $d = 1$ ), compactly support wavelets (see [5]), and actually any orthonormal wavelet in  $L^2(\mathbb{R}^d)$  with mild decay conditions (see [32, 28]).

In the following result we prove that wavelet admissible basis are also unconditional basis of weighted Orlicz spaces  $L^\Phi(w)$ , for appropriate  $w$ , since the norm can be characterize in terms of a square function. Without loss of generality we assume  $L = 1$  in the rest of this work.

**Theorem 2.2.** *Let  $L^\Phi(w)$  be a weighted Orlicz space, with the Boyd indices satisfying  $0 < i_\varphi \leq I_\varphi < 1$ , and  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  an admissible wavelet basis. Then, if  $w \in A_{p^\Phi}(\mathbb{R}^d)$ , where  $p^\Phi = \frac{1}{I_\varphi}$ , we have*

$$\|f(\cdot)\|_{L^\Phi(w)} \simeq \|S_\psi f(\cdot)\|_{L^\Phi(w)}, \quad \text{for all } f \in L^\Phi(w). \quad (2.14)$$

For the proof we shall use the following extrapolation theorem adapted to our situation.

**Theorem 2.3.** ([4]) *Let  $\mathcal{F}$  be a family of couples of measurable non-negative functions  $(f, g)$ . Suppose that for some  $1 \leq p_0 < \infty$ , and every weight  $w \in A_{p_0}(\mathbb{R}^d)$ ,*

$$\int_{\mathbb{R}^d} f(x)^{p_0} w(x) dx \leq C \int_{\mathbb{R}^d} g(x)^{p_0} w(x) dx, \quad \text{for all } (f, g) \in \mathcal{F}. \quad (2.15)$$

*Then, if  $L^\Phi(w)$  is a weighted Orlicz space such that the Boyd indices satisfies,  $0 < i_\varphi \leq I_\varphi < 1$  and  $w \in A_{p^\Phi}(\mathbb{R}^d)$ ,  $p^\Phi = \frac{1}{I_\varphi}$ , we have*

$$\|f\|_{L^\Phi(w)} \leq C \|g\|_{L^\Phi(w)}, \quad \text{for all } (f, g) \in \mathcal{F}. \quad (2.16)$$

*Proof.* ( **of Theorem 2.2** ) It is proved in [12] (see also [1]) that

$$\|f\|_{L^p(w)} \simeq \|S_\psi(f)\|_{L^p(w)}, \quad (2.17)$$

for all  $1 < p < \infty$  and  $w \in A_p$ . We consider the family  $\mathcal{F} = \{(|f|, S_\psi(f)) : S_\psi(f) \in L^p(w)\}$ . From the equivalence (2.17), we obtain

$$\int_{\mathbb{R}^d} |f(x)|^p w(x) dx \leq C_1 \int_{\mathbb{R}^d} |S_\psi(f)|^p w(x) dx$$

for all  $1 < p < \infty$  and  $w \in A_p(\mathbb{R}^d)$ . Then, by Theorem 2.3 we obtain

$$\|f\|_{L^\Phi(w)} \leq C_1 \|S_\psi(f)\|_{L^\Phi(w)}$$

when  $w \in A_{p^\Phi}$ . The other inequality is proved similarly taking  $\mathcal{F} = \{(S_\psi(f), |f|), f \in L^p(w)\}$ .  $\square$

**2.3. Greedy basis and democracy.** We defined in the introduction the notion of **greedy basis** in a quasi-normed Banach space  $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$ . We also mentioned the result of Konyagin and Temlyakov [21] characterizing greedy bases as those which are unconditional and **democratic**. For simplicity, given a basis  $\mathcal{B} = \{e_j : j \in \mathbb{N}\}$  in  $\mathbb{B}$  we shall denote the (normalized) characteristic function of a finite set of indices  $\Gamma \in \mathbb{N}$  by

$$\tilde{\mathbf{1}}_\Gamma = \tilde{\mathbf{1}}_\Gamma^{\mathbb{B}, \mathcal{B}} = \sum_{j \in \Gamma} \frac{e_j}{\|e_j\|_{\mathbb{B}}}.$$

The basis  $\mathcal{B}$  is **democratic** in  $\mathbb{B}$  if there exists  $C \geq 1$  such that

$$\|\tilde{1}_\Gamma\|_{\mathbb{B}} \leq C \|\tilde{1}_{\Gamma'}\|_{\mathbb{B}} \quad (2.18)$$

for all finite sets of indices  $\Gamma, \Gamma' \subset \mathbb{N}$  with  $\#\Gamma = \#\Gamma'$  (the symbol  $\#\Gamma$  denotes the cardinality of the set  $\Gamma$ ). Quite often one can show democracy by finding a function  $h : \mathbb{N} \rightarrow \mathbb{R}^+$  for which

$$\frac{1}{C}h(\#\Gamma) \leq \|\tilde{1}_\Gamma\|_{\mathbb{B}} \leq Ch(\#\Gamma), \quad \forall \Gamma \subset \mathbb{N}, \text{ finite}. \quad (2.19)$$

In the case of wavelet bases, many classical function and distribution spaces satisfy (2.19) with  $h(\#\Gamma) = (\#\Gamma)^{\frac{1}{p}}$ . Indeed, this is the situation for the Lebesgue spaces  $L^p(\mathbb{R}^d)$  when  $1 < p < \infty$ , for the Hardy spaces  $H^p(\mathbb{R}^d)$ ,  $0 < p \leq 1$  and for the Sobolev spaces  $\dot{W}^{s,p}(\mathbb{R}^d)$ ,  $1 < p < \infty$  (see [17]), and more generally for the family of Triebel-Lizorkin spaces  $\dot{F}_{p,r}^s(\mathbb{R}^d)$  with  $0 < p < \infty$ ,  $s \in \mathbb{R}$ ,  $0 < r \leq \infty$  (under the usual smoothness assumptions, and with the standard modification of the basis in the case of inhomogeneous spaces; see [13]). Thus, wavelet bases are democratic and hence greedy in all these spaces.

The Haar system is not greedy in rearrangement invariant spaces defined in  $[0, 1]$  other than  $L^p[0, 1]$  (see [33]). Moreover, wavelet bases are not democratic in other classical spaces, such as  $BMO$ , the Besov spaces  $\dot{B}_{p,q}^\alpha(\mathbb{R}^d)$  with  $p \neq q$ , Orlicz spaces  $L^\Phi(\mathbb{R}^d)$  distinct from  $L^p(\mathbb{R}^d)$ , and as we shall see below, weighted Orlicz spaces  $L^\Phi(w)$  distinct from  $L^p(w)$ .

**Definition 2.4.** Let  $\mathcal{B}$  be a collection of elements in a quasi-Banach space  $\mathbb{B}$ . The right-democracy function associated with  $\mathcal{B}$  is defined by

$$h_r(N; \mathbb{B}, \mathcal{B}) = \sup_{\text{Card}(\Gamma)=N} \|\tilde{1}_\Gamma\|_{\mathbb{B}}; \quad (2.20)$$

analogously, the left-democracy function associated with  $\mathcal{B}$  is defined by

$$h_l(N; \mathbb{B}, \mathcal{B}) = \inf_{\text{Card}(\Gamma)=N} \|\tilde{1}_\Gamma\|_{\mathbb{B}} \quad (2.21)$$

Observe that a basis  $\mathcal{B}$  is democratic in  $\mathbb{B}$  if and only if  $h_r(N; \mathbb{B}, \mathcal{B}) \leq Ch_l(N; \mathbb{B}, \mathcal{B})$  for all  $N \geq 1$  and some  $C > 0$ .

We want to show that, in general, admissible wavelet bases are not democratic in weighted Orlicz spaces. In order to do so one needs to estimate  $\|\tilde{1}_\Gamma\|_{L^\Phi(w)}$  in terms of  $\#\Gamma$ . This can be done when  $\Gamma$  is a collection of pairwise disjoint dyadic cubes  $\{Q_j\}_{j=1}^N$ , such that  $w(Q_j) \approx \tau$ , for any  $\tau > 0$ .

We state and prove the following results.

**Lemma 2.5.** Let  $w \in A_\infty(\mathbb{R}^d)$  be a weight. If  $\{Q_k\}_{k=-\infty}^\infty$  is a family of dyadic cubes such that  $Q_k \subset Q_{k+1}$  and  $|Q_{k+1}| = 2^d|Q_k|$  for all  $k \in \mathbb{Z}^+$ , then

$$\lim_{k \rightarrow \infty} w(Q_k) = \infty \quad \text{and} \quad \lim_{k \rightarrow -\infty} w(Q_k) = 0. \quad (2.22)$$

*Proof.* Because  $w \in A_\infty$ , if  $k \geq 0$ , by (2.3) we obtain

$$\frac{w(Q_0)}{w(Q_k)} \leq C_w^2 \left( \frac{|Q_0|}{|Q_k|} \right)^\delta = C_w^2 \left( \frac{1}{2^{kd}} \right)^\delta.$$

Then,  $w(Q_k) \geq (C_w^2)^{-1} 2^{kd\delta} w(Q_0)$  and  $\lim_{k \rightarrow \infty} w(Q_k) = \infty$ . On the other hand, if  $k \leq 0$ , by (2.3) we obtain

$$\frac{w(Q_k)}{w(Q_0)} \leq C_w^2 \left( \frac{|Q_k|}{|Q_0|} \right)^\delta = C_w^2 2^{kd\delta}.$$

Then,  $w(Q_k) \leq C_w^2 2^{kd\delta} w(Q_0)$  and  $\lim_{k \rightarrow -\infty} w(Q_k) = 0$ .  $\square$

**Lemma 2.6.** *Let  $w \in A_\infty(\mathbb{R}^d)$  be a weight. Given  $\tau > 0$  there exists a pairwise disjoint sequence of cubes  $\{R_j\}_{j=1}^\infty \subset \mathcal{D}$  such that*

$$C\tau < w(R_j) \leq \tau$$

where  $C > 0$  is a constant depending only on  $w$ .

*Proof.* Let  $Q_k = [0, 2^k)^d$ ,  $k \in \mathbb{Z}$ . By lemma 2.5 there exists  $k_1 \in \mathbb{Z}$  such that

$$w(Q_{k_1}) \leq \tau < w(Q_{k_1+1}). \quad (2.23)$$

Choose  $R_1 = Q_{k_1}$ . We have  $w(R_1) = w(Q_{k_1}) \leq \tau$ . On the other hand, by (2.3), we obtain

$$\frac{w(Q_{k_1})}{w(Q_{k_1+1})} \geq C_w^1 \left( \frac{|Q_{k_1}|}{|Q_{k_1+1}|} \right)^p = C_w^1 2^{-dp},$$

so that

$$w(R_1) = w(Q_{k_1}) \geq C_w^1 2^{-dp} w(Q_{k_1+1}) > C_w^1 2^{-dp} \tau.$$

Thus, we can take  $C = C_w^1 2^{-dp}$ .

Suppose we have chosen disjoint cubes  $R_1, R_2, \dots, R_{m-1}$  such that  $C\tau < w(R_j) \leq \tau$  for all  $j = 1, 2, \dots, m-1$ . Without loss of generality we can assume that all the  $R_j$  are contained in the positive cone of  $\mathbb{R}^d$ , that is, the set of points of  $\mathbb{R}^d$  with non-negative coordinates.

Choose  $Q_0 = 2^{k_m} [0, 1)^d$ ,  $k_m \in \mathbb{Z}$ , such that  $R_j \subset Q_0$  for all  $j = 1, 2, \dots, m-1$ . Consider the increasing family of dyadic cubes given by  $Q_k = 2^{k_m+k} [0, 1)^d$ ,  $k = 0, 1, 2, \dots$ . Let  $\tilde{Q}_k$ ,  $k = 1, 2, \dots$ , be a dyadic cube contained in  $Q_k$  such that  $|\tilde{Q}_k| = \frac{|Q_k|}{2^d}$  and  $\tilde{Q}_k \cap Q_{k-1} = \emptyset$ . If  $w(\tilde{Q}_k) \leq \tau$  for all  $k = 1, 2, 3, \dots$  by (2.3) we obtain

$$\frac{w(\tilde{Q}_k)}{w(Q_k)} \geq C_w^1 \left( \frac{|\tilde{Q}_k|}{|Q_k|} \right)^p = C_w^1 2^{-dp}.$$

Thus,  $w(Q_k) \leq (C_w^1)^{-1} 2^{dp} \tau$  for all  $k = 1, 2, \dots$  contradicting lemma 2.5. Thus, there exists  $k_m^0 \in \mathbb{Z}$  such that  $w(\tilde{Q}_{k_m^0}) > \tau$ . Consider a family of descendants of the dyadic cube  $\tilde{Q}_{k_m^0}$ . By lemma 2.5, there exists  $\tilde{Q}_{k_m}$ ,  $\tilde{\tilde{Q}}_{k_m} \in \mathcal{D}$  such that

$$w(\tilde{\tilde{Q}}_{k_m}) \leq \tau < w(\tilde{Q}_{k_m}) \quad (2.24)$$

and  $|\tilde{\tilde{Q}}_{k_m}| = \frac{|\tilde{Q}_{k_m}|}{2^d}$ . Choose  $R_m = \tilde{\tilde{Q}}_{k_m}$ . Since (2.24) is the same relation as (2.23) it follows that

$$C_w^1 2^{-dp} \tau < w(R_m) \leq \tau.$$

Observe that  $R_m$  has been chosen in the positive cone of  $\mathbb{R}^d$  and is disjoint to  $R_1, \dots, R_{m-1}$ .  $\square$



**Proposition 2.7.** *Let  $L^\Phi(w)$  be a weighted Orlicz space with Boyd indices  $0 < i_\varphi \leq I_\varphi < 1$ ,  $w \in A_{p^\Phi}$  a weight in  $\mathbb{R}^d$ , and let  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  be an admissible wavelet basis.*

*i) If  $\Gamma = \{Q_1, \dots, Q_N\} \subset \mathcal{D}$  is a pairwise disjoint family then*

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \approx \left\| \sum_{Q \in \Gamma} \frac{\chi_Q(\cdot)}{\varphi(w(Q))} \right\|_{L^\Phi(w)}. \quad (2.25)$$

*ii) Moreover, for any  $\tau > 0$ , there exist a family of disjoint cubes  $\Gamma = \{R_1, R_2, \dots, R_N\} \subset \mathcal{D}$ , such that*

$$\|\tilde{1}_\Gamma\|_{L^\Phi} \approx \frac{\varphi(N\tau)}{\varphi(\tau)} \quad (2.26)$$

*Proof.* *i)* For a single element of the basis  $\mathcal{B}$  we have, by (2.14) that

$$\|\psi_Q\|_{L^\Phi(w)} \simeq \left\| \left( \frac{\chi_Q(\cdot)}{|Q|} \right)^{\frac{1}{2}} \right\|_{L^\Phi(w)} = \frac{\|\chi_Q(\cdot)\|_{L^\Phi(w)}}{|Q|^{\frac{1}{2}}} = \frac{\varphi(w(Q))}{|Q|^{\frac{1}{2}}}. \quad (2.27)$$

By (2.14) again

$$\begin{aligned} \|\tilde{1}_\Gamma\|_{L^\Phi(w)} &\simeq \left\| \left( \sum_{Q \in \Gamma} \frac{1}{\|\psi_Q\|_{L^\Phi(w)}^2} \chi_Q |Q|^{-1} \right)^{\frac{1}{2}} \right\|_{L^\Phi(w)} \approx \left\| \left( \sum_{Q \in \Gamma} \frac{\chi_Q}{\varphi(w(Q))^2} \right)^{\frac{1}{2}} \right\|_{L^\Phi(w)} = \\ &= \left\| \sum_{Q \in \Gamma} \frac{\chi_Q}{\varphi(w(Q))} \right\|_{L^\Phi(w)}, \end{aligned}$$

where in the last equality we have used that the cubes in  $\Gamma$  are pairwise disjoint.

*ii)* The existence of the family  $\Gamma = \{R_1, R_2, \dots, R_N\} \subset \mathcal{D}$  is proved in Lemma 2.6 where it is shown that  $w(R_j) \approx \tau$ ,  $j = 1, 2, \dots, N$ . In this situation

$$\begin{aligned} \|\tilde{1}_\Gamma\|_{L^\Phi(w)} &= \left\| \sum_{j=1}^N \frac{\chi_{R_j}(\cdot)}{\varphi(w(R_j))} \right\|_{L^\Phi(w)} \approx \frac{1}{\varphi(\tau)} \left\| \bigcup_{j=1}^N \chi_{R_j} \right\|_{L^\Phi(w)} \\ &= \frac{1}{\varphi(\tau)} \varphi\left(w\left(\bigcup_{j=1}^N R_j\right)\right) \approx \frac{\varphi(N\tau)}{\varphi(\tau)}. \end{aligned} \quad (2.28)$$

□

**Remark 2.8.** *It follows from part ii) of Proposition 2.7 that for admissible wavelet basis  $\mathcal{B}$*

$$h_r(N; L^\Phi(w), \mathcal{B}) \gtrsim \sup_{\tau > 0} \frac{\varphi(N\tau)}{\varphi(\tau)} = h_\varphi^+(N)$$

and

$$h_l(N; L^\Phi(w), \mathcal{B}) \lesssim \inf_{\tau > 0} \frac{\varphi(N\tau)}{\varphi(\tau)} = h_\varphi^-(N).$$

Thus, if  $h_\varphi^+(N)$  and  $h_\varphi^-(N)$  are not comparable for  $N \geq 1$  it follows that admissible wavelet bases are non democratic in weighted Orlicz spaces. On the other hand, if wavelet admissible bases are democratic in  $L^\Phi(w)$ ,  $h_\varphi^+(N) \leq h_\varphi^-(N)$ , and Lemma 5.2 in [14] shows that  $\varphi(t) \approx t^\alpha$  for some  $\alpha \in (0, 1)$ ; thus, the only democratic weighted Orlicz spaces are the spaces  $L^p(w)$  for some  $p = \frac{1}{\alpha} \in (1, \infty)$ .

### 3. LEFT AND RIGHT DEMOCRACY FUNCTIONS FOR WEIGHTED ORLICZ SPACES

Our main theorem in this section shows that  $h_r(N; L^\Phi(w), \mathcal{B}) \lesssim h_\varphi^+(N)$  and  $h_l(N; L^\Phi, \mathcal{B}) \gtrsim h_\varphi^-(N)$  (see theorem 3.1 below) giving us together with remark 2.8 a complete description (up to multiplicative constants) of the left and right democracy functions of wavelet basis on weighted Orlicz spaces.

**Theorem 3.1.** *Let  $L^\Phi(w)$ , be a weighted Orlicz space with Boyd indices satisfying  $0 < i_\varphi \leq I_\varphi < 1$ ,  $w \in A_{p^\Phi}$  a weight in  $\mathbb{R}^d$ , and let  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  be an admissible wavelet basis. Then for all  $\Gamma \subset \mathcal{D}$*

$$h_\varphi^-(\#\Gamma) \lesssim \|\tilde{1}_\Gamma\|_{L^\Phi(w)} \lesssim h_\varphi^+(\#\Gamma). \quad (3.1)$$

This, together with Remark 2.8 gives

$$h_r(N; \mathcal{B}, L^\Phi(w)) \approx h_\varphi^+(N) \quad \text{and} \quad h_l(N; \mathcal{B}, L^\Phi(w)) \approx h_\varphi^-(N),$$

which is Theorem 1.1.

The rest of this section is devoted to prove Theorem 3.1. We first present a very simple argument for the case of pairwise disjoint cubes.

**3.1. Proof of Theorem 3.1: The case of disjoint cubes.** We assume first that  $\Gamma = \{Q_1, \dots, Q_N\}$  consists of pairwise disjoint cubes. Let  $\lambda = h_\varphi^+(N)$ , so that  $\varphi(N\omega(Q)) \leq \lambda\varphi(\omega(Q))$ ,  $\forall Q \in \Gamma$ . Therefore, since the elements of  $\Gamma$  are disjoint, and  $\Phi$  is increasing

$$\begin{aligned} \int_{\mathbb{R}^d} \Phi\left(\frac{\sum_{j=1}^N \frac{\chi_{Q_j}(x)}{\varphi(\omega(Q_j))}}{\lambda}\right) w(x) dx &= \sum_{j=1}^N \Phi\left(\frac{1}{\lambda\varphi(\omega(Q_j))}\right) \omega(Q_j) \\ &\leq \sum_{j=1}^N \Phi\left(\frac{1}{\varphi(N\omega(Q_j))}\right) \omega(Q_j) = \sum_{j=1}^N \Phi\left(\Phi^{-1}\left(\frac{1}{N\omega(Q_j)}\right)\right) \omega(Q_j) = 1 \end{aligned}$$

Then by (2.25) and (2.4) we have

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \simeq \left\| \sum_{j=1}^N \frac{\chi_{Q_j}(\cdot)}{\varphi(\omega(Q_j))} \right\|_{L^\Phi(w)} \leq h_\varphi^+(N).$$

The lower estimate is obtained in a similar way.

**3.2. Proof of Theorem 3.1: The general case.** In the case of disjoint cubes just considered we have two important features. First, Proposition 2.7 allows us to “linearize” the square function in (2.14). Second, for the estimates obtained in the previous argument it is crucial that the sets involved are disjoint. For general families of cubes we are going to follow the same scheme. First we “linearize” the square function and we dominate this by an expression involving only disjoint subsets from the elements of  $\Gamma$ .

**Linearization of the square function.** Given a finite set  $\Gamma \subset \mathcal{D}$ , we denote

$$S_\Gamma(x) = \left( \sum_{Q \in \Gamma} \frac{\chi_Q(x)}{\varphi(\omega(Q))^2} \right)^{\frac{1}{2}}, \quad (3.2)$$

so that by (2.14) and (2.27), we have  $\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \simeq \|S_\Gamma(\cdot)\|_{L^\Phi(w)}$ . For every  $x \in \bigcup_{Q \in \Gamma} Q$ , we define  $Q_x$  as the smallest (hence unique) cube in  $\Gamma$  containing  $x$ . It is clear that

$$S_\Gamma(x) \geq \frac{\chi_{Q_x}(x)}{\varphi(w(Q_x))}, \quad \forall x \in \bigcup_{Q \in \Gamma} Q, \quad (3.3)$$

since the left hand side contains at least the cube  $Q_x$  (and possible more). We now show that the reverse inequality holds. Indeed, if we enlarge the sum to include all dyadic cubes containing  $Q_x$  we have

$$S_\Gamma(x)^2 = \sum_{Q \in \Gamma} \frac{\chi_Q(x)}{\varphi(w(Q))^2} \leq \sum_{\substack{Q \supset Q_x \\ Q \in \mathcal{D}}} \frac{1}{\varphi(w(Q))^2} \leq \sum_{j=0}^{\infty} \frac{1}{\varphi(w(Q_x^j))^2},$$

where  $Q_x^j$  denotes the unique cube of measure  $2^{jd}|Q_x|$  containing  $Q_x$ . Now since  $Q_x = Q_x^0 \subset Q_x^1 \subset Q_x^2 \subset \dots$  we can use (2.3) to obtain

$$\frac{w(Q_x)}{w(Q_x^j)} \leq C_w^2 \left( \frac{|Q_x|}{|Q_x^j|} \right)^\delta = C_w^2 2^{-jd\delta}.$$

Hence,

$$w(Q_x^j) \geq (C_w^2)^{-1} w(Q_x) 2^{jd\delta},$$

and

$$\varphi(w(Q_x^j)) \geq \varphi((C_w^2)^{-1} w(Q_x) 2^{jd\delta}).$$

Since  $i_\varphi > 0$ , by (2.10) we can choose  $0 < \epsilon < i_\varphi$  and find a  $C_\epsilon > 0$  such that  $\varphi((C_w^2)^{-1} 2^{jd\delta} w(Q_x)) \geq C_\epsilon ((C_w^2)^{-1} 2^{jd\delta})^{(i_\varphi - \epsilon)} \varphi(w(Q_x))$ .

Thus,

$$S_\Gamma(x)^2 \leq \frac{(C_w^2)^{(i_\varphi - \epsilon)} C_\epsilon}{(\varphi(w(Q_x)))^2} \sum_{j=0}^{\infty} 2^{-jd\delta(i_\varphi - \epsilon)} \lesssim \frac{\chi_{Q_x}(x)}{(\varphi(w(Q_x)))^2}.$$

This and (3.3) show that

$$S_\Gamma(x) \simeq \frac{\chi_{Q_x}(x)}{\varphi(w(Q_x))}. \quad (3.4)$$

Observe from (3.4) that  $S_\Gamma(x) \simeq S_{\Gamma_{\min}}(x)$ , where  $\Gamma_{\min}(x)$  denotes the family of minimal cubes in  $\Gamma$ , that is,

$$\Gamma_{\min} = \left\{ Q_x : x \in \bigcup_{Q \in \Gamma} Q \right\}.$$

**3.3. Shaded and Lighted Cubes.** Shaded and lighted cubes were introduced in [14]. We recall the definitions. Given a fixed  $\Gamma \subset \mathcal{D}$ , for any  $Q \in \Gamma$  we define the **Shade** of  $Q$  as the union of all cubes from  $\Gamma$  strictly contained in  $Q$

$$\text{Shade}(Q) = \bigcup \left\{ R : R \in \Gamma, R \subsetneq Q \right\}.$$

We define the **Light** of  $Q$  as  $\text{Light}(Q) = Q \setminus \text{Shade}(Q)$ . It is clear that  $Q \in \Gamma_{\min}$ , if and only if,  $\text{Light}(Q) \neq \emptyset$ , and moreover

$$\bigcup_{Q \in \Gamma} Q = \bigcup_{Q \in \Gamma_{\min}} \text{Light}(Q).$$

Therefore, by (3.4) we can write

$$S_\Gamma(x) \simeq \sum_{Q \in \Gamma_{\min}} \frac{\chi_{\text{Light}(Q)}(x)}{\varphi(w(Q))}, \quad (3.5)$$

where in the last sum there is at most one non-zero term for each  $x$ . We shall classify the cubes as shaded if the shade is a big portion of the cube or lighted if this does not happen. Precisely, a cube  $Q \in \Gamma$  is called **shaded** if  $|\text{Shade}(Q)| > \frac{2^d-1}{2^d}|Q|$ , and we write  $\Gamma_s$  for the collection of cubes from  $\Gamma$  that are shaded. A cube  $Q$  from  $\Gamma$  is called **lighted** if it is not shaded, that is, if  $|\text{Light}(Q)| \geq \frac{1}{2^d}|Q|$ . We write  $\Gamma_L$  for the collection of all cubes from  $\Gamma$  that are lighted.

**Remark 3.2.** Observe that  $\Gamma_L \subset \Gamma_{\min}$  and by Lemma 4.3 in [14] we have

$$\frac{2^d-1}{2^d}(\#\Gamma) \leq (\#\Gamma_L) \leq (\#\Gamma_{\min}) \leq (\#\Gamma), \quad \forall \Gamma \subset \mathcal{D}$$

Now we shall conclude the proof of theorem 3.1.

*Proof.* ( of Theorem 3.1) By (2.14) and (3.5) we know that

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \simeq \left\| \sum_{Q \in \Gamma_{\min}} \frac{\chi_{\text{Light}(Q)}(x)}{\varphi(w(Q))} \right\|_{L^\Phi(w)}. \quad (3.6)$$

Thus it is enough to estimate the quantity in the right side of (3.6). We let  $\lambda = h_\varphi^+(\#\Gamma_{\min})$  so that  $\varphi(w(Q)\#\Gamma_{\min}) \leq \lambda\varphi(w(Q))$  for all  $Q \in \Gamma_{\min}$ . Since  $\{\text{Light}(Q) : Q \in \Gamma_{\min}\}$  is a pairwise disjoint collection and  $\Phi$  is increasing, we have

$$\begin{aligned} \int_{\mathbb{R}^d} \Phi\left(\frac{\sum_{Q \in \Gamma_{\min}} \frac{\chi_{\text{Light}(Q)}(x)}{\varphi(w(Q))}}{\lambda}\right) w(x) dx &= \sum_{Q \in \Gamma_{\min}} \Phi\left(\frac{1}{\lambda\varphi(w(Q))}\right) w(\text{Light}(Q)) \\ &\leq \sum_{Q \in \Gamma_{\min}} \Phi\left(\frac{1}{\varphi(w(Q)\#\Gamma_{\min})}\right) w(Q) = \sum_{Q \in \Gamma_{\min}} \Phi\left(\Phi^{-1}\left(\frac{1}{w(Q)\#\Gamma_{\min}}\right)\right) w(Q) = 1. \end{aligned}$$

Hence by, (3.6), Remark 3.2 and since  $h_\varphi^+$  is non decreasing, we have

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \lesssim h_\varphi^+(\#\Gamma_{\min}) \lesssim h_\varphi^+(\#\Gamma).$$

For the left inequality, by (3.6) and using that  $\Gamma_L \subset \Gamma_{\min}$ , we can write

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \gtrsim \left\| \sum_{Q \in \Gamma_L} \frac{\chi_{\text{Light}(Q)}(x)}{\varphi(w(Q))} \right\|_{L^\Phi(w)}.$$

Now let  $\lambda < h_\varphi^-(2^{-dp}C_w^1(\#\Gamma_L))$  so that  $\lambda\varphi(w(Q)) < \varphi(w(Q)2^{-dp}C_w^1(\#\Gamma_L))$  for all  $Q \in \Gamma_L$ . Using (2.3), and since  $|\text{Light}(Q)| > 2^{-d}|Q|$  for  $Q \in \Gamma_L$ , we deduce, with  $p = p^\Phi$ , that

$$\begin{aligned} \int_{\mathbb{R}^d} \Phi\left(\frac{\sum_{Q \in \Gamma_L} \frac{\chi_{\text{Light}(Q)}(x)}{\varphi(w(Q))}}{\lambda}\right) w(x) dx &= \sum_{Q \in \Gamma_L} \Phi\left(\frac{1}{\lambda\varphi(w(Q))}\right) w(\text{Light}(Q)) \\ &> \sum_{Q \in \Gamma_L} \Phi\left(\frac{1}{\varphi(2^{-dp}C_w^1 w(Q)(\#\Gamma_L))}\right) C_w^1 2^{-dp} w(Q) \end{aligned}$$

$$= \sum_{Q \in \Gamma_L} \Phi \left( \Phi^{-1} \left( \frac{1}{2^{-dp} C_w^1(Q) (\#\Gamma_L)} \right) \right) 2^{-dp} w(Q) C_w^1 = 1.$$

Then by (2.4), and Remark 3.2 we obtain

$$\|\tilde{1}_\Gamma\|_{L^\Phi(w)} \geq h_\varphi^-(2^{-dp} C_w^1(\#\Gamma_L)) \geq h_\varphi^-(C_w^1(2^d - 1) 2^{-d(p+1)}(\#\Gamma)).$$

Now using (2.10) it can be shown that

$$h_\varphi^-(C_w^1(2^d - 1) 2^{-d(p+1)}(\#\Gamma)) \geq C h_\varphi^-(\#\Gamma).$$

□

If  $\Phi(t) = t^p$ , from Theorem 3.1 and Example 2.1 we deduce that admissible wavelet bases are democratic in weighted Lebesgue spaces  $L^p(w)$ .

**Corollary 3.3.** *Let  $\Phi(t) = t^p$ ,  $1 < p < \infty$ ,  $w \in A_p$  a weight in  $\mathbb{R}^d$ , and  $\mathcal{B} = \{\psi_Q : Q \in \mathcal{D}\}$  an admissible wavelet basis. Then*

$$h_r(N; \mathcal{B}, L^p(w)) \approx h_l(N; \mathcal{B}, L^p(w)) \approx N^{\frac{1}{p}} \quad (3.7)$$

#### 4. INCLUSIONS FOR $N$ -TERM APPROXIMATION SPACES OF $L^\Phi(w)$ .

In this section we investigate Jackson and Bernstein type inequalities and the corresponding inclusions for the  $N$ -term approximation spaces  $\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w))$ ,  $\alpha > 0$ ,  $0 < q \leq \infty$ ,  $w \in A_{p,\Phi}(\mathbb{R}^d)$ , where the error of approximation is measured in  $L^\Phi(w)$  (see 1.2). These inclusions are given in terms of the discrete Lorentz spaces  $\Lambda_\eta^q$  (see definition and properties of this spaces in subsection 4.1, bellow).

**4.1. Sequence spaces in  $\mathcal{D}$ .** We recall the definition of some classical sequence spaces over the index set  $\mathcal{D}$  of all dyadic cubes on  $\mathbb{R}^d$ . All of them are subspaces of  $c_o$  and therefore for each sequence  $\{s_Q\}_{Q \in \mathcal{D}}$  we can find an enumeration of the index set  $\mathcal{D} = \{Q_k\}_{k=1}^\infty$  so that  $|s_{Q_1}| \geq |s_{Q_2}| \geq \dots$  and in addition  $\lim_{k \rightarrow \infty} s_{Q_k} = 0$ . We shall always assume that  $\{s_{Q_k}\}_{k \geq 1}$  corresponds to such ordering, which coincides with the *non-increasing rearrangement*  $\mathbf{s}^*$  of the sequence  $\mathbf{s}$ .

Let  $\eta = \{\eta(k)\}_{k \geq 1}$  be a fixed positive increasing sequence so that  $\lim_{k \rightarrow \infty} \eta(k) = \infty$  and  $\eta$  is *doubling* (i.e.  $\eta(2k) \leq C\eta(k)$ ,  $k \geq 1$ ). Then, for each  $0 < q \leq \infty$  we define a *weighted discrete Lorentz space* by

$$\Lambda_\eta^q = \left\{ \mathbf{s} \in c_o : \|\mathbf{s}\|_{\Lambda_\eta^q} = \left[ \sum_{k \geq 1} (\eta(k) |s_{Q_k}|)^q \frac{1}{k} \right]^{\frac{1}{q}} < \infty \right\}.$$

Note that for  $q = \infty$  one writes  $\|\mathbf{s}\|_{\Lambda_\eta^\infty} = \sup_k \eta(k) |s_{Q_k}|$ . These are quasi-Banach rearrangement invariant spaces, which are Banach when  $q \geq 1$  and  $\{\frac{\eta(k)^q}{k}\}_k$  is non-increasing ([3], p. 28). When  $q = 1$  or  $q = \infty$  we shall write, respectively,  $\Lambda_\eta$  and  $\mathbb{M}_\eta$  (the latter called Marcinkiewicz space). The particular case  $\eta(k) = k^{\frac{1}{\tau}}$  gives the classical (discrete) Lorentz space  $\Lambda_\eta^q = \ell^{\tau,q}(\mathcal{D})$ . The spaces  $\Lambda_\eta^q$  for general  $\eta$ , and in particular, their interpolation properties, have been studied, e.g., in [3, 24, 27]. In our applications we use the sequences  $\{k^\alpha h_\varphi^\pm(k)\}_{k \geq 1}$ , for  $\alpha > 0$ , which always satisfy the required assumptions.

Given a fixed sequence space  $\mathfrak{s}$  as above, we define a new sequence space  $\mathfrak{s}(L^\Phi(w))$  isomorphic to  $\mathfrak{s}$ , by

$$\mathfrak{s}(L^\Phi(w)) = \left\{ f = \sum_{Q \in \mathcal{D}} \langle f, \psi_Q \rangle \psi_Q \in L^\Phi(w) : \{ \|\langle f, \psi_Q \rangle \psi_Q\|_{L^\Phi(w)} \}_Q \in \mathfrak{s} \right\},$$

with  $\|f\|_{\mathfrak{s}(L^\Phi(w))} = \left\| \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} \right\|_{\mathfrak{s}}$ . Such definitions appear naturally in relation with approximation when the basis is not normalized (see, e.g., [13]).

**4.2. Jackson type inequalities.** In order to obtain the left embedding of the inclusions of approximation spaces given in Theorem 1.2, we start by proving some inequalities of Jackson Type.

**Proposition 4.1.** *Let  $\Phi$  be a Young function so that  $0 < i_\varphi \leq I_\varphi < 1$ ,  $w \in A_{p^\Phi}$  a weight in  $\mathbb{R}^d$ , and let  $\alpha > 0$ . Let  $\mathcal{B}$  be an admissible wavelet basis. Then, there exists  $C > 0$  such that for every  $f \in \mathbb{M}_{k^\alpha h_\varphi^+(k)}(\mathcal{B}, L^\Phi(w))$  we have*

$$\|f - G_{N-1}(f)\|_{L^\Phi(w)} \leq CN^{-\alpha} \|f\|_{\mathbb{M}_{k^\alpha h_\varphi^+(k)}(\mathcal{B}, L^\Phi(w))}, \quad \forall N \geq 1. \quad (4.1)$$

*Proof.* By the triangle inequality and (1.3) we have

$$\begin{aligned} \|f - G_{N-1}(f)\|_{L^\Phi(w)} &= \left\| \sum_{k \geq N} \langle f, \psi_{Q_k} \rangle \psi_{Q_k} \right\|_{L^\Phi(w)} \leq \sum_{j=0}^{\infty} \left\| \sum_{2^j N \leq k < 2^{j+1} N} \langle f, \psi_{Q_k} \rangle \psi_{Q_k} \right\|_{L^\Phi} \\ &\leq \sum_{j=0}^{\infty} \left\| \langle f, \psi_{Q_{2^j N}} \rangle \psi_{Q_{2^j N}} \right\|_{L^\Phi(w)} \left\| \sum_{2^j N \leq k < 2^{j+1} N} \frac{\psi_{Q_k}}{\|\psi_{Q_k}\|_{L^\Phi(w)}} \right\|_{L^\Phi(w)} \\ &\lesssim \sum_{j=0}^{\infty} \|\langle f, \psi_{2^j N} \rangle \psi_{2^j N}\|_{L^\Phi(w)} h_\varphi^+(2^j N) \end{aligned} \quad (4.2)$$

where in the last inequality we have used Theorem 3.1. Now using that  $\frac{h_\varphi^+(k)}{k}$  is non-increasing (this follows from the fact that  $\frac{\varphi(t)}{t}$  is non-increasing for all  $t > 0$ , see [2]) and the definition of the Marcinkiewicz space we have

$$\begin{aligned} \sum_{j=0}^{\infty} \|\langle f, \psi_{Q_{2^j N}} \rangle \psi_{Q_{2^j N}}\|_{L^\Phi(w)} h_\varphi^+(2^j N) &= \sum_{j=0}^{\infty} \sum_{2^{j-1} N \leq k < 2^j N} \|\langle f, \psi_{Q_{2^j N}} \rangle \psi_{Q_{2^j N}}\|_{L^\Phi(w)} \frac{h_\varphi^+(2^j N)}{2^{j-1} N} \\ &\leq 2 \sum_{k > \frac{N}{2}} \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} \frac{h_\varphi^+(k)}{k} \leq C \|f\|_{\mathbb{M}_{k^\alpha h_\varphi^+(k)}(\mathcal{B}, L^\Phi(w))} \sum_{k > \frac{N}{2}} k^{-\alpha} \frac{1}{k} \\ &\leq CN^{-\alpha} \|f\|_{\mathbb{M}_{k^\alpha h_\varphi^+(k)}(\mathcal{B}, L^\Phi(w))}. \end{aligned} \quad (4.3)$$

□

The previous result can be translated as the following inclusion for approximation spaces

$$\mathbb{M}_{k^\alpha h_\varphi^+(k)}(\mathcal{B}, L^\Phi(w)) \hookrightarrow \mathcal{A}_\infty^\alpha(\mathcal{B}, L^\Phi(w)). \quad (4.4)$$

**4.3. Bernstein type inequalities.** Bernstein type estimates are useful to obtain the right hand inclusions for approximation spaces of Theorem 1.2.

**Proposition 4.2.** *Let  $\Phi$  be a Young function such that  $0 < i_\varphi \leq I_\varphi < 1$ ,  $w \in A_{p,\Phi}$  a weight in  $\mathbb{R}^d$ , and let  $\alpha > 0$ . Let  $\mathcal{B}$  be an admissible basis. Then, there exists  $C > 0$  so that, for all  $N \geq 1$ , and all  $f \in \Sigma_N$*

$$\|f\|_{\Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w))} \leq CN^\alpha \|f\|_{L^\Phi(w)}. \quad (4.5)$$

*Proof.* Let  $f = \sum_{j=1}^N \langle f, \psi_{Q_j} \rangle \psi_{Q_j} \in \Sigma_N$ , written in such a way that  $\|\langle f, \psi_{Q_1} \rangle \psi_{Q_1}\|_{L^\Phi(w)} \geq \|\langle f, \psi_{Q_2} \rangle \psi_{Q_2}\|_{L^\Phi(w)} \geq \dots$ . For  $1 \leq k \leq N$ , using Theorem 3.1

$$\begin{aligned} \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} h_\varphi^-(k) &\leq C \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} \left\| \sum_{j=1}^k \frac{\psi_{Q_j}}{\|\psi_{Q_j}\|_{L^\Phi(w)}} \right\|_{L^\Phi(w)} \\ &\leq C \|G_N(f)\|_{L^\Phi(w)}. \end{aligned} \quad (4.6)$$

By (4.6) we have

$$\begin{aligned} \|f\|_{\Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w))} &= \sum_{k=1}^N k^\alpha h_\varphi^-(k) \|\langle f, \psi_{Q_k} \rangle \psi_{Q_k}\|_{L^\Phi(w)} \frac{1}{k} \leq C \|G_N(f)\|_{L^\Phi(w)} \sum_{k=1}^N \frac{k^\alpha}{k} \\ &\leq C' N^\alpha \|f\|_{L^\Phi(w)}. \end{aligned}$$

□

As before, the above result can be stated as an inclusion for approximation spaces. Below, the number  $\rho_\alpha \in (0, 1]$  is chosen so that the quasi-normed space  $\Lambda_{k^\alpha h_\varphi^-(k)}$  satisfies the  $\rho_\alpha$ -triangle inequality, that is,

$$\|\mathbf{s}_1 + \mathbf{s}_2\|_{\Lambda_{k^\alpha h_\varphi^-(k)}^{\rho_\alpha}} \leq \|\mathbf{s}_1\|_{\Lambda_{k^\alpha h_\varphi^-(k)}^{\rho_\alpha}} + \|\mathbf{s}_2\|_{\Lambda_{k^\alpha h_\varphi^-(k)}^{\rho_\alpha}}. \quad (4.7)$$

**Corollary 4.3.** *Let  $\alpha > 0$ . Then, with the same hypothesis as in Proposition 4.2, we have*

$$\mathcal{A}_{\rho_\alpha}^\alpha(\mathcal{B}, L^\Phi(w)) \hookrightarrow \Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w)) \quad (4.8)$$

*Proof.* The argument for (4.8) is standard (see, e.g., [7]). It suffices to prove that

$$\|f\|_{\Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w))} \leq C \|f\|_{\mathcal{A}_{\rho_\alpha}^\alpha(\mathcal{B}, L^\Phi(w))}, \quad \forall f \in \Sigma_N, \quad N \geq 1$$

with a constant  $C > 0$  independent of  $N$  and one obtains the desired inclusion by letting  $N \rightarrow \infty$ . We also assume that  $N = 2^J$ . Now, write  $f = \sum_{j=0}^J [f^{(j)} - f^{(j-1)}]$ , where by convection  $f^{(J)} = f$ ,  $f^{(-1)} = 0$  and  $f^{(j)} \in \Sigma_{2^j}$  is so that  $\|f - f^{(j)}\|_{L^\Phi(w)} \leq 2\sigma_{2^j}(f)_{L^\Phi(w)}$ ,  $0 \leq j < J$ . Then applying, (4.7) and Proposition 4.2 to  $f^{(j)} - f^{(j-1)} \in \Sigma_{2^{j+1}}$  we obtain

$$\begin{aligned} \|f\|_{\Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w))} &\leq \left[ \sum_{j=0}^J \|f^{(j)} - f^{(j-1)}\|_{\Lambda_{k^\alpha h_\varphi^-(k)}^{\rho_\alpha}(\mathcal{B}, L^\Phi(w))} \right]^{\frac{1}{\rho_\alpha}} \\ &\leq C \left[ \sum_{j=0}^J 2^{j\alpha\rho} \|f^{(j)} - f^{(j-1)}\|_{L^\Phi(w)} \right]^{\frac{1}{\rho}}. \end{aligned}$$

Now, by assumption, for  $1 \leq j \leq J$

$$\|f^{(j)} - f^{(j-1)}\|_{L^\Phi(w)} \leq \|f^{(j)} - f\|_{L^\Phi(w)} + \|f - f^{(j-1)}\|_{L^\Phi(w)} \leq 4\sigma_{2^{j-1}}(f)_{L^\Phi(w)}.$$

On the other hand, for  $j = 0$  we have

$$\|f^{(0)} - f^{(-1)}\|_{L^\Phi(w)} = \|f^{(0)}\|_{L^\Phi(w)} \leq \|f^{(0)} - f\|_{L^\Phi(w)} + \|f\|_{L^\Phi(w)} \leq 2\sigma_1(f)_{L^\Phi(w)} + \|f\|_{L^\Phi(w)}.$$

Hence,

$$\|f\|_{\Lambda_{k^\alpha h_\varphi^-(k)}(\mathcal{B}, L^\Phi(w))} \leq C \left[ \|f\|_{L^\Phi(w)} + \sum_{j=0}^{J-1} (2^{j\alpha} \sigma_{2^j}(f)_{L^\Phi(w)})^\rho \right]^{\frac{1}{\rho}} \approx \|f\|_{\mathcal{A}_\rho^\alpha(\mathcal{B}, L^\Phi(w))}.$$

□

Finally, using real interpolation we can obtain inclusions for the whole family of approximation spaces  $\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w))$ ,  $0 < q \leq \infty$ . For this we consider the interpolation properties of the sequence spaces  $\Lambda_\eta^q$ , namely,

$$(\Lambda_{k^{\alpha_0 \eta(k)}}^r, \Lambda_{k^{\alpha_1 \eta(k)}}^r)_{\alpha, q} = \Lambda_{k^{\alpha \eta(k)}}^q, \quad \alpha = (1 - \theta)\alpha_0 + \theta\alpha_1, \quad (4.9)$$

for all  $0 < q, r \leq \infty$ ,  $0 < \theta < 1$  (see, e.g., [27] Proposition 6.2, [24], Theorem 3).

**Theorem 4.4.** *Let  $\Phi$  be a Young function such that  $0 < i_\varphi \leq I_\varphi < 1$ ,  $w \in A_{p^\Phi}$  a weight in  $\mathbb{R}^d$ ,  $\alpha > 0$ , and  $0 < q \leq \infty$ . Let  $\mathcal{B}$  be an admissible wavelet basis. Then*

$$\Lambda_{k^\alpha h_\varphi^+(k)}^q(\mathcal{B}, L^\Phi(w)) \hookrightarrow \mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) \hookrightarrow \Lambda_{k^\alpha h_\varphi^-(k)}^q(\mathcal{B}, L^\Phi(w)). \quad (4.10)$$

*Proof.* Let  $\alpha_0 < \alpha < \alpha_1$ , so that  $\alpha = (\alpha_0 + \alpha_1)/2$ . Then, for every  $0 < q, r \leq \infty$  we have (see, e.g., [7])

$$\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) = (\mathcal{A}_r^{\alpha_0}(\mathcal{B}, L^\Phi(w)), \mathcal{A}_r^{\alpha_1}(\mathcal{B}, L^\Phi(w)))_{\frac{1}{2}, q}.$$

Letting  $r = \min(\rho_{\alpha_0}, \rho_{\alpha_1})$  and using (4.8)

$$\begin{aligned} \mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) &= (\mathcal{A}_r^{\alpha_0}(\mathcal{B}, L^\Phi(w)), \mathcal{A}_r^{\alpha_1}(\mathcal{B}, L^\Phi(w)))_{\frac{1}{2}, q} \\ &\hookrightarrow (\Lambda_{k^{\alpha_0 h_\varphi^-(k)}}(\mathcal{B}, L^\Phi(w)), \Lambda_{k^{\alpha_1 h_\varphi^-(k)}}(\mathcal{B}, L^\Phi(w)))_{\frac{1}{2}, q} \\ &= \Lambda_{k^\alpha h_\varphi^-(k)}^q(\mathcal{B}, L^\Phi(w)), \end{aligned}$$

where the last equality follows from (4.9). Similarly, by (4.4)

$$\begin{aligned} \mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) &= (\mathcal{A}_\infty^{\alpha_0}(\mathcal{B}, L^\Phi(w)), \mathcal{A}_\infty^{\alpha_1}(\mathcal{B}, L^\Phi(w)))_{\frac{1}{2}, q} \\ &\leftrightarrow (\mathbb{M}_{k^{\alpha_0 h_\varphi^+(k)}}(\mathcal{B}, L^\Phi(w)), \mathbb{M}_{k^{\alpha_1 h_\varphi^+(k)}}(\mathcal{B}, L^\Phi(w))) \\ &= \Lambda_{k^\alpha h_\varphi^+(k)}^q(\mathcal{B}, L^\Phi(w)). \end{aligned}$$

□

We now prove that the inclusions (4.10) are optimal. To state the Theorem we write  $\mathcal{D}$  for the class of sequences  $\eta = \{\eta(k)\}_{k=1}^\infty$  that are increasing, and doubling.

**Theorem 4.5.** *Same hypothesis as in Theorem 4.4. For fixed  $\alpha > 0$  and  $q$ ,  $0 < q \leq \infty$ , the inclusions given in (4.10) are best possible in the scale of weighted Lorentz spaces  $\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))$  where  $\eta \in \mathcal{D}$*



*Proof.* Suppose  $\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w)) \hookrightarrow \mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w))$ . We want to prove that  $h_\varphi^+(N) \lesssim \eta(N)$  for all  $N = 1, 2, \dots$ . By definition of  $h_\varphi^+(N)$  we can choose  $\tau = \tau(N) > 0$  such that

$$\frac{\varphi(N\tau)}{\varphi(\tau)} \leq h_\varphi^+(N) \leq 2 \frac{\varphi(N\tau)}{\varphi(\tau)}. \quad (4.11)$$

By Lemma 2.6 we can choose a sequence of pairwise disjoint cubes  $\Gamma = \{R_j\}_{j=1}^{2N}$  such that  $w(R_j) \approx \tau$ . Let  $\tilde{1}_\Gamma = \sum_{j=1}^{2N} \frac{\psi_{R_j}}{\|\psi_{R_j}\|_{L^\Phi(w)}}$ . By Theorem 2.2,  $\|f\|_{L^\Phi(w)}$  is equivalent to the lattice norm  $\|S_\psi(f)\|_{L^\Phi(w)}$ ; thus there exists  $\Gamma' \subset \Gamma$  with  $|\Gamma'| = N$  such that  $\sigma_N(\tilde{1}_\Gamma) \approx \|\tilde{1}_{\Gamma'}\|_{L^\Phi(w)}$  (see (2.6) in [13]). Thus, by (2.28) and (4.11)

$$\sigma_N(\tilde{1}_\Gamma)_{L^\Phi(w)} \approx \|\tilde{1}_{\Gamma'}\|_{L^\Phi(w)} \approx \frac{\varphi(N\tau)}{\varphi(\tau)} \approx h_\varphi^+(N).$$

Hence,

$$\|\tilde{1}_\Gamma\|_{\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w))} \geq \left( \sum_{k=N/2}^N k^{\alpha q} \sigma_k(\tilde{1}_\Gamma)^q \frac{1}{k} \right)^{\frac{1}{q}} \gtrsim \sigma_N(\tilde{1}_\Gamma) N^\alpha \approx N^\alpha h_\varphi^+(N). \quad (4.12)$$

On the other hand

$$\|\tilde{1}_\Gamma\|_{\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))} = \left( \sum_{k=1}^{2N} (k^\alpha \eta(k))^q \frac{1}{k} \right)^{\frac{1}{q}} \lesssim \eta(2N) N^\alpha \lesssim \eta(N) N^\alpha \quad (4.13)$$

by the doubling property of  $\eta$ . The inequalities (4.12) and (4.13) together with our assumption imply the desired result.

Suppose now that  $\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w)) \hookrightarrow \Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))$ . We want to prove that  $\eta(k) \leq h_\varphi^-(N)$  for all  $N = 1, 2, \dots$ . Let  $\Gamma \subset \mathcal{D}$  with  $|\Gamma| = N$ . Write  $\tilde{1}_\Gamma = \sum_{Q \in \Gamma} \frac{\psi_Q}{\|\psi_Q\|_{L^\Phi(w)}}$ . Since  $\sigma_k(\tilde{1}_\Gamma) \leq \|\tilde{1}_\Gamma\|_{L^\Phi(w)}$  for all  $k = 1, 2, \dots, N$ , our hypothesis imply

$$\|\tilde{1}_\Gamma\|_{\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))} \lesssim \|\tilde{1}_\Gamma\|_{\mathcal{A}_q^\alpha(\mathcal{B}, L^\Phi(w))} \lesssim N^\alpha \|\tilde{1}_\Gamma\|_{L^\Phi(w)}. \quad (4.14)$$

On the other hand

$$\|\tilde{1}_\Gamma\|_{\Lambda_{k^\alpha \eta(k)}^q(\mathcal{B}, L^\Phi(w))} \geq \left( \sum_{k=\frac{N}{2}}^N (\eta(k) k^\alpha)^q \frac{1}{k} \right)^{\frac{1}{q}} \gtrsim N^\alpha \eta(N/2) \gtrsim N^\alpha \eta(N) \quad (4.15)$$

since  $\eta$  is doubling. By (4.14) and (4.15) we have  $\eta(N) \lesssim \|\tilde{1}_\Gamma\|_{L^\Phi(w)}$  for all  $\Gamma \subset \mathcal{D}$ , with  $|\Gamma| = N$ . Taking the infimum over all  $\Gamma \in \mathcal{D}$ , with  $|\Gamma| = N$ , we obtain  $\eta(N) \leq h_l(N; \mathcal{B}, L^\Phi(w)) \approx h_\varphi^-(N)$  by Theorem 3.1.  $\square$

5. APPROXIMATION SPACES FOR  $L^p(w)$ 

Corollary 1.3 is now an easy consequence of Theorem 4.4 and Corollary 3.3. The rest of this section is devoted to prove Theorem 1.4 ( see Theorem 5.4 bellow).

The approximation spaces  $\mathcal{A}_\tau^\gamma(\mathcal{B}, L^p(w))$  can also be identified with weighted Besov spaces. Our definition of weighted Besov spaces is borrowed from [29, 30], and it is modeled on the corresponding definition of Besov spaces without weights developed in [26] (see also [9] and [10]).

We say that a function  $\varphi \in \mathcal{S}(\mathbb{R}^d)$  belongs to the class of admissible kernels if  $\text{Supp}\widehat{\varphi} \subset \{\xi \in \mathbb{R}^d : \frac{1}{2} < |\xi| < 2\}$  and  $|\widehat{\varphi}(\xi)| \geq c > 0$ , if  $\frac{3}{5} < |\xi| < \frac{5}{3}$ . Set  $\varphi_k(x) = 2^{kd}\varphi(2^kx)$  for  $k \in \mathbb{Z}$ .

Let  $\alpha \in \mathbb{R}$ ,  $1 \leq p < \infty$ ,  $0 < q \leq \infty$ ,  $\varphi$  admissible kernel, and  $w$  an  $A_p$  weight on  $\mathbb{R}^d$ . The homogeneous weighted Besov space  $\dot{B}_{p,q}^\alpha(w)$  is the set of all tempered distributions  $f \in \mathcal{S}'/\mathcal{P}$  (modulo polynomials) such that

$$\|f\|_{\dot{B}_{p,q}^\alpha(w)} = \left[ \sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\varphi_k * f\|_{L^p(w)})^q \right]^{1/q} < \infty. \quad (5.1)$$

This definition depends initially of the choice of admissible  $\varphi$ . It can be proved (see Theorem 1.8 in [29] or [30]) that this is independent of the choice of admissible  $\varphi$ . Also, the spaces  $\dot{B}_{p,q}^\alpha(w)$  are (quasi)-Banach spaces (see section 4.4 of [30]).

Let  $\Psi = \{\psi^l : l = 1, 2, \dots, 2^{d-1}\}$  be an orthonormal wavelet family in  $L^2(\mathbb{R}^d)$  constructed from the 1-dimensional Lemarié-Meyer wavelets (see [16, 23, 25]). Write  $s_Q^l = \langle f, \psi_Q^l \rangle$ ,  $Q \in \mathcal{D}$ ,  $l = 1, 2, \dots, 2^{d-1}$  for the wavelet coefficients.

**Proposition 5.1.** (see Theorem 10.2 in [29] or Theorem 6.2 in [30]).

Let  $\alpha \in \mathbb{R}$ ,  $0 < q \leq \infty$ ,  $1 \leq p < \infty$  and let  $w$  be an  $A_p$  weight in  $\mathbb{R}^d$ . Let  $\Psi$  be a family of Lemarié-Meyer wavelets as defined above. Then

$$\|f\|_{\dot{B}_{p,q}^\alpha(w)} \approx \sum_{l=1}^{2^{d-1}} \left[ \sum_{j \in \mathbb{Z}} \left( \sum_{|Q|=2^{-jd}} (|Q|^{-\frac{\alpha}{d}-\frac{1}{2}} |s_Q^l| w(Q)^{\frac{1}{p}})^\tau \right)^{\frac{q}{\tau}} \right]^{\frac{1}{q}}. \quad (5.2)$$

**Remark 5.2.** It is also proved in Theorem 10.2 of [29] and Theorem 6.2 of [30] that the condition  $w$  doubling, that is, there exists  $C > 0$  such that

$$\int_{B_{2\delta}(z)} w(x) dx \leq C \int_{B_\delta(z)} w(x) dx, \quad \forall z \in \mathbb{R}^d \text{ and } \forall \delta > 0,$$

is sufficient to guarantee the equivalence (5.2).

**Remark 5.3.** Equivalence (5.2) also holds for the family  ${}_N\Psi = \{N\psi^l : l = 1, \dots, 2^{d-1}\}$  constructed from the 1-dimensional Daubechies compactly supported wavelets (see [5]), provided  $N$  is sufficiently large (see [29]).

**Theorem 5.4.** Let  $\gamma > 0$ ,  $1 < p < \infty$ . We have

$$\mathcal{A}_\tau^{\gamma/d}(\Psi, L^p(w)) = l^\tau(\Psi, L^p(w)) = \dot{B}_{\tau,\tau}^\gamma(w^{\frac{\tau}{p}}), \quad \text{whenever } \frac{1}{\tau} = \frac{\gamma}{d} + \frac{1}{p}, \quad (5.3)$$

for all  $w \in A_\tau(\mathbb{R}^d)$  and all orthonormal wavelet families  $\Psi$  for which (5.2) holds for  $\dot{B}_{\tau,\tau}^{\gamma/d}(w)$ .

For the proof we shall need the following lemma:

**Lemma 5.5.** *Let  $w \in A_r$  be a weight in  $\mathbb{R}^d$ ,  $r \geq 1$ ,  $0 < \delta < 1$ , and  $u(x) = w(x)^\delta$ . Then  $u \in A_r$ , and  $w_Q \approx (u_Q)^{\frac{1}{\delta}}$ , where*

$$w_Q = \frac{1}{|Q|} \int_Q w(x) dx.$$

*Proof.* When  $r > 1$ , since  $w \in A_r$  and  $0 < \delta < 1$ , using Jensen's inequality we have

$$\begin{aligned} & \left( \frac{1}{|Q|} \int_Q u(x) dx \right) \left( \frac{1}{|Q|} \int_Q u^{1-r'}(x) dx \right)^{r-1} \\ &= \left( \frac{1}{|Q|} \int_Q w(x)^\delta dx \right) \left( \frac{1}{|Q|} \int_Q w(x)^{\delta(1-r')} dx \right)^{r-1} \\ &\leq \left[ \left( \frac{1}{|Q|} \int_Q w(x) dx \right) \left( \frac{1}{|Q|} \int_Q w(x)^{1-r'} dx \right)^{r-1} \right]^\delta \leq C_w^\delta. \end{aligned}$$

Thus,  $u \in A_r$ . Now we shall to prove the equivalence  $w_Q \approx (u_Q)^{\frac{1}{\delta}}$ . On the one hand, using Jensen's inequality with  $\delta < 1$ , we have

$$(u_Q)^{\frac{1}{\delta}} = \left( \frac{1}{|Q|} \int_Q w(x)^\delta dx \right)^{\frac{1}{\delta}} \leq \left( \frac{1}{|Q|} \int_Q w(x) dx \right) = w_Q.$$

On the other hand, since  $h(t) = t^{-(r-1)\delta}$ ,  $t > 0$ , is a convex function, using again Jensen's inequality we have

$$\left( \frac{1}{|Q|} \int_Q w^{1-r'}(x) dx \right)^{-(r-1)\delta} \leq \left( \frac{1}{|Q|} \int_Q w(x)^\delta dx \right) = u_Q.$$

From the condition  $w \in A_r$  it follows that

$$w_Q = \left( \frac{1}{|Q|} \int_Q w(x) dx \right) \leq C_w \left( \frac{1}{|Q|} \int_Q w^{1-r'}(x) dx \right)^{-(r-1)} \leq C(u_Q)^{\frac{1}{\delta}}.$$

For  $r = 1$ , on the one hand, since  $w \in A_1$ , for almost all  $x \in Q$ , using again Jensen's inequality we have

$$\left( \frac{1}{|Q|} \int_Q u(x) dx \right) = \left( \frac{1}{|Q|} \int_Q w^\delta(x) dx \right) \leq \left( \frac{1}{|Q|} \int_Q w(x) dx \right)^\delta \leq C w(x)^\delta = C u(x).$$

Thus  $u \in A_1$ . On the other hand we can use again Jensen's inequality and obtain,  $(u_Q)^{\frac{1}{\delta}} \leq w_Q$ . Moreover, the condition  $w \in A_1$  implies that

$$w_Q = \frac{1}{|Q|} \int_Q w(x) dx \leq C \operatorname{ess}_Q \inf w = C(\operatorname{ess}_Q \inf u)^{\frac{1}{\delta}} \leq C \left( \frac{1}{|Q|} \int_Q u(x) dx \right)^{\frac{1}{\delta}} = C(u_Q)^{\frac{1}{\delta}}$$

□

*Proof. (of Theorem 5.4).* The first equality in (5.3) follows from Corollary 1.3 (with  $\tau = q$  and  $\alpha = \frac{\gamma}{d}$ ). For the second equality, observe that for a single element of the basis  $\Psi$ , we have

$$\|\psi_Q\|_{L^p(w)} \approx \left\| \left( \frac{\chi_Q(x)}{|Q|} \right)^{\frac{1}{2}} \right\|_{L^p(w)} = |Q|^{-\frac{1}{2}} \|\chi_Q(x)\|_{L^p(w)} = |Q|^{-\frac{1}{2}} \left( \int_Q w(x) dx \right)^{\frac{1}{p}}$$

$$= |Q|^{-\frac{1}{2}} w(Q)^{\frac{1}{p}}. \quad (5.4)$$

Let  $u(x) = w(x)^{\frac{\tau}{p}}$ . By Lemma 5.5 with  $r = \tau$  and  $\delta = \frac{\tau}{p} < 1$  we deduce that  $u \in A_\tau \subset A_p$  and  $w_Q^{\frac{1}{p}} \approx (u_Q)^{\frac{1}{\tau}}$ . Thus, since  $w(Q) = |Q|w_Q$  we obtain

$$\begin{aligned} \|f\|_{\ell^\tau(\Psi, L^p(w))} &= \left\| \left\| \langle f, \psi_{Q_k} \rangle \psi_{Q_k} \right\|_{L^p(w)} \right\|_{\ell^\tau} \approx \left( \sum_{Q \in \mathcal{D}} (|Q|^{-\frac{1}{2} + \frac{1}{p}} |\langle f, \psi_Q \rangle| (w_Q)^{\frac{1}{p}})^\tau \right)^{\frac{1}{\tau}} \\ &\approx \left( \sum_{Q \in \mathcal{D}} (|Q|^{-\frac{1}{2} + \frac{1}{p}} |\langle f, \psi_Q \rangle| (u_Q)^{\frac{1}{\tau}})^\tau \right)^{\frac{1}{\tau}} \\ &= \left( \sum_{Q \in \mathcal{D}} (|Q|^{-\frac{1}{2} - \frac{\gamma}{d} + \frac{1}{p}} |\langle f, \psi_Q \rangle| (u_Q)^{\frac{1}{\tau}})^\tau \right)^{\frac{1}{\tau}} \\ &= \left( \sum_{Q \in \mathcal{D}} (|Q|^{-\frac{1}{2} - \frac{\gamma}{d}} |\langle f, \psi_Q \rangle| (u(Q))^{\frac{1}{\tau}})^\tau \right)^{\frac{1}{\tau}} \approx \|f\|_{\dot{B}_{\tau, \tau}^{\gamma}(w^{\frac{\tau}{p}})}. \end{aligned}$$

□

As a corollary we prove a non-trivial interpolation result.

**Corollary 5.6.** *Let  $\gamma > 0$ ,  $1 < p < \infty$ ,  $\frac{1}{\tau} = \frac{\gamma}{d} + \frac{1}{p}$ , and  $w \in A_\tau(\mathbb{R}^d)$ . Let  $\Psi$  be an orthonormal wavelet family for which (5.2) holds for the Besov spaces involved in this Corollary. For  $0 < \theta < 1$  we have*

$$(L^p(w), \dot{B}_{\tau, \tau}^{\gamma}(w^{\tau/p}))_{\theta, \tau\theta} = \dot{B}_{\tau\theta, \tau\theta}^{\theta\gamma}(w^{\tau\theta/p})$$

where  $\frac{1}{\tau\theta} = \frac{\theta\gamma}{d} + \frac{1}{p}$ .

*Proof.* If  $\Phi(t) = t^p$ , use Proposition 4.1, the continuous embedding  $l^\tau \hookrightarrow l^{\tau, \infty}$ , and Theorem 5.4 to obtain, for all  $N = 1, 2, \dots$

$$\begin{aligned} \sigma_N(f)_{L^p(w)} &\leq CN^{-(\frac{1}{\tau} - \frac{1}{p})} \|f\|_{l^{\tau, \infty}(\Psi, L^p(w))} \leq CN^{-(\frac{1}{\tau} - \frac{1}{p})} \|f\|_{l^\tau(\Psi, L^p(w))} \\ &\leq CN^{-(\frac{1}{\tau} - \frac{1}{p})} \|f\|_{\dot{B}_{\tau, \tau}^{\gamma}(w^{\tau/p})}, \end{aligned} \quad (5.5)$$

whenever  $\frac{1}{\tau} = \frac{\gamma}{d} + \frac{1}{p}$ .

From Theorem 5.4, and Proposition 4.2 we obtain, for all  $g \in \Sigma_N$ ,  $N = 1, 2, 3, \dots$

$$\|g\|_{\dot{B}_{\tau, \tau}^{\gamma}(w^{\tau/p})} \leq C \|g\|_{l^\tau(\Psi, L^p(w))} \leq CN^{\frac{1}{\tau} - \frac{1}{p}} \|g\|_{L^p(w)}, \quad (5.6)$$

whenever  $\frac{1}{\tau} = \frac{\gamma}{d} + \frac{1}{p}$ . From (5.5), (5.6), and the general theory developed by R. DeVore and V. A. Popov (see Theorem 3.1 in [7]) we deduce

$$\mathcal{A}_q^{\frac{\theta\gamma}{d}}(\Psi, L^p(w)) = (L^p(w), \dot{B}_{\tau, \tau}^{\gamma}(w^{\tau/p}))_{\theta, q} \quad (5.7)$$

whenever  $0 < q \leq \infty$  and  $0 < \theta < 1$ . We use again Theorem 5.4 to obtain

$$\mathcal{A}_{\tau\theta}^{\frac{\theta\gamma}{d}}(\Psi, L^p(w)) = \dot{B}_{\tau\theta, \tau\theta}^{\theta\gamma}(w^{\tau\theta/p})$$

when  $\frac{1}{\tau\theta} = \frac{\theta\gamma}{d} + \frac{1}{p}$ . The result follows from (5.7) with  $q = \tau\theta$ . □

## REFERENCES

- [1] H.A. AIMAR, A.L. BERNARDIS, AND F.J. MARTÍN-REYS, *Multiresolution Approximation and Wavelet Bases of Weighted Lebesgue Spaces*, J. Fourier Anal. and Appl., 9, n 5, (2003), 497-510.
- [2] C. BENETT, AND R.C. SHARPLEY, *Interpolation of Operators*, Pure and Appl. Math. 129, Academic Press, 1988.
- [3] M.J. CARRO, J. RAPOSO, AND J. SORIA, *Recent developments in the theory of Lorentz spaces and weighted inequalities*, Mem. Amer. Math. Soc. 107 (2007).
- [4] D. CRUZ-URIBE, J.M. MARTELL, AND C. PÉREZ, *Weights, Extrapolation and Theory of Rubio de Francia*, Preprint, (2008).
- [5] I. DAUBECHIES, *Orthonormal bases of compactly supported wavelets*, Comm. Pure Appl. Math., 2, (1986), 1–18.
- [6] R. DEVORE, G. PETROVA, AND V. TEMLYAKOV, *Best basis selection for approximation in  $L^p$* , Found. Comput. Math. 3, (2003), 161–185.
- [7] R. DEVORE AND V.A. POPOV, *Interpolation spaces and nonlinear approximation*, Function spaces and applications (Lund, 1986), Lecture Notes in Math., 1302, Springer, Berlin, (1988), 191–205.
- [8] S.J. DILWORTH, N.J. KALTON, D. KUTZAROVA, AND V.N. TEMLYAKOV, *The Thresholding Greedy Algorithm, Greedy Bases, and Duality*, Constructive Approximation, 19, (2003), 575–597.
- [9] M. FRAZIER AND B. JAWERTH, *Decomposition of Besov Spaces*, Indiana Univ. Math. J., 34, (1985), 777–799.
- [10] M. FRAZIER AND B. JAWERTH, *A discrete transform and decomposition of distribution spaces*, J. Func. Anal., 93, (1990), 34–170.
- [11] J. GARCIA-CUERVA AND J.L. RUBIO DE FRANCIA, *Weighted Norm Inequalities and Related Topics*, North Holland Mathematical Studies 116, Elsevier Sciences Publishers, 1985.
- [12] J. GARCÍA-CUERVA AND J.M. MARTELL, *Wavelet characterization of weighted spaces*, J. Geom. Anal. 11, no. 2,(2001), 241–262.
- [13] G. GARRIGÓS AND E. HERNÁNDEZ, *Sharp Jackson and Bernstein Inequalities for  $n$ -term Approximation in Sequence Spaces with Applications*, Indiana Univ. Math. J., 53 (2004), 1739–1762.
- [14] G. GARRIGÓS, E.HERNÁNDEZ, AND J.M. MARTELL, *Wavelets, Orlicz and Greedy bases*, Appl. Compt. Harmon. Anal., 24, (2008), 70–93.
- [15] R. GRIBONVAL, AND M. NIELSEN, *Some Remarks on non-linear approximation with Schauder bases*, East J. Approx. 7, (3), (2001), 267–285.
- [16] E. HERNÁNDEZ AND G.WEISS, *A first course on wavelets*, CRC Press, Boca Raton FL, 1996.
- [17] C. HSIAO, B. JAWERTH, B.J. LUCIER, AND X.M. YU, *Near optimal compression of almost optimal wavelet expansions*, Wavelets: mathematics and applications, Stud. Adv. Math., CRC, Boca Raton, FL, (1994), 425–446.
- [18] A. KAMONT AND V. N. TEMLYAKOV, *Greedy approximation and the multivariate Haar system*, Studia Math. 161, no. 3,(2004), 199–223.
- [19] G. KERKIACHARYAN, AND D. PICARD, *Nonlinear Approximation and Mckenhaupt Weights*, Constr. Approx. 20, (2006), 123–156.
- [20] V. KOKILASHVILI, AND M. KREBEC, *Weighted Inequalities in Lorentz and Orlicz spaces*, World Scientific, Singapore, 1991.
- [21] S.V. KONYAGIN AND V.N. TEMLYAKOV, *A remark on greedy approximation in Banach spaces*. East J. Approx., 5, (1999), 365–379.
- [22] S. KREIN, J. PETUNIN, AND E. SEMENOV, *Interpolation of Linear Operators*, Translations Math. Monographs, vol. 55, Amer. Math. Soc., Providence, RI, 1982.
- [23] P.G. LEMARIÉ AND Y. MEYER, *Ondelettes et bases hilbertiennes*, Rev. Mat. Iberoamericana, 2, no. 1-2, (1986), 1–18.
- [24] C. MERUCI, *Applications of interpolation with a function parameter to Lorentz, Sobolev and Besov spaces*, in : M. Cwikel, J. Peetre (Eds.), *Interpolation spaces and Allied Topics in Analysis*, Lund, 1983, in: Lecture Notes in Math., vol.1070, Springer, Berlin, (1984), pp. 183–201-
- [25] Y. MEYER, *Ondelettes et opérateurs, I: Ondelettes*, Hermann, Paris, 1990. [English translation: *Wavelets and operators*, Cambridge University Press 1992.]

- [26] J. PEETRE, *New Thoughts on Besov Spaces*, Duke Univ. Math. Series, Durham, N.C., (1976).
- [27] L. PERSON, *Interpolation with a parameter function*, Math. Scand. 59, (1986), 199–222.
- [28] W. POMPE, *Unconditional biorthogonal bases in  $L^p(\mathbb{R}^d)$* , Colloq. Math., 92, (2002), 19–34.
- [29] S. ROUDENKO, *Matrix-weighted Besov Spaces*, Trans. Amer. Math. Soc., 355 (1), (2003), 273–314.
- [30] S. ROUDENKO, *The Theory of Function Spaces with Matrix Weights*, Ph. D. Thesis, Department of Mathematics, Michigan State University, (2002).
- [31] V.N. TEMLYAKOV, *The best  $m$ -term approximation and Greedy Algorithms*, Advances in Comp. Math., 8, (1998), 249–265.
- [32] P. WOJTASZCZYK, *Wavelets as unconditional bases in  $L_p(\mathbb{R})$* , J. Fourier Anal. Appl. 5, no. 1, (1999), 73–85.
- [33] P. WOJTASZCZYK, *Greediness of the Haar system in rearrangement invariant spaces*, in Approximation and Probability, Banach Center Publications, 72, T. Figiel and A. Kamont Eds., Warszawa (2006), 385–395.

MARIA DE NATIVIDADE, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN

*E-mail address:* maria.denatividade@uam.es