# GENERIC TWISTED T-ADIC EXPONENTIAL SUMS OF BINOMIALS

CHUNLEI LIU AND CHUANZE NIU

ABSTRACT. The twisted *T*-adic exponential sum associated to  $x^d + \lambda x$  is studied. If  $\lambda \neq 0$ , then an explicit arithmetic polygon is proved to be the Newton polygon of the twisted *C*-function of the T-adic exponential sum. It gives the Newton polygons of the *L*-functions of twisted *p*-power order exponential sums.

#### 1. INTRODUCTION TO TWISTED EXPONENTIAL SUMS

Let W be a Witt ring scheme of Witt vectors,  $\mathbb{F}_q$  the field of characteristic p with q elements,  $\mathbb{Z}_q = W(\mathbb{F}_q)$ , and  $\mathbb{Q}_q = \mathbb{Z}_q[\frac{1}{p}]$ .

Let  $\Delta \supseteq \{0\}$  be an integral convex polytope in  $\mathbb{R}^n$ , and I the set of vertices of  $\Delta$  different from the origin. Let

$$f(x) = \sum_{v \in \Delta} (a_v x^v, 0, 0, \cdots) \in W(\mathbb{F}_q[x_1^{\pm 1}, \cdots, x_n^{\pm 1}]) \text{ with } \prod_{v \in I} a_v \neq 0,$$

where  $x^v = x_1^{v_1} \cdots x_n^{v_n}$  if  $v = (v_1, \cdots, v_n) \in \mathbb{Z}^n$ .

Let T be a variable. Let  $\mu_{q-1}$  be the group of q-1-th roots of unity in  $\mathbb{Z}_q$ and  $\chi = \omega^{-u}$  with  $u \in \mathbb{Z}^n/(q-1)$  a fixed multiplicative character of  $(\mathbb{F}_q^{\times})^n$ into  $\mu_{q-1}$ , where  $\omega : x \to \hat{x}$  is the Teichmüller character.

## Definition 1.1. The sum

$$S_{f,u}(k,T) = \sum_{x \in \mathbb{F}_{q^k}^{\times}} \chi(\operatorname{Norm}_{\mathbb{F}_{q^k}/\mathbb{F}_q}(x))(1+T)^{Tr_{\mathbb{Q}_{q^k}}/\mathbb{Q}_p}(f(\hat{x})) \in \mathbb{Z}_q[[T]]$$

is called a twisted T-adic exponential sum. And the function

$$L_{f,u}(s,T) = \exp(\sum_{k=1}^{\infty} S_{f,u}(k,T) \frac{s^k}{k}) \in 1 + s\mathbb{Z}_q[[T]][[s]]$$

is called an L-function of a twisted T-adic exponential sum.

**Definition 1.2.** The function

$$C_{f,u}(s,T) = \exp(\sum_{k=1}^{\infty} -(q^k - 1)^{-1} S_{f,u}(k,T) \frac{s^k}{k}),$$

is called a C-function of a twisted T-adic exponential sums.

The L-function and C-function determine each other :

$$L_{f,u}(s,T) = \prod_{i=0}^{n} C_{f,u}(q^{i}s,T)^{(-1)^{n-i+1}\binom{n}{i}},$$

and

$$C_{f,u}(s,T) = \prod_{j=0}^{\infty} L_{f,u}(q^j s,T)^{(-1)^{n-1}\binom{n-j+1}{j}}.$$

By the last identity, one sees that

$$C_{f,u}(s,T) \in 1 + s\mathbb{Z}_q[[T]][[s]].$$

The *T*-adic exponential sums were first introduced by Liu-Wan [9]. We view  $L_{f,u}(s,T)$  and  $C_{f,u}(s,T)$  as power series in the single variable *s* with coefficients in the *T*-adic complete field  $\mathbb{Q}_q((T))$ . The *C*-function  $C_{f,u}(s,T)$  was shown to be *T*-adic entire in *s* by Liu-Wan [9] for u = 0 and Liu [6] for all u.

Let  $\zeta_{p^m}$  be a primitive  $p^m$ -th root of unity, and  $\pi_m = \zeta_{p^m} - 1$ . Then  $L_{f,u}(s, \pi_m)$  is the *L*-function of the *p*-power order exponential sums  $S_{f,u}(k, \pi_m)$  studied by Adolphson-Sperber [1–4] for m = 1, by Liu-Wei [8] and Liu [6] for  $m \geq 1$ .

Let  $C(\Delta)$  be the cone generated by  $\Delta$ . There is a degree function deg on  $C(\Delta)$  which is  $\mathbb{R}_{\geq 0}$ -linear and takes values 1 on each co-dimension 1 face not containing 0. For  $a \notin C(\Delta)$ , we define deg $(a) = +\infty$ . Write  $C_u(\Delta) = C(\Delta) \cap (u + (q - 1)\mathbb{Z}^n)$  and  $M_u(\Delta) = \frac{1}{q-1}C_u(\Delta)$ . Let b be the least positive integer such that  $p^b u \equiv u \pmod{q-1}$ . Order elements of  $\bigcup_{i=0}^{b-1} M_{p^i u}(\Delta)$  so that deg $(x_1) \leq \deg(x_2) \leq \cdots$ .

**Definition 1.3.** The infinite u-twisted Hodge polygon  $H_{\Delta,u}^{\infty}$  of  $\Delta$  is the convex function on  $[0, +\infty]$  with initial point 0 which is linear between consecutive integers and whose slopes are

$$\frac{\deg(x_{bi+1}) + \deg(x_{bi+2}) + \dots + \deg(x_{b(i+1)})}{b}, i = 0, 1, \dots$$

Write NP for the short of Newton polygon. Liu [7] proved the following Hodge bound for the C-function  $C_{f,u}(s,T)$ .

Theorem 1.4. We have

$$T - adic \ NP \ of \ C_{f,u}(s,T) \ge ord_p(q)(p-1)H^{\infty}_{\Delta,u}$$

In the rest of this paper,  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots) \in W(\mathbb{F}_q[x])$ where  $\lambda \in \mathbb{F}_q^{\times}$  and  $\Delta = [0, d]$ . We fix  $0 \leq u \leq q - 1$ . We shall study the twisted exponential sum of f(x).

Let  $a = \log_p q$ . Write  $u = u_0 + u_1 p + \dots + u_{a-1} p^{a-1}$  with  $0 \le u_i \le p-1$ . Then we have

$$\frac{u}{q-1} = -(u_0 + u_1 p + \cdots), \ u_i = u_{b+i}.$$

Write  $p^{i}u = q_{i}(q-1) + s_{i}$  for  $i \in \mathbb{N}$  with  $0 \leq s_{i} < q-1$ , then  $s_{a-l} = u_{l} + u_{l+1}p + \cdots + u_{a+l-1}p^{a-1}$  for  $0 \leq l \leq a-1$  and  $s_{i} = s_{b+i}$ .

**Lemma 1.5.** The infinite u-twisted Hodge polygon  $H_{[0,d],u}^{\infty}$  is the convex function on  $[0, +\infty]$  with initial point 0 which is linear between consecutive integers and whose slopes are

$$\frac{u_0 + u_1 + \dots + u_{b-1}}{bd(p-1)} + \frac{l}{d}, \ l = 0, 1, \dots$$

*Proof.* Observe that  $\frac{s_i}{q-1} + l$  with  $0 \le i \le b-1$  is just a permutation of  $x_{bl+1}, x_{bl+2}, \cdots, x_{b(l+1)}$ . The lemma follows.

**Definition 1.6.** The arithmetic polygon  $P_{\{1,d\},u}$  is the convex function on  $\mathbb{N}$  which is linear between consecutive integers with initial point 0 and whose slopes are

$$\omega(n) = \frac{(p-1)n + \frac{1}{b}\sum_{i=0}^{b-1} u_i}{d} + (d-1)\frac{1}{b}\sum_{i=0}^{b-1} (\{\frac{pn+u_i}{d}\} - \{\frac{n}{d}\}),$$

where  $\{\cdot\}$  is the fractional part of a real number.

The main results of this paper are the following.

Theorem 1.7. We have

$$P_{\{1,d\},u} \ge (p-1)H^{\infty}_{[0,d],u}.$$

Moreover, they coincide at the point d.

**Theorem 1.8.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1.$ Then

$$T - adic NP \text{ of } C_{f,u}(s,T) = ord_p(q)P_{\{1,d\},u}$$

**Theorem 1.9.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1$ , and  $m \ge 1$ . Then

$$\pi_m - adic \ NP \ of \ C_{f,u}(s,\pi_m) = ord_p(q)P_{\{1,d\},u}$$

**Theorem 1.10.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1,$ and  $m \ge 1$ . Then

$$\pi_m - adic \ NP \ of \ L_{f,u}(s,\pi_m) = ord_p(q)P_{\{1,d\},u} \ on \ [0,p^{m-1}d].$$

The q-adic Newton polygon of  $L_{f,0}(s, \pi_m)$  with m = 1 was considered by Roger Yang [11] and Zhu [15]. In the first paper, the author got all slopes of the Newton polygon for  $p \equiv -1 \pmod{d}$  and the first slope for all p. In Zhu's paper, she pointed that if  $\lambda \neq 0$ , the Newton polygon are all generic, and the Newton polygon goes to Hodge as p goes to infinity, hence proved Wan's conjecture [14] for this certain case. S. Sperber [13] also studied the Newton polygon of  $L_{f,0}(s, \pi_m)$  when f is of degree 3 and m = 1.

## 2. Twisted T-Adic Dwork's trace formula

In this section we review the twisted T-adic analogy of Dwork's theory on exponential sums.

Let  $E(x) = \exp(\sum_{i=0}^{\infty} \frac{x^{p^i}}{p^i}) = \sum_{n=0}^{\infty} \lambda_n x^n$  be the Artin-Hasse series. Define a new T-adic uniformizer  $\pi$  of the *T*-adic local ring  $\mathbb{Q}_p[[T]]$  by the formula  $E(\pi) = 1 + T.$ 

Recall  $C_u = C_u([0,d]) = \{v \in \mathbb{N} | v \equiv u \pmod{q-1}\}$ . Write

$$L_u = \{\sum_{v \in C_u} b_v \pi^{\frac{v}{d(q-1)}} x^{\frac{v}{q-1}} : b_v \in \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]] \}$$

and

$$B_u = \{\sum_{v \in C_u} b_v \pi^{\frac{v}{d(q-1)}} x^{\frac{v}{q-1}} : b_v \in \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]]; \operatorname{ord}_{\pi} b_v \to \infty, v \to \infty\}.$$

Define  $\psi_p: L_u \to L_{p^{-1}u}$  by  $\psi_p(\sum_{v \in C_u} b_v x^v) = \sum_{v \in C_{p^{-1}u}} b_{pv} x^v$ . Then the map

 $\psi_p \circ E_f$  sends  $L_u$  to  $B_{p^{-1}u}$ , where  $E_f(x) = E(\pi x^d)E(\pi \lambda x) = \sum_{n=0}^{\infty} \gamma_n x^n$ .

The Galois group  $\operatorname{Gal}(\mathbb{Q}_q/\mathbb{Q}_p)$  can act on  $B_u$  by fixing  $\pi^{\frac{1}{d(q-1)}}$  and x. Let  $\sigma \in \operatorname{Gal}(\mathbb{Q}_q/\mathbb{Q}_p)$  be the Frobenius element such that  $\sigma(\zeta) = \zeta^p$  if  $\zeta$  is a (q-1)-th root of unity. The operator  $\Psi = \sigma^{-1} \circ \psi_p \circ E_f(x)$  sends  $B_u$ to  $B_{p^{-1}u}$ , hence  $\Psi$  operators on  $B = \bigoplus_{i=0}^{b-1} B_{p^i u}$ . We call it Dwork's *T*-adic

semi-linear operator because it is semi-linear over  $\mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]]$ . We have  $\Psi^a = \psi^a_p \circ \prod_{i=0}^{a-1} E_f^{\sigma^i}(x^{p^i})$ . It follows that  $\Psi^a$  operates on  $B_u$ and is linear over  $\mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]]$ . Moreover, one can show that  $\Psi$  is completely continuous in sense of Serre [12], so det $(1-\Psi^a s|B_u/\mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$  and det $(1-\Psi^a s|B_u/\mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$  $\Psi s|B/\mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]])$  are well defined. Now we state the twisted *T*-adic Dwork's trace formula [6].

### Theorem 2.1. We have

$$C_{f,u}(s,T) = \det(1 - \Psi^a s | B_u / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]]).$$

## 3. The twisted Dwork semi-linear operator

In this section, we study the twisted Dwork semi-linear operator  $\Psi$ . Recall that  $\Psi = \sigma^{-1} \circ \psi_p \circ E_f$ ,  $E_f(x) = E(\pi x^d) E(\pi \lambda x) = \sum_{n=0}^{\infty} \gamma_n x^n$ , where

$$\gamma_n = \sum_{di+j=n; i,j \ge 0} \pi^{i+j} \lambda_i \lambda_j \hat{\lambda}^j.$$

We see  $B = \bigoplus_{i=1}^{b} B_{p^{i}u}$  has a basis represented by  $\{x^{\frac{s_{i}}{q-1}+j}\}_{1 \leq i \leq b, j \in \mathbb{N}}$  over  $\mathbb{Z}_{q}[[\pi^{\frac{1}{d(q-1)}}]]$ . We have

$$\Psi(x^{\frac{s_i}{q-1}+j}) = \sigma^{-1} \circ \psi_p(\sum_{l=0}^{\infty} \gamma_l x^{\frac{s_i}{q-1}+j+l}) = \sum_{l=0}^{\infty} \gamma_{pl+u_{b-i}-j}^{\sigma^{-1}} x^{l+\frac{s_{i-1}}{q-1}}.$$

For  $1 \leq k, i \leq b$  and  $l, j \in \mathbb{N}$ , define

$$\gamma_{(\frac{s_k}{q-1}+l,\frac{s_i}{q-1}+j)} = \begin{cases} \gamma_{pl+u_{b-i}-j}, & k=i-1;\\ 0, & otherwise. \end{cases}$$

Let  $\xi_1, \dots, \xi_a$  be a normal basis of  $\mathbb{Q}_q$  over  $\mathbb{Q}_p$  and write

$$\xi_v^{\sigma^{-1}} \gamma_{(\frac{s_k}{q-1}+l,\frac{s_i}{q-1}+j)}^{\sigma^{-1}} = \sum_{w=1}^a \gamma_{(w,\frac{s_k}{q-1}+l)(v,\frac{s_i}{q-1}+j)} \xi_w.$$

It is easy to see  $\gamma_{(w,\frac{s_k}{q-1}+l)(v,\frac{s_i}{q-1}+j)} = 0$  for any w and v if  $k \neq i-1$ . Define the *i*-th submatrix by

$$\Gamma^{(i)} = (\gamma_{(w, \frac{s_{i-1}}{q-1} + l)(v, \frac{s_i}{q-1} + j)})_{1 \le w, v \le a; l, j \in \mathbb{N}},$$

then the matrix of the operator  $\Psi$  on B over  $\mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]]$  with respect to the basis  $\{\xi_v x^{\frac{s_i}{q-1}+j}\}_{1 \le i \le b, 1 \le v \le a; j \in \mathbb{N}}$  is

$$\Gamma = \begin{pmatrix} 0 & \Gamma^{(1)} & 0 & \cdots & 0 \\ 0 & 0 & \Gamma^{(2)} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \Gamma^{(b-1)} \\ \Gamma^{(b)} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

Hence by a result of Li and Zhu [5], we have

$$\det(1 - \Psi s | B / \mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]]) = \det(1 - \Gamma s) = \sum_{n=0}^{\infty} (-1)^{bn} C_{bn} s^{bn},$$

with  $C_n = \sum \det(A)$  where A runs over all principle  $n \times n$  submatrix of  $\Gamma$ .

For every principle submatrix A of  $\Gamma$ , write  $A^{(i)} = A \cap \Gamma^{(i)}$  as the submatrix of  $\Gamma^{(i)}$ . For a principle  $bn \times bn$  submatrix A of  $\Gamma$ , by linear algebra, if one of  $A^{(i)}$  is not  $n \times n$  submatrix of  $\Gamma^{(i)}$  then at least one row or column of A are 0 since A is principle.

Let  $\mathcal{A}_n$  be the set of all  $bn \times bn$  principle submatrix A of  $\Gamma$  with  $A^{(i)}$  all  $n \times n$  submatrix of  $\Gamma^{(i)}$  for each  $1 \leq i \leq b$ . Then we have

$$C_{bn} = \sum_{A \in \mathcal{A}_n} \det(A) = \sum_{A \in \mathcal{A}_n} (-1)^{n^2(b-1)} \prod_{i=1}^b \det(A^{(i)}).$$

Let  $O(\pi^{\alpha})$  denotes any element of  $\pi$ -adic order  $\geq \alpha$ .

Lemma 3.1. We have

$$\gamma_n = \pi^{[\frac{n}{d}] + d\{\frac{n}{d}\}} \lambda_{[\frac{n}{d}]} \lambda_{d\{\frac{n}{d}\}} \hat{\lambda}^{d\{\frac{n}{d}\}} + O(\pi^{[\frac{n}{d}] + d\{\frac{n}{d}\} + 1}).$$

*Proof.* This follows from the fact that  $i + j \ge \lfloor \frac{n}{d} \rfloor + d \lfloor \frac{n}{d} \rfloor$  if di + j = n.  $\Box$ 

**Corollary 3.2.** For any  $1 \le i \le b$  and  $1 \le w, u \le a$ , we have

$$\gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_i}{q-1}+j)} = O(\pi^{[\frac{p_i+u_{b-i}-j}{d}]+d\{\frac{p_i+u_{b-i}-j}{d}\}}).$$

Proof. By

$$\xi_{v}^{\sigma^{-1}} \gamma_{(\frac{s_{i-1}}{q-1}+l,\frac{s_{i}}{q-1}+j)}^{\sigma^{-1}} = \xi_{v}^{\sigma^{-1}} \gamma_{pl-j+u_{b-i}}^{\sigma^{-1}} = \sum_{w=1}^{a} \gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_{i}}{q-1}+j)} \xi_{w},$$
  
Follows from last lemma.

this follows from last lemma.

**Theorem 3.3.** Let  $R_i$  be a finite subset of  $\{1, 2, \dots, a\} \times (\frac{s_i}{q-1} + \mathbb{N})$  of cardinality an for each  $1 \leq i \leq b, \tau$  a permutation of  $\bigcup_{i=1}^{b} R_i$ . If  $p > 2(d-1)^2 + 1$ , then

$$\sum_{i=1}^{b} \sum_{l \in R_{i}} \left( \left[ \frac{p\phi_{i}(l) + u_{b-i} - \phi_{i(\tau(l))}(\tau(l))}{d} \right] + d\left\{ \frac{p\phi_{i}(l) + u_{b-i} - \phi_{i(\tau(l))}(\tau(l))}{d} \right\} \right)$$
$$\geq abP_{\{1,d\},u}(n) + \sum_{i=1}^{b} \sharp\{l \in R_{i} | \phi_{i}(l) > n - 1\},$$

where i(l) = i if  $l \in R_i$  and  $\phi_i$  is the projection  $\{1, 2, \cdots, a\} \times (\frac{s_i}{q-1} + \mathbb{N}) \to \mathbb{N}$ such that

$$\phi_i((\cdot, \frac{s_i}{q-1}+l)) = l.$$

*Proof.* We have

$$\sum_{i=1}^{b} \sum_{l \in R_{i}} \left( \left[ \frac{p\phi_{i}(l) + u_{b-i} - \phi_{i(\tau(l))}(\tau(l))}{d} \right] + d\left\{ \frac{p\phi_{i}(l) + u_{b-i} - \phi_{i(\tau(l))}(\tau(l))}{d} \right\} \right)$$

$$= \sum_{i=1}^{b} \sum_{l \in R_{i}} \frac{(p-1)\phi_{i}(l) + u_{b-i}}{d} + (d-1) \sum_{i=1}^{b} \sum_{l \in R_{i}} \left( \left\{ \frac{p\phi_{i}(l) + u_{b-i}}{d} \right\} - \left\{ \frac{\phi_{i}(l)}{d} \right\} \right)$$

$$+ (d-1) \sum_{i=1}^{b} \sum_{l \in R_{i}} 1_{\left\{ \frac{p\phi_{i}(l) + u_{b-i}}{d} \right\}} < \left\{ \frac{\phi_{i(\tau(l))}(\tau(l))}{d} \right\}.$$
Observe that  $\#\{l \in R: |\phi_{i}(l) > n-1\} = \#\{l \notin R: |\phi_{i}(l) \le n-1\}$  we have

Observe that  $\sharp \{l \in R_i | \phi_i(l) > n-1\} = \sharp \{l \notin R_i | \phi_i(l) \le n-1\}$ , we have  $(n-1)\phi(l) + u_l$ 

$$\sum_{l \in R_i} \frac{(p-1)\phi_i(l) + u_{b-i}}{d}$$
$$= \left(\sum_{l \in \phi_i^{-1}(A_n)} + \sum_{l \in R_i, \phi_i(l) > n-1} + \sum_{\phi_i(l) \le n-1, l \notin R_i}\right) \frac{(p-1)\phi_i(l) + u_{b-i}}{d}$$

$$\geq a \sum_{l=0}^{n-1} \frac{(p-1)l + u_{b-i}}{d} + \frac{p-1}{d} \sharp \{l \in R_i | \phi_i(l) > n-1\}.$$

Similarly, dividing the sum into three disjoint parts as above, we have

$$\sum_{l \in R_i} \left\{ \frac{p\phi_i(l) + u_{b-i}}{d} \right\} - \left\{ \frac{\phi_i(l)}{d} \right\}$$
$$\geq a \sum_{l=0}^{n-1} \left\{ \left\{ \frac{pl + u_{b-i}}{d} \right\} - \left\{ \frac{l}{d} \right\} \right\} - \frac{2(d-1)^2}{d} \sharp \{ l \in R_i | \phi_i(l) > n-1 \}.$$

The theorem follows from  $p > 2(d-1)^2 + 1$ .

**Theorem 3.4.** If  $p > 2(d-1)^2 + 1$ , then we have

$$ord_{\pi}(C_{abn}) \ge abP_{\{1,d\},u}(n).$$

In particular,

$$C_{abn} = \pm \operatorname{Norm}(\prod_{i=1}^{o} \det((\gamma_{pl-j+u_{b-i}})_{l,j\in A_n})) + O(\pi^{abP_{\{1,d\},u}(n) + \frac{1}{d(q-1)}}),$$

where Norm is the norm map from  $\mathbb{Q}_q(\pi^{\frac{1}{d(q-1)}})$  to  $\mathbb{Q}_p(\pi^{\frac{1}{d(q-1)}})$ .

*Proof.* Let  $A \in \mathcal{A}_{an}$ ,  $R_i$  the set of rows of  $A^{(i)}$  as the submatrix of  $\Gamma^{(i)}$ ,  $\tau$  a permutation of  $\bigcup_i R_i$ . By the above corollary and Theorem 3.3, we have

$$\operatorname{ord}_{\pi}(C_{abn}) \ge \sum_{i=1}^{b} \sum_{l \in R_{i}} \lceil \frac{p\phi_{i}(l) + u_{b-i} - \phi_{i(\tau(l))}(\tau(l))}{d} \rceil \ge abP_{\{1,d\},u}(n).$$

Moreover the strict inequality holds if there exist  $1 \leq i \leq b$  such that  $R_i \neq \phi_i^{-1}(A_n)$ , hence

$$C_{abn} = \prod_{i=1}^{b} \det(\gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_{i}}{q-1}+j)})_{1 \le w, v \le a; l, j \in A_n} + O(\pi^{abP_{\{1,d\},u}(n) + \frac{1}{d(q-1)}}).$$

Therefore the theorem follows from the following.

**Lemma 3.5.** For finite subset  $A \subset \mathbb{N}$ , we have

$$\det((\gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_{i}}{q-1}+j)})_{1 \le w, v \le a; l, j \in A}) = \pm \operatorname{Norm}(\det(\gamma_{(\frac{s_{i-1}}{q-1}+l,\frac{s_{i}}{q-1}+j)})_{l, j \in A})),$$
  
where Norm is the norm map from  $\mathbb{Q}_{q}(\pi^{\frac{1}{d(q-1)}})$  to  $\mathbb{Q}_{p}(\pi^{\frac{1}{d(q-1)}}).$ 

Proof. Let  $V = \bigoplus_{j \in A} \mathbb{Q}_q(\pi^{\frac{1}{d(q-1)}})e_j$  be a vector space over  $\mathbb{Q}_q(\pi^{\frac{1}{d(q-1)}})$ , and let F be the linear operator on V whose matrix with respect to the basis  $\{e_j\}$  is  $(\gamma_{(\frac{s_{i-1}}{q-1}+l,\frac{s_i}{q-1}+j)})_{l,j \in A}$ , and let  $\sigma$  act on V coordinate-wise. It is easy to see that  $(\gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_i}{q-1}+j)})_{1 \leq w,v \leq a; l,j \in A}$  is the matrix of  $\sigma^{-1} \circ F$  over  $\mathbb{Q}_p(\pi^{\frac{1}{d(q-1)}})$  with respect to the basis  $\{\xi_u e_j\}_{1 \leq u \leq a; l \in A}$ . Therefore  $\det((\gamma_{(w,\frac{s_{i-1}}{q-1}+l)(v,\frac{s_i}{q-1}+j)})_{1 \leq w,v \leq a; l,j \in A}) = \det(\sigma^{-1} \circ F)$ 

$$= \pm \det(F \mid \mathbb{Q}_p(\pi^{\frac{1}{d(q-1)}})) = \pm \operatorname{Norm}(\det(F)).$$

The lemma is proved.

# 4. HASSE POLYNOMIAL

In this section,  $1 \le n \le d-1$ . We shall study  $\det((\gamma_{pl-j+u_{b-i}})_{0\le l,j\le n-1}))$ . Definition 4.1. For any  $1 \le n \le d-1$  and  $1 \le i \le b$ , we define

$$S_{n,i} = \{\tau \in S_n | \tau(l) \le d\{\frac{pl + u_{b-i}}{d}\}\},$$

$$H_{n,u}^{(i)}(y) = \sum_{\tau \in S_{n,i}} sgn(\tau) \prod_{l=0}^{n-1} \lambda_{\lfloor \frac{pl + u_{b-i} - \tau(l)}{d} \rfloor} \lambda_{d\{\frac{pl + u_{b-i} - \tau(l)}{d}\}} y^{d\{\frac{pl + u_{b-i} - \tau(l)}{d}\}}.$$

The u-twisted Hasse polynomial  $H_{n,u}(y)$  of  $\{1,d\}$  at n is defined by

$$H_{n,u}(y) = \prod_{i=1}^{b} H_{n,u}^{(i)}(y) \pmod{p}.$$

The u-twisted Hasse polynomial  $H_u(y)$  of  $\{1, d\}$  is defined by  $H_u = \prod_{n=1}^{d-1} H_{n,u}$ .

Lemma 4.2. We have

$$H_{n,u}^{(i)}(y) = y^{\sum_{l=0}^{n-1} (d\{\frac{pl+u_{b-i}}{d}\}-l)} \sum_{\tau \in S_{n,i}} sgn(\tau) \prod_{l=0}^{n-1} \lambda_{\lfloor \frac{pl+u_{b-i}}{d}\rfloor} \lambda_{d\{\frac{pl+u_{b-i}}{d}\}-\tau(l)}.$$

*Proof.* The lemma follows from

$$\sum_{l=0}^{n-1} d\{\frac{pl+u_{b-i}-\tau(l)}{d}\} = \sum_{l=0}^{n-1} (d\{\frac{pl+u_{b-i}}{d}\}-l),$$

for any  $\tau \in S_{n,i}$ .

**Theorem 4.3.** For  $1 \le n \le d-1$  and  $p > 2(d-1)^2 + 1$ , we have

$$C_{abn} = \pm Norm(H_{n,u}(\hat{\lambda}))\pi^{abP_{\{1,d\},u}(n)} + O(\pi^{abP_{\{1,d\},u}(n) + \frac{1}{d(q-1)}}).$$

*Proof.* We have

$$\det(\gamma_{pl-j+u_{b-i}})_{0 \le l,j \le n-1} = \sum_{\tau \in S_n} \operatorname{sgn}(\tau) \prod_{l=0}^{n-1} \gamma_{pl-\tau(l)+u_{b-i}}$$

For any  $\tau \in S_n$ , we have

$$\sum_{l=0}^{n-1} \operatorname{ord}_{\pi} \gamma_{pl+u_{b-i}-\tau(l)} \ge \sum_{l=0}^{n-1} \frac{pl+u_{b-i}-\tau(l)}{d} + (d-1)\{\frac{pl+u_{b-i}-\tau(l)}{d}\}$$
$$\ge \sum_{l=0}^{n-1} \frac{(p-1)l+u_{b-i}}{d} + (d-1)\sum_{l=0}^{n-1} \{\frac{pl+u_{b-i}}{d}\} - \{\frac{l}{d}\} + 1_{\tau(l) > d\{\frac{pl+u_{b-i}}{d}\}})$$

$$\geq \sum_{l=0}^{n-1} \frac{(p-1)l + u_{b-i}}{d} + (d-1)\sum_{l=0}^{n-1} \left\{ \frac{pl + u_{b-i}}{d} \right\} - \left\{ \frac{l}{d} \right\} \right),$$

with the equalities holding if and only if  $\tau \in S_{n,i}$ . The lemma follows.  $\Box$ 

**Theorem 4.4.** The Hasse polynomial  $H_u$  is nonzero.

*Proof.* By Lemma 4.2, we see  $H_{n,u}^{(i)}$  is a monomial, so is  $H_u$ . It suffices to show for any  $1 \le i \le b$  and  $1 \le n \le d$ , the coefficients of  $H_{n,u}^{(i)}$ 

$$f_{n,p}^{(i)} = \sum_{\tau \in S_{n,i}} \operatorname{sgn}(\tau) \prod_{l=0}^{n-1} \lambda_{\lfloor \frac{pl+u_{b-i}}{d} \rfloor} \lambda_{d\{\frac{pl+u_{b-i}}{d}\} - \tau(l)} \in \mathbb{Z}_p^{\times}$$

Write  $u_{n,i} = \prod_{l=0}^{n-1} \left[\frac{pl+u_{b-i}}{d}\right]! \left(d\left\{\frac{pl+u_{b-i}}{d}\right\}\right)! \in \mathbb{Z}_p^{\times} \text{ and } \alpha_l = d\left\{\frac{pl+u_{b-i}}{d}\right\}$ , it suffices to show  $u_{n,i} f_{n,p}^{(i)} \in \mathbb{Z}_p^{\times}$ .

$$u_{n,i}f_{n,p}^{(i)} = \sum_{\tau \in S_{n,i}} \operatorname{sgn}(\tau) \prod_{l=0}^{n-1} \alpha_l(\alpha_l - 1) \cdots (\alpha_l - \tau(l) + 1)$$
$$= \sum_{\tau \in S_n} \operatorname{sgn}(\tau) \prod_{l=0}^{n-1} \alpha_l(\alpha_l - 1) \cdots (\alpha_l - \tau(l) + 1) = \det M,$$

where

$$M = \begin{pmatrix} 1 & \alpha_0 & \alpha_0(\alpha_0 - 1) & \cdots & \alpha_0 \cdots (\alpha_0 - n + 2) \\ 1 & \alpha_1 & \alpha_1(\alpha_1 - 1) & \cdots & \alpha_1 \cdots (\alpha_1 - n + 2) \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & \alpha_{n-1} & \alpha_{n-1}(\alpha_{n-1} - 1) & \cdots & \alpha_{n-1} \cdots (\alpha_{n-1} - n + 2) \end{pmatrix}.$$

It is easy to see that M is equivalent to a Vandermonde matrix. So

$$u_{n,i}f_{n,p}^{(i)} = \det M = \prod_{0 \le l < j \le n-1} (\alpha_l - \alpha_j) \in \mathbb{Z}_p^{\times},$$

the theorem is proved.

**Theorem 4.5.** For  $1 \le n \le d-1$  and  $p > 2(d-1)^2 + 1$ , we have

$$prd_{\pi}(C_{abn}) = abP_{\{1,d\},u}(n).$$

*Proof.* The theorem follows from the last two theorems and Lemma 4.2.  $\Box$ 

### 5. Proof of the main theorems

In this section we prove the main theorems of this paper. Firstly we prove Theorem 1.7, which says that

$$P_{\{1,d\},u} \ge (p-1)H^{\infty}_{[0,d],u}$$

with equality holding at the point d.

*Proof of Theorem 1.7.* It need only to show this for  $n \leq d$ . The inequality follows from the following

$$\sum_{l=0}^{n-1} \{ \frac{pl+u_i}{d} \} \ge \sum_{l=0}^{n-1} \{ \frac{l}{d} \}.$$

Moreover, the equality above holds if n = d, the theorem follows.

**Lemma 5.1.** The *T*-adic Newton polygon of det $(1 - \Psi^a s^a | B / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$ coincides with that of det $(1 - \Psi s | B / \mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]])$ 

*Proof.* The lemma follows from the following:

$$\prod_{\zeta^a=1} \det(1 - \Psi \zeta s | B / \mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]]) = \det(1 - \Psi^a s^a | B / \mathbb{Z}_p[[\pi^{\frac{1}{d(q-1)}}]])$$
$$= \operatorname{Norm}(\det(1 - \Psi^a s^a | B / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])),$$

where Norm is the norm map from  $\mathbb{Q}_q[[\pi^{\frac{1}{d(q-1)}}]]$  to  $\mathbb{Q}_p[[\pi^{\frac{1}{d(q-1)}}]]$ .

**Lemma 5.2.** The *T*-adic Newton polygon of  $C_{f,u}(s,T)^b$  coincides with that of det $(1 - \Psi^a s | B / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$ .

*Proof.* Let  $\operatorname{Gal}(\mathbb{Q}_q/\mathbb{Q}_p)$  act on  $\mathbb{Z}_q[[T]][[s]]$  by fixing s and T, then we have  $C_{pu,f}(s,T) = C_{f,u}(s,T)^{\sigma}.$ 

Therefore the lemma follows from the following

$$\prod_{i=0}^{b-1} C_{f,u}(s,T)^{\sigma^{i}} = \prod_{i=0}^{b-1} C_{p^{i}u,f}(s,T)$$
$$= \prod_{i=0}^{b-1} \det(1 - \Psi^{a}s|B_{p^{i}u}/\mathbb{Z}_{q}[[\pi^{\frac{1}{d(q-1)}}]]) = \det(1 - \Psi^{a}s|B/\mathbb{Z}_{q}[[\pi^{\frac{1}{d(q-1)}}]]).$$

**Theorem 5.3.** The T-adic Newton polygon of  $C_{f,u}(s,T)$  is the lower convex closure of the points

$$(n, \frac{1}{b} ord_T(C_{abn})), n = 0, 1, \cdots$$

*Proof.* By Lemma 5.1, we see the T-adic Newton polygon of the power series  $det(1 - \Psi^a s^a | B / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$  is the lower convex closure of the points

$$(n, \operatorname{ord}_T C_n), \ n = 0, 1, \cdots$$

It is clear that  $(n, \operatorname{ord}_T(c_n))$  is not a vertex of that polygon if  $a \nmid n$ . So that Newton polygon is the lower convex closure of the points

$$(an, \operatorname{ord}_T(c_{an})), n = 0, 1, \cdots$$

Hence the T-adic Newton polygon of det $(1 - \Psi^a s | B / \mathbb{Z}_q[[\pi^{\frac{1}{d(q-1)}}]])$  is the convex closure of the points

$$(n, \text{ord}_T C_{an}), \ n = 0, 1, \cdots$$

By Lemma 5.2, the T-adic Newton polygon of  $C_{f,u}(s,T)^b$  is the lower convex closure of the points

$$(n, \operatorname{ord}_T C_{an}), n = 0, 1, \cdots,$$

hence the closure of the points

$$(bn, \operatorname{ord}_T C_{abn}), n = 0, 1, \cdots$$

It follows that the *T*-adic Newton polygon of  $C_{f,u}(s,T)$  is the convex closure of the points

$$(n, \frac{1}{b} \operatorname{ord}_T(C_{abn})), \ n = 0, 1, \cdots$$

The theorem is proved.

**Lemma 5.4.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1.$ Then

$$T - adic NP \text{ of } C_{f,u}(s,T) \geq ord_p(q)P_{\{1,d\},u}$$

*Proof.* By the last theorem, we see that the *T*-adic Newton polygon of  $C_{f,u}(s,T)$  is the convex closure of the points

$$(n, \frac{1}{b} \operatorname{ord}_T(C_{abn})), \ n = 0, 1, \cdots$$

The theorem now follows from Theorem 3.4.

**Theorem 5.5.** Let A(s,T) be a T-adic entrie series in s with unitary constant term. If  $0 \neq |t|_p < 1$ , then

$$t - adic \ NP \ of \ A(s, t) \ge T - adic \ NP \ of \ A(s, T),$$

where NP is the short for Newton polygon. Moreover, the equality holds for one t if and only if it holds for all t.

*Proof.* The reader may refer [10] and we omit the proof here.

**Theorem 5.6.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1$ . If the equality

$$\pi_m - adic \ NP \ of \ C_{f,u}(s,\pi_m) = \operatorname{ord}_p(q) P_{\{1,d\},u}$$

holds for one  $m \ge 1$ , then it holds for all  $m \ge 1$ , and we have

$$T - adic \ NP \ of \ C_{f,u}(s,T) = \operatorname{ord}_p(q) P_{\{1,d\},u}.$$

Proof. It follows from Lemma 5.4 and the last theorem.

**Theorem 5.7.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1$ . Then

$$\pi_m - adic \ NP \ of \ C_{f,u}(s,\pi_m) = \operatorname{ord}_p(q) P_{\{1,d\},u}$$

if and only if

$$\pi_m - adic \ NP \ of \ L_{f,u}(s,\pi_m) = \operatorname{ord}_p(q) P_{\{1,d\},u} \ on \ [0, p^{m-1}d].$$

*Proof.* Assume that  $L_{f,u}(s, \pi_m) = \prod_{i=1}^{p^{m-1}d} (1 - \beta_i s)$ . Then

$$C_{f,u}(s,\pi_m) = \prod_{j=0}^{\infty} L_{f,u}(q^j s,\pi_m) = \prod_{j=0}^{\infty} \prod_{i=1}^{p^{m-1}d} (1-\beta_i q^j s).$$

Therefore the slopes of the q-adic Newton polygon of  $C_{f,u}(s, \pi_m)$  are the numbers

$$j + \operatorname{ord}_q(\beta_i), \ 1 \le i \le p^{m-1} \operatorname{Vol}(\Delta), j = 0, 1, \cdots$$

Since

$$w(n+m^{m-1}d) = p^m - p^{m-1} + w(n),$$

then the slopes of  $P_{\{1,d\},u}$  are the numbers

$$j(p^m - p^{m-1}) + w(i), \ 1 \le i \le p^{m-1}d, j = 0, 1, \cdots$$

It follows that

$$\pi_m$$
 - adic NP of  $C_{f,u}(s,\pi_m) = \operatorname{ord}_p(q) P_{\{1,d\},u}$ 

if and only if

$$\pi_m - \text{adic NP of } L_{f,u}(s, \pi_m) = \operatorname{ord}_p(q) P_{\{1,d\},u} \text{ on } [0, p^{m-1}d].$$

We now prove Theorems 1.8, 1.9 and 1.10. By the above theorems, it suffices to prove the following.

**Theorem 5.8.** Let  $f(x) = (x^d, 0, 0, \dots) + (\lambda x, 0, 0, \dots), \ p > 2(d-1)^2 + 1$ . Then

$$\pi_1 - adic \ NP \ of \ L_{f,u}(s, \pi_1) = \operatorname{ord}_p(q) P_{\{1,d\},u} \ on \ [0,d]$$
  
if and only if  $H((a_i)_{0 \le i \le d}) \ne 0$ .

*Proof.* By a result of Liu [7], the q-adic Newton polygon of  $L_{f,u}(s, \pi_1)$  coincides with  $H^{\infty}_{[0,d],u}$  at the point d. By Theorem 1.7,  $P_{\{1,d\},u}$  coincide with  $(p-1)H^{\infty}_{[0,d],u}$  at the point d, It follows that the  $\pi_1$ -adic Newton polygon of  $L_{f,u}(s, \pi_1)$  coincides with  $\operatorname{ord}_p(q)p_{\triangle}$  at the point d. Therefore it suffices to show that

$$\pi_1$$
 - adic NP of  $L_{f,u}(s,\pi_1) = \operatorname{ord}_p(q) P_{\{1,d\},u}$  on  $[0,d-1]$ 

if and only if  $H((a_u)_{u\in\triangle}) \neq 0$ .

From the identity

$$C_{f,u}(s,\pi_1) = \prod_{j=0}^{\infty} L_{f,u}(q^j s,\pi_1),$$

and the fact the q-adic orders of the reciprocal roots of  $L_{f,u}(s,\pi_1)$  are no greater than 1, we infer that

 $\pi_1$  - adic NP of  $L_{f,u}(s,\pi_1) = \pi_1$  - adic NP of  $C_{f,u}(s,\pi_1)$  on [0,d-1].

Therefore it suffices to show that

$$\pi_1$$
 - adic NP of  $C_{f,u}(s, \pi_1) = \operatorname{ord}_p(q) P_{\{1,d\},u}$  on  $[0, d-1]$ 

if and only if  $H((a_i)_{0 \le i \le d}) \ne 0$ . The theorem now follows from the *T*-adic Dwork trace formula and Theorems 4.5.

#### References

- A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: cohomology and estimates, Ann. Math., 130 (1989), 367-406.
- [2] A. Adolphson and S. Sperber, Newton polyhedra and the degree of the L-function associated to an exponential sum, Invent. Math. 88(1987), 555-569.
- [3] A. Adolphson and S. Sperber, On twisted exponential sums, Math. Ann., 290 (1991), 713-726.
- [4] A. Adolphson and S. Sperber, Twisted exponential sums and Newton polyhedra, J. reine angew. Math., 443 (1993), 151-177.
- [5] Hanfeng Li and Hui June Zhu, Zeta functions of totally ramified p-covers of the projective line, Rend. Sem. Mat. Univ. Padova, Vol. 113(2005), 203-225.
- [6] C.Liu, The L-functions of twisted Witt coverings, J. Number Theory, 125(2007), 267-284.
- [7] C.Liu, T-adic exponential sums under base change, arXiv:0909.1111.
- [8] C. Liu and D. Wei, The L-functions of Witt coverings, Math. Z., 255 (2007), 95-115.
- C. Liu and D. Wan, T-adic exponential sums, Algebra & Number Theory, Vol. 3, No. 5 (2009), 489-509.
- [10] C. Liu, Wenxin Liu and Chuanze Niu, T-adic exponential sums in one variable, arXiv:0901.0354v5.
- [11] Roger Yang, Newton polygons of L-functions of polynomials of the form  $x^d + \lambda x$ , Finite Fields Appl. 9(2003), no. 1, 59-88.
- [12] J-P. Serre, Endomorphismes complétement continus des espaces de Banach p-adiques, Publ. Math., IHES., 12(1962), 69-85.
- [13] S. Sperber, On the p-adic theory of exponential sums, Amer. J. Math. 108 (1986) 255C296.
- [14] D. Wan, Variation of p-adic Newton polygons for L-functions of exponential sums, Asian J. Math., Vol 8, 3(2004), 427-474.
- [15] J. H. Zhu, p-adic variation of L functions of one variable exponential sums, I. Amer. J. Math., 125 (2003), 669-690.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI 200240, P.R. CHINA, E-MAIL: CLLIU@SJTU.EDU.CN

School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P.R. China, E-Mail: Niuchuanze@mail.bnu.edu.cn