Middle-Four Maps and Net Categories

BRIAN DAY *

April 2, 2009

Abstract

We briefly relate the existence of a middle-four interchange map in a category with two monoidal structures, to the standard Cockett and Seely notion of a weakly distributive category.

1 Introduction

From a middle-four map we shall obtain an example of a structure (on the category of I_{\oplus} -bimodules) with two weak distributivity maps:

$$d^{t}: A \otimes (B \oplus C) \longrightarrow (A \otimes B) \oplus C$$
$$d^{r}: (B \oplus C) \otimes A \longrightarrow B \oplus (C \otimes A)$$

satisfying the eleven (planar) conditions (2)-(4) and (7)-(14) of [1], but not necessarily [1] (1), (5), and (6). We call such a structure a "net" category (see [2]) with the view that it is a proof-net category ([2]) without negation, or, equivalently, a (planar) weakly distributive category [1] without the unit objects.

Moreover, we can talk about a middle-four category with negation, which satisfies four more suitable axioms (c.f. [1] §4 (15)-(18)), so that, on passing to the category of I_{\oplus} -bimodules, we get a proof-net category with a middle-four map.

2 Middle-Four Categories

First, we consider a category V with two monoidal structures $(\otimes, I_{\otimes}, a, l, r)$ and (\oplus, I_{\oplus}) , where we have assumed that the second structure is strictly monoidal. Suppose this category is equipped with a natural middle-four transformation:

 $m: (A \oplus B) \otimes (C \oplus D) \longrightarrow (A \otimes C) \oplus (B \otimes D)$

^{*}Department of Mathematics, Macquarie University, NSW 2109, Australia.

and maps:

$$\mu: I_{\oplus} \otimes I_{\oplus} \longrightarrow I_{\oplus}$$
$$\eta: I_{\otimes} \longrightarrow I_{\oplus}$$

satisfying the following four axioms:

$\mathbf{M1}$

 $(a \oplus a)m(m \otimes 1) = m(1 \otimes m)a : ((U \oplus V) \otimes (W \oplus X)) \otimes (Y \oplus Z) \longrightarrow (U \otimes (W \otimes Y)) \oplus (V \otimes (X \otimes Z))$

M2

$(m\oplus 1)m = (1\oplus m)m$

from

 to

$(U \otimes X) \oplus (V \otimes Y) \oplus (W \oplus Z)$

 $(U \oplus V \oplus W) \otimes (X \oplus Y \oplus Z)$

and, if we put $I=I_{\otimes}$ and $R=I_{\oplus},$ we want (R,μ,η) to be a $\otimes\text{-monoid in }V$ such that both

M3

 $(1 \oplus (\mu \otimes 1))m(m \otimes 1) = m$

 $((A \oplus R) \otimes (B \oplus R)) \otimes (C \oplus D)$

from

 to

 $((A \otimes B) \otimes C) \oplus (R \otimes D)$

M4

 $((1\otimes\mu)\oplus1)m(1\otimes m)=m$

from

$$(B\oplus C)\otimes ((R\oplus A)\otimes (R\oplus B))$$


 to

$$(B\otimes R)\oplus (C\otimes (A\otimes D))$$

Moreover, a middle-four category with negation has maps γ and $\tau,$ where

$$\gamma: A \otimes A^{\perp} \longrightarrow R$$
$$\tau: I \longrightarrow A^{\perp} \oplus A$$

satisfying four axioms similar to:



commutes (cf. [1] §4).

3 The Category of *R*-bimodules as a net category

The category V(R) of R-bimodules in V has two induced (coherent) tensor products. These are:

(1) $A \otimes B$ with action:

$$\begin{array}{ccc} R \otimes (A \otimes B) \otimes R \xrightarrow{\sim} (R \otimes A) \otimes (B \otimes R) \\ & \xrightarrow{\sim} (R \otimes A \otimes I) \otimes (I \otimes B \otimes R) \\ & \xrightarrow{(1 \otimes \eta) \otimes (\eta \otimes 1)} (R \otimes A \otimes R) \otimes (R \otimes B \otimes R) \\ & \xrightarrow{\alpha \otimes \beta} A \otimes B \end{array}$$

which is unitless, and

(2) $A \oplus B$ with action

$$R\otimes (A\oplus B)\otimes R \xrightarrow{\overset{\mathbf{x}_{m^{\prime\prime}}}{\longrightarrow}} (R\otimes A\otimes R)\oplus (R\otimes B\otimes R) \xrightarrow{\alpha\oplus\beta} A\oplus B$$

which has the unit R.


Furthermore, it is easily checked that our basic transformation

$$m: (A \oplus B) \otimes (C \oplus D) \longrightarrow (A \otimes C) \oplus (B \otimes D)$$


becomes an R-bimodule homomorphism. Thus, on putting B = R and C = R respectively, and using the bimodule actions, we get two natural transformations:

$$d^{l}: A \otimes (B \oplus C) \longrightarrow (A \otimes B) \oplus C$$
$$d^{r}: (B \oplus C) \otimes A \longrightarrow B \oplus (C \otimes A)$$

of *R*-bimodules. We now check some [1] axioms on d^l and d^r . Axioms [1] (7) and (8) are trivial. Axiom [1] (9):



where $\alpha(1 \otimes \alpha)a = \alpha(\mu \otimes 1)$ sine *D* is an *R*-bimodule. Axiom [1] (10) is similar, but uses M1 and M4. Axiom [1] (11) uses M2 and the action (*) of *R* on $A \oplus B$.



Axiom [1] (12) is similar and still uses M2, while Axiom [1](13) uses M2, and [1] (14) uses M1.

Thus, from any category with two monoidal structures related by a middlefour interchange transformation satisfying the standard axioms for such a map, we obtain an example of a net-category (that is, a proof-net category without negation) which is also equipped with a middle-four interchange map.

Communication on this article can be made through Micah McCurdy (Macquarie University, micah.mccurdy@gmail.com) who kindly typed this manuscript. I would also like to thank Ross Street for a helpful comment on the middle-four map.

References

- [1] Cockett, J.R.B. and R.A.G. Seely, *Weakly Distributive Categories*, Journal of Pure and Applied Algebra v114 n2 (1997).
- [2] Dosen, K., and Z. Petric, "Proof-net categories", (Preprint) Math. Inst. Serbian Academy of Sciences and Arts (2005), arXiv:math/0503301v3