

THE PROGENITORS OF TYPE IA SUPERNOVAE:

I. Are they Supersoft Sources?

R. DiStefano

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

ABSTRACT

In a canonical model, the progenitors of Type Ia supernovae (SNe Ia) are accreting, nuclear-burning white dwarfs (NBWDs), which explode when the white dwarf (WD) reaches the Chandrasekhar mass, M_C . Such massive NBWDs are hot ($kT \sim 100$ eV), luminous ($L \sim 10^{38}$ erg s $^{-1}$), and are potentially observable as luminous supersoft X-ray sources (SSSs). During the past several years, surveys for soft X-ray sources in external galaxies have been conducted. This paper shows that the results falsify the hypothesis that a large fraction of progenitors are NBWDs which are presently observable as SSSs. The data also place limits on sub- M_C models. While $\hat{1}$ progenitors may pass through one or more phases of SSS activity, these phases are far shorter than the time needed to accrete most of the matter that brings them close to M_C .

Subject headings: binaries:close; ISM:planetary nebulae:general; stars:evolution; X-rays:stars; X-rays:general

1. Introduction

Type Ia supernovae (SNe Ia) help us to explore the history of the cosmos (see, e.g., Riess et al. 2007; Kuznetsova et al. 2008). Yet we still know little about their progenitors. There are two classes of progenitor models. (See Kotak 2009 and Branch et al. 1995 for reviews.) In “single degenerate” models, the progenitors are accreting white dwarfs (WDs). In “double-degenerate” models, the explosion is initiated through mass transfer between and/or the merger of two WDs. This paper is devoted to searching for the progenitors.

In §2 we show that, if the single-degenerate model is dominant, then galaxies such as our own and M31 contain on the order of 1000 nuclear-burning WDs (NBWDs) that will explode within 10^5 years. In §3 we show that these progenitors could be hot and bright enough to be detected as supersoft x-ray sources (SSSs) in many external galaxies. We then show that existing data from *Chandra* place strict limits on the numbers of Type Ia progenitors that

are detected as SSSs. In section §4 we discuss the implications. These are explored more fully, in the context of binary models for $\hat{1}$ progenitors, in the companion paper (DiStefano 2009).

2. The Numbers of Accreting WD Progenitors

In single-degenerate models, a WD must accrete and retain matter. Let M_0 represent the initial mass of the WD. An explosion occurs when the mass has increased to a value M_f . We start with models in which M_f is the Chandrasekhar mass, $M_C \approx 1.4 M_\odot$. The WD must have a donor that can contribute the requisite mass. In addition, the rate of mass infall, \dot{M}_{in} , must be large enough that accreted matter can be burned and retained. The rate of genuine mass increase, \dot{M}_{WD} , is determined by the rate of infall to the WD and a retention factor $\beta(M_{WD}, \dot{M}_{in})$.

$$\dot{M}_{WD} = \beta(M_{WD}, \dot{M}_{in}) \dot{M}_{in} \quad (1)$$

The top panel of Figure 1 illustrates that mass retention requires mass donation rates in a narrow range between $\dot{M}_{in,min}(M_{WD})$ (roughly $10^{-7} M_\odot \text{ yr}^{-1}$ for a solar-mass CO WD), and $\dot{M}_{in,max}(M_{WD})$, which is generally a few times larger than $\dot{M}_{in,min}(M_{WD})$. This range corresponds to rates of infall for which accreting matter can undergo quasisteady nuclear burning.¹ For $\dot{M}_{in} > \dot{M}_{in,max}$, not all of the incoming matter can be burned, and β declines. For $\dot{M}_{in} < \dot{M}_{in,min}$, incoming matter accumulates and burns episodically. For rates just below $\dot{M}_{in,min}(M_{WD})$, the explosions are frequent (separated by decades) and weak, corresponding to recurrent novae (RNe); β can be close to unity. The lower the rate of infall, the more violent are the explosions produced by nuclear burning, and the more matter is ejected from the system. In some cases, more matter may be ejected than is accumulated between novae; β is negative. Most of the accreting WDs already identified are cataclysmic variables with accretion rates typically two orders of magnitude or more below the rate required for steady burning. Binaries can produce the requisite high rates, however (van den Heuvel et al. 1992).

The duration, τ_{acc} , of the accretion phase is

$$\tau_{acc} = \int_{M_0}^{M_f} \frac{dM}{\dot{M}_{in} \beta(M_{WD}, \dot{M}_{in})} \quad (2)$$

There may be long periods in which the mass transfer period is too low (or too high) for mass retention. In this paper we are concerned primarily with the interval during which the

¹Note that even within the so-called *steady-burning region*, β need not be equal to unity. For example, some of the infalling mass may be driven from the white dwarf in the form of winds.

value of \dot{M}_{in} is large enough that infalling mass is burned and retained. The approximate duration of this interval can be written as follows.

$$\tau_{acc} = 5 \times 10^5 \text{ yrs} \left(\frac{\Delta M}{0.4 M_{\odot}} \right) \left(\frac{8 \times 10^{-7} M_{\odot} \text{ yr}^{-1}}{\beta \dot{M}_{in}} \right). \quad (3)$$

where the value of $\beta \dot{M}_{in}$ represents an average during the epoch when the WD's mass is changing.

The rate of SNe Ia in galaxies is roughly 0.3 per century per $10^{10} L_{\odot}$ in blue luminosity (Cappellaro et al. 1993; Turatto et al. 1994). In a galaxy with blue luminosity L_B , the number, N_{acc} , of accreting progenitors within $0.4 M_{\odot}$ of M_C is roughly

$$N_{acc} = 1500 \left(\frac{\Delta M}{0.4 M_{\odot}} \right) \left(\frac{8 \times 10^{-7} M_{\odot} \text{ yr}^{-1}}{\beta \dot{M}_{in}} \right) \left(\frac{L_B}{10^{10} L_{\odot}} \right). \quad (4)$$

To determine the expected range of values of N_{acc} , we assess the range of likely values of each factor. The amount of mass, ΔM , that the white dwarf must gain is constrained from below by the requirement that the accretor must start as a CO white dwarf, while white dwarfs born with the highest mass start as O-Ne-Mg white dwarfs, which undergo accretion-induced collapse (AIC) upon achieving M_C (see, e.g., Nomoto et al. 1979). The value of ΔM can therefore not be smaller than $0.15 - 0.2 M_{\odot}$. Furthermore, the numbers of high-mass white dwarfs are constrained by two considerations. First the initial mass function favors low-mass stars (Miller & Scalo 1979; Salpeter 1955). Second, low-mass stars produce low-mass white dwarfs (Kalirai et al. 2008; Catalán et al. 2009; Dobbie et al. 2006; Williams 2007; Weideman & Koester 1983). Therefore ΔM is almost certainly in the range: $0.2 - 0.8 M_{\odot}$, consistent with N_{acc} in the range $750 - 2250$. In order for the average value of ΔM to be small enough to place N_{acc} below 750, the relationship between a star and the mass and/or composition of the remnant would have to be different from current models. Alternatively, part of the answer could be that O-Ne-Mg white dwarfs produce $\hat{1}e$. Either case would represent a shift in our understanding of fundamental astrophysics.

If the average value of $\beta \dot{M}_{in}$ is larger than $8 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$, this would reduce the value of N_{acc} . Higher rates of nuclear burning will, however, produce luminosities that exceed the Eddington luminosity. It is more likely that the average rate at which mass is accreted is actually smaller, producing larger values of N_{acc} . Note in addition that calculations by different groups agree on the general range of \dot{M}_{in} needed for high-mass white dwarfs to burn accreting matter (see, e.g., Nomoto 1982; Iben 1982, Fujimoto 1982; Shen & Bildsten 2008 and references therein).

The third factor influencing the value of N_{acc} is proportional to L_B . This factor is included because observations find that the rate of Type Ia supernova scales with L_B . As

more $\hat{1}\text{e}$ are discovered in more galaxies, this rate is becoming better measured. (See, e.g., Dilday et al. 2008; Graham et al. 2008; Kuznetsova et al. 2008; Poznanski et al. 2008; Panagia et al. 2007.) It is unlikely that the estimated rates will change enough to reduce or enhance the predicted values of N_{acc} by a factor as large as 2.

Therefore, taking into account the range of possible values of the factors in Equation 4, we find that for spiral galaxies similar to the Milky Way and M31 (and even for ellipticals, with a lower rate of SNe Ia), N_{acc} cannot be much smaller than several hundred, and is likely to be significantly larger. The assumptions that lead to these predictions are that (1) the single-degenerate channel produces the majority of $\hat{1}\text{e}$, and (2) nuclear burning of accreted material is needed to prevent explosive mass loss.

3. Detecting the Progenitors

3.1. The High Luminosity of Progenitors

The crucial point is that nuclear-burning, either episodic or quasisteady, appears to be necessary for the retention of matter by WDs. The burning of matter releases a great deal of energy.

$$L_{NB} = 2.4 \times 10^{38} \text{ erg s}^{-1} \left(\frac{\eta}{0.007} \right) \left(\frac{f}{0.75} \right) \left(\frac{\beta \dot{M}_{in}}{8 \times 10^{-7} M_{\odot} \text{ yr}^{-1}} \right) \quad (5)$$

In this equation, L_{NB} is the power released through the burning of accreted material; f is the fraction of the incoming matter burned, and η is the efficiency.

Consider a WD with accretion rate in the steady-burning regime. Because the mass infall rate is large, the accretion luminosity is also large, although typically less than 10% of the power provided by nuclear burning. If, however, the accretion rate lies slightly below the steady-burning regime, then nuclear burning will occur only episodically, during recurrent novae. In contrast, the accretion luminosity, which can be $\sim 10^{37} \text{ erg s}^{-1}$ for high-mass white dwarfs, could make the system appear bright even between nova explosions. Therefore, particularly when nuclear burning is occurring, but even when mass infall rates are slightly below the nuclear-burning limit, accreting WDs that are Type Ia progenitors should be highly luminous (second panel of Figure 1). We address the issue of the detectability of the accretion luminosity in the companion paper (Di Stefano 2009).




Fig. 1.— **Properties of NBWDs:** green point correspond to systems in which the rate of mass infall is in the range necessary for quasisteady nuclear burning. In each panel, the green points lie between two curves; the upper (lower) curves correspond to the maximum (minimum) rate of infall consistent with quasisteady nuclear burning. **Top panel:** \dot{M}_{in} is plotted against the white dwarf mass, M_{WD} . For values of \dot{M}_{in} below the steady-burning region, novae are expected. The repetition frequency is highest for \dot{M}_{in} just below the steady-burning limit, so that the novae are recurrent novae and a large fraction of the accreted matter can be retained. For even lower accretion rates, the novae are classical novae, blasting accreted material from the binary at intervals $> 10^4$ years. Just above the steady burning region is the region in which mass comes in too quickly for it all to be burned upon arrival. Unless mass transfer can be turned off during an interval in which all excess matter is burned, the value of β must be smaller than unity. **Second panel:** The logarithm of the luminosity versus M_{WD} . The positions of the green points are computed by assuming that 0.75 of the infalling matter is burned with an efficiency of 0.007. The red points show the accretion luminosity, which was computed to be: $\dot{M}_{in} M_{WD} G / R_{WD}$. Although more than an order of magnitude lower than the luminosity derived from nuclear burning, accretion by high-mass white dwarfs is detectable in nearby external galaxies. **Third panel:** kT versus M_{WD} . The value of T was computed by assuming an effective radius equal to $1.5 R_{WD}$. This panel illustrates that the effective temperature is a strong function of the WD mass. For WDs with mass near M_C , kT is large enough to make x-ray detection efficiency comparable to what it would be for a canonical (i.e., hard) x-ray source.

3.2. Effective Temperatures: Detection at X-ray Wavelengths

As the third panel of Figure 1 shows, effective values of kT for NBWDs are expected to be in the range of tens of eV. The x-ray missions *Einstein* and *ROSAT* had good low-energy sensitivity and could detect soft x-ray sources in the Local Group. By the mid-1990s, more than 30 sources with the properties expected for NBWDs had been discovered. (See Greiner 2000 for a catalog and for references.) The soft emitters were dubbed luminous supersoft x-ray sources (SSSs). A natural conjecture is that the SSSs are in fact NBWDs. Some nearby SSSs are known to be hot white dwarfs post-nova, or are pre-WDs still ensconced in planetary nebulae. The binary properties of roughly half of the SSSs with optical IDs suggests that they might be WDs accreting at high-enough rates to burn matter in a quasisteady fashion. Soon after their discovery, it was conjectured that some SSSs could be progenitors of AIC (van den Heuvel et al. 1992) or of 1e (Rappaport, Di Stefano & Smith 1994).

As Figure 1 demonstrates, the temperature is higher for WDs with higher mass. This is because higher-mass WDs (1) are smaller, and (2) require a higher infall rate, and therefore a higher rate of nuclear burning. In principle, NBWDs with mass close to M_C can have values of kT near or even slightly above 100 eV.

Di Stefano & Rappaport (1994) modeled the gas distribution of several galaxies to determine what fraction of SSSs could be detected as a function of L and T . While only roughly a percent of low- L /low- T SSSs could be detected by *ROSAT*, almost all SSSs associated with NBWDs with mass near M_C could be detected in M31, because they are both hot and bright. Figure 2 demonstrates that, with *Chandra*, for typical amounts of absorption, we can expect to have a complete census of NBWDs with masses within a few tenths of M_C in M31, M101, M83, and M51. In fact, with typical values of $N_H \sim 10^{21} \text{ cm}^{-2}$, the *Chandra* exposures we analyzed should allow us to detect and identify all NBWDs with M_{WD} larger than $0.8 - 0.9 M_\odot$ ($1.2 - 1.3 M_\odot$) in M31 (M81 and M83). Elliptical galaxies are good places in which to hunt for bright SSSs because they have relatively small amounts of gas and dust.

If, therefore, (1) the class of Type Ia progenitors consists primarily of accreting white dwarfs with mass close to M_C , and (2) these white dwarfs are burning the accreted matter in a quasi-steady manner, and (3) this produces luminosities and temperatures in the expected regime, then these galaxies should each contain at least several hundred hot, bright, and detectable SSSs.

3.3. Observations

3.3.1. Individual Galaxies

Table 1 shows the numbers of soft x-ray sources in each of six galaxies that have been carefully studied (Di Stefano et al. 2003a, 2003b, 2004a, 2004b). M101, M83, and M51 are spiral galaxies. M104 is a bulge-dominated spiral, while both NGC4472 and NGC4697 are elliptical galaxies. Two categories of soft sources are listed.

(1) **SSSs** refer to sources with values of kT below 150 eV. These should encompass the majority of NBWDs, including all of the Type Ia progenitors. In each case, the numbers of SSSs actually detected are smaller than the numbers of predicted Type Ia progenitors by more than an order of magnitude. Furthermore, only a fraction of the detected SSSs are candidate progenitors of $\hat{1}e$. This is because luminosities and temperatures of many of the detected SSSs are so low that, if they correspond to NBWDs, the mass of the white dwarf is much smaller than M_C (see, e.g., Pence et al. 2001). On the other hand, some SSSs are too luminous to correspond to NBWDs. Some ultraluminous SSSs, with x-ray luminosities above 10^{39} erg s $^{-1}$, may be accreting black holes (Kong & Di Stefano 2005; Mukai et al. 2005; Kong et al. 2004).

(2) **Quasisoft sources (QSSs)** are sources that are slightly harder than SSSs, but which exhibit little or no emission above 2 keV (Di Stefano & Kong 2004; Di Stefano et al. 2004a; Di Stefano et al. 2004b). With effective values of kT above ~ 150 eV, some could be NBWDs with mass near M_C : the addition of a small hard component and/or high levels of absorption could cause such a source to have a QSS spectrum. Nevertheless, many QSSs are too hard to be associated with NBWDs. At most a small fraction could be progenitors of Type Ia SNe.

Results: At most, a modest fraction of all SSSs and QSSs could be progenitors of $\hat{1}e$. Yet, even if the total of all SSSs *and* QSSs in each galaxy were progenitors, they would

Table 1: Soft Sources in External Galaxies

Galaxy	SSSs	QSSs	Other Sources
M101	42	21	24
M83	28	26	74
M51	15	21	56
M104	5	17	100
NGC4472	5	22	184
NGC4697	4	15	72

constitute less than a tenth of the total NBWDs with mass near M_C needed to provide the observed rate of $\dot{1}\text{e}$. Therefore, even if some of the detected SSSs and QSS are progenitors, there is a clear and large mismatch, with at least 90–99% of the actively accreting progenitors not appearing as SSSs or QSSs. Figure 2 also indicates that the observations already place limits on the numbers of lower-mass NBWDs in each of several galaxies.

It may be important to note, however, that there is an interesting trend in the data. That is, both the rate of $\dot{1}\text{e}$ and the number of SSSs are significantly larger in late-type galaxies than in early-type galaxies.

If the rate of mass transfer is high enough to support quasisteady-nuclear burning, then the character of the disk is remarkable, because the total accretion energy is only a small fraction of the energy provided to the disk by the white dwarf through irradiation and heating. The inner disk itself becomes a source of supersoft x-rays. At x-ray wavelengths, the effect is therefore to make the source marginally brighter, but to likely keep it in the SSS regime. Should it b

3.3.2. *A Large Sample of Galaxies:*

An automated source detection and identification process was employed by Liu (2008) to study x-ray sources in fields containing 383 nearby galaxies. SSSs and QSSs were identified using the same algorithm used for the galaxies in Table 1. Liu found that only 2.6% of all sources bright enough for spectral classification are SSS. For every SSS there are roughly four QSSs.² The combination of SSSs and QSSs constitute about 13% of all x-ray sources. Yet, the total numbers of x-ray sources per galaxy were generally several times smaller than the expected population of nuclear-burning Type Ia progenitors. Thus, the discrepancy between the numbers of progenitors and the total numbers of soft sources per galaxy is typically larger than an order of magnitude. Furthermore, a significant fraction of the soft sources have x-ray luminosities above $10^{39} \text{ erg s}^{-1}$, and are therefore not good candidates for white dwarf accretors.

Liu created merged images to achieve effective exposure times far longer than those utilized in Figure 2. The total exposure time of M101, for example, is roughly ten times the value used to create Figure 2. This means that Liu’s catalog can be used to place stringent limits on sub-Chandrasekhar models (with white dwarf masses as low as $0.8 - 0.9 M_\odot$) as

²The high ratio of QSSs to SSSs may reflect the prevalence of older stellar populations among the galaxies in Liu’s survey.

well as on Chandrasekhar-mass models.

Note that, in addition to the observations of the previous subsection, and the work of Liu (2008), many studies of extragalactic x-ray sources have identified SSSs. See, e.g., Sarazin et al.(2001); Swartz et al.(2002); Pence et al.(2001); Jenkins et al.(2004); Fabbiano et al.(2003). We focus on the Liu (2008) paper because it is comprehensive and uses uniform detection and selection procedures. Note, however, that the results of the individual studies are consistent with those of Liu: the total numbers of bright SSSs per galaxy is small, consistent with the source counts presented in Table 1.

3.3.3. M31

M31 is a special case because of its proximity. A source with x-ray luminosity ($0.2 - 10$ keV) of $10^{38} \text{erg s}^{-1}$, possibly corresponding to a NBWD with mass within roughly $0.15 M_{\odot}$ of M_C , would produce a count rate of 0.04 s^{-1} in ACIS-S, or 0.014 s^{-1} in ACIS-I. Since most exposures of M31 are longer than 5 ksec, these sources are readily detected and their spectra can be classified as SSS. A source which exhibits a small hard component might appear as a QSS, producing an even higher count rate. DiStefano et al. (2004b) published a list of 42 SSSs and QSSs drawn from three regions in the disk and also from the galactic bulge. Of these, only 4 sources (3 of them in the bulge), had count rates above 0.01 s^{-1} . Most of the detected soft sources are too dim to be high-mass NBWDs. These observations cover only a fraction of the galaxy. (Note that most of the SSSs are in the center, which was well covered, so the total number of sources we expect does not scale with the area covered.) Orio (2006) identified SSSs and QSSs in M31 by using deep *XMM-Newton* observations that covered the entire body of the galaxy. Using x-ray spectral information alone, Orio found significant contamination from supernova remnants. The data contained evidence for 15 SSSs and 18 QSSs that were not associated with known supernova remnants. In addition, Orio found that about one in ten nova outbursts in M31 produces SSS-behavior post nova. The link to novae has been studied in more detail using additional data sets (Pietsch et al. 2006; Henze et al. 2009). della Valle & Livio (1996) have demonstrated that only $\sim 10\%$ of all novae are likely to be RNe, retaining most of the matter they accrete. Most novae, therefore, are not $\hat{1}$ progenitor candidates. In summary, the total number of SSSs detected in M31 is smaller than the requisite number of progenitors, and many of the ones that are detected are not good candidates for Type Ia progenitors.

In fact in M31, where typical values of N_H are $\sim 10^{21} \text{ cm}^{-2}$, we likely have a complete census of NBWDs with mass greater than roughly $0.9 M_{\odot}$. This places strong limits on sub-Chandrasekhar models.

3.3.4. *The Milky Way*

Only a handful of SSSs have been discovered in the Milky Way. This small number is consistent with the fact that absorption by the interstellar medium (ISM) in the Galactic disk can obscure the soft radiation emanating from l-m NBWDs. If, however, the Galaxy contains 500 – 1000 high-mass NBWDs, 5 – 10 of them should lie within roughly a kpc. If these have values of kT in the range 75 – 100 eV and luminosities of $\sim 10^{38}$ erg s $^{-1}$, they will be easily detected by *ROSAT*, *Chandra*, and *XMM-Newton*.

For *Chandra* count rates we can scale from the numbers given in the paragraph on M31; the *XMM-Newton* count rates would be ~ 4 times higher. While the *ROSAT* count rates would be smaller than the *Chandra* rates by a factor of ~ 2 , the *All-Sky Survey* could have discovered all high-mass NBWDs within a kpc, even along directions in the sky in which the exposure times were shortest. The persistent SSSs discovered by *ROSAT* were dimmer and cooler than high-mass NBWDs (Greiner 2000). The failure to detect bright, hot SSSs and/or QSSs in the neighborhood (< 1 kpc) of the Sun is an independent indication that such sources do not comprise the majority of $\hat{1}$ progenitors.

3.3.5. *Pre- and Post-explosion Comparisons*

The most direct way to determine if $\hat{1}$ e are generated by SSSs is to check pre-explosion x-ray images. Voss & Nelemans (2008) found 4 galaxies that had been observed by *Chandra* prior to a $\hat{1}$ explosion. There was no sign of an x-ray source in the vicinities of 3 of the supernovae. The detection limits were below those expected for high-mass NBWDs. In one case, there was a nearby source, but it was 1.15'' from the supernova (~ 80 pc in projection; Roelofs et al. 2008), and was neither an SSS nor a QSS.

4. Conclusions

4.1. SSSs for the Long Term?

The discovery of SSSs seemed to provide hope that Type Ia progenitors could be identified in a fairly straightforward way, through the signatures of their soft x-ray emission. While some bright SSSs may be progenitors of Type Ia supernovae, we can now say conclusively that the majority of the progenitors do not appear as bright SSSs during intervals long enough ($\sim 10^5$ yrs) to allow quasisteady burning of the necessary amounts of accreting matter. The discrepancy is at least an order of magnitude, and may be as much as two




Fig. 2.— Each panel refers to a quasi-steady NBWD with the listed mass; L and T were taken from Iben (1982). Within each panel, each curve corresponds to a single galaxy (M31, green; M101, red; M83, magenta; M51, cyan; NGC 4697, blue); the galaxy’s distance and the longest ACIS-S exposure we have analyzed were used to compute the numbers of counts. We assumed the sensitivity of *Chandra*’s ACIS-S as computed by PIMMS for AO4. The horizontal dot-dashed line shown on each panel corresponds to 14 counts, the number needed to ascertain whether a given source can be classified as an SSS (see DiStefano & Kong 2003).

orders of magnitude.

In addition, we find that existing data already place restrictions on sub-Chandrasekhar models (Figure 2). These restrictions will be tightened as more data are analyzed, and additional exposures with *Chandra* and *XMM-Newton* become available.

4.2. SSSs for the Short Term?

Although we have ruled out the possibility that the majority of $\hat{1}$ progenitors appear as SSSs during much of the all-important mass-gaining phase, there is almost certainly a tight link between SSSs and $\hat{1}$ progenitors. While there are several ways to “hide” the soft x-ray emission from NBWDs (see the companion paper, DiStefano 2009), it is difficult to construct a model in which a white dwarf burns a tenth of a solar mass or more, yet never exhibits at least brief SSS or QSS phases. For example, some progenitors may experience short periods of SSS behavior during the time when the system approaches or passes the minimum accretion rate needed for steady burning. This can happen both as mass transfer ramps up and also as it ramps down. In addition, a small fraction of binaries in which the white dwarf reaches M_C can sustain mass transfer rates in the steady burning regime for most of the mass-transfer phase. (See, e.g., van den Heuvel et al. 1992; Rappaport, Di Stefano & Smith 1994; Di Stefano & Nelson 1996; Yungelson et al. 1996; Lü et al. 2006; Han & Podsiadlowski 2006.)

It is therefore possible that a portion of the observed SSSs and QSSs are CO white dwarfs on their way to the Chandrasekhar mass. It is important to identify candidates. Given the expected temperature range, the NBWDs that are $\hat{1}$ progenitors could appear as very hot SSSs. If there is a small hard component and/or if absorption is important, they could instead appear as the coolest of QSSs. Wherever possible, these candidates should be monitored. Those that are not high-mass NBWDs will be interesting anyway, because they are unusual states of neutron-star or black-hole accretors. For example, some could be black holes of $\sim 50 - 100 M_\odot$ in thermal dominant states (Di Stefano et al. 2004a).

The candidates that turn out to be NBWDs could be near explosion. The likelihood that an individual source is about to “go off” is small. Even a white dwarf within $0.01 M_\odot$ may take more than 1000 years to achieve M_C . Nevertheless, by identifying a large pool of candidates, drawn from the hundreds of galaxies observed with *Chandra* and *XMM-Newton*, we increase the chance of having a useful pre-explosion image of a $\hat{1}$ that explodes within

the next decade.³

4.3. A Multiwavelength Guide

The restrictions on the numbers of SSSs we have derived above do not necessarily translate into restrictions on the numbers of NBWDs. In fact, theoretical considerations predict the presence of a population of high-mass NBWDs. First, starting with a large population of binaries, a subset will pass through a phase in which a high-mass white dwarf accretes matter at a high rate (see, e.g., Rappaport, Di Stefano & Smith 1994; Yungelson et al. 1996). Second, we know that $\hat{1e}$ occur, and that every evolutionary pathway to explosion passes through channels that involve high accretion rates onto a white dwarf with mass larger than $\sim 0.8 M_{\odot}$. Certainly a long phase of nuclear-burning is required in single-degenerate models (Equation 3). This would be true in sub-Chandrasekhar models as well. Furthermore, we show in the companion paper that even many γ -d binaries require a prior single-degenerate phase during which nuclear burning is expected (Di Stefano 2009).

We have also found that the numbers of NBWDs needed per galaxy is large. Thus, at any given time, galaxies such as the Milky Way house ~ 1000 NBWDs with masses near M_C . In sub-Chandrasekhar models, or γ -d models the average mass of the NBWDs may be smaller, but nevertheless larger than $0.8 M_{\odot}$. This increases our chances of establishing the existence of a population of progenitors in M31 and in other nearby galaxies. Perhaps most exciting, the large number guarantees that some progenitors that are actively accreting NBWDs must lie within a kpc of Earth. If we can identify these, we will be able to make detailed studies of the progenitors and the processes that allow high-mass white dwarfs to grow in mass.

In fact, we have almost certainly detected NBWDs that are $\hat{1}$ progenitors, in the Milky Way and in nearby galaxies. These must be bright systems. Since the bulk of the energy produced by nuclear burning does not escape at x-ray wavelengths, it must be emitted in other portions of the electromagnetic spectrum. Radiation from the white dwarf could, for example, be reprocessed by winds from the binary. In this case, the spectrum would be shifted toward longer wavelengths, and the progenitors could be highly luminous in the IR. Alternatively, if the photospheres of massive NBWDs tend to be large, compared with size of the white dwarf, the power can be emitted primarily in the UV. If progenitors pass through phases of SSS behavior, even with a low duty cycle, they may be associated with large, distinctive ionization nebulae (Rappaport et al. 1994; Chiang & Rappaport 1996;

³In the companion paper we also discuss the discovery of progenitors at other wavelengths (IR, optical, and UV) in the context of binary models.

Di Stefano, Paerels & Rappaport 1995). In the companion paper (Di Stefano 2009) we link the possible signatures of nuclear burning to the characteristics of the binary models for $\hat{1}$ progenitors.

Summary: Nuclear burning appears to be required in order for Type Ia supernovae to occur. As the companion paper shows, even many $\hat{1}$ progenitors of $\hat{1}$ e are expected to pass through an earlier single-degenerate phase in which nuclear burning occurs. We have demonstrated that the majority of the progenitors do not appear as SSSs or QSSs during most of the all-important nuclear-burning phase. The energy must escape in other wavebands, making it possible to detect and identify dozens of progenitors in our own Galaxy, and comparable or perhaps even larger numbers in M31 and other nearby galaxies. Given the long duration of the requisite nuclear-burning phases, it would be surprising if some progenitors do not appear as SSSs or QSSs, at least during part of their evolution. It is therefore still interesting to explore the connection between soft x-ray sources and $\hat{1}$ progenitors, even as we extend searches to bright IR and UV sources, and large ionization nebulae, so as to find evidence of the centrally-important nuclear-burning phase. The bottom line is that, although $\hat{1}$ e are rare and therefore tend to occur at great distance from us, thousands of the progenitors are nearby, and are luminous enough that we can study them in detail well before they explode.

Acknowledgements: It is a pleasure to acknowledge helpful conversations and comments from many of the participants (especially Ed van den Heuvel, Lev Yungelson, and Jim Liebert), of the KITP conference and workshop on *Accretion and Explosion* held at UC Santa Barbara in 2007. This work was supported in part by an LTSA grant from NASA and by funding from the Smithsonian Institution.

REFERENCES

- Branch, D., Livio, M., Yungelson, L. R., Boffi, F. R., & Baron, E. 1995, *PASP*, 107, 1019
- Cappellaro, E., Turatto, M., Benetti, S., Tsvetkov, D. Y., Bartunov, O. S., & Makarova, I. N. 1993, *A&A*, 273, 383
- Catalán, S., Isern, J., García-Berro, E., & Ribas, I. 2009, *Journal of Physics Conference Series*, 172, 012007
- Chiang, E., & Rappaport, S. 1996, *ApJ*, 469, 255
- della Valle, M., & Livio, M. 1996, *ApJ*, 473, 240

- Dilday, B., et al. 2008, ApJ, 682, 262
- Di Stefano, R., 2009, *The progenitors of Type Ia Supernovae: II. A Guide to Discovery*, in preparation.
- Di Stefano, R., Primini, F. A., Liu, J., Kong, A., & Patel, B. 2009, arXiv:0909.2046
- Di Stefano, R., Primini, F. A., Kong, A. K. H., & Russo, T. 2004a, arXiv:astro-ph/0405238
- Di Stefano, R., et al. 2004b, ApJ, 610, 247
- Di Stefano, R., & Kong, A. K. H. 2004, ApJ, 609, 710
- Di Stefano, R., Friedman, R., Kundu, A., & Kong, A. K. H. 2003a, arXiv:astro-ph/0312391
- Di Stefano, R. et al. 2003b, ApJ, 599, 1067
- Di Stefano, R., & Kong, A. K. H. 2003b, ApJ, 592, 884
- Di Stefano, R., & Nelson, L. A. 1996, Supersoft X-Ray Sources, 472, 3
- Di Stefano, R., Paerels, F., & Rappaport, S. 1995, ApJ, 450, 705
- Di Stefano, R., & Rappaport, S. 1994, ApJ, 437, 733
- Dobbie, P. D., et al. 2006, MNRAS, 369, 383
- Fabbiano, G., King, A. R., Zezas, A., Ponman, T. J., Rots, A., & Schweizer, F. 2003, ApJ, 591, 843
- Fujimoto, M. Y. 1982, ApJ, 257, 767
- Graham, M. L., et al. 2008, AJ, 135, 1343
- Greiner, J. 2000, New Astronomy, 5, 137
- Han, Z., & Podsiadlowski, P. 2006, MNRAS, 368, 1095
- Henze, M., et al. 2009, A&A, 500, 769
- Iben, I., Jr. 1982, ApJ, 259, 244
- Jenkins, L. P., Roberts, T. P., Warwick, R. S., Kilgard, R. E., & Ward, M. J. 2005, MNRAS, 357, 401

- Kalirai, J. S., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., & Richer, H. B. 2008, *ApJ*, 676, 594
- Kong, A. K. H., & Di Stefano, R. 2005, *ApJ*, 632, L107
- Kong, A. K. H., Di Stefano, R., & Yuan, F. 2004, *ApJ*, 617, L49
- Kong, A. K. H., & Di Stefano, R. 2003, *ApJ*, 590, L13
- Kotak, R. 2009, *Astronomical Society of the Pacific Conference Series*, 401, 150
- Kuznetsova, N., et al. 2008, *ApJ*, 673, 981
- Liu, J. 2008, arXiv:0811.0804
- Lü, G., Yungelson, L., & Han, Z. 2006, *MNRAS*, 372, 1389
- Miller, G. E., & Scalo, J. M. 1979, *ApJS*, 41, 513
- Mukai, K. et al. 2005, *ApJ*, 634, 1085
- Nomoto, K. 1982, *ApJ*, 253, 798
- Nomoto, K., Miyaji, S., Sugimoto, D., & Yokoi, K. 1979, *IAU Colloq. 53: White Dwarfs and Variable Degenerate Stars*, 56
- Orio, M. 2006, *ApJ*, 643, 844
- Paczynski, B. 1971, *ARA&A*, 9, 183
- Panagia, N., Della Valle, M., & Mannucci, F. 2007, *The Multicolored Landscape of Compact Objects and Their Explosive Origins*, 924, 373
- Pence, W. D., Snowden, S. L., Mukai, K., & Kuntz, K. D. 2001, *ApJ*, 561, 189
- Pietsch, W. et al. 2005, *A&A*, 442, 879
- Poznanski, D., et al. 2007, *MNRAS*, 382, 1169
- Rappaport, S., Di Stefano, R., & Smith, J. D. 1994, *ApJ*, 426, 692
- Rappaport, S., Chiang, E., Kallman, T., & Malina, R. 1994, *ApJ*, 431, 237
- Riess, A. G., et al. 2007, *ApJ*, 659, 98
- Roelofs, G., Bassa, C., Voss, R., & Nelemans, G. 2008, *MNRAS*, 391, 290

- Sarazin, C. L., Irwin, J. A., & Bregman, J. N. 2001, *ApJ*, 556, 533
- Shen, K. J., & Bildsten, L. 2008, *ApJ*, 678, 1530
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Swartz, D. A., Ghosh, K. K., Suleimanov, V., Tennant, A. F., & Wu, K. 2002, *ApJ*, 574, 382
- Turatto, M., Cappellaro, E., & Benetti, S. 1994, *AJ*, 108, 202
- van den Heuvel, E.P.J., Bhattacharya, D., Nomoto, K., & Rappaport, S.A. 1992, *A&A*, 262, 97.
- Voss, R., & Nelemans, G. 2008, *Nature*, 451, 802
- Weidemann, V., & Koester, D. 1983, *A&A*, 121, 77
- Williams, K. A. 2007, 15th European Workshop on White Dwarfs, 372, 85