

A Spin Chain for the Symmetric Product CFT₂

Ari Pakman^{*1}, Leonardo Rastelli^{†2}, and Shlomo S. Razamat^{‡2}

¹ *Department of Physics,
Brown University,
Providence, RI 02912, USA*

² *C.N. Yang Institute for Theoretical Physics,
Stony Brook University,
Stony Brook, NY 11794-3840, USA*

ABSTRACT:

We consider “gauge invariant” operators in $\text{Sym}^N T^4$, the symmetric product orbifold of N copies of the $2d$ supersymmetric sigma model with T^4 target. We discuss a spin chain representation for single-cycle operators and study their two point functions at large N . We perform systematic calculations at the orbifold point (“tree level”), where non-trivial mixing is already present, and some sample calculations to first order in the blow-up mode of the orbifold (“one loop”).

KEYWORDS: CFT, AdS/CFT.

*Email: ari_pakman@brown.edu

†Email: leonardo.rastelli@stonybrook.edu

‡Email: razamat@max2.physics.sunysb.edu

Contents

1. Introduction	2
2. Definition of the spin chain	4
2.1 Generic gauge invariant state	4
2.2 Towards a spin chain	7
3. Impurities and tree level computations	11
3.1 Fermionic Impurities	12
3.2 J^- impurities	17
3.3 Fractional moded J^- impurities	18
3.4 ∂X impurities	21
4. The spin chain at one loop	22
4.1 The $\mathcal{N} = (4,4)$ supersymmetry algebra and the interaction term	22
4.2 The map to the covering surface in presence of the interactions	25
4.3 Evaluating $t_L(t)$	29
4.4 Non-renormalization of the chiral vacuum at one loop	33
4.5 Comments on one loop with impurities	37
5. Summary and Discussion	39
A. A short explanation of the diagrams	40
B. Diagonalization of the mixing of two J^- impurity states	42
C. Some details of the one loop map	42
D. Arutyunov-Frolov map	44
E. The one loop correlator of bare twist fields	46

1. Introduction

In gauge/string dualities, it is desirable to fix the bulk-to-boundary dictionary as precisely as possible. In general this is a daunting task, but there are some very (super)symmetric examples where integrability makes the problem more tractable. In the maximally supersymmetric AdS_5/CFT_4 duality, the discovery of integrability both in the string sigma model [1] and in the large N field theory [2] has triggered spectacular progress in calculating and matching the spectrum of states. See for instance [3, 4, 5, 6, 7] for reviews of this very vast literature. More recently this program has been extended to the AdS_4/CFT_3 duality [8, 9], see *e.g.* [10, 11, 12, 13]. Curiously another classic duality, the AdS_3/CFT_2 correspondence [14], has been largely left out from these developments. While on the string theory side there is evidence for integrability [15, 16], much less is known on the field theory side. This predicament is the background motivation for this paper.

To describe the peculiarities of AdS_3/CFT_2 , let us briefly review how integrability works in the higher-dimensional cases. In $4d$ Yang-Mills (and also in $3d$ Chern-Simons coupled to matter), the elementary gauge invariant states at large N are single-trace operators, which can be viewed as spin chain systems with periodic boundary conditions. The dilatation operator of $\mathcal{N} = 4$ SYM is identified with an integrable spin chain Hamiltonian. The spin chain Hamiltonian is of nearest-neighbor type to lowest order in planar perturbation theory, and it becomes more and more non-local to higher orders. In fact at higher orders the Hamiltonian is not explicitly known and the most effective tool is instead the S-matrix of asymptotic magnon excitations propagating on the spin chain [17]. An analogous S-matrix can be defined for the light-cone string sigma model, and when phrased in this language the string theory analysis and field theory analysis become largely isomorphic. The AdS/CFT S-matrix is fixed by symmetries and various consistency requirements and is the main input in an asymptotic Bethe ansatz [18], which in principle allows to calculate the dimension of all “long” operators for arbitrary coupling. More recently a conjecture has been formulated that extends the Bethe ansatz to arbitrary finite-size operators [19].

The same ideas and techniques should apply at least to the string side of the AdS_3/CFT_2 duality. However the real interest would be to develop the string and field theory side simultaneously and understand their dictionary. The CFT_2 side is the less understood and it is the focus of our work. In a certain (possibly singular) region of its moduli space, the theory is believed to be described by a deformation of $\text{Sym}^N T^4$, see *e.g.* [20, 21]. This is the best analogy we have for the weakly coupled region of Yang-Mills or Chern-Simons theory. However the notion of a spin chain is less obvious than in a gauge theory, and perturbative computations (in conformal perturbation theory, infinitesimally away from the orbifold point) take a qualitatively different form. Following the ideas of BMN [22] a spin chain language for symmetric product orbifolds was put forward by several authors [23, 24, 25]. In [26] (see

also [27]) the giant magnon solutions on the gravity side were mapped to certain excitations above the chiral vacuum in the symmetric product orbifold. A dynamical spin chain picture based on the symmetry properties of the theory was suggested and an all-order dispersion relation for the magnon proposed using a central extension of the symmetry algebra. However the magnon S-matrix has not been computed yet. While on the sigma model side this is in principle a straightforward calculation, on the CFT_2 side even setting up the question is non-trivial and seems to require the proper construction of a spin chain in “position space”, which would allow a concrete definition of asymptotic magnon excitations.

In this paper, which builds upon our recent work [28, 29], we set the stage for a systematic discussion of “gauge invariant” states in symmetric product orbifolds. We will study a specific “position space” spin chain picture for single-cycle operators, which are analogous to single-trace operators in gauge theory. We start in section 2 by reviewing basic facts about $\text{Sym}^N T^4$. We discuss the most generic gauge invariant operators and introduce a spin chain interpretation for single-cycle operators, which can be regarded as the elementary building blocks at large N . In section 3 we define several different ways to introduce “impurities” on the chain and compute two point functions of states with impurities at the orbifold point. By analogy with the standard gauge theory case we will refer to these as “tree level” calculations, since they are the closest we can get to free field theory calculations. The analogy is however far from perfect, since even at the orbifold point correlators of twisted fields are non-trivial (there is no simple sense in which Wick theorem applies). Unlike the gauge theory spin chain, we encounter large mixing already at “tree level”: for example two states that differ by the position of one impurity are in general not orthogonal, even at large N . In section 4 we turn on an exactly marginal deformation that preserves $(4, 4)$ supersymmetry and discuss computations at leading non-trivial order in the deformation (“one loop”). Because of the complication of tree level mixing, the fundamental question of whether the one-loop Hamiltonian is “local” is difficult to answer. We find however some encouraging hints. In section 5 we summarize and discuss our results. Several technical appendices complement the text.

We end this introduction with a brief recapitulation of the current evidence for the holographic correspondence between $\text{Sym}^N T^4$ and type IIB string theory on $AdS_3 \times S^3 \times T^4$ [14]. See [30, 31, 32, 33] for reviews. The early checks of this duality included comparison of the moduli spaces [34, 21], the spectra of both theories [35, 36, 37, 38], and the symmetries [39, 40, 41]. Recently much progress was made in comparing correlation functions. The structure constants of single-cycle operators in the chiral ring of the symmetric product were computed early on in [42] and, for a subset of these operators they were extended in [43, 44] to the full $1/2$ BPS $SU(2)$ multiplet. These three-point functions were exactly reproduced in the string theory/supergravity dual [45, 46, 47, 48] (see also [49, 50, 51]), which also predicts some correlators not yet computed in the symmetric product [47]. The bulk-boundary agreement

of these computations, performed far apart in the moduli space [34, 21], is explained by a non-renormalization theorem proved in [52]. The latter also holds for extremal correlators, a large class of which was computed in the symmetric product orbifold in [29]. For examples of explicit computations of correlators in symmetric product orbifolds see [53, 54, 43, 44, 28, 55, 56]. The AdS_3/CFT_2 duality was also discussed in the pp-wave limit [23, 24, 25, 57].

2. Definition of the spin chain

2.1 Generic gauge invariant state

We are interested in classifying and studying gauge invariant operators in symmetric product orbifolds. For a general discussion of symmetric product orbifolds we refer the reader to [28] and references therein. The specific theory of our interest will be $\text{Sym}^N T^4$. This theory has the following matter content: 4 real left/right mover fermions and 4 real bosons, each coming in N copies. The different copies of the fields are identified under the action of the group of permutations S_N . In analogy to gauge theory we will refer to the index I of the copies of the fields as the “color” index.

The four real holomorphic fermions of T^4 can be combined, in each copy I , into two complex fermions ψ_I^1, ψ_I^2 , (with $I = 1 \dots N$) and bosonized as

$$\psi_I^1 = e^{i\phi_I^1}, \tag{2.1}$$

$$\psi_I^2 = e^{i\phi_I^2}, \quad I = 1, \dots, N. \tag{2.2}$$

In each copy I , we pair the four real bosons into two complex bosons X_α^I ($\alpha = 1, 2$).

The basic observables of a symmetric product orbifold are the twist fields $\sigma_{[g]}$, labeled by a conjugacy class $[g]$ of the permutation group. “Gauge invariant” twist fields $\sigma_{[g]}$ can be constructed from “gauge non-invariant” ones, σ_g , associated to a group element $g \in S_N$ and not to a conjugacy class. The operator $\sigma_g(z, \bar{z})$ is defined as a “defect” imposing the following monodromies on the different copies of the fields

$$X_I^i(e^{2\pi i} z) \sigma_g(0) = X_{g(I)}^i(z) \sigma_g(0), \tag{2.3}$$

and similarly for the fermionic fields. Gauge invariant operators are obtained by averaging over the group orbit,

$$\sigma_{[g]} \equiv \mathcal{A}_{[g]}(N) \sum_{h \in S(N)} \sigma_{h^{-1}gh}, \tag{2.4}$$

where $\mathcal{A}_{[g]}(N)$ is an appropriate normalization, see *e.g.* [28].

The theory has $\mathcal{N} = 4$ supersymmetry. We will review the realization of the algebra in terms of the fields of the theory in section 4.1. A notable set of gauge invariant states is

the set of (anti)chiral states under some $\mathcal{N} = 2$ subalgebra. Let us discuss these first (see *e.g.* [42, 44, 29]). The $U(1)$ current of a $\mathcal{N} = 2$ subalgebra of supersymmetry we will use is

$$J = \frac{1}{2} \sum_{I=1}^N \left(\psi_I^1 \psi_I^{1\dagger} + \psi_I^2 \psi_I^{2\dagger} \right) = \frac{i}{2} \sum_{I=1}^N \left(\partial \phi_I^1 + \partial \phi_I^2 \right). \quad (2.5)$$

We define the gauge-non-invariant chiral operators associated to the single-cycle $g = (12 \dots n)$,

$$o_{(12 \dots n)}^{(0,0)} = e^{i \frac{n-1}{2n} \sum_{I=1}^n (\phi_I^1 + \phi_I^2 + \bar{\phi}_I^1 + \bar{\phi}_I^2)} \sigma_{(12 \dots n)}, \quad (2.6)$$

$$o_{(12 \dots n)}^{(a=1, \bar{a}=1)} = e^{i \frac{n+1}{2n} \sum_{I=1}^n (\phi_I^1 + \bar{\phi}_I^1) + i \frac{n-1}{2n} \sum_{I=1}^n (\phi_I^2 + \bar{\phi}_I^2)} \sigma_{(12 \dots n)}, \quad (2.7)$$

$$o_{(12 \dots n)}^{(a=2, \bar{a}=2)} = e^{i \frac{n-1}{2n} \sum_{I=1}^n (\phi_I^1 + \bar{\phi}_I^1) + i \frac{n+1}{2n} \sum_{I=1}^n (\phi_I^2 + \bar{\phi}_I^2)} \sigma_{(12 \dots n)}, \quad (2.8)$$

$$o_{(12 \dots n)}^{(2,2)} = e^{i \frac{n+1}{2n} \sum_{I=1}^n (\phi_I^1 + \phi_I^2 + \bar{\phi}_I^1 + \bar{\phi}_I^2)} \sigma_{(12 \dots n)}. \quad (2.9)$$

The gauge invariant operators are obtained by summing over the group orbit,

$$O_n^{(0,0)} = \frac{1}{\sqrt{n N!(N-n)!}} \sum_{h \in S(N)} o_{h^{-1}(12 \dots n)h}^{(0,0)}, \quad (2.10)$$

$$O_n^{(a, \bar{a})} = \frac{1}{\sqrt{n N!(N-n)!}} \sum_{h \in S(N)} o_{h^{-1}(12 \dots n)h}^{(a, \bar{a})}, \quad (2.11)$$

$$O_n^{(2,2)} = \frac{1}{\sqrt{n N!(N-n)!}} \sum_{h \in S(N)} o_{h^{-1}(12 \dots n)h}^{(2,2)}. \quad (2.12)$$

The conformal dimensions and charges are

$$\Delta_n^0 = Q_n^0 = \frac{n-1}{2}, \quad (2.13)$$

$$\Delta_n^a = Q_n^a = \frac{n}{2}, \quad (2.14)$$

$$\Delta_n^2 = Q_n^2 = \frac{n+1}{2}, \quad (2.15)$$

and similarly for the antiholomorphic sector. The antichiral operators $O_n^{(0,0)\dagger}$, $O_n^{(a, \bar{a})\dagger}$, $O_n^{(2,2)\dagger}$ are obtained by reversing the sign in the exponents in (2.6)-(2.9).¹

Let us now build a generic gauge invariant state. We will denote schematically all the fields in copy I as χ_I . The generic state invariant under the action of the permutation group S_N is built from

$$o_g = \mathcal{G} [\chi_I | g \cdot I = I] \mathcal{F} [\chi_I | g \cdot I \neq I] \sigma_g. \quad (2.16)$$

¹One can also define operators with different left and right properties.

Here g is some group element of S_N . We will refer to the twist field σ_g as *bare* twist field to emphasize that it is an operator without any \mathcal{F} and \mathcal{G} dressing. The function $\mathcal{G}[\chi_I|g \cdot I = I]$ commutes with the bare twist field by definition and is arbitrary. However, we have to demand that $\mathcal{F}[\chi_I|g \cdot I \neq I]$ also commutes with σ_g and this implies certain non trivial restrictions. In the example of chiral fields (2.6)-(2.9) the fermionic dressing is the \mathcal{F} part and there is no \mathcal{G} part. A set of gauge invariant states is then given by

$$\mathcal{O}_{[g]} = \sum_{\beta \in S_N} o_{\beta g \beta^{-1}}, \quad (2.17)$$

and the most generic gauge invariant state is built from linear combinations of these states. The ultimate goal is to find the spectrum of conformal dimensions of the gauge invariant operators after turning on interaction terms. While the general problem is hugely complicated, it should simplify in the large N limit, which corresponds to the classical limit on the string theory side. The theory is expected to be integrable in this limit.

The states $\mathcal{O}_{[g]}$ are classified according to the conjugacy class $[g]$, which is specified by the cycle structure of the group element, *i.e.* the number of single-cycles and their length. States with different cycle structure are exactly orthogonal, even at finite N , since for $\langle o_g o_h \rangle$ to be non zero the product gh has to be the identity. It is common (see *e.g.* [23, 24, 26]) to draw an analogy between multi-cycle states in the symmetric product orbifold and multi-trace states in gauge theories. Both, single traces and single-cycles, have a natural cyclic structure and are bounded in size by the number of colors N . (In the symmetric product this is a strict bound, while in a gauge theory it is the statement that a single trace operator longer than N can be re-written as a linear combination of shorter multi-traces.) An important rationale for the analogy is the fact that in both cases single (multiple) trace/cycle states correspond to single (multiple) gravitons. This is confirmed for single gravitons by the bulk-boundary agreement of three-point functions in both cases [58, 45, 46, 47, 48]. While multi-trace states in the gauge theory are not exactly orthogonal at finite N (unlike multi-cycle states in the symmetric product), they become orthogonal at large N (if the length of the traces is kept fixed), which is the relevant limit for our discussion.

Since the cycle structure of $[g]$ is specified by a partition of n , with $n \leq N$, we can conveniently associate $[g]$ to a Young tableau. If $n = \sum_k n_k$ where n_k is the number of cycles of length k , we draw a tableau with n_k columns of length k , that is we associate to each single-cycle of length k a column with k boxes. The untwisted sector states of the orbifold correspond to columns of the Young tableaux with a single box, and therefore we can represent an untwisted state which involves k colors as a single row with k boxes. This is illustrated in figure 1. The total number n of boxes in the Young tableaux is bounded by N , and this is related to the stringy exclusion principle [35]. We can use a similar Young tableaux representation for multi-trace states in a gauge theory, associating single-traces of length k

to columns of length k , in keeping with our analogy between single-cycles and single-traces. A column with a single box corresponds to the trace of a single field in the adjoint of $U(N)$, *i.e.* to its $U(1)$ part. In some rough sense the untwisted states of the symmetric product are analogous to states in the $U(1)$ decoupled sector of the gauge theory. Unlike the symmetric product case, in the gauge theory there is no upper bound on the total number of boxes.²




Figure 1: (1) Young tableau representation of a state in the untwisted sector, $\mathcal{O} = \sum_{I \neq J \neq K \neq L} e^{i(\chi_I + \chi_J + \chi_K + \chi_L)}$. Here \mathcal{O} has only a \mathcal{G} part. (2) Representation of a state in a twisted sector, $\mathcal{O} = \sum_{I \neq J \neq K \neq L} e^{i(\chi_I + \chi_J + \chi_K + \chi_L)} \sigma_{(IJKL)}$. Here there is only an \mathcal{F} part. (3) Representation of the state $\mathcal{O} = \sum_{I \neq J \neq K} e^{i(\chi_I + \chi_J + \chi_K)} \sigma_{(IJ)}$. The \mathcal{G} dressing is $\mathcal{G} = e^{i\chi_K}$, and the \mathcal{F} dressing is $\mathcal{F} = e^{i(\chi_I + \chi_J)}$. (4) Representation of the state $\mathcal{O} = \sum_{I \neq J \neq K \neq L} e^{i(\chi_I + \chi_J + \chi_K + \chi_L)} \sigma_{(IJ)} \sigma_{(KL)}$. There is only \mathcal{F} dressing, $\mathcal{F} = e^{i(\chi_I + \chi_J + \chi_K + \chi_L)}$.

In what follows we consider the strict infinite N limit of the symmetric product. In this limit correlators of generic gauge invariant operators factorize into correlators of single-cycle operators with trivial $\mathcal{G}[\chi_I]$ part, and untwisted correlators. Single-cycle operators with trivial $\mathcal{G}[\chi_I]$ are elementary building blocks, and at large N form a close sector under the action of the dilatation operator, even away from the orbifold point. They are expected to map to single closed string states on the dual side.

2.2 Towards a spin chain

As familiar, in a gauge theory we can represent single-trace operators as spin chains with periodic boundary conditions. The simplest example is the $SU(2)$ sector of $\mathcal{N} = 4$ SYM, which consists of operators made of two adjoint scalars, Z and X , which are thought as spin

²At finite N , a useful orthogonal basis for Yang-Mills operators built out of a single adjoint scalar Z is given by the Schur polynomial basis [59]. Schur polynomials are also naturally represented by Young tableaux, which should not be confused with the way we use Young tableaux to represent multi traces. In the Schur basis, a column of the Young tableau of length $k \lesssim N$ is associated to a subdeterminant and is holographic to sphere giant gravitons [60], while a row is holographic to AdS giant gravitons [61]. See [62, 63] for the issue of giant gravitons in $AdS_3 \times S^3 \times T^4$.

up and spin down. The operator with only Z s, $\text{Tr}Z^n$, is the vacuum of the spin chain (of length n); replacing Z with X at various sites we introduce “impurities”. In some special gauge theories, *e.g.* $\mathcal{N} = 4$ SYM [2], the one-loop dilatation operator turns out to be the nearest-neighbor Hamiltonian of an integrable spin chain – for example, it is the Heisenberg spin chain in the $SU(2)$ sector. At higher orders the Hamiltonian becomes very complicated, and it has proved more fruitful to phrase the integrability structure in terms of the S-matrix of asymptotic magnon excitations of the spin chain.

We would like to uncover the analogous notions for symmetric product orbifolds in general and in particular for $\text{Sym}^N T^4$. We assume the analogy between a single-trace operator in a gauge theory and a single-cycle twist field in the symmetric product. Next we should specify what we mean by an individual “site” of the single-cycle twist field. In Yang-Mills the sites are identified with the elementary adjoint fields of the composite operator. In the symmetric product, it was suggested by several authors [23, 24, 26] to decompose a given n -cycle element of S_N as a product of transpositions, and to consider this decomposition as a collection of $n - 1$ sites. For example

$$(1\ 2\ 3\ 4\ 5) = (1\ 5)(5\ 4)(4\ 3)(3\ 2), \quad (2.18)$$

$$= (2\ 3)(3\ 4)(4\ 5)(5\ 1). \quad (2.19)$$

There are however some difficulties with this identification. First, as we sprinkle the different sites with impurities, the dressed transpositions will generally have singularities in the OPE with one another, and some prescription must be specified in recombining the transpositions into a single-cycle. More fundamentally, as is clear from the above example, the decomposition of a single-cycle into transpositions is not unique and it breaks the explicit cyclic structure of the single-cycle. Given a single-cycle operator there is no canonical gauge invariant way to specify the sites of the associated spin chain.

To avoid these problems we propose to identify the sites of the spin chain with the “colors” permuted by the cycle. This gives the sites a natural cyclic ordering. There *is* a natural gauge invariant way to act on a single site of the chain, as follows (see [26] for similar manipulations). We insert an impurity at a site I by inserting an operator \hat{P}_I that depends on the elementary fields χ_I of that site (=color I),

$$\hat{P}_I \sigma_{(12\dots n)}, \quad (2.20)$$

where σ_n is the gauge non-invariant twist field. Then we sum cyclically over the n sites of the chain,

$$\hat{P}\sigma_{(12\dots n)} \equiv \sum_{I=1}^n \hat{P}_I \sigma_{(12\dots n)}, \quad (2.21)$$

and finally we sum over all relabelings of the colors to get the desired gauge invariant operator (as in (2.17)),

$$\sum_{h \in S_N} \sum_{I=1}^n \hat{P}_I \sigma_{h(12\dots n)h^{-1}}. \quad (2.22)$$

In many concrete cases it will be useful to represent \hat{P}_I as a contour integral

$$\hat{P}_I = \oint \frac{dz}{2\pi i} P_I(z) \quad (2.23)$$

$$\hat{P} \tilde{o}_n(0) \equiv \sum_{I=1}^n \oint \frac{dz}{2\pi i} P_I(z) \tilde{o}_n(0). \quad (2.24)$$

Here $\tilde{o}_n(z)$ is a generic gauge non-invariant operator in the twisted sector associated to the single-cycle $(12\dots n)$. It is most convenient to discuss symmetric product orbifolds on a covering surface where the twist fields disappear and the fields are single-valued. The exact covering map, $z(t)$, is determined by the correlator being evaluated. The covering map has the property that near the location of single-cycle twist field, say $z = 0$ and $t = 0$, it behaves as $z(t) \sim a t^n$. On the covering surface the action of an operator on a single color can be elegantly written as

$$\hat{P} \tilde{o}_n(0) \sim \oint \frac{dt}{2\pi i} \left(\frac{dt}{dz} \right)^{\delta-1} P(t) \tilde{o}_n^{(t)}(0). \quad (2.25)$$

where δ is the conformal dimension of $P(z)$ and the superscript t in $\tilde{o}_n^{(t)}$ denotes that the operator is lifted to the covering surface, on which the bare twist field σ_n disappears. The fact that we can easily lift the definition of the impurities to the covering surface will give us powerful computational tools to evaluate correlators.

To introduce two impurities at different sites of the chain one proceeds as follows. The action of operator $\hat{P}^{(1)}$ and $\hat{P}^{(2)}$ on two different sites, I and $I + L$, becomes

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} P_I^{(1)}(z) P_{I+L}^{(2)}(z) \tilde{o}_n \rightarrow \oint \frac{dt}{2\pi i} \left(\frac{dt}{dz} \right)^{\delta_1-1} \left(\frac{dt_L}{dz} \right)^{\delta_2} P^{(1)}(t) P^{(2)}(t_L(t)) \tilde{o}_n^{(t)}. \quad (2.26)$$

where the ordering on the color is defined by the cyclic ordering inferred from the structure of the twist field. The function $t_L(t)$ satisfies the following

$$z(t) = z(t_L(t)), \quad (2.27)$$

and for t in the vicinity of the insertion of the chiral field $t = 0$ we have

$$t_L(t) \sim e^{\frac{2\pi i}{n} L t}. \quad (2.28)$$




Figure 2: A vertex corresponding to a seven-cycle operator \mathcal{O}_7 . The red circles are color loops. We can think of the red dots as sites of a spin chain.

Near the pre-image of a twist field on the covering surface if a position t corresponds on the base to color I then the position $e^{\frac{2\pi i}{n}L}t$ corresponds to color $I + L$. The function $t_L(t)$ is crucial to define the action of operators on different sites of the chain and we will discuss several explicit examples in the following sections.

The generalization to states with many impurities is straightforward. The most general single-cycle state with trivial \mathcal{G} part is a linear combination of states of the form

$$\hat{o}_n = \oint \frac{dt}{2\pi i} \left(\frac{dt}{dz} \right)^{-1} \prod_{I=1}^n \left(\frac{dt_{L=I}}{dz} \right)^{\delta_I} \prod_{I=1}^n P^{(I)}(t_{L=I}), \quad (2.29)$$

with $t_{L=1} \equiv t$.

We can represent the spin chain graphically, using a diagrammatic language for symmetric product orbifolds introduced in [28] and briefly reviewed in appendix A. In this language a single-cycle twist field is represented as a loop with $2n$ vertices, see figure 2. The vertices are of two types, “color” and “non-color” vertices. In the limit of large length of the chain we will depict it as a horizontal line. The “Feynman diagrams” for correlators are obtained by gluing together appropriately the vertices of loops corresponding to different twist fields. An example of a tree level two point function is depicted in figure 3.

The next step is to settle on a choice of vacuum for the spin chain and to identify its basic excitations, *i.e.* the impurities. A natural choice is to identify the vacuum as the chiral state with lowest dimension for a given length n [23, 24, 26], namely the state

$$o_n \equiv o_n^{(0,0)}. \quad (2.30)$$

The other chiral primary fields (2.10)-(2.12) can be obtained from o_n by the action of some operators. The chiral state satisfies $\Delta - J = 0$ and the basic impurities can be defined as operators acting on a single site of a chain and having minimal $\Delta - J$. In the next section we




Figure 3: A qualitative picture of the two point diagram. The two solid lines correspond to dressed twist field. We have taken the length of the cycles to be large and have drawn the circles corresponding to twist fields as infinite lines. The diagrams of [28] are obtained by identifying the vertices according to the dashed lines.

will discuss such excitations. After the action of the impurities on o_n , we should symmetrize over the orbit of the S_N group as in (2.10).

Before going into a detailed analysis of the different types of impurities, let us briefly review the situation on the dual string side [22, 23, 24, 25]. In the light-cone gauge, the asymptotic excitations of the (massive) worldsheet sigma model fall into two classes: there are excitations related to the $AdS_3 \times S^3$ portion of the geometry, which are in one-to-one correspondence with the generators of the superconformal algebra left unbroken by the light-cone gauge choice, and excitations related to the T^4 . By contrast in $AdS_5 \times S^5$ all excitations are of the first type – they are associated with the unbroken superconformal generators. We will indeed find that in $\text{Sym}^N T^4$ the spectrum of impurities is in a sense richer than in $\mathcal{N} = 4$ SYM: besides “universal” impurities associated with the symmetry generators, there are impurities associated to non-trivial primaries of T^4 , which are perhaps more elementary. In the CFT_2 the holomorphic objects with lowest dimension are the fermions ($\Delta = \frac{1}{2}$), while the symmetry generators start at $\Delta = 1$ (the R -symmetry currents). Single-fermion impurities in the spin chain of $\text{Sym}^N T^4$ have no direct analogue in $\mathcal{N} = 4$ SYM.

3. Impurities and tree level computations

In this section we consider different types of impurities on the vacuum represented by the chiral operator o_n . A natural choice is to create impurities by acting with symmetry generators on the sites of the spin chain [23, 24, 26], *e.g.* the modes of the R -current and the susy generators. However, the smallest-dimensional fields in the orbifold are the fermions and thus the most fundamental impurities one can build include these objects.³ In what follows we will study impurities generated by modes of the fermions (ψ^a and $\psi^{a\dagger}$), and by modes

³One can always restrict to states with impurities generated only by symmetry currents, as these form a closed sector of the theory.

of the bosons (∂X^i). We also discuss impurities generated by modes of the R -current J^- , which is a quadratic composite of the fermions. The discussion will focus mostly on states with two impurities at a distance L , and in each case we check whether states with different L 's are orthogonal at tree level - as is the case in $\mathcal{N} = 4$ SYM in the limit of large N . In symmetric product CFTs at tree level the only covering surfaces contributing to two point functions have topology of a sphere.⁴ Thus, the full N dependence is given by a simple overall combinatorial factor [28]. In particular the large N considerations do not play any role in tree level computations. As we will see, some types of two-impurity states are orthogonal, some are not and some are orthogonal only when the length n of the spin chain (not to be confused with the number N) is taken to be large. These results illustrate the lack of a sharp analogy between the symmetric product and $\mathcal{N} = 4$ SYM.

3.1 Fermionic Impurities

We first introduce fermionic impurities by directly constructing the states in the bosonized language for the fermions. We then show that this construction can be recast as the action of a certain current algebra on the chiral vacuum.

Direct Construction

The states with a single impurity are defined as the following currents acting on the chiral vacuum o_n ,

$$A^a = \sum_{I=1}^n \oint \frac{dz}{2\pi i} \frac{1}{z} e^{-i\phi_I^a(z)}, \quad a = 1, 2 \quad (3.1)$$

$$= \sum_{I=1}^n \psi_{I,-1/2}^a, \quad (3.2)$$

$$B^a = \sum_{I=1}^n \oint \frac{dz}{2\pi i} \frac{1}{z^2} e^{i\phi_I^a(z)}, \quad a = 1, 2 \quad (3.3)$$

$$= \sum_{I=1}^n \psi_{I,-3/2}^{a\dagger}, \quad (3.4)$$

and analogously for the anti holomorphic sector. Note that these impurities have different conformal dimensions and thus are orthogonal at tree level. Strictly speaking this definition is correct only for odd n , as otherwise the fermions are antiperiodic when rotated around the twist field [24]. The currents A^a increase the dimension by $1/2$ and decrease the charge by

⁴This can be deduced for instance from the Riemann-Hurwitz relation determining the genus of a covering surface by noting that the number of colors is uniquely determined to be equal to the size of the cycles, see *e.g.* [28].

1/2, while the currents B^a increase the dimension by 3/2 and increase the charge by 1/2, so both types of impurities satisfy $\Delta - J = 1$.

Note that we could also act on o_n with the following current

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} \frac{1}{z} e^{i\phi_I^a(z)} = \sum_{I=1}^n \psi_{I,-1/2}^{a\dagger}, \quad (3.5)$$

which has $\Delta - J = 0$. However, this gives us just the fermionic chiral state $o_n^{(a,0)}$, with $(\Delta, \bar{\Delta}) = (\frac{n}{2}, \frac{n-1}{2})$, and thus we should not treat this as an impurity.

Let us consider the A type impurities in more detail. The current generating two impurities is defined by

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} e^{-i\phi_I^a(z)} e^{-i\phi_{I+L}^b(z)} \quad (3.6)$$

$$= \sum_{I=1}^n [\psi_I^a \psi_{I+L}^b]_0, \quad (3.7)$$

where $[\dots]_0$ denotes the zero mode in the mode expansion of the normal product inside the square brackets. On the covering surface this becomes

$$\oint \frac{dt}{2\pi i} \sqrt{\frac{\partial t_L}{\partial t}} (t - t_L)^{\delta_{a,b}} e^{-i\phi^a(t) - i\phi^b(t_L)}. \quad (3.8)$$

Note that this expression is single valued and well defined, as near $t = 0$ the function $t_L(t)$ by definition behaves as $t_L(t) \sim e^{\frac{2\pi i}{n} L} t$ and thus we do not cross the branch cut of the square root. This impurity does not change the conformal dimension of the chiral operator but reduces its charge by one unit, and thus has $\Delta - J = 1$. For $L > 0$, this current creates a two-impurity state with minimal $\Delta - J$ and this is in contrast to $\mathcal{N} = 4$ SYM where such states do not exist.

We define a state with two impurities by acting with the above current on a state o_n to obtain⁵

$$\begin{aligned} |L\rangle &= \oint \frac{dt}{2\pi i} \sqrt{\frac{\partial t_L}{\partial t}} (t - t_L)^{\delta_{a,b}} e^{-i\phi^a(t) - i\phi^b(t_L)} e^{i\frac{n-1}{2}(\phi^1(t) + \phi^2(t))} e^{i\frac{n-1}{2}(\bar{\phi}^1(\bar{t}) + \bar{\phi}^2(\bar{t}))} |0\rangle, \quad (3.9) \\ &= \oint \frac{dt}{2\pi i} \sqrt{\frac{\partial t_L}{\partial t}} \frac{(t - t_L)^{\delta_{a,b}}}{t^{\frac{n-1}{2}} t_L^{\frac{n-1}{2}}} e^{-i\phi^a(t) - i\phi^b(t_L) + i\frac{n-1}{2}(\phi^1(t) + \phi^2(t))} e^{i\frac{n-1}{2}(\bar{\phi}^1(\bar{t}) + \bar{\phi}^2(\bar{t}))} |0\rangle. \end{aligned}$$

Note that taking $L = 0$ for $a = b$ we get zero, but if $a \neq b$ we get single impurity created by J_0^- .

⁵Omitting an overall rescaling factor which should be correctly regularized in any correlator, *e.g.* following Lunin-Mathur procedure [43].

The generalization to an operator creating an even number, $2l + 2$, of impurities, *i.e.* a bosonic chain, is

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} z^l \prod_{j=1}^{2l+2} e^{-i\phi_{I+L_j}^{b_j}(z)}, \quad (3.10)$$

with $L_j < L_{j+1}$, and $L_1 = 0$. On the covering surface this becomes

$$\oint \frac{dt}{2\pi i} z(t)^l \sqrt{z'(t)} \prod_{j=2}^{2l+2} \frac{1}{\sqrt{z'(t_{L_j})}} \prod_{j \neq j'} (t_{L_j} - t_{L_{j'}})^{\delta_{b_j, b_{j'}}} e^{-i \sum_{j=1}^{2l+2} \phi^{b_j}(t_{L_j})}. \quad (3.11)$$

These states have $\Delta - J = l + 1$.

The generalization to an operator creating an odd number, $2l + 1$, of impurities, *i.e.* a fermionic chain, is

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} z^{l-1} \prod_{j=1}^{2l+1} e^{-i\phi_{I+L_j}^{b_j}(z)}, \quad (3.12)$$

with $L_j < L_{j+1}$, and $L_1 = 0$. On the covering surface this becomes

$$\oint \frac{dt}{2\pi i} z(t)^{l-1} \sqrt{z'(t)} \prod_{j=2}^{2l+1} \frac{1}{\sqrt{z'(t_{L_j})}} \prod_{j \neq j'} (t_{L_j} - t_{L_{j'}})^{\delta_{b_j, b_{j'}}} e^{-i \sum_{j=1}^{2l+1} \phi^{b_j}(t_{L_j})}. \quad (3.13)$$

This definition holds for odd n . These states have $\Delta - J = l + 1$. We note that there are bosonic and fermionic chains with same $\Delta - J$. Note also that $\Delta - J$ for the chain is roughly twice the number of the impurities.

Let us compute two point functions of two-impurity states at tree level. Putting the twist fields at 0 and ∞ , both in the base and the covering sphere, the map to the covering surface is

$$z(t) = t^n. \quad (3.14)$$

In particular this implies that

$$t_L(t) = e^{\frac{2\pi i}{n} L} t. \quad (3.15)$$

We define

$$p = e^{\frac{2\pi i}{n} L}, \quad q = e^{\frac{2\pi i}{n} M}, \quad (3.16)$$

to obtain

$$\langle M|L \rangle \sim p(1-p)^{\delta_{ab}}(1-q)^{\delta_{a'b'}} \oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \frac{t^{1-n+\delta_{ab}} t'^{n-1+\delta_{a'b'}-\delta_{ab'}-\delta_{a'b'}-\delta_{aa'}-\delta_{bb'}}{(1-\frac{t}{t'})^{\delta_{aa'}}(1-p\frac{t}{t'})^{\delta_{ba'}}(q-\frac{t}{t'})^{\delta_{ab'}}(q-p\frac{t}{t'})^{\delta_{bb'}}}. \quad (3.17)$$

In order to get simple poles for both contour integrals above we have to satisfy

$$2 - n + \delta_{ab} + \alpha = 0, \quad n + \delta_{a'b'} - \delta_{ab'} - \delta_{a'b} - \delta_{aa'} - \delta_{bb'} - \alpha = 0, \quad (3.18)$$

where α is the expansion order in t/t' of the denominator. This implies the selection rule

$$\delta_{ab} + \delta_{a'b'} - \delta_{ab'} - \delta_{a'b} - \delta_{aa'} - \delta_{bb'} + 2 = 0, \quad (3.19)$$

which implies either $a = b = a' = b'$ or $a = a' \neq b = b'$ ($a = b' \neq b = a'$). Evaluating the integrals by residues we obtain, up to an overall constant, for the first case,

$$\langle M|L \rangle \sim -n \delta_{L,M} + n \delta_{L,-M}, \quad (3.20)$$

and for the second case

$$\langle M|L \rangle \sim -1 + n \delta_{L,M}, \quad (\langle M|L \rangle \sim -1 + n \delta_{L,-M}). \quad (3.21)$$

In the large n limit, *i.e.* long spin chain, all these two-impurities states are orthogonal.

As another example of tree level properties let us consider the two point functions of states with two ψ_1 and two ψ_2 impurities. The two point function is given by

$$\langle \{M_i\} | \{L_i\} \rangle \sim \oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \frac{p\hat{p}_1\hat{p}_2(1-p)(\hat{p}_1-\hat{p}_2)(1-q)(\hat{q}_1-\hat{q}_2)t^{5-2n}t'^{2n-7}}{(1-\frac{t}{t'})(1-p\frac{t}{t'})(q-\frac{t}{t'})(q-p\frac{t}{t'})(\hat{q}_1-\hat{p}_1\frac{t}{t'})(\hat{q}_1-\hat{p}_2\frac{t}{t'})(\hat{q}_2-\hat{p}_1\frac{t}{t'})(\hat{q}_2-\hat{p}_2\frac{t}{t'})}. \quad (3.22)$$

Here the ψ_1 impurities are at sites $e^{\frac{2\pi i}{n}L} = 1, p$ and $e^{\frac{2\pi i}{n}M} = 1, q$. The ψ_2 impurities are at $e^{\frac{2\pi i}{n}L} = \hat{p}_1, \hat{p}_2$ and $e^{\frac{2\pi i}{n}M} = \hat{q}_1, \hat{q}_2$. For generic values of M_i and L_i the above two point function is exactly zero. It is non vanishing when at least one of the L_i and M_i overlap. For instance if $p = q$ the above is equal to

$$\frac{2n\hat{p}_1\hat{p}_2(\hat{p}_1-\hat{p}_2)(\hat{q}_1-\hat{q}_2)}{(\hat{p}_1-\hat{q}_1)(\hat{q}_1-\hat{p}_2)(\hat{p}_1-\hat{q}_2)(\hat{p}_2-\hat{q}_2)}. \quad (3.23)$$

This expression scales as n . However, if say \hat{p}_1 is close enough to \hat{q}_1 , *i.e.* $L_i - M_j \ll n$, then the scaling is enhanced to n^2 . Farther, if all the impurities are “approximately” aligned in the above sense then the behavior is enhanced to n^3 .

In general, we can say that the states with impurities are orthogonal until at least one pair of impurities aligns on the two chains. Then, even in the large n limit, if the rest of the impurities are approximately aligned, *i.e.* the difference in their position is much less than n , the two point function is not zero at tree level and the states mix. This fact has to be contrasted with YM . There two states with impurities which are not aligned do not mix because of the large N suppression. However, the large N limit in symmetric product orbifold does not rule out such mixings.

Current Algebra Construction

In this subsection we present an algebraic approach to the impurities introduced above. For simplicity, we focus on two-impurity states with $a = b$ and we omit the labels a, b from the fermions.

The exponent that dresses the spin field of the chiral vacuum o_n in (2.6) is invariant under Z_n action of the twist field. In the same spirit, we define the following Z_n -invariant currents,

$$P_L^- \equiv \sum_{I=1}^n \psi_I^\dagger \psi_{I+L}^\dagger = \sum_{r \in \mathbb{Z}} \frac{P_{L,r}^-}{z^{m+1}}, \quad (3.24)$$

$$P_L^+ \equiv \sum_{I=1}^n \psi_I \psi_{I+L} = \sum_{r \in \mathbb{Z}} \frac{P_{L,r}^+}{z^{m+1}}, \quad (3.25)$$

$$N_L \equiv \sum_{I=1}^n \psi_I \psi_{I+L}^\dagger + \psi_{I+L} \psi_I^\dagger = \sum_{r \in \mathbb{Z}} \frac{N_{L,r}}{z^{m+1}}, \quad (3.26)$$

which have dimension $\Delta = 1$ and charges ± 1 and 0 under the $U(1)$ current J . Since these currents are Z_n -invariant they have integer-moded expansions near the chiral vacuum o_n , even though the modes cannot be expressed easily in terms of the modes of the fermions ψ_I . Note the properties

$$P_0^\pm = 0, \quad (3.27)$$

$$P_{-L}^\pm = -P_L^\pm, \quad (3.28)$$

$$N_L = N_{-L} = N_L^\dagger. \quad (3.29)$$

Using the same contour integral arguments as in the previous section, it is easy to verify that the chiral vacuum

$$|n\rangle \equiv o_n|0\rangle \quad (3.30)$$

is a highest weight state for the currents P_L^\pm, N_L , satisfying

$$P_{L,m}^+ |n\rangle = 0 \quad m \geq -1, \quad (3.31)$$

$$P_{L,m}^- |n\rangle = 0 \quad m \geq 1, \quad (3.32)$$

$$N_{L,m} |n\rangle = 0 \quad m \geq 1, \quad (3.33)$$

$$N_{L,0} |n\rangle = c_L |n\rangle, \quad (3.34)$$

with

$$c_L = 1 - n\delta_{L,0}. \quad (3.35)$$

In this language, the two-impurities states (3.9) are given by

$$|L\rangle = P_{L,0}^- |n\rangle, \quad (3.36)$$

and acting with higher modes we can build all the module of the algebra. Using

$$\psi_i(z)\psi_j^\dagger(w) \sim \frac{\delta_{ij}}{z-w}, \quad (3.37)$$

we can obtain the algebra satisfied by the currents,

$$P_{L_1}^+ P_{L_2}^- \sim \frac{\delta_{L_1, L_2} n}{(z-w)^2} + \frac{N_{L_1+L_2} - N_{L_1-L_2}}{z-w}, \quad (3.38)$$

or equivalently

$$\left[P_{L_1, r}^+ P_{L_2, s}^- \right] = nr \delta_{L_1, L_2} \delta_{r+s, 0} + N_{L_2+L_1, r+s} - N_{-L_2-L_1, r+s}. \quad (3.39)$$

Using now these commutation relations, we can compute the norm

$$\langle L_1 | L_2 \rangle = \langle n | P_{L_1, 0}^+ P_{L_2, 0}^- | n \rangle \quad (3.40)$$

$$= \langle n | N_{L_2+L_1, 0} - N_{-L_2-L_1, 0} | n \rangle \quad (3.41)$$

$$= c_{L_2+L_1} - c_{L_2-L_1} \quad (3.42)$$

$$= n \delta_{L_1, L_2} - n \delta_{L_1, -L_2} \quad (3.43)$$

which coincides, up to an overall constant which is not determined by the covering surface method, with (3.20). Impurities with $a \neq b$ can be obtained in a similar way, and the currents (3.24)-(3.26) can be easily generalized to the cases of three and more impurities. It would be interesting to compute the commutators for these cases and study the representation theory of these algebras.

3.2 J^- impurities

In this section we discuss impurities built from modes of J^- . Since the latter is a quadratic combination of the fermions, $\psi^1 \psi^2$, the state with a single J^- impurity is a special case of the fermionic two-impurities state we discussed in the previous section.⁶

But considering J^- as the basic impurity, we can build a state with two or more impurities of type J^- as follows. Denoting by $J_{(k)}^-$ the J^- current acting on k th copy, a two-impurities state is obtained from the zero mode of the combined operator (assuming that the current acts on state at $z = 0$ for simplicity)

$$\sum_{k=1}^n J_{(k)}^- J_{(k+L)}^- \rightarrow \sum_{k=1}^n \oint \frac{dz}{2\pi i} z e^{-i\phi_k^1(z) - i\phi_k^2(z)} e^{-i\phi_{k+L}^1(z) - i\phi_{k+L}^2(z)}. \quad (3.44)$$

⁶More generally, the two fermions ψ^a have same quantum numbers as the two supersymmetry generators broken by the vacuum, the quadratic combination $\psi^1 \psi^2$ is the R -current J^- , and the combination $\psi^1 \psi^{\dagger 2}$ has the same quantum numbers as the derivative ∂ .

Lifting this to the covering surface we obtain (assuming that the image of the twist field on the covering surface is $t = 0$)

$$\begin{aligned} \sum_{k=1}^n J_{(k)}^- J_{(k+L)}^- &\rightarrow \oint \frac{dt}{2\pi i} \frac{\partial t_L}{\partial z} z(t) e^{-i\phi^1(t)-i\phi^2(t)} e^{-i\phi^1(t_L)-i\phi^2(t_L)} \\ &= \oint \frac{dt}{2\pi i} \frac{\partial t_L}{\partial z} z(t) (t-t_L)^2 e^{-i\phi^1(t)-i\phi^2(t)-i\phi^1(t_L)-i\phi^2(t_L)}. \end{aligned} \quad (3.45)$$

The state is obtained by acting with this operator on the chiral vacuum o_n to obtain

$$|L\rangle_{J^-} = \oint \frac{dt}{2\pi i} \frac{\partial t_L}{\partial z} z(t) \frac{(t-t_L)^2}{t^{n-1} t_L^{n-1}} e^{-i\phi^1(t)-i\phi^2(t)-i\phi^1(t_L)-i\phi^2(t_L)+i\frac{n-1}{2}(\phi^1+\phi^2)} e^{i\frac{n-1}{2}(\bar{\phi}^1+\bar{\phi}^2)} |0\rangle. \quad (3.46)$$

Let us now compute the two point a two point function of the two-impurities states above. Using the map to the covering surface (3.14) the two point function can be evaluated to be proportional to

$${}_{J^-} \langle M|L \rangle_{J^-} \sim \frac{1}{n^2} \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} \frac{p^2 (1-p)^2 (1-q)^2 t^{5-2n} (t')^{2n-7}}{(1-\frac{t}{p'})^2 (1-p\frac{t}{p'})^2 (q-\frac{t}{p'})^2 (q-p\frac{t}{p'})^2}, \quad (3.47)$$

where $p = e^{\frac{2\pi i}{n}L}$, $q = e^{\frac{2\pi i}{n}M}$ as before. Performing explicitly the computation one obtains

$${}_{J^-} \langle M|L \rangle_{J^-} \sim \begin{cases} L \neq \pm M & \frac{4}{n} - \frac{1}{n} \frac{1}{\sin^2 \frac{\pi(L-M)}{n}} - \frac{1}{n} \frac{1}{\sin^2 \frac{\pi(L+M)}{n}} \\ L = \pm M & \frac{17-12n+4n^2}{3n} - \frac{5}{4n} \frac{1}{\sin^2 \frac{\pi L}{n}} - \frac{1}{4n} \frac{1}{\cos^2 \frac{\pi L}{n}} \\ L = \pm M = \pm \frac{n}{2} & \frac{16(2-3n+n^2)}{3n} \end{cases} \quad (3.48)$$

Note thus that our states are not orthogonal at tree level. In appendix B we comment on the diagonalization of these states.

Let us consider the large n limit. In this case if $L \neq \pm M$ the two point function scales as n^{-1} . However, the $L = \pm M$ case scales as n . Thus, naively in the large n limit the states with impurities are approximately orthogonal and the mixing matrix is proportional to the identity with coefficient scaling as n . However note that if $L - M \ll n$ for instance the term $\frac{1}{n} \frac{1}{\sin^2 \frac{\pi(L-M)}{n}}$ will scale as n and thus will not be subleading. Thus, even in the large n limit chains with close L and M mix.

3.3 Fractional moded J^- impurities

In the previous sections we defined a spin chain in ‘‘position space’’ using the functions $t_L(t)$, but we can also define states in ‘‘momentum space’’. One way to do so would be to Fourier

transform the states we defined above. However, in orbifold theories there is a natural definition using fractional modes of the fields. For concreteness, let us discuss here the fractional modes of the R current J^- . In the presence of twist n field we define [44]

$$J_{-m/n}^-(z) \equiv \oint \frac{dz}{2\pi i} \sum_{k=1}^n J_{(k)}^- e^{-\frac{2\pi i m(k-1)}{n}} z^{-\frac{m}{n}}. \quad (3.49)$$

This operator is gauge invariant and has dimension $\frac{m}{n}$. Note that here $\Delta - J \neq 1$. We can refer to the quantum number m as a ‘‘momentum’’ variable.

The fractional moded operators are lifted in a very simple manner to the covering surface

$$J_{-m/n}^- \rightarrow \oint \frac{dt}{2\pi i} z(t)^{-\frac{m}{n}} J^-(t). \quad (3.50)$$

Note that these operators on the covering surface become just the integer modes of J^- as $z(t) \sim t^n$ near $t = 0$. We can act with these operators on the chiral primary o_n to obtain a general state of the form

$$\prod_{k=1}^s J_{-m_k/n}^- o_n. \quad (3.51)$$

Note that if $\sum_{k=1}^s m_k = 0$ we have a state with charge shifted by s and unshifted dimension, implying $\Delta - J = s$.

Let us consider a state with two impurities

$$\begin{aligned} \hat{o}_k &\equiv J_{k/n}^- J_{-k/n}^- o_n = \\ &\oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \left(\frac{z(t)}{z(t')} \right)^{-\frac{k}{n}} \frac{(t-t')^2}{t^{n-1} t'^{n-1}} e^{-i\phi^1(t)-i\phi^2(t)-i\phi^1(t')-i\phi^2(t')+i\frac{n-1}{2}(\phi^1+\phi^2)} e^{i\frac{n-1}{2}(\bar{\phi}^1+\bar{\phi}^2)}. \end{aligned} \quad (3.52)$$

Note that the ‘‘momentum’’ k is bounded. The map near zero behaves as $z(t) \sim t^n$ and for $k \geq n-1$ the t' integration does not have a pole and thus vanishes. For $k \leq 1-n$ the t integration vanishes for the same reason. Of course we can take k to be non negative as $J_{k/n}^-$ commutes with $J_{-k/n}^-$ and thus the independent values of k are $0, \dots, n-2$.

Let us compute two point function of states with two impurities,

$$\begin{aligned} \langle \hat{o}_k | \hat{o}_m \rangle &\sim \frac{\oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \oint \frac{ds'}{2\pi i} \oint \frac{ds}{2\pi i} \left(\frac{t}{t'} \right)^{-k} \left(\frac{s}{s'} \right)^{-m} \frac{(t-t')^2}{t^{n-1} t'^{n-1}} \frac{(s-s')^2}{s^{1-n} s'^{1-n}}}{1} \\ &\quad \frac{1}{(t-s)^2 (t'-s')^2 (t-s')^2 (t'-s)^2}. \end{aligned} \quad (3.53)$$

We assume that the contours are ordered as $|t'| < |t| < |s'| < |s|$ and write

$$\langle \hat{o}_k | \hat{o}_m \rangle \sim \oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \oint \frac{ds'}{2\pi i} \oint \frac{ds}{2\pi i} \frac{t'^{k+1-n} t^{-k-n+3} (1-\frac{t'}{t})^2 s^{n-m-3} s'^{m+m-5} (1-\frac{s'}{s})^2}{(1-\frac{t}{s})^2 (1-\frac{t'}{s'})^2 (1-\frac{t}{s'})^2 (1-\frac{t'}{s})^2} \quad (3.54)$$

This can be evaluated to give

$$\langle \hat{o}_k | \hat{o}_m \rangle \sim \begin{cases} k = m & (n-2)^2 - (k-1)^2 + \delta_{k,0} (n-1)^2 \\ k > m & -2(n-k-1) \\ m > k & -2(n-m-1) \end{cases} \quad (3.55)$$

We see that these operators also mix at tree level. However, in the strict infinite n limit rescaling the operators with $1/n$ we get that this basis is orthogonal. To get an orthonormal basis we have to rescale with $\frac{1}{n} \frac{1}{\sqrt{1-\frac{k^2}{n^2}}}$. Thus defining $p = k/n$ we get an orthonormal set of states labeled by p such that $|p| - 1$ is finite.

The operators in “position space” we defined in previous section using $t_L(t)$ can be obtained from states built using fractional modes as follows. Let us consider the following operator

$$\begin{aligned} \hat{o}_L^{pos} &\equiv \sum_{m=0}^{\infty} e^{\frac{2\pi i m L}{n}} \hat{o}_m \rightarrow \\ &\oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} \frac{1}{1 - e^{\frac{2\pi i L}{n}} \left(\frac{z(t')}{z(t)}\right)^{\frac{1}{n}}} \frac{(t-t')^2}{t^{n-1} t'^{n-1}} e^{-i\phi^1(t) - i\phi^2(t) - i\phi^1(t') - i\phi^2(t') + i\frac{n-1}{2}(\phi^1 + \phi^2)} e^{i\frac{n-1}{2}(\bar{\phi}^1 + \bar{\phi}^2)}, \end{aligned} \quad (3.56)$$

where we assumed that $|z(t)| > |z(t')|$. We have interchanged the order of taking the contour integral and performing the infinite sum. These two limits do not commute. Note that if we were to truncate the sum above at finite m the pole in the integrand coming from summing up the geometric series would not have appeared. This is a sign of the fact that $J_{k/n}^-$ commutes with $J_{-k/n}^-$. However, upon changing the limits we develop a pole and the two currents effectively cease to commute. The commutator term which we develop is exactly the operator we defined in the previous section. We can deform the t integration to two contours, one around t' and the other around 0. Thus, computing it with residue theorem we get

$$\begin{aligned} o_L^{pos} &\equiv \hat{o}_L^{pos} + \sum_{m=-\infty}^{-1} e^{\frac{2\pi i m L}{n}} \hat{o}_m = \sum_{m=-\infty}^{\infty} e^{\frac{2\pi i m L}{n}} \hat{o}_m \rightarrow \\ &\oint \frac{dt'}{2\pi i} z(t_L) \frac{dt_L}{dz} \frac{(t_L - t')^2}{t_L^{n-1} t'^{n-1}} e^{-i\phi^1(t_L) - i\phi^2(t_L) - i\phi^1(t') - i\phi^2(t') + i\frac{n-1}{2}(\phi^1 + \phi^2)} e^{i\frac{n-1}{2}(\bar{\phi}^1 + \bar{\phi}^2)}, \end{aligned} \quad (3.57)$$

where t_L is defined by

$$z(t_L)^{1/n} = e^{\frac{2\pi i L}{n}} z(t)^{1/n}. \quad (3.58)$$

This is the operator we defined in the previous section.

3.4 ∂X impurities

We can add impurities using the modes of ∂X^i , $\partial X^{\dagger i}$ and the antiholomorphic counterparts. A chain with l impurities is defined as

$$\sum_{I=1}^n \oint \frac{dz}{2\pi i} \frac{1}{z} \prod_{j=1}^l \partial X_{I+L_j}^{i_j} o_n. \quad (3.59)$$

The bosons do not carry R -charge. However, they carry a charge under the outer automorphism $SU(2)$. As we will see in what follows the interactions are invariant under this $SU(2)$ and thus we can use it as a selection rule. The above state has $\Delta - J = l$. On the covering surface it becomes

$$\oint \frac{dt}{2\pi i} \frac{1}{z(t)} \prod_{k=2}^l \frac{\partial t_{L_j}}{\partial z} \prod_{j=1}^l \partial X^{i_j}(t_{L_j}) o_n. \quad (3.60)$$

Let us consider the two point function in the simplest case when all the bosons are of the same kind. This case is simple because there are no contractions between the ∂X s of the chain. The tree level two point function is

$$\begin{aligned} & \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} z(t')^{2l-1} \prod_{k=2}^l \frac{\partial t'_{M_j}}{\partial z} \frac{1}{z(t)} \prod_{k=2}^l \frac{\partial t_{L_j}}{\partial z} \sum_{\alpha \in S_l} \prod_{i=1}^l \frac{1}{(t_{L_i} - t'_{M_{\alpha(i)}})^2} = \\ & \frac{1}{n^{2l-2}} \prod_{k=2}^l q_l p_l \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} t'^{m(2l-1)-(n-1)(l-1)-2l} t^{-n-(n-1)(l-1)} \sum_{\alpha \in S_l} \prod_{i=1}^l \frac{1}{(p_i \frac{t}{t'} - q_{\alpha(i)})^2} \\ & \frac{1}{n^{2l-2}} \prod_{k=2}^l q_l p_l \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} t'^{ml-l-1} t^{-nl+l-1} \sum_{\alpha \in S_l} \prod_{i=1}^l \frac{1}{(p_i \frac{t}{t'} - q_{\alpha(i)})^2}, \end{aligned} \quad (3.61)$$

where S_l is the group of permutations of l objects. Taking a state with two impurities, *i.e.* $l = 2$, we get that the two point function is

$$\mathcal{I}_{LM} = \begin{cases} L \neq \pm M & -\frac{1}{n} \frac{1}{\sin^2 \frac{\pi(L-M)}{n}} - \frac{1}{n} \frac{1}{\sin^2 \frac{\pi(L+M)}{n}} \\ L = \pm M & \frac{4n^2-1}{3n} - \frac{1}{4n} \frac{1}{\sin^2 \frac{\pi L}{n}} - \frac{1}{4n} \frac{1}{\cos^2 \frac{\pi L}{n}} \\ L = \pm M = \pm \frac{n}{2} & \frac{4n^2+4}{3n} \end{cases} \quad (3.62)$$

Thus we see that this chain is not orthogonal at tree level, much as the chain built from R currents. The basic objects in the theory are in a sense the fermions and these objects are “composite”s as far as the quantum numbers go.

4. The spin chain at one loop

In this section we discuss the structure of the “one loop” computation of anomalous dimensions of states with impurities. First, we discuss the symmetry algebra of the theory and the structure of the interaction terms. Then, we discuss the map to the covering surface and the function $t_L(t)$ at one loop. Next, we perform the one loop computation of the vanishing anomalous dimension of the vacuum. Finally, we comment on the one loop computation of states with impurities.

4.1 The $\mathcal{N} = (4, 4)$ supersymmetry algebra and the interaction term

Let us first discuss the supersymmetry generators.⁷ The left-moving supercharges are given by

$$\begin{aligned} G^a &= \sqrt{2} \begin{bmatrix} i\psi_I^1 \\ -\psi_I^{2\dagger} \end{bmatrix} \partial X_I^{1\dagger} + \sqrt{2} \begin{bmatrix} i\psi_I^2 \\ \psi_I^{1\dagger} \end{bmatrix} \partial X_I^{2\dagger}, \\ \hat{G}^a &= \sqrt{2} \begin{bmatrix} i\psi_I^{1\dagger} \\ \psi_I^2 \end{bmatrix} \partial X_I^1 + \sqrt{2} \begin{bmatrix} i\psi_I^{2\dagger} \\ -\psi_I^1 \end{bmatrix} \partial X_I^2. \end{aligned} \quad (4.1)$$

In the above expressions a summation over the copy index I is implied. In the bosonized language this becomes

$$\begin{aligned} G^a &= \sqrt{2} \begin{bmatrix} ie^{i\phi_I^1} \\ -e^{-i\phi_I^2} \end{bmatrix} \partial X_I^{1\dagger} + \sqrt{2} \begin{bmatrix} ie^{i\phi_I^2} \\ e^{-i\phi_I^1} \end{bmatrix} \partial X_I^{2\dagger}, \\ \hat{G}^a &= \sqrt{2} \begin{bmatrix} ie^{-i\phi_I^1} \\ e^{i\phi_I^2} \end{bmatrix} \partial X_I^1 + \sqrt{2} \begin{bmatrix} ie^{-i\phi_I^2} \\ -e^{i\phi_I^1} \end{bmatrix} \partial X_I^2. \end{aligned} \quad (4.2)$$

The global symmetry of the theory is $SU(2)_R \times SU(2)_I$, where $SU(2)_I$ acts on the generators in the following way

$$\begin{aligned} [J_I^i, J_I^j] &= i\epsilon_{ijk} J_I^k, & \mathcal{G} &= (G^1, \hat{G}^2), & \mathcal{G}^\dagger &= \begin{pmatrix} \hat{G}^1 \\ G^2 \end{pmatrix} \\ [J_I^i, \mathcal{G}^a] &= \frac{1}{2} \mathcal{G}^b \sigma_{ba}^i, & [J_I^i, \mathcal{G}^a] &= -\frac{1}{2} \sigma_{ab}^i \mathcal{G}^{b\dagger}. \end{aligned} \quad (4.3)$$

From here we see that under the J_3 charge G^1 and G^2 have charge $+\frac{1}{2}$ while \hat{G}^1 and \hat{G}^2 have charge $-\frac{1}{2}$. The $SU(2)_R$ acts in the following way,

$$[J_R^i, J_R^j] = i\epsilon^{ijk} J_R^k, \quad [J_R^i, G^a] = -\frac{1}{2} (\sigma^i)^a_b G^b, \quad [J_R^i, \hat{G}_a] = -\frac{1}{2} \hat{G}_b (\sigma^i)^b_a. \quad (4.4)$$

⁷For more details see *e.g.* [32, 46].

From here we get that generators G^1 and \hat{G}^2 have charge $+\frac{1}{2}$ while G^2 and \hat{G}^1 have charge $-\frac{1}{2}$. Thus we can write the following

$$G^1 = G^{++}, \quad G^2 = G^{-+}, \quad \hat{G}^1 = G^{--}, \quad \hat{G}^2 = G^{+-}, \quad (4.5)$$

where the first sign is the $SU(2)_R$ charge and the second is the $SU(2)_I$ charge. We can repeat the same procedure for the anti-holomorphic fields and denote the antiholomorphic super-charges as \tilde{G}^{ab} , $a = \pm$ and $b = \pm$.

We are ready now to discuss the deformation away from the orbifold point. In a unitary $N = 4$ theory, all marginal deformations that preserve the $N = 4$ symmetry are obtained from chiral fields with dimension $\Delta = 1/2$ [64], such as our twist-two fields O_2 . The explicit form of the deformation we will use is

$$I_{int} = \int d^2u \mathcal{O}_2 + c.c., \quad (4.6)$$

$$\mathcal{O}_2 = \lambda \left[G_{-1/2}^{--} \tilde{G}_{-1/2}^{-+} - G_{-1/2}^{-+} \tilde{G}_{-1/2}^{--} \right] O_2. \quad (4.7)$$

This deformation has charge zero under both $SU(2)_I$ and $SU(2)_R$. Each one of the two terms above can be turned on with separate coupling with the price of breaking the global $SU(2)_I$ symmetry.

We will perform one loop computations on the covering surface using stress-energy tensor method. Let us thus lift the deformation explicitly to the covering surface. The explicit computation of the first term in (4.7) reads

$$\begin{aligned} & G_{-1/2}^{--} \tilde{G}_{-1/2}^{-+} O_2(u, \bar{u}) = \quad (4.8) \\ & \oint \frac{dz}{2\pi i} \oint \frac{d\bar{z}'}{2\pi i} \left(\sqrt{2} i e^{-i\phi_I^1} \partial X_I^1 + \sqrt{2} i e^{-i\phi_I^2} \partial X_I^2 \right) (z) \left(-\sqrt{2} e^{-i\bar{\phi}_J^1} \bar{\partial} X_J^{1\dagger} + \sqrt{2} e^{-i\bar{\phi}_J^2} \bar{\partial} X_J^{2\dagger} \right) (\bar{z}') \times \\ & \times \frac{1}{\sqrt{2N!(N-2)!}} \sum_{h \in S_N} e^{i\frac{1}{4} \sum_{I'=h-1, h-2} (\phi_{I'}^1 + \phi_{I'}^2 + \bar{\phi}_{I'}^1 + \bar{\phi}_{I'}^2)} \sigma_{h^{-1}(12)h}(u, \bar{u}) \end{aligned}$$

On the covering surface up to overall constants the integrand of the above expression becomes

$$\begin{aligned} & \frac{2i \left(\frac{\partial z}{\partial t} \right)^{-\frac{3}{2}} \left(\frac{\partial \bar{z}'}{\partial \bar{t}'} \right)^{-\frac{3}{2}}}{(x-t)^{\frac{1}{2}} (\bar{x}-\bar{t}')^{\frac{1}{2}}} \left[\begin{aligned} & e^{-i\phi^2(t)-i\bar{\phi}^1(\bar{t}')+\frac{i}{2}(\phi^1(x)+\phi^2(x)+\bar{\phi}^1(\bar{x})+\bar{\phi}^2(\bar{x}))} : \partial X^2 \bar{\partial} X^{2\dagger} - \\ & - : e^{-i\phi^1(t)-i\bar{\phi}^2(\bar{t}')+\frac{i}{2}(\phi^1(x)+\phi^2(x)+\bar{\phi}^1(\bar{x})+\bar{\phi}^2(\bar{x}))} : \partial X^1 \bar{\partial} X^{1\dagger} + \\ & + : e^{-i\phi^1(t)-i\bar{\phi}^1(\bar{t}')+\frac{i}{2}(\phi^1(x)+\phi^2(x)+\bar{\phi}^1(\bar{x})+\bar{\phi}^2(\bar{x}))} : \partial X^1 \bar{\partial} X^{2\dagger} - \\ & - : e^{-i\phi^2(t)-i\bar{\phi}^2(\bar{t}')+\frac{i}{2}(\phi^1(x)+\phi^2(x)+\bar{\phi}^1(\bar{x})+\bar{\phi}^2(\bar{x}))} : \partial X^2 \bar{\partial} X^{1\dagger} \end{aligned} \right] \quad (4.9) \end{aligned}$$

where we assumed that the insertion at u on the base sphere is mapped to $t = x$ on the covering. We also dropped overall x dependent factors coming from conformal transformations

of the operators as these will cancel out in the stress-energy method of computing correlators. We will drop these kind of terms everywhere in what follows. Near twist two field the map has the property

$$z(t) \sim v(x) + \frac{1}{2} a(x) (t - x)^2, \quad (4.10)$$

and thus the contour integrals in (4.8) can be lifted to the covering surface as

$$\oint \frac{dz}{2\pi i} \oint \frac{d\bar{z}'}{2\pi i} \frac{1}{(x-t)^{\frac{1}{2}}(\bar{x}-\bar{t}')^{\frac{1}{2}}} \left(\frac{\partial z}{\partial t}\right)^{-\frac{3}{2}} \left(\frac{\partial \bar{z}'}{\partial \bar{t}'}\right)^{-\frac{3}{2}} \rightarrow \oint \frac{dt}{2\pi i} \oint \frac{d\bar{t}'}{2\pi i} \frac{1}{(x-t)(\bar{x}-\bar{t}')}. \quad (4.11)$$

Using Cauchy theorem we obtain

$$\begin{aligned} G_{-1/2}^{--} \tilde{G}_{-1/2}^{-+} O_2 \sim & \quad (4.12) \\ 2i \left[e^{\frac{i}{2}(\phi^1 - \phi^2 - \bar{\phi}^1 + \bar{\phi}^2)} \partial X^2 \bar{\partial} X^{2\dagger} - e^{\frac{i}{2}(-\phi^1 + \phi^2 + \bar{\phi}^1 - \bar{\phi}^2)} \partial X^1 \bar{\partial} X^{1\dagger} + \right. \\ & \left. + e^{\frac{i}{2}(-\phi^1 + \phi^2 - \bar{\phi}^1 + \bar{\phi}^2)} \partial X^1 \bar{\partial} X^{2\dagger} - e^{\frac{i}{2}(\phi^1 - \phi^2 + \bar{\phi}^1 - \bar{\phi}^2)} \partial X^2 \bar{\partial} X^{1\dagger} \right] (x, \bar{x}). \end{aligned}$$

Performing the same computation for all the terms in (4.6) we finally get

$$\begin{aligned} \mathcal{O}^\dagger + \mathcal{O} \sim & -4i \text{Re}[\lambda] \left(e^{-\frac{i}{2}(\phi^1 - \phi^2 - \bar{\phi}^1 + \bar{\phi}^2)} \partial X^{2\dagger} \bar{\partial} X^2 - e^{-\frac{i}{2}(\bar{\phi}^1 - \bar{\phi}^2 - \phi^1 + \phi^2)} \bar{\partial} X^{2\dagger} \partial X^2 - \right. \\ & \left. - e^{-\frac{i}{2}(-\phi^1 + \phi^2 + \bar{\phi}^1 - \bar{\phi}^2)} \partial X^{1\dagger} \bar{\partial} X^1 + e^{-\frac{i}{2}(-\bar{\phi}^1 + \bar{\phi}^2 + \phi^1 - \phi^2)} \bar{\partial} X^{1\dagger} \partial X^1 \right) + \\ & + 4\text{Im}[\lambda] \left(e^{-\frac{i}{2}(-\phi^1 + \phi^2 - \bar{\phi}^1 + \bar{\phi}^2)} \partial X^{1\dagger} \bar{\partial} X^2 + e^{-\frac{i}{2}(\bar{\phi}^1 - \bar{\phi}^2 + \phi^1 - \phi^2)} \bar{\partial} X^{2\dagger} \partial X^1 - \right. \\ & \left. - e^{-\frac{i}{2}(\phi^1 - \phi^2 + \bar{\phi}^1 - \bar{\phi}^2)} \partial X^{2\dagger} \bar{\partial} X^1 - e^{-\frac{i}{2}(-\bar{\phi}^1 + \bar{\phi}^2 - \phi^1 + \phi^2)} \bar{\partial} X^{1\dagger} \partial X^2 \right). \end{aligned} \quad (4.13)$$

By computing *OPEs* of the above interactions one can verify that there are no dimension (1, 1) contact terms. The conformal dimension on the covering surface of the above operator is $(\frac{5}{4}, \frac{5}{4})$ and one obtains dimension (1, 1) on the base by using the fact that for the above operator $\Delta_{base} = \Delta_n + \frac{\Delta_{cover}}{n}$, where n is the size of the twist field and Δ_n is the dimension of the bare twist field.⁸

⁸Note that the interaction has a very simple ‘‘local’’ form on the covering surface. However, this does not imply that on the base surface the interaction has a simple form.

4.2 The map to the covering surface in presence of the interactions

To determine the two point functions of states with impurities in presence of the interactions we have to lift the computation to the covering surface and specify the position of the impurities using the function $t_L(t)$. After turning on the interaction term (4.6) in principle, a dressed n -cycle twist field mixes with dressed $n - 1$ cycle twist field already in order λ^1 . However, the mixing of two dressed n cycle fields occurs only at even orders in $|\lambda|$. We will restrict our explicit discussion to the latter case, *i.e.* we will study the mixing matrix to the first non trivial order in λ between chains of same length. The reason for this is that we will discuss only computations with fermionic impurities, and the former mixing is absent for these as the X correlator becomes a one point function on the covering surface and thus vanishes. Using the same techniques as will be used below one can also perform calculations of two point functions of chains of different lengths. In what follows we first discuss in detail the map to the covering surface in presence of two twist-two interactions, the first non-trivial order contributing to mixing of two chains of same length. Then we briefly comment on generalizations to maps with more interaction terms. Finally, in the next section we discuss the properties of $t_L(t)$.

Let us specify the map to the covering surface of two twist n fields in presence of two twist two interactions. The map we construct has twist n fields at 0 and ∞ on the base and the covering. One twist two field is at $z = 1$ on the base and $t = 1$ on the covering and the other one is at $z = u$ on the base and $t = x$ on the cover. The relevant map is given by [28]

$$z(t; x) = \left(\frac{f_2(1)}{f_1(1)} \right) \frac{f_1(t)}{f_2(t)}, \quad (4.14)$$

where

$$f_1(t) = t^n \left(1 - \frac{n(1+x) \mp \sqrt{n^2(1-x)^2 + 4x}}{2x(n+1)} t \right), \quad (4.15)$$

$$f_2(t) = 1 - \frac{n(1+x) \pm \sqrt{n^2(1-x)^2 + 4x}}{2x(n-1)} t.$$

We have two choices of the sign before the square root. The map with the $-$ in f_1 and $+$ in f_2 will be called map a , and the map with the $+$ in f_1 and $-$ in f_2 will be called map b in what follows. We will also write (4.14) as

$$z(t; x) = C t^n \frac{t - t_0}{t - t_\infty}. \quad (4.16)$$

For map a we have

$$t_0 = \frac{2(n+1)x}{n(1+x) - \sqrt{n^2(x-1)^2 + 4x}} \sim x \left(1 + \frac{1}{n} \right) + O(1/n^2), \quad (4.17)$$

$$t_\infty = \frac{2(n-1)x}{n(1+x) + \sqrt{n^2(x-1)^2 + 4x}} \sim 1 - \frac{1}{n} + O(1/n^2),$$

and for map b

$$\begin{aligned} t_0 &= \frac{2(n+1)x}{n(1+x) + \sqrt{n^2(x-1)^2 + 4x}} \sim 1 + \frac{1}{n} + O(1/n^2), \\ t_\infty &= \frac{2(n-1)x}{n(1+x) - \sqrt{n^2(x-1)^2 + 4x}} \sim x \left(1 - \frac{1}{n}\right) + O(1/n^2). \end{aligned} \quad (4.18)$$

Additional useful identities are

$$\frac{1}{1-t_\infty} - \frac{1}{1-t_0} = n, \quad t_0 t_\infty = x. \quad (4.19)$$

The parameter x is fixed by demanding that $z(x; x) = u$, *i.e.*

$$u = \frac{1}{2}x^{n-1} \left(2x + n^2(x-1)^2 - n(x-1)\sqrt{n^2(1-x)^2 + 4x} \right), \quad (4.20)$$

which can be more conveniently written as

$$u^2 + x^{2n} - ux^{n-1} [n^2(x-1)^2 + 2x] = 0. \quad (4.21)$$

Given u we have $2n$ solutions to this equation. Using the diagrammatic language of [28] these can be represented as $2n$ different diagrams. The diagrams split into two groups, with $n+1$ and $n-1$ diagrams in each group. The groups differ by the behavior of $x(u)$ in the OPE limits $u=0$ and $u=\infty$. For the first group we have

$$u \rightarrow \infty \quad : \quad u \sim n^2 x^{n+1}, \quad u \rightarrow 0 \quad : \quad u \sim \frac{1}{n^2} x^{n+1}, \quad (4.22)$$

and for the second group

$$u \rightarrow \infty \quad : \quad u \sim \frac{1}{n^2} x^{n-1}, \quad u \rightarrow 0 \quad : \quad u \sim n^2 x^{n-1}. \quad (4.23)$$

Moreover, in the OPE limit $u=1$ we again have different behaviors for the two groups of solutions,

$$u-1 \sim 2n(x-1), \quad u-1 \sim \frac{n(n^2-1)}{24}(x-1)^3, \quad (4.24)$$

for the first and the second groups respectively. The diagrams of the first group appear in figure 4 and the diagrams of the second group appear in figure 5. There are three diagrams with a non trivial OPE limit near $u=1$ for the second group and one diagram with non trivial behavior for the first one, as is clear from (4.24). In $u \rightarrow 1$ limit the non trivial OPE manifests itself in the diagrams having a shared color between the two interactions. The space of possible values of parameter x , the “moduli space” space of maps [28], consists of two copies of a sphere glued along a branch cut between

$$x_+ = 1 - \frac{2}{n^2} \left(1 - \sqrt{1-n^2} \right), \quad x_- = 1 - \frac{2}{n^2} \left(1 + \sqrt{1-n^2} \right), \quad (4.25)$$




Figure 4: The diagrams of map b in the vicinity of $x = 0$. The bottom diagram is the generic one, there are $n - 2$ of these.

as can be seen from (4.20). Note that in the large n limit x_+ and x_- are both very near $x = 1$, and effectively the two spheres pinch away.⁹ Note also that one moves between maps a and b by crossing with x the branch cut between x_+ and x_- and thus map a and map b correspond to the two copies of the moduli space. In particular map a roughly corresponds to the group of $n - 1$ diagrams and map b to the group of $n + 1$ diagrams. One way to establish this fact is to compute the four point function, *i.e.* two chiral fields with two interactions, for the two maps and look at the behavior in the OPE limits. The details of this computation can be found in appendix E. The schematic picture of the moduli space is depicted in figure 6.

⁹Note that $|x_+| = |x_-| = 1$.




Figure 5: The diagrams of map a in the vicinity of $x = 0$. The bottom diagram is the generic one, there are $n - 2$ of these.

It is possible to choose another parametrization of the moduli spaces which combine the two copies of the moduli space [53]. We present this parametrization in appendix D. However, the parametrization with the two copies appearing in this section will be used in what follows. Additional details on the map needed in the following sections are collected in appendix C.

Finally, let us comment on higher loop maps. Adding more interactions it becomes harder to determine the exact map to the covering surface. However, in the leading $1/n$ orders the map is essentially simple. Let us take $2k$ interaction terms, *i.e.* $2k$ twist two fields. The generic map takes the following form

$$z(t) = C t^n \frac{\prod_{j=1}^k (t - t_0^j)}{\prod_{j=1}^k (t - t_\infty^j)}. \quad (4.26)$$

Differentiating this map we obtain that the twist fields are located at the solutions of the following equation

$$\prod_{j=1}^k (t - t_0^j) \prod_{j=1}^k (t - t_\infty^j) + \frac{t}{n} \sum_{i=0}^k \left[\prod_{i \neq j} (t - t_0^j) \prod_j (t - t_\infty^j) - \prod_{i \neq j} (t - t_\infty^j) \prod_j (t - t_0^j) \right] = 0. \quad (4.27)$$




Figure 6: The schematic structure of the moduli space for $x < 1$. On the left we have map a and on the right map b . In this figure $n = 4$. Each region of the moduli space corresponds to a diagram. The dashed blue line is the branch cut connecting the two copies of the moduli space.

To the leading $1/n$ order the second term is vanishing and we get that all the maps are given by different assignments of the positions of twist fields x_i ($i = 1 \dots 2k$ and $x_0 = 1$) to t_∞^i and t_0^i . For example, in the two interaction case we discussed above in detail we had two possibilities, $(t_0 = 1, t_\infty = x)$ and $(t_0 = x, t_\infty = 1)$. The subleading behavior is also easy to obtain.

4.3 Evaluating $t_L(t)$

Let us compute the function $t_L(t)$ for the covering map of the previous section. The inverse (to (4.16)) map near $t = 0$ is given by the following expansion

$$t = \sum_{k=1}^{\infty} b_k z^{\frac{k}{n}}. \quad (4.28)$$

Then by definition $t_L(t)$ is given by

$$t_L = \sum_{k=1}^{\infty} b_k z^{\frac{k}{n}} p^k, \quad p = e^{\frac{2\pi i}{n} L}. \quad (4.29)$$

We want to understand the properties of this function. First, again by definition

$$z(t) = z(t_L(t)), \quad (4.30)$$

i.e. the points $t_L(t)$ and t correspond to the same position z on the base sphere but represent different colors. This implies the following equality,

$$t^n \frac{t - t_0}{t - t_\infty} = t_L^n \frac{t_L - t_0}{t_L - t_\infty}, \quad (4.31)$$

which has $n + 1$ solutions, and thus there are only $n + 1$ functions satisfying (4.30). One solution is trivial $t_L(t) = t$, but the others are not. Out of n non-trivial solutions $n - 1$ correspond to the different choices of p , *i.e.* $p = e^{\frac{2\pi i}{n} j}$ with $j = 1 \dots n - 1$. We are interested in the solution with $j = L$. The one extra solution satisfies $t_L(0) = t_0$.

Using the above one can derive the following useful identities,

$$(t - 1)(t - x) = (t - t_0)(t - t_\infty) + \frac{t}{n}(t_0 - t_\infty), \quad (4.32)$$

$$\frac{\partial t_L}{\partial t} = \frac{t_L}{t} \frac{t_L - t_0}{t_L - x} \frac{t_L - t_\infty}{t_L - 1} \frac{t - x}{t - t_\infty} \frac{t - 1}{t - t_0}.$$

The function $t_L(t)$ is a solution to a polynomial equation (4.31) and thus it is clear that $t_L(t)$ might have branch cuts and indeed it does. By definition t_0 is an image of $z = 0$, and thus for the non trivial solutions of interest to us we have $t_L(t_0) = 0$. This implies, using (4.31), the following

$$t_L \sim \left[\frac{t_\infty}{t_0} \frac{t_0^n}{t_0 - t_\infty} \right]^{1/n} (t - t_0)^{\frac{1}{n}}. \quad (4.33)$$

In analogous way $t_L(t_\infty) = \infty$, and we have

$$t_L \sim [t_\infty^n (t_\infty - t_0)]^{1/n} \frac{1}{(t - t_\infty)^{\frac{1}{n}}}. \quad (4.34)$$

From here we see that $t = t_\infty$ and $t = t_0$ are branch points of order n and are connected by a branch cut. This means that the n non-trivial solution of (4.31) are different branches of the same function.

Looking for zeros of $\frac{dt}{dt_L}$ we find that there are additional branch points of order 2 at t s satisfying

$$t_L(\hat{t}_1) = 1, \quad t_L(\hat{t}_x) = x, \quad (4.35)$$

with $\hat{t}_1 \neq 1$ and $\hat{t}_x \neq x$. There are $n - 1$ solutions of each type. These solutions are distributed between the different Riemann sheets. The exact way one distributes these branch point among the sheets depends, by definition, on the choice of the branch cuts. One can make the following choice (see figure 7) . On each sheet with $t_L(0) = 0$ there are exactly two solutions for \hat{t}_1 and two solutions for \hat{t}_x , with two branch cuts connecting \hat{t}_1 and \hat{t}_x . On a single sheet with $t_L(0) = t_0$ we have a single pair of solutions to (4.35). Essentially here $\{\hat{t}_1 = 1, \hat{t}_x = x\}$ and thus we do not have an additional cut.

We can think of (4.31) as an equation defining a Riemann surface *i.e.* a map between a sphere and another surface. The other surface has two connected components. The first one is a sphere and corresponds to $t_L(t) = t$. The second one corresponds to a genus $n - 1$ surface. The genus can be calculated through Riemann-Hurwitz formula by noting that, as

shown above, we have two ramification points of order n , $2(n-1)$ ramification points of order 2 and the number of sheets is n . The freedom of distributing the ramification points among the sheets translates to the freedom of dividing the Riemann surface into Riemann sheets.




Figure 7: A schematic example of a consistent division of the Riemann surface to sheets. The different sheets represented by horizontal lines and the cuts by the vertical line. The special sheet is the bottom one.

Let us note the following relations

$$\begin{aligned} p z^{1/n} &= t_L e^{\frac{1}{n} \ln Q(t_L)}, \\ p z^{1/n} &= p t e^{\frac{1}{n} \ln Q(t)}, \end{aligned} \quad (4.36)$$

where we define

$$z(t) = C t^n \frac{t - t_0}{t - t_\infty} \equiv t^n Q(t). \quad (4.37)$$

Using this one can write

$$t_L = p t e^{\frac{1}{n} \ln \frac{Q(t)}{Q(t_L)}}. \quad (4.38)$$

The expression (4.38) can be consistently expanded in $1/n$ away from the branch cuts, say around $t = 0$. To leading orders in $1/n$ gives

$$t_L = p t \left(1 + \frac{1}{n} \ln \frac{Q(t)}{Q(pt)} + \frac{2\pi i}{n} l \dots \right). \quad (4.39)$$

The term proportional to l takes into account possible crossing of a branch cut of the \ln , and in vicinity of $t = 0$ by definition $l = 0$. For our specific map $\ln \frac{Q(t)}{Q(pt)}$ in the large n limit evaluates to

$$\ln \frac{Q(t)}{Q(pt)} = (-)^{\sigma+1} \ln \left[\frac{(t-x)(pt-1)}{(pt-x)(t-1)} \right], \quad (4.40)$$

where $\sigma = 1$ for map a and $\sigma = 0$ for map b . Note that this result makes sense if t is away from x and 1, thus we have to be either sufficiently close to 0 or ∞ .¹⁰

Let us compute the points \hat{t}_x and \hat{t}_1 in the limit $1 \ll L \ll n$. Near $t_L = 1$ we define $t_L = 1 + \epsilon$ and $t = \chi$. From here we obtain that

$$1 + \epsilon = \left(1 + \frac{2\pi i}{n}L\right)\chi e^{\frac{1}{n} \ln \frac{\chi - t_0}{\chi - t_\infty} \frac{1 + \epsilon - t_\infty}{1 + \epsilon - t_0}}. \quad (4.41)$$

Remembering the behavior of $t_{0,\infty}$ for the two maps and assuming that $\chi \sim 1 + \delta$ we deduce

$$\begin{aligned} (a) & : \quad \epsilon n - \ln(1 + \epsilon n) = 2\pi i L + \delta n - \ln(1 + \delta n), \\ (b) & : \quad \epsilon n + \ln(1 - \epsilon n) = 2\pi i L + \delta n + \ln(1 - \delta n). \end{aligned} \quad (4.42)$$

The points with $\epsilon = 0$ are \hat{t}_1 by definition. We get two such points on every sheet and the solutions are

$$\begin{aligned} (a) & : \quad \hat{t}_1 = 1 - \frac{2\pi i}{n}L + \frac{1}{n} \ln(-2\pi i L) \\ & \quad \hat{t}'_1 = 1 - \frac{2\pi i}{n}(L + 1) + \frac{1}{n} \ln(-2\pi i L) \\ (b) & : \quad \hat{t}_1 = 1 - \frac{2\pi i}{n}L - \frac{1}{n} \ln(2\pi i L) \\ & \quad \hat{t}'_1 = 1 - \frac{2\pi i}{n}(L - 1) - \frac{1}{n} \ln(2\pi i L), \end{aligned} \quad (4.43)$$

where the second solution comes from encircling once the branch point of the log on the *l.h.s* of (4.42). The branch point is at $t_L = t_\infty$ for map a and at $t_L = t_0$ for map b . Note that on generic sheets there are no points such that $t_L(t) = t_{0,\infty}$ and these points are “swallowed” by the branch cuts.

Near $t_L = x$ we define $t_L = x(1 + \epsilon)$ and $t = \chi$. We obtain that

$$x(1 + \epsilon) = \left(1 + \frac{2\pi i}{n}L\right)\chi e^{\frac{1}{n} \ln \frac{\chi - t_0}{\chi - t_\infty} \frac{x(1 + \epsilon) - t_\infty}{x(1 + \epsilon) - t_0}}. \quad (4.44)$$

Assuming that $\chi \sim x(1 + \delta)$ one deduces

$$\begin{aligned} (a) & : \quad \epsilon n + \ln(1 - \epsilon n) = 2\pi i L + \delta n + \ln(1 - \delta n), \\ (b) & : \quad \epsilon n - \ln(1 + \epsilon n) = 2\pi i L + \delta n - \ln(1 + \delta n). \end{aligned} \quad (4.45)$$

The points with $\epsilon = 0$ are \hat{t}_x by definition. We get two such points on every sheet and the

¹⁰In large n limit we have $t_0 \sim x^\sigma$ and $t_\infty \sim x^{1-\sigma}$.

solutions are

$$\begin{aligned}
(a) \quad & \hat{t}_x = x \left(1 - \frac{2\pi i}{n} L - \frac{1}{n} \ln(2\pi i L) \right) \\
& \hat{t}'_x = x \left(1 - \frac{2\pi i}{n} (L + 1) - \frac{1}{n} \ln(2\pi i L) \right) \\
(b) \quad & \hat{t}_x = x \left(1 - \frac{2\pi i}{n} L + \frac{1}{n} \ln(-2\pi i L) \right) \\
& \hat{t}'_x = x \left(1 - \frac{2\pi i}{n} (L - 1) + \frac{1}{n} \ln(-2\pi i L) \right),
\end{aligned} \tag{4.46}$$

Note also that a loop around the two branch cuts between the images of $t_L = 1, x$ under $t_L(t)$




Figure 8: A contour around the two order two cuts on the t sheet transforms to a contour around the order n cut on the t_L sheet. The two green segments on the right are on different sheets and thus the integral over the green loop is in principal not vanishing. This illustration is for map a , and for map b t_0 has to be interchanged with t_∞ .

maps to a loop around the cut between t_∞ and t_0 . This fact is illustrated in figure 8. Thus, a contour integral of a function of t and t_L around the cut between t_∞ and t_0 , integrated over t , can be brought to an integral over t_L and around the two additional cuts. Equivalently, it is equal to an integral around the two additional cuts with L traded for $-L$. This fact will play a role in what follows.

4.4 Non-renormalization of the chiral vacuum at one loop

As a first step toward discussing the one loop structure of the two point functions of states with impurities we will compute first correction to the two point functions of operators without impurities in the deformed symmetric product CFT. These two point functions are protected [52] and thus the corrections will vanish.

Let us discuss the general structure of the one loop computation. The first correction to the two-point function is given by

$$\int d^2z_2 d^2z_3 \langle \mathcal{O}_1(z_1) \mathcal{O}_2(z_2) \mathcal{O}_3(z_3) \mathcal{O}_4(z_4) \rangle, \tag{4.47}$$

where \mathcal{O}_2 and \mathcal{O}_3 are the interaction vertices. Using $\Delta_1 = \Delta_4 = \Delta$ and $\Delta_2 = \Delta_3 = 1$, global conformal invariance fixes the form

$$\langle \mathcal{O}_1(z_1) \mathcal{O}_2(z_2) \mathcal{O}_3(z_3) \mathcal{O}_4(z_4) \rangle = \frac{G(u, \bar{u})}{|z_{24}|^4 |z_{13}|^4 |z_{14}|^{4\Delta-4}}, \quad (4.48)$$

where

$$u = \frac{z_{12} z_{34}}{z_{13} z_{24}}. \quad (4.49)$$

We can change now the integration variables in (4.47) to u, z_3 , and using

$$\left| \frac{\partial(u, z_3)}{\partial(z_2, z_3)} \right| = \left| \frac{z_{34} z_{14}}{z_{13} z_{24}^2} \right|, \quad (4.50)$$

eq. (4.47) becomes

$$\frac{1}{|z_{14}|^{4\Delta}} \int d^2 z_3 \frac{|z_{14}|^2}{|z_{31}|^2 |z_{34}|^2} \int d^2 u G(u, \bar{u}). \quad (4.51)$$

The first integral is divergent and must be regulated as

$$\int d^2 z_3 \frac{|z_{14}|^2}{|z_{31}|^2 |z_{34}|^2} = \int \frac{d^2 w}{|w|^2 |w+1|^2} = 2\pi \log \Lambda, \quad (4.52)$$

where $w = z_{31}/z_{14}$, and $\frac{1}{\Lambda}$ is a cutoff in $|w|$. So the expression (4.47) is finally

$$2\pi \frac{\log \Lambda}{|z_{14}|^{4\Delta}} \int d^2 u G(u, \bar{u}), \quad (4.53)$$

and we expect the integral over u, \bar{u} to give a finite contribution.

We will now explicitly compute $G(u, \bar{u})$. We do the computation using stress-energy tensor technique. On the base sphere the two chiral operators are at $z = 0, \infty$ and the two interactions at $z = 1, u$. On the covering sphere the two chiral operators are at $t = 0$ and $t = \infty$ and the two interactions are at $t = x$ and $t = 1$. The stress-energy tensor is given by

$$T(z) = - \sum_{i=1}^2 \left[\partial X_I^i(z) \partial X_I^{i\dagger}(z) + \frac{1}{2} \partial \phi_I^i(z) \partial \phi_I^i(z) \right] \quad (4.54)$$

$$= - \lim_{w \rightarrow z} \sum_{i=1}^2 \left[\partial X_I^i(z) \partial X_I^{i\dagger}(w) + \frac{1}{2} \partial \phi_I^i(z) \partial \phi_I^i(w) + \frac{6N}{(z-w)^2} \right]. \quad (4.55)$$

The bosonic correlator is given by

$$\begin{aligned} I_b &= \sum_{i=1}^2 \frac{\langle \partial X_I^i(z) \partial X_I^{i\dagger}(w) o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j}{\langle o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j} = \\ &= -4 \frac{t'_{j,I}(z) t'_{j,I}(w)}{(t_{j,I}(z) - t_{j,I}(w))^2} - 2 (t'_{j,I}(z))^2 \frac{(x-t_1)^2}{(t-1)^2 (t-x)^2}. \end{aligned} \quad (4.56)$$

The fermionic contribution is

$$\begin{aligned}
I_f &= \sum_{i=1}^2 \frac{\langle \partial \phi_I^i(z) \partial \phi_I^i(w) o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j}{\langle o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j} = \\
&= -2 \frac{t'_{j,I}(z) t'_{j,I}(w)}{(t_{j,I}(z) - t_{j,I}(w))^2} - \frac{1}{2} (t'_{j,I}(z))^2 \left[\left(\frac{1}{t-1} - \frac{1}{t-x} \right)^2 + \frac{(n-1)^2}{t^2} \right].
\end{aligned} \tag{4.57}$$

Then we define

$$g_j(z, u) = \frac{\langle T(z) o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j}{\langle o_n(\infty) \mathcal{I}(1) \mathcal{I}(u) o_n(0) \rangle_j}, \tag{4.58}$$

and obtain

$$\begin{aligned}
g_j(z, u) &= \frac{6}{12} \sum_{I=1}^{\tilde{n}} \{t_{j,I}, z\} + \sum_{I=1}^{\tilde{n}} \frac{(t'_{j,I}(z))^2}{2} \left(2 \frac{(x-1)^2}{(t-1)^2(t-x)^2} + \right. \\
&\quad \left. + \frac{1}{2} \left[\left(\frac{1}{t-1} - \frac{1}{t-x} \right)^2 + \frac{(n-1)^2}{t^2} \right] \right),
\end{aligned} \tag{4.59}$$

where $\{t, z\}$ is the Schwartz derivative,

$$\{t, z\} = \frac{t'''}{t'} - \frac{3}{2} \left(\frac{t''}{t'} \right)^2 = \left(\frac{t''}{t'} \right)' - \frac{1}{2} \left(\frac{t''}{t'} \right)^2, \tag{4.60}$$

The number \tilde{n} is the number of active colors in the vicinity of the field at $z = u$, *i.e.* in our case $\tilde{n} = 2$. Finally, remembering that

$$\partial_u \ln G_j(u) = \{g_j(z, u)\}_{\frac{1}{z-u}}, \tag{4.61}$$

we obtain the differential equation for $G(u)$

$$\begin{aligned}
\partial_u \ln G(x(u)) &= \left\{ \left[\left(\frac{t''}{t'} \right)' - \frac{1}{2} \left(\frac{t''}{t'} \right)^2 \right] + (t')^2 \left(2 \frac{(x-1)^2}{(t-1)^2(t-x)^2} + \right. \right. \\
&\quad \left. \left. + \frac{1}{2} \left[\left(\frac{1}{t-1} - \frac{1}{t-x} \right)^2 + \frac{(n-1)^2}{t^2} \right] \right) \right\}_{\frac{1}{z-u}}.
\end{aligned} \tag{4.62}$$

Note also that in this case we can write

$$v'(x) \partial_u \ln G(x(u)) = \partial_x \ln G(x(u)), \tag{4.63}$$

as the only dependence on u on the right hand side of the differential equation is through $x(u)$. Using the explicit map (see appendix C for details) this equation can be integrated to obtain

$$\begin{aligned}
G_0^a(x) &= C x^{2-n} (x-1)^{-4} \left(1 + x + \sqrt{n^2(x-1)^2 + 4x} \right) \\
&\quad \left(n^2(x-1) - 2x - n\sqrt{n^2(x-1)^2 + 4x} \right)^{-\frac{1}{2}} \left(n^2(x-1) + 2 + n\sqrt{n^2(x-1)^2 + 4x} \right)^{\frac{3}{2}},
\end{aligned} \tag{4.64}$$

$$G_0^b(x) = 4(1-n^2)^2 C x^{1-n} (x-1)^{-2} \left(1+x+\sqrt{n^2(x-1)^2+4x}\right)^{-1} \quad (4.65)$$

$$\left(n^2(x-1)-2x-n\sqrt{n^2(x-1)^2+4x}\right)^{\frac{1}{2}} \left(n^2(x-1)+2+n\sqrt{n^2(x-1)^2+4x}\right)^{-\frac{3}{2}},$$

where C is an overall constant not fixed by our method of computation. Note that the second expression is equal to the first one after we change the sign in front of the square roots. Thus, we can only take the first expression and remember that x takes values in a double cover of a sphere. The OPE behavior on the two covers is different. Note also the only singularities of the four point function occur when $x = 0, 1$ or $x = \infty$.

A simple check of this equation is to take $x \rightarrow 1$ in $G_0^b(x)$. In this limit $u \rightarrow 1$ and corresponds to an OPE limit of the two twist-two interactions sharing both colors and thus annihilating each other, see equation (4.24). The four point function scales as $G(u) \sim \frac{1}{(1-u)^2}$ as expected as the interactions have dimension one.

We can easily integrate (4.64) over the moduli space of maps

$$\begin{aligned} \sum_{j=1}^{2n} \int d^2u G(x_j(u), \bar{x}_j(\bar{u})) &= \int d^2x |v'(x)|^2 G(x)G(\bar{x}) = \int d^2x |v'(x)G(x)|^2 = \quad (4.66) \\ &= 2|n(n^2-1)C|^2 \int d^2x \left| \frac{\sqrt{2x+n(1+x)\sqrt{n^2(x-1)^2+4x}+n^2(1+x^2)}}{(x-1)^2\sqrt{n^2(x-1)^2+4x}} \right|^2 = \\ &= \sqrt{2}|n(n^2-1)C|^2 \int d^2x \left| \frac{n(x+1)+\sqrt{n^2(x-1)^2+4x}}{(x-1)^2\sqrt{n^2(x-1)^2+4x}} \right|^2, \end{aligned}$$

where the integration over x presumes integration over the double cover. The sum in the first line is over all the solution to (4.21), *i.e.* over all the diagrams in figures 4 and 5.

We are interested in working in large n limit.¹¹ In this limit we obtain the following simple equation

$$\sum_{j=1}^{2n} \int d^2u G(x_j(u), \bar{x}_j(\bar{u})) \sim |2n^3 C|^2 \int d^2x \left[\left| \frac{x}{(x-1)^3} \right|^2 + \left| \frac{1}{(x-1)^3} \right|^2 \right], \quad (4.67)$$

where on the *r.h.s* the two terms come from the two different maps. Using the following

$$\int d^2z z^a \bar{z}^{\bar{a}} (1-z)^b (1-\bar{z})^{\bar{b}} = \pi \frac{\Gamma(1+a)\Gamma(1+b)\Gamma(-\bar{a}-\bar{b}-1)}{\Gamma(-\bar{a})\Gamma(-\bar{b})\Gamma(a+b+2)}, \quad (4.68)$$

and plugging $a = \bar{a} = 1, 0$, $b = \bar{b} = -3$ we get that (4.67) vanishes and thus the chiral operators do not acquire an anomalous dimension.

¹¹Note that one can repeat the following with finite n . Finite n result can be found in appendix D using an alternative parametrization of the covering map.

Let us compute the first $1/n$ correction. We have

$$\begin{aligned} \delta_1 = & -\frac{1}{n} \delta_0 + \sqrt{2} |n C|^2 \int d^2x \left| \frac{2x}{(x-1)^3} \right|^2 \left(\frac{1}{(x-1)} - \frac{2x}{(x-1)^2} + c.c. \right) - \\ & - \sqrt{2} |n C|^2 \int d^2x \left| \frac{2}{(x-1)^3} \right|^2 \left(\frac{x}{(x-1)} + \frac{2x}{(x-1)^2} + c.c. \right), \end{aligned}$$

and again using (4.68) this is vanishing.

We have mentioned that there are no dimension $(1, 1)$ contact terms and one can verify this from the above explicit expression by taking appropriate *OPE* limits. In appendix D we explicitly show this in a different setup.

4.5 Comments on one loop with impurities

We are now ready to add impurities to the computation. In previous sections we have computed the two point function of operators with impurities but without the interactions, and two points of chiral operators without impurities but with interaction insertions. In this section we will combine these results to discuss the general structure of the computation of anomalous dimensions of the spin chain in one loop. For simplicity we will discuss only states with two holomorphic fermionic impurities of type *A*, *i.e.* one loop correction to (3.20).

We add impurities by “dressing” the chiral vacuum with contour integrals. It is convenient to keep the contour integrals and to compute the six point function of the four fermionic dressings and the two chiral fields first. The dressings are located at t, t_L and t', t'_M . Note that because the dressing is given in terms of untwisted sector fields this six point function is simply given by a product of the one loop vacuum result we obtained in previous section and the free field contractions of the dressing fermions and the fermions appearing in the interaction vertices. This free field computation gives the following result

$$\begin{aligned} G_1^{(a/b)}(t, t_L, t', t'_M, x) \equiv & (t - t_L)(t' - t'_M) \sqrt{\frac{\partial t_L}{\partial t} \frac{\partial t'_M}{\partial t'}} \frac{t^{\frac{1}{2}(1-n)} t_L^{\frac{1}{2}(1-n)} t'^{\frac{1}{2}(n-1)} t'_M^{\frac{1}{2}(n-1)}}{(t-t')(t_L-t')(t-t'_M)(t_L-t'_M)} \times \\ & \times \left[\sqrt{\frac{t-1}{t-x} \frac{t_L-1}{t_L-x} \frac{t'-x}{t'-1} \frac{t'_M-x}{t'_M-1}} + \sqrt{\frac{t-x}{t-1} \frac{t_L-x}{t_L-1} \frac{t'-1}{t'-x} \frac{t'_M-1}{t'_M-x}} \right], \end{aligned} \quad (4.69)$$

where the index (a/b) refers to the map with which we evaluate t_L and t'_M . The terms in the second line come from contractions with the interactions. Thus, to leading order in $1/n$, the two point function with impurities is given by

$$\begin{aligned} \langle M|L \rangle_{1-loop} = & |2n^3 C|^2 \int d^2x \left\{ \left| \frac{x}{(x-1)^3} \right|^2 \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} G_1^{(a)}(t, t_L, t', t'_M, x) + \right. \\ & \left. + \left| \frac{1}{(x-1)^3} \right|^2 \oint \frac{dt}{2\pi i} \oint \frac{dt'}{2\pi i} G_1^{(b)}(t, t_L, t', t'_M, x) \right\}. \end{aligned} \quad (4.70)$$

Let us discuss the structure of the above computation. First, we have to evaluate the following contour integrals

$$\oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} (t - t_L)(t' - t'_M) \sqrt{\frac{\partial t_L}{\partial t} \frac{\partial t'_M}{\partial t'}} \frac{t^{\frac{1}{2}(1-n)} t_L^{\frac{1}{2}(1-n)} t'^{\frac{1}{2}(n-1)} t'_M^{\frac{1}{2}(n-1)}}{(t-t')(t_L-t')(t-t'_M)(t_L-t'_M)} \times$$

$$\times \left[\sqrt{\frac{t-1}{t-x} \frac{t_L-1}{t_L-x} \frac{t'-x}{t'-1} \frac{t'_M-x}{t'_M-1}} + \sqrt{\frac{t-x}{t-1} \frac{t_L-x}{t_L-1} \frac{t'-1}{t'-x} \frac{t'_M-1}{t'_M-x}} \right] \quad (4.71)$$

and then integrate over x . Using the relations (4.32) the above can be written as

$$\oint \frac{dt'}{2\pi i} \oint \frac{dt}{2\pi i} (t - t_L)(t' - t'_M) \frac{t^{-n} t'^{n-1} t_L}{(t-t')(t_L-t')(t-t'_M)(t_L-t'_M)} \times$$

$$\times \frac{t_L - t_0}{t - t_0} \frac{t'_M - t_\infty}{t' - t_\infty} \left[\frac{t-1}{t_L-x} \frac{t'-x}{t'_M-1} + \frac{t-x}{t_L-1} \frac{t'-1}{t'_M-x} \right]. \quad (4.72)$$

Note that the integrand is not a meromorphic function as $t_L(t)$ and $t'_M(t')$ have branch cuts. The t contour is around zero and there are no branch cuts in the vicinity of the origin. The t' contour is around infinity. Thus, deforming this integral towards the origin we will encounter the branch cuts. This contour can be split into an integral around the branch cuts and an integral around the origin. Let us analyze the latter part first.

Note that for the t' integral to have a simple pole at $t' = 0$ we have to expand the denominator in the first line of (4.72) at least to order $n-3$ in t . The reason is as follows. Taking $t \rightarrow 0$ the integral scales as t^{n-4} . Thus, we at least have to get the above mentioned negative power of t' as the other terms in the integrand give positive powers of t' . However, expanding this denominator to order $n-3$ also gives us a simple pole in t . Thus, all the other terms have to be expanded to zeroth order. This in particular implies that the residue will be independent of x . Thus, the integral over the moduli space x will vanish as it does for the chiral operators. We deduce that the only non zero contribution to the one loop comes from t' integrals around the cuts.

There are three cuts for any given M : one cut of order n running between $t' = t_0$ and $t' = t_\infty$ and two cuts of order two running between points satisfying $t'_M(t') = 1, x$. As we do not know explicitly the functions t_L and t'_M the evaluation of these contour integrals is a complicated task. In principle, one can perform a consistent expansion in $1/n$ of these functions and try to evaluate the integrals. We leave the detailed investigation of these issues for future results. However, just from the generic structure of $t_L(t)$ discussed in section 4.3 we can learn that the two point function has the following structure.

$$\langle M|L \rangle_{1-loop} = f_a(L|M) - f_a(L|M+1) + f_b(L|M) - f_b(L|M-1), \quad (4.73)$$

$$+ f_a(L|-M) - f_a(L|-M) + f_b(L|-M) - f_b(L|-M-1),$$

where a, b label the two different maps. The first line above comes from the two order two cuts and the second line from the order n cut. We have assumed the large n limit and thus that the contour integrals around the two order two cuts are equal up to a sign. Of course we can also write another expression by exchanging M and L . This expression has a structure of “nearest neighbor interactions”. For this to hold precisely the functions $f_{a/b}(L|M)$ have to be proportional to $\delta_{L,M}$. To determine whether this is true an explicit computation has to be performed. However, in general we have mixing already at tree level and thus we do not have a reason to expect the above precise property to hold. This observation can be generalized to any type of impurities. Adding more interaction vertices we will have more cuts and thus smearing of this nearest neighbor feature.

5. Summary and Discussion

We have investigated a spin chain picture for the single-cycle gauge invariant states of $\text{Sym}^N T^4$. The ground state of a chain of length n is given by the chiral state $O_n^{(0,0)}$. This state is a bare twist field permuting n copies of T^4 appropriately dressed with fermions to render the conformal dimension equal to the R -charge. One considers the different copies (“colors”) the bare twist field permutes as the sites of the spin chain. In the vacuum state all the “colors” entering the bare twist field have the same dressing. Note that the state with lowest conformal dimension is the bare twist field and not the chiral state one builds from it and which we take as the vacuum of the spin chain. This is just the first of many qualitative differences between the symmetric product orbifolds and the gauge theories. In some sense the ground state is a “Dirac sea” of fermions [24] and the impurities are the excitations of this sea.

A natural set of impurities is introduced by changing the dressing of single colors permuted by the twist field. The fields with the lowest conformal dimension in the theory are the fermions and the basic impurities are given by dressing a site of the spin chain with these. We have explicitly shown that states with the same quantum numbers but with different impurities mix already at tree level. The technical reason for this is that on the covering surface the different copies permuted by the twist fields are identified and there is no suppression of contractions of fields sitting at different sites of the chain. This is in sharp contrast with the $1/N$ suppression of such contractions in a gauge theory. Of course, one can always diagonalize the states at tree level. An important open problem is to find whether there is a natural and economic way to describe the orthogonal basis of impurities. Ultimately one would like to find an operational definition of asymptotic states.

The generators of the superconformal algebra have an explicit realization in terms of the basic fields of the theory. Some of these generators, *e.g.* the R -charge, are quadratic combinations of the fields. Thus, again unlike the gauge theory case, the symmetry generators

are not the most fundamental impurities. Excitations of the spin chain generated by the symmetry algebra can be regarded as “composites” of more fundamental impurities.

We have also discussed higher “loop” computations. These are given by turning on an appropriate twist two interaction and expanding in the coupling constant. We have discussed the first non trivial, “one loop”, order in this expansion. The computation is also restricted to the leading order in $1/N$, *i.e.* the covering surface is a sphere. The explicit evaluation of the one loop reduces to a free field computation on a covering surface. The fact that the theory is interacting is manifested in two ways. First, the twist two field insertions give rise to a non trivial covering map. Moreover, the definition of the impurities in terms of contour integrals when lifted to the covering surface involves a function, $t_L(t)$, which is sensitive to the presence of the interaction terms. We have shown that the one loop computation of a two point function amounts to evaluating certain contour integrals around branch cuts of $t_L(t)$. This system of branch cuts has in some sense a “nearest neighbor” structure. This might be an indication of the local (nearest neighbor) nature of the one-loop Hamiltonian acting on the appropriate basis of (tree level) orthogonal impurities.

The main difficulty of performing explicit higher loop computations in gauge theories is the fast growth in the number of Feynman diagrams. In symmetric product CFTs the different “diagrams” are the different maps to the covering surface. As we have shown in section 4.2 *all* the maps relevant to our problem can be explicitly computed at least in the limit of large size of the spin chain. It would appear that higher loop calculations are not much more complex than one loop calculations, at least for chains of large size. So we may hope that a thorough understanding of one loop would lead to an all loop result.

Acknowledgements

We thank Antal Jevicki, Samir Mathur, and Luca Mazzucato for useful discussions. The work of AP was supported in part by DOE grant DE-FG02-91ER40688 and NSF grant PHY-0643150. The work of LR and SSR is supported in part by a Junior Investigator Award of the Department of Energy and by a Grant of the National Science Foundation. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

A. A short explanation of the diagrams

In this appendix we give a lightening summary of the diagrammatic technique for symmetric product orbifolds introduced in [28]. One way to compute a correlator of twist fields is to lift

the computation to the covering surface where the fields are single valued. There are however different ways to lift a given set of twist fields, *i.e.* given set of branching points. The different maps correspond to different ways to choose common indices, *i.e.* common “colors”, for the twist fields. To obtain a “gauge” invariant result one has to sum over all such maps, *i.e.* inequivalent choices of the color assignments to the twist fields. One can think of this sum as a formal sum over diagrams, much like the correlators in a gauge theory are sums over Feynman diagrams.

For simplicity we take all the twist fields to correspond to single-cycles. There are two ways to define the diagrams which are graph theoretic dual of each other. In the body of the paper we use diagrams in which each twist field corresponds to a loop. A cycle of length n corresponds to a loop connecting $2n$ vertices. See example in figure 9. There are two types of vertices, color and non-color ones, and their position is alternated along each loop. One can assign numbers to the color vertices and then the cycle-structure can be read off each loop by reading counter clockwise these numbers. The diagrams are obtained by gluing the loops in all possible ways modulo two rules. The first rule is that in each diagram the number of color vertices should equal the number of non color ones. The second rule is defined as follows. Each vertex defines a partial cyclic ordering on the loops, the color vertices by going around them counter clockwise and the non-color ones by going around them clockwise. The second rule is that all these orderings should be compatible with each other and also compatible with the radial ordering of the positions of the twist fields. The claim is that there is one to one correspondence between diagrams satisfying these conditions and the different maps contributing to a correlator of twist fields.




Figure 9: A diagram contributing to the correlator $\langle O_{[3]}(z_a)O_{[2]}(z_b)O_{[2]}(z_c)O_{[3]}(z_d) \rangle$ with $z_a < z_b < z_c < z_d$. The diagram corresponds to the following choice of (color) indices $(132)_a(24)_b(34)_c(241)_d$. The (solid) red dots are the color vertices and the white dots are the non color ones. Each loop (letter) corresponds to a twist field: going around the loop counterclockwise one reads off the color indices of the corresponding cyclic permutation.

We can think of the loop corresponding to a single-cycle as a spin chain, with the color vertices being the sites of the chain. The case with two twist-two interaction terms all the possible diagrams are depicted in figures 4 and 5. It is easy to check explicitly that the two basic rules are satisfied.

B. Diagonalization of the mixing of two J^- impurity states

In this appendix we describe the diagonalization procedure for the tree level mixing matrix (3.48). The diagonalization matrix can be conveniently written as a product of three matrices DVU . The non unitary matrix U is the following Fourier-like transform

$$U^k_L = \cos \frac{2\pi k L}{n}, \quad (\text{B.1})$$

where k labels the new states and we can think of it as a “momentum” variable. The unitary matrix V is given by

$$V = \begin{pmatrix} (\lfloor \frac{n}{2} \rfloor)^{-\frac{1}{2}} & (\lfloor \frac{n}{2} \rfloor)^{-\frac{1}{2}} & (\lfloor \frac{n}{2} \rfloor)^{-\frac{1}{2}} & (\lfloor \frac{n}{2} \rfloor)^{-\frac{1}{2}} & \dots & (\lfloor \frac{n}{2} \rfloor)^{-\frac{1}{2}} \\ 0 & \dots & 0 & 0 & \alpha_2 & -\alpha_2 \\ 0 & \dots & 0 & 2\alpha_3 & -\alpha_3 & -\alpha_3 \\ 0 & \dots & 3\alpha_4 & -\alpha_4 & -\alpha_4 & -\alpha_4 \\ & & \dots & & & \\ \lfloor \frac{n}{2} \rfloor \alpha_{\lfloor \frac{n}{2} \rfloor} & \dots & -\alpha_{\lfloor \frac{n}{2} \rfloor} & -\alpha_{\lfloor \frac{n}{2} \rfloor} & -\alpha_{\lfloor \frac{n}{2} \rfloor} & -\alpha_{\lfloor \frac{n}{2} \rfloor} \end{pmatrix}, \quad \alpha_k = \frac{1}{\sqrt{k(k-1)}}. \quad (\text{B.2})$$

The matrices are $\lfloor \frac{n}{2} \rfloor \times \lfloor \frac{n}{2} \rfloor$ dimensional. After diagonalizing we also have to rescale the states with the inverse of the eigenvalues. The eigenvalues of the state k is

$$k \neq 1 : \epsilon_k = \frac{6n^2 - 16n + 6}{n^2} - \frac{8}{n^2} (k-1)k, \quad k = 1 : \epsilon_1 = 4(n-2). \quad (\text{B.3})$$

That is we define a matrix D by

$$D_{kk'} = \sqrt{\epsilon_k} \delta_{kk'}. \quad (\text{B.4})$$

The matrix \mathcal{I} transforms as $(D^{-1} V (U^\dagger)^{-1}) \mathcal{I} (U^{-1} V^\dagger D^{-1}) = Id$. Note that after acting only with U the small momentum states are approximately orthogonal in the large n limit, a fact illustrated in figure 10.

C. Some details of the one loop map

In this appendix we give some formulae needed for the one loop computation in section 4.4. Lets us find the expansion of $z - u$ in terms of $y = t - x$, *i.e.* the inverse map. We can write




Figure 10: The expansion coefficients for two diagonal states, *i.e.* on the x axis are the L states and the y axis is the coefficient of each state in the “momentum” state decomposition. The two states are obtained by taking $k = 98$ (left) and $k = 10$ (right) with $n = 201$. The blue dots are the expansion coefficients and the red dots are the coefficients only of U . Note that for small k these are approximately identical.

the expansion as

$$y = \sum_{k=1}^{\infty} c_k (z - u)^{k/2}. \quad (\text{C.1})$$

To obtain the coefficients of the inverse map we first write the following

$$\ln \frac{z(t)}{u} = n \ln \frac{t}{x} + \ln \frac{t - t_0}{x - t_0} - \ln \frac{t - t_\infty}{x - t_\infty}, \quad (\text{C.2})$$

and expand both sides to get

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \left(\frac{z - u}{u} \right)^k = (t - x)^2 \sum_{k=0}^{\infty} a_k (t - x)^k, \quad (\text{C.3})$$

with

$$a_k = \frac{(-1)^{k+1}}{k + 2} \left(\frac{n}{x^{k+2}} + \frac{1}{(x - t_0)^{k+2}} - \frac{1}{(x - t_\infty)^{k+2}} \right). \quad (\text{C.4})$$

The expansion coefficients are related as

$$c_1^2 = \frac{1}{u(x) a_0}, \quad c_2 = -\frac{a_1}{2u(x) a_0^2}, \quad c_3 = -\frac{2a_0^3 - 5a_1^2 + 4a_0 a_2}{8u(x)^2 a_0^4 c_1}. \quad (\text{C.5})$$

The above results are needed to write down the differential equation (4.62). We get for the quantities appearing in this equation,

$$\begin{aligned}\left(\frac{t''}{t'}\right)' &= \frac{1}{2} \frac{1}{(z-u)^2} - \frac{c_2}{c_1} \frac{1}{2} (z-u)^{-3/2} + \dots \\ \left(\frac{t''}{t'}\right)^2 &= \frac{1}{4} \frac{1}{(z-u)^2} - \frac{c_2}{c_1} (z-u)^{-3/2} + 3 \left[\frac{c_2^2}{c_1^2} - \frac{c_3}{c_1} \right] (z-u)^{-1} + \dots\end{aligned}\tag{C.6}$$

From here we obtain

$$\left(\frac{t''}{t'}\right)' - \frac{1}{2} \left(\frac{t''}{t'}\right)^2 = \frac{3}{8} \frac{1}{(z-u)^2} - \frac{3}{2} \left[\frac{c_2^2}{c_1^2} - \frac{c_3}{c_1} \right] (z-u)^{-1} + \dots\tag{C.7}$$

We will also need the following

$$\begin{aligned}(t')^2 &= \frac{c_1^2}{4} (z-u)^{-1} + c_1 c_2 (z-u)^{-1/2} + \frac{1}{2} (2c_2^2 + 3c_1 c_3) \dots \\ \frac{1}{t-x} &= \frac{1}{c_1} (z-u)^{-1/2} - \frac{c_2}{c_1^2} + \frac{c_2^2 - c_1 c_3}{c_1^3} (z-u)^{1/2} + \dots \\ \frac{1}{t-x+a} &= \frac{1}{a} - \frac{c_1}{a^2} (z-u)^{1/2} - \left[\frac{c_2}{a^2} - \frac{c_1^2}{a^3} \right] (z-u) + \frac{-c_1^3 + 2ac_1 c_2 - a^2 c_3}{a^4} (z-u)^{3/2} + \dots,\end{aligned}\tag{C.8}$$

where a is some complex number.

D. Arutyunov-Frolov map

Note that the one loop map of section 4.2, although useful to understand the structure of different diagrams, is complicated as it contains square roots. Essentially, the map can be recast in much simpler form, as has been done by Arutyunov and Frolov in [53]. The simplification occurs if we map the twist at $z = \infty$ to $t = \infty$ and the twist $z = 0$ to $t = 0$ as before, but we map the additional image of $z = 0$ to $t = x - 1$ (instead of mapping the twist at $z = 1$ to $t = 1$ as was done in section 4.2). Then the map is given by the following

$$z(t) = t^n \frac{t - t_0}{t - t_\infty} \frac{t_1 - t_\infty}{t_1^n (t_1 - t_0)},\tag{D.1}$$

where we have

$$t_0 = x - 1, \quad t_\infty = x - \frac{x}{x+n},\tag{D.2}$$

$$t_1 = \frac{1-n}{n} + x - \frac{n+1}{n} \frac{x}{x+n}.\tag{D.3}$$

With these definition the point $z = u$ maps to $t = x$ with the following relation between the two

$$u = v(x) = \frac{x^{n-1} (x+n)^{n+1}}{(x-1)^{n+1} (x+n-1)^{n-1}}.\tag{D.4}$$

Note, in contrast to (4.20), that there are no square roots in this expression. Let us understand the moduli space in the new coordinates. The images of different OPE limits are

$$\begin{aligned}
u \rightarrow 0 & : & u & \sim \frac{n^{n+1}}{(n-1)^{n-1}} x^{n-1}, & u & \sim \frac{n^{n-1}}{(-n-1)^{n+1}} (x+n)^{n+1}, & (D.5) \\
u \rightarrow \infty & : & u & \sim \frac{(1+n)^{n+1}}{(n)^{n-1}} \frac{1}{(x-1)^{n+1}}, & u & \sim \frac{(n-1)^{n-1}}{n^{n+1}} \frac{1}{(x+n-1)^{n-1}}, \\
u \rightarrow 1 & : & u-1 & \sim -\frac{64}{3} \frac{n}{(n^2-1)^2} \left(x - \frac{1-n}{2}\right)^3, & x & \rightarrow \infty
\end{aligned}$$

In the limit $u \rightarrow 1$ we wrote down only those solutions contributing to the OPE limit, *i.e.* also $t_1 \rightarrow x$.

We can calculate the first correction to the two point functions of the chiral states using this map to obtain

$$G_0(x) = C x^{2-n} (x-1)^{3+n} (n+x)^{1-n} (x+n-1)^n \left(x + \frac{n-1}{2}\right)^{-4}, \quad (D.6)$$

where C is an overall constant not fixed by our method of computation. A simple check of this equation is to take $x \rightarrow \infty$. In this limit $u \rightarrow 1$ and corresponds to an OPE limit of the two interactions in the way that they share all the colors and thus annihilate each other, see equation (D.5). The four point function scales as $G(u) \sim \frac{1}{(1-u)^2}$ as expected as the interactions have dimension one. Computing the coefficient of the subleading term in the *OPE*, *i.e.* the term coming from dimension one operator, we find that it is zero. Thus there is no contribution from possible contact terms from the untwisted sector. In another limit, $x \rightarrow \frac{1-n}{2}$, we have $u \rightarrow 1$ and the leading singularity is -4 . This gives the conformal dimension of the operator of the leading *OPE* singularity to be $\Delta = \frac{2}{3}$. This is the dimension of bare twist three field. Farther, expanding to subleading order, *i.e.* $(x + \frac{n-1}{2})^{-3}$, we get that the coefficient is zero. This subleading order corresponds to $\Delta = 1$. Thus, the vanishing of this coefficient implies either that the correlator of the chiral states with the contact, twist three, terms vanishes, or that the contact terms simply do not exist.

We can easily integrate the above expression over the location of the interaction

$$\begin{aligned}
\sum_{j=1}^{2n} \int d^2u G(x_j(u), \bar{x}_j(\bar{u})) &= \int d^2x |v'(x)|^2 G(x) G(\bar{x}) = \int d^2x |v'(x) G(x)|^2 = & (D.7) \\
&= |nC|^2 \int d^2x \left| \frac{(x-1)(n+x)}{(n-1+2x)^2} \right|^2
\end{aligned}$$

We now make the following change of variables

$$y = -4 \frac{(x-1)(n+x)}{(1+n)^2}, \quad (D.8)$$

we obtain

$$|nC|^2 \int d^2x \left| \frac{(x-1)(n+x)}{(n-1+2x)^2} \right|^2 = \left(\frac{n^2(1+n)|C|}{16} \right)^2 \int d^2y \frac{|y|^2}{|1-y|^3} \quad (\text{D.9})$$

Using the following

$$\int d^2z z^a \bar{z}^{\bar{a}} (1-z)^b (1-\bar{z})^{\bar{b}} = \pi \frac{\Gamma(1+a)\Gamma(1+b)\Gamma(-\bar{a}-\bar{b}-1)}{\Gamma(-\bar{a})\Gamma(-\bar{b})\Gamma(a+b+2)}, \quad (\text{D.10})$$

and plugging $a = \bar{a} = 1$, $b = \bar{b} = -\frac{3}{2}$ we get that (4.67) is vanishing and thus the chiral operators do not acquire an anomalous dimension.

E. The one loop correlator of bare twist fields

Using map a and computing the four point function of twist fields without the dressings we obtain

$$\begin{aligned} \ln G(u(x)) = \frac{1}{8n} \bigg(& -2n \ln(x-1) - (2+5(n-1)n) \ln x - 2n \ln(1+x+\sqrt{n^2(x-1)^2+4x}) + \\ & + (2+n(2n-3)) \ln(n^2(x-1)-2x-n\sqrt{n^2(x-1)^2+4x}) + \\ & + (2+n(3+2n)) \ln(2+n(n(x-1)+\sqrt{n^2(x-1)^2+4x})) \bigg). \end{aligned} \quad (\text{E.1})$$

Remembering that the conformal dimension of a twist field is

$$\Delta_n = \frac{1}{4} \left(n - \frac{1}{n} \right), \quad (\text{E.2})$$

we can check different OPE limits. For instance as $u \rightarrow 1$ and $x \rightarrow 1$ we get that

$$\ln G(u) \sim -\frac{1}{4} \ln(x-1) \sim -\frac{1}{12} \ln(u-1), \quad (\text{E.3})$$

and we get also

$$\frac{1}{12} = \Delta_2 + \Delta_2 - \Delta_3. \quad (\text{E.4})$$

Taking the limit $u \rightarrow 0$ and $x \rightarrow 0$ we get

$$\ln G(u) \sim -\frac{2+5(n-1)n}{8n} \ln x \sim -\frac{2+5(n-1)n}{8n(n-1)} \ln u, \quad (\text{E.5})$$

which is to be understood as

$$\frac{2+5(n-1)n}{8n(n-1)} = \Delta_n + \Delta_2 - \Delta_{n-1}. \quad (\text{E.6})$$

In the same way for map b we obtain

$$\begin{aligned} \ln G(u(x)) = & -\frac{1}{8n} \left(6n \ln(x-1) + (n+n^2-2) \ln x - 2n \ln(1+x + \sqrt{n^2(x-1)^2 + 4x}) + \right. \\ & + (2+n(2n-3)) \ln(n^2(x-1) - 2x - n\sqrt{n^2(x-1)^2 + 4x}) + \\ & \left. + (2+n(3+2n)) \ln(2+n(n(x-1) + \sqrt{n^2(x-1)^2 + 4x}) \right). \end{aligned} \quad (\text{E.7})$$

The different OPE limits give here the following. As $u \rightarrow 1$ and $x \rightarrow 1$ we get that

$$\ln G(u) \sim -\frac{3}{4} \ln(x-1) \sim -\frac{3}{4} \ln(u-1), \quad (\text{E.8})$$

which is consistent with

$$\frac{3}{4} = \Delta_2 + \Delta_2, \quad (\text{E.9})$$

the two interactions annihilate each other as they share both their colors. Taking the limit $u \rightarrow 0$ and $x \rightarrow 0$ we get

$$\ln G(u) \sim -\frac{n^2+n-2}{8n} \ln x \sim -\frac{n^2+n-2}{8n(n+1)} \ln u, \quad (\text{E.10})$$

and we get also

$$\frac{n^2+n-2}{8n(n+1)} = \Delta_n + \Delta_2 - \Delta_{n+1}. \quad (\text{E.11})$$

Thus this computation gives us the identification between the two maps and the two copies of the moduli space.

References

- [1] I. Bena, J. Polchinski, and R. Roiban, *Hidden symmetries of the AdS(5) x S**5 superstring*, *Phys. Rev.* **D69** (2004) 046002, [[hep-th/0305116](#)].
- [2] J. A. Minahan and K. Zarembo, *The Bethe-ansatz for N = 4 super Yang-Mills*, *JHEP* **03** (2003) 013, [[hep-th/0212208](#)].
- [3] N. Beisert, *The dilatation operator of N = 4 super Yang-Mills theory and integrability*, *Phys. Rept.* **405** (2005) 1–202, [[hep-th/0407277](#)].
- [4] J. Plefka, *Spinning strings and integrable spin chains in the AdS/CFT correspondence*, *Living Rev. Rel.* **8** (2005) 9, [[hep-th/0507136](#)].
- [5] J. A. Minahan, *A brief introduction to the Bethe ansatz in N=4 super- Yang-Mills*, *J. Phys.* **A39** (2006) 12657–12677.
- [6] N. Dorey, *Notes on integrability in gauge theory and string theory*, *J. Phys.* **A42** (2009) 254001.

- [7] G. Arutyunov and S. Frolov, *Foundations of the $AdS_5 \times S^5$ Superstring. Part I*, *J. Phys.* **A42** (2009) 254003, [0901.4937].
- [8] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *$N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091, [0806.1218].
- [9] O. Aharony, O. Bergman, and D. L. Jafferis, *Fractional M2-branes*, *JHEP* **11** (2008) 043, [0807.4924].
- [10] J. A. Minahan and K. Zarembo, *The Bethe ansatz for superconformal Chern-Simons*, *JHEP* **09** (2008) 040, [0806.3951].
- [11] D. Gaiotto, S. Giombi, and X. Yin, *Spin Chains in $N=6$ Superconformal Chern-Simons-Matter Theory*, *JHEP* **04** (2009) 066, [0806.4589].
- [12] D. Bak and S.-J. Rey, *Integrable Spin Chain in Superconformal Chern-Simons Theory*, *JHEP* **10** (2008) 053, [0807.2063].
- [13] N. Gromov and P. Vieira, *The all loop AdS_4/CFT_3 Bethe ansatz*, *JHEP* **01** (2009) 016, [0807.0777].
- [14] J. M. Maldacena, *The large n limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [hep-th/9711200].
- [15] B. Chen, Y.-L. He, P. Zhang, and X.-C. Song, *Flat currents of the Green-Schwarz superstrings in $AdS(5) \times S^{*1}$ and $AdS(3) \times S^{*3}$ backgrounds*, *Phys. Rev.* **D71** (2005) 086007, [hep-th/0503089].
- [16] I. Adam, A. Dekel, L. Mazzucato, and Y. Oz, *Integrability of type II superstrings on Ramond-Ramond backgrounds in various dimensions*, *JHEP* **06** (2007) 085, [hep-th/0702083].
- [17] M. Staudacher, *The factorized S-matrix of CFT/AdS*, *JHEP* **05** (2005) 054, [hep-th/0412188].
- [18] N. Beisert, B. Eden, and M. Staudacher, *Transcendentality and crossing*, *J. Stat. Mech.* **0701** (2007) P021, [hep-th/0610251].
- [19] N. Gromov, V. Kazakov, and P. Vieira, *Integrability for the Full Spectrum of Planar AdS/CFT* , 0901.3753.
- [20] N. Seiberg and E. Witten, *The $d1/d5$ system and singular cft*, *JHEP* **04** (1999) 017, [hep-th/9903224].
- [21] F. Larsen and E. J. Martinec, *$U(1)$ charges and moduli in the $d1-d5$ system*, *JHEP* **06** (1999) 019, [hep-th/9905064].
- [22] D. E. Berenstein, J. M. Maldacena, and H. S. Nastase, *Strings in flat space and pp waves from $N = 4$ super Yang Mills*, *JHEP* **04** (2002) 013, [hep-th/0202021].
- [23] J. Gomis, L. Motl, and A. Strominger, *pp-wave / $CFT(2)$ duality*, *JHEP* **11** (2002) 016, [hep-th/0206166].

- [24] O. Lunin and S. D. Mathur, *Rotating deformations of $AdS(3) \times S(3)$, the orbifold CFT and strings in the pp-wave limit*, *Nucl. Phys.* **B642** (2002) 91–113, [[hep-th/0206107](#)].
- [25] E. Gava and K. S. Narain, *Proving the pp-wave / CFT(2) duality*, *JHEP* **12** (2002) 023, [[hep-th/0208081](#)].
- [26] J. R. David and B. Sahoo, *Giant magnons in the D1-D5 system*, *JHEP* **07** (2008) 033, [[0804.3267](#)].
- [27] B.-H. Lee, R. R. Nayak, K. L. Panigrahi, and C. Park, *On the giant magnon and spike solutions for strings on $AdS_3 \times S^3$* , *JHEP* **06** (2008) 065, [[0804.2923](#)].
- [28] A. Pakman, L. Rastelli, and S. S. Razamat, *Diagrams for Symmetric Product Orbifolds*, *JHEP* **10** (2009) 034, [[0905.3448](#)].
- [29] A. Pakman, L. Rastelli, and S. S. Razamat, *Extremal Correlators and Hurwitz Numbers in Symmetric Product Orbifolds*, *Phys. Rev.* **D80** (2009) 086009, [[0905.3451](#)].
- [30] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, *Large n field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183–386, [[hep-th/9905111](#)].
- [31] R. Dijkgraaf, *On the d1-d5 conformal field theory*, *Class. Quant. Grav.* **17** (2000) 1035–1048.
- [32] J. R. David, G. Mandal, and S. R. Wadia, *Microscopic formulation of black holes in string theory*, *Phys. Rept.* **369** (2002) 549–686, [[hep-th/0203048](#)].
- [33] E. Martinec, *The d1-d5 system*, <http://hamilton.uchicago.edu/~ejm/japan99.ps>.
- [34] R. Dijkgraaf, *Instanton strings and hyperkaehler geometry*, *Nucl. Phys.* **B543** (1999) 545–571, [[hep-th/9810210](#)].
- [35] J. M. Maldacena and A. Strominger, *$AdS(3)$ black holes and a stringy exclusion principle*, *JHEP* **12** (1998) 005, [[hep-th/9804085](#)].
- [36] J. de Boer, *Six-dimensional supergravity on $s^{**3} \times ads(3)$ and 2d conformal field theory*, *Nucl. Phys.* **B548** (1999) 139–166, [[hep-th/9806104](#)].
- [37] D. Kutasov, F. Larsen, and R. G. Leigh, *String theory in magnetic monopole backgrounds*, *Nucl. Phys.* **B550** (1999) 183–213, [[hep-th/9812027](#)].
- [38] R. Argurio, A. Giveon, and A. Shomer, *Superstrings on $ads(3)$ and symmetric products*, *JHEP* **12** (2000) 003, [[hep-th/0009242](#)].
- [39] A. Giveon, D. Kutasov, and N. Seiberg, *Comments on string theory on $ads(3)$* , *Adv. Theor. Math. Phys.* **2** (1998) 733–780, [[hep-th/9806194](#)].
- [40] A. Giveon and A. Pakman, *More on superstrings in $ads(3) \times n$* , *JHEP* **03** (2003) 056, [[hep-th/0302217](#)].
- [41] S. K. Ashok, R. Benichou, and J. Troost, *Asymptotic Symmetries of String Theory on $AdS_3 \times S^3$ with Ramond-Ramond Fluxes*, *JHEP* **10** (2009) 051, [[0907.1242](#)].

- [42] A. Jevicki, M. Mihailescu, and S. Ramgoolam, *Gravity from cft on $s^{**n}(x)$: Symmetries and interactions*, *Nucl. Phys.* **B577** (2000) 47–72, [[hep-th/9907144](#)].
- [43] O. Lunin and S. D. Mathur, *Correlation functions for $m(n)/s(n)$ orbifolds*, *Commun. Math. Phys.* **219** (2001) 399–442, [[hep-th/0006196](#)].
- [44] O. Lunin and S. D. Mathur, *Three-point functions for $m(n)/s(n)$ orbifolds with $n = 4$ supersymmetry*, *Commun. Math. Phys.* **227** (2002) 385–419, [[hep-th/0103169](#)].
- [45] M. R. Gaberdiel and I. Kirsch, *Worldsheet correlators in $ads(3)/cft(2)$* , *JHEP* **04** (2007) 050, [[hep-th/0703001](#)].
- [46] A. Dabholkar and A. Pakman, *Exact chiral ring of $ads(3)/cft(2)$* , *Adv. Theor. Math. Phys.* **13** (2009) 409–462, [[hep-th/0703022](#)].
- [47] A. Pakman and A. Sever, *Exact $n=4$ correlators of $ads(3)/cft(2)$* , *Phys. Lett.* **B652** (2007) 60–62, [[arXiv:0704.3040](#) [[hep-th](#)]].
- [48] M. Taylor, *Matching of correlators in AdS_3/CFT_2* , *JHEP* **06** (2008) 010, [[0709.1838](#)].
- [49] G. Giribet, A. Pakman, and L. Rastelli, *Spectral Flow in $AdS(3)/CFT(2)$* , *JHEP* **06** (2008) 013, [[0712.3046](#)].
- [50] G. Giribet and L. Nicolas, *Comment on three-point function in $AdS(3)/CFT(2)$* , *J. Math. Phys.* **50** (2009) 042304, [[0812.2732](#)].
- [51] C. A. Cardona and C. A. Nunez, *Three-point functions in superstring theory on $AdS_3 \times S^3 \times T^4$* , *JHEP* **06** (2009) 009, [[0903.2001](#)].
- [52] J. de Boer, J. Manschot, K. Papadodimas, and E. Verlinde, *The chiral ring of AdS_3/CFT_2 and the attractor mechanism*, *JHEP* **03** (2009) 030, [[0809.0507](#)].
- [53] G. E. Arutyunov and S. A. Frolov, *Virasoro amplitude from the $s(n) r^{**24}$ orbifold sigma model*, *Theor. Math. Phys.* **114** (1998) 43–66, [[hep-th/9708129](#)].
- [54] G. E. Arutyunov and S. A. Frolov, *Four graviton scattering amplitude from $s(n) r^{**8}$ supersymmetric orbifold sigma model*, *Nucl. Phys.* **B524** (1998) 159–206, [[hep-th/9712061](#)].
- [55] S. G. Avery, B. D. Chowdhury, and S. D. Mathur, *Emission from the $D1D5$ CFT*, *JHEP* **10** (2009) 065, [[0906.2015](#)].
- [56] S. G. Avery and B. D. Chowdhury, *Emission from the $D1D5$ CFT: Higher Twists*, [0907.1663](#).
- [57] Y. Hikida and Y. Sugawara, *Superstrings on PP -wave backgrounds and symmetric orbifolds*, *JHEP* **06** (2002) 037, [[hep-th/0205200](#)].
- [58] S.-M. Lee, S. Minwalla, M. Rangamani, and N. Seiberg, *Three-point functions of chiral operators in $d = 4$, $n = 4$ sym at large n* , *Adv. Theor. Math. Phys.* **2** (1998) 697–718, [[hep-th/9806074](#)].
- [59] S. Corley, A. Jevicki, and S. Ramgoolam, *Exact correlators of giant gravitons from dual $N = 4$ SYM theory*, *Adv. Theor. Math. Phys.* **5** (2002) 809–839, [[hep-th/0111222](#)].

- [60] J. McGreevy, L. Susskind, and N. Toumbas, *Invasion of the giant gravitons from anti-de Sitter space*, *JHEP* **06** (2000) 008, [[hep-th/0003075](#)].
- [61] A. Hashimoto, S. Hirano, and N. Itzhaki, *Large branes in AdS and their field theory dual*, *JHEP* **08** (2000) 051, [[hep-th/0008016](#)].
- [62] O. Lunin, S. D. Mathur, and A. Saxena, *What is the gravity dual of a chiral primary?*, *Nucl. Phys.* **B655** (2003) 185–217, [[hep-th/0211292](#)].
- [63] S. Raju, *Counting giant gravitons in ads_3* , [arXiv:0709.1171](#) [[hep-th](#)].
- [64] N. Berkovits and C. Vafa, *$N=4$ topological strings*, *Nucl. Phys.* **B433** (1995) 123–180, [[hep-th/9407190](#)].