A Combinatorial Enumeration of Distances for Calculating Energy in Molecular Conformational Space.

Jacques Gabarro-Arpa

November 8, 2018

Ecole Normale Supérieure de Cachan - LBPA CNRS UMR 8113 - 61, Avenue du Président Wilson - 94235 Cachan cedex - France.

e-mail: jga@gtran.org

Abstract

In previous works it was shown that protein 3D-conformations could be encoded into discrete sequences called dominance partition sequences (DPS), that generated a linear partition of molecular conformational space into regions of molecular conformations that have the same DPS. In this work we describe procedures for building in a cubic lattice the set of 3Dconformations that are compatible with a given DPS. Furthermore, this set can be structured as a graph upon which a combinatorial algorithm can be applied for computing the mean energy of the conformations in a cell.

1 Introduction

In previous papers [1-5] we have built a series of mathematical tools for studying the multidimensional molecular conformational space of proteins, with the aim of understanding the dynamical states of proteins by building a complete energy surface.

In this approach, the 3D-structures of protein molecules are encoded into a linear sequence of numbers called **dominance partition sequences** (DPS) [1-5], there are three of these sequences one for each coordinate x, y and z. For a molecule of N atoms assigning to each atom a number in the range 1 - N, then for a coordinate c the DPS is : the sequence of atom numbers sorted in ascending order of the value of the c-coordinate of their respective atoms.

A typical 3D-DP sequence may be something like

 $\{\{(5)(3)(1)(4)(2)\}_x, \{(2)(4)(3)(1)(5)\}_y, \{(1)(3)(5)(2)(4)\}_z\}$

which means that

 $x_5 < x_3 < x_1 < x_4 < x_2, \qquad y_2 < y_4 < y_3 < y_1 < y_5, \qquad z_1 < z_3 < z_5 < z_2 < z_4$

A simple example of DPS can be extracted from Fig. 1 where the α -carbon skeleton of the pancreatic trypsin inhibitor protein [6,7] is shown




Figure 1

Numbered α -carbon chain stereoview of the pancreatic trypsin inhibitor protein.

with the atoms positions numbered. The DPSs for this protein conformation are :

 $\{ \{(58)(49)(29)(48)(57)(27)(28)(31)(30)(52) (32)(47)(53)(50)(19)(26)(21)(56)(51)(24) \\ (33)(20)(23)(55)(46)(25)(22)(34)(54)(18) (1)(45)(17) (5) (6)(35)(44) (2) (8)(43) \\ (16) (9)(11) (7) (3)(10)(36) (4)(37)(42) (15)(12)(41)(40)(14)(38)(13)(39)\}_x, \\ \{(15)(16)(17)(14)(18)(37)(36)(13)(19)(38) (34)(35)(12)(11)(39)(20)(33)(46)(10)(40) \\ (32)(47)(21)(45)(44) (9)(48)(31)(41)(22) (49)(50)(43)(42) (8)(51)(30)(23)(24)(52) \\ (7)(29)(54)(53) (5)(27)(55)(26)(25) (4) (6)(28)(57)(56)(58) (3) (2) (1)\}_y, \\ \{(26)(27)(10) (8)(25) (7)(24)(11) (6) (9) (12)(28)(13)(33)(34)(15)(31)(32)(29)(17) \\ (14)(23)(36)(41)(35) (3)(40) (5)(22)(16) (30) (4)(39)(43) (1)(18)(21)(19)(42)(20) \\ (44)(38) (2)(37)(55)(48)(45)(51)(52)(57) (56)(47)(46)(54)(49)(53)(58)(50)\}_z \}$

The main tool developped [3-5] in this approach can be described as a **fluctuation amplifier**: the small movements of a molecular system, which are essentially sampled with the current computer simulating tools, are encoded by means of a simple combinatorial structure, from which we can generate the complete set of DPSs corresponding to realizable 3D-conformations that arise from the combination of these movements. In the preceeding papers [3-5] it was described how to build a graph whose nodes are the cells that are visited by the system in its thermal wandering, with edges towards the adjacent cells.

As it was it was suggested in [1] molecular 3D-conformations are constrained in a small fraction of the cell volume, for this formalism to be useful the conformational volume inside a cell has to be probed and the mean energy of its conformations must be evaluated. The problem adressed in the present work is how from a DPS code the set of 3D-conformations it encodes can be reconstructed.

2 A procedure for embedding molecular conformations in a cubic 3D spatial lattice

We start by describing a procedure for embedding the molecular 3D-structures in a cubic lattice, this can be done using empirical data sampled from molecular dynamics simulations [7]

Procedure 1

- 1. First we determine the dimensions of the lattice by taking as reference the mean bond length between C_{α} carbons, which is : $3.58\text{\AA} < 3.86\text{\AA} < 4.13\text{\AA}$. In the example developped here we set this length arbitrarily to 20 lattice units, which gives a lattice spacing of 0.19\AA .
- 2. From the range of variation extracted from molecular dynamics any segment between to lattice points with a length range between $3.58 \times 20/3.86$ and $4.13 \times 20/3.86$ is potentially C_{α} - C_{α} bond segment. The set of valid lattice bond segments, modulo a lattice translation along the

x, y and z axes, is the set of segments starting at the origin and ending in any lattice point that lies between two spheres of radius $3.58 \times 20/3.86$ and $4.13 \times 20/3.86$ respectively. This gives a total of 1883 primary segments, i.e. modulo a reflection through the xy, xz and yz planes.

3. Next we determine the range of variation for the bond angles, which is greater than that for the bond length and varies considerably along the C_{α} chain. For each bond angle $A_{\alpha_1\alpha_2\alpha_3}$ along the main chain we determine two integer numbers : the floored minimum $\lfloor min(A_{\alpha_1\alpha_2\alpha_3}) \rfloor$ and the ceiled maximum range $\lceil max(A_{\alpha_1\alpha_2\alpha_3}) \rceil$ respectively. These divide the range $0^{\circ} - 360^{\circ}$ in a number of intervals which in our case give $71^{\circ}-74^{\circ}-75^{\circ}-76^{\circ}-77^{\circ}-78^{\circ}-79^{\circ}-80^{\circ}-81^{\circ}-82^{\circ}-89^{\circ}-$

each $A_{\alpha_1\alpha_2\alpha_3}$ along the backbone has a range spanning a given interval set from (2).

- 4. The set of allowed bond angles formed by pairs of primary segments are classified according to the following :
 - (a) Their sign vector : from a broken line formed by two primary segments with end coordinates $v1 = \{x_0, y_0, z_0\}$ and $v2 = \{x_1, y_1, z_1\}$, a set of three sign vectors can be generated

 $\{ sign(-x_0), sign(-x_1), sign(x_0 - x_1) \} \\ \{ sign(-y_0), sign(-y_1), sign(y_0 - y_1) \} \\ \{ sign(-y_0), sign(-y_1), sign(y_0 - y_1) \}$

where the function sign(x) returns the symbols +,- and 0 if x is positive, negative or zero¹. In order to reduce the size of data we introduce the constraint that the first three signs of each vector must be +, from these all other sign classes can be generated by reflection symmetry through the planes perpendicular to x, y and z. A total of 328 sign classes are thus generated.

- (b) Their bond angle interval : pairs of primary bond segments within a sign class are sorted in angular interval subclasses according to their bond angle.
- 5. Next lattice embedded molecular 3D-conformations of the protein backbone can be generated by joining lattice bond segments with the correct angular interval and the correct sign matrix between the atoms.

3 The graph of lattice points

Each bond between two atoms in the molecular backbone can be approximated by a lattice primary segment, and each pair of consecutive bonds can be approximated by a pair of segments having an angle within the bond angle dynamic range (2). In our approach molecular backbone 3D-conformations are characterized by the dynamic range of their bond lengths and bond angles, and by the sign matrix or equivalently the dominance partition sequence. We have seen in the previous section how to embed the molecular backbone in a discrete lattice, the problem we try to solve here

¹The sign matrix and DPSs are equivalent since $sign(x_0 - x_1) = +$ means that $x_0 > x_1[1]$.

is how to enumerate the finite set of lattice 3D-conformations that fulfill the DPS constraints of a cell or set of cells.

For this we need to build a graphical structure that we call the **graph of lattice points** in three steps

Procedure 2

- 1. we arbitrarily set the coordinates of the first C_{α} (the **root** node) in the backbone to $\{0, 0, 0\}$ to avoid translation ambiguities.
- 2. we choose all segment pairs that have the same sign vectors as the first 3 C_{α} s and angular value within the interval limits allowed for the first bond angle.
- 3. building the $(n+1)^{th}$ lattice bond level in the C_{α} backbone is done from each individual n^{th} level bond segment by joining it with the second segments of those lattice bond pairs that :
 - (a) the first segment is the same as the n^{th} bond,
 - (b) the angular value of the pair is within the range of the $(n+1)^{th}$ bond angles.
- 4. for each level the nodes of the graph are the lattice points at the upper end of a bond segment. There will be two arcs between any two points in two consecutive levels if they are connected by a bond segment: a forward arc from the node in the n^{th} level towards the one in the $(n+1)^{th}$ level if the two are connected by a bond segment, there is also a reverse arc from the $(n+1)^{th}$ level towards the n^{th} level towards the n^{th}
- 5. The root node has an arc towards every node in the upper level and reciprocally.

4 An algorithm for determining the set of inter atomic distances and weights

In force fields currently used in molecular dynamics simulations [8] atoms are represented by pointlike structures and conformational energy is calculated with a Hamiltonian which is a sum of two kinds of terms a) **local**: which involve groups of 2, 3 and 4 consecutive atoms needed to calculate the bond, bond angle and torsion angle energies; b) **non-local** involving pairs of atoms lying anywhere in the structure, these are needed for calculating the electrostatic and van der Waals energy terms.

Usually energy is calculated for only one conformation at a time, not here : in this work we intend to calculate the energy for a great number of structures simultaneously, as each term in the hamiltonian may appear in many structures, we want to calculate it only once. For this we need to know in how many 3D-structures, or equivalently for how many paths in the graph of lattice points, our term arises. For the non-local terms we need further to enumerate the whole set of inter-atomic distances between pairs of atoms, this is possible on a lattice because squared distances are integer numbers.

We first start by describing an algorithm that assigns two weights to each node : for a given node n the lower weight (LW_n) is the number of downward paths starting at the node and ending at the root node, the upper weight (UW_n) is the number of upward paths starting at the node and ending at ending at some node in the top level. The algorithm is as follows :

Procedure 3

- 1. For the root node we set its lower weight value to 1.
- 2. We go to the next level.
- 3. For each node n in the level we set the LW_n value to the sum of its downward link nodes LWs.
- 4. If the top level has not been reached we go to step 2.
- 5. Otherwise for every node in the top level we set the upper weight to 1.
- 6. We go to the previous level.
- 7. For each node n in the level we set the UW_n value to the sum of its upward link nodes UW_s .
- 8. If the root level has not been reached we go step 6.
- 9. Otherwise we terminate the procedure.

Thus two consecutive lattice points a and b will contribute the quantity $E_{bond}(d_{a,b}) \times LW_a \times UW_b$ to the global bond energy, where $d_{a,b}$ is the length of the lattice segment between a and b and LW_a and UW_b are the lower and upper weights of a and b respectively. Similarly for the bond and torsion energy of consecutive lattice points a, b, c and d we have $E_{angle}(\theta_{a,b,c}) \times LW_a \times UW_c$ and $E_{torsion}(\phi_{a,b,c,d}) \times LW_a \times UW_d$ respectively. These quantities can be calculated using a variant of the basic algorithm described in procedure 1.

To determine the weights for distances between arbitrary pairs of nodes we need three new data structures : an integer flag variable in every node, a sorted table of distances and a register array whose lengh is the number of levels.

Procedure 4

- 1. We set the flag in every node to 0.
- 2. For the N_{top} nodes in the top level.
- 3. For every node in the top level
 - (a) We assign to each one a number n_{top} from 1 to N_{top} .
 - (b) We enter the node in the top level of the array register.
 - (c) We follow every downward link in succession to the previous level.
- 4. For each node thus reached
 - (a) We set its flag to the top level number and we enter the node in the corresponding level of the array register.
 - (b) We follow every downward link in succession.
 - (c) Upon reaching the root level or a node whose flag value is equal to the current n_{top}
 - i. We calculate the distances between all nodes in the array register from the current to the top level.
 - ii. Every calculated distance $d_{a,b}$ between nodes a and b in levels $L_a > L_b$ is assigned an upper and lower weight : UW_a and LW_b . The array $\{d_{a,b}, LW_b, UW_a\}$ is searched in the table of distances

- A. If not found it is entered in the table together with the number $count_{a,b}$ which is set to 1.
- B. If the array already exists in the table $count_{a,b}$ is increased by 1.
- iii. We return to the previous node and from there we follow the next downward link to a node in the lower level.
- (d) Upon reaching a node in level L_v whose flag value n > 0 is different from the current n_{top}
 - i. We continue the downward exploration of the graph setting the flag of every node to n_{top} .
 - ii. We upon reaching the root node we do not compute distances between nodes in levels L_v or lower.
- 5. The procedure terminates when all downward links in every top node have been explored.

Each distance $d_{a,b}$ from the table of distances will have a weight $W_{a,b} = UW_a + LW_b + count_{a,b}$ which will we used to compute the van der Waals and electrostatic terms of the hamiltonian $E_{vdW}(d_{a,b}) \times W_{a,b}$ and $E_{elec}(d_{a,b}) \times W_{a,b}$ respectively.

5 Discussion

The procedures discussed in this work make possible removal of two important hurdles of the formalism in its way towards practical applications :

- 1. The construction of realistic 3D-conformations with a given partition sequence. As was discussed in [1], a lot of structural information disappears when replacing the coordinates of a molecule by the inverse sequences of its DPSs, however though the molecule appears heavily deformed all the secondary structure motifs : α -helices, β -sheets, turns, ... together with the overall 3D folding arrangement can still be recognized. This means that the $3 \times (N-1)$ -dimensional volume² of a cell in conformational space is big enough to allow for very lean codes, but it still exerts a constraint that keeps the 3D-structures close to the real ones.
- 2. The combinatorial structures described above for building molecular conformations can be transformed to calculate their mean energy. Which is essential for comparing the results from this formalism with experimental data.

It is also quite conceivable that in some practical situations the procedures decribed above may overwhelmed by the amount of data generated, in this case the combinatorial structures they generate will have to be pruned in order to make them useful. This question will be adresses when submitting this model of conformational molecular space to phenomenological tests.

²For a molecule with N atoms.

References

- J. Gabarro-Arpa, A central partition of molecular conformational space. I. Basic structures, Comp. Biol. Chem. 27, 153-159 (2003).
- [2] J. Gabarro-Arpa, A central partition of molecular conformational space. II. Embedding 3Dstructures, in *Proceedings of the 26th Conference of the IEEE EMBS*, San Francisco, pp. 3007-3010, 2004.
- [3] J. Gabarro-Arpa, A central partition of molecular conformational space. III. Combinatorial determination of the volume spanned by a molecular system in conformational space, J. Math. Chem. 42, 691-706 (2006).
- [4] J. Gabarro-Arpa, A central partition of molecular conformational space. IV. Extracting information from the graph of cells, J. Math. Chem. 44, 872-883 (2008).
- [5] J. Gabarro-Arpa, A central partition of molecular conformational space. V. The hypergraph of 3D partition sequences, arXiv: 0812-1.844v1 (2008).
- [6] M. Marquart, J. Walter, J. Deisenhofer, W. Bode, R. Huber, The geometry of the reactive site and of the peptide groups in trypsin, trypsinogen and its complexes with inhibitors, *Acta Crystallogr. Sect. B* 39, 480–490 (1983).
- [7] J. Gabarro-Arpa, R. Revilla, Clustering of a molecular dynamics trajectory with a Hamming distance, Comp. and Chem. 24, 693-698 (2000).
- [8] W. Wang, O. Donini, C.M. Reyes, P.A. Kollman, Biomolecular simulations: recent developments in force fields, simulations of enzyme catalysis, protein-ligand, protein-protein, and proteinnucleic acid noncovalent interactions, Ann. Rev. Biophys. Biomol. Struct. 30, 211-243 (2001).