# THE CALIFORNIA PLANET SURVEY II. A SATURN-MASS PLANET ORBITING THE M DWARF GL 649<sup>1</sup>

John Asher Johnson<sup>2</sup>, Andrew W. Howard<sup>3,4</sup>, Geoffrey W. Marcy<sup>3</sup>, Brendan P. Bowler<sup>5</sup>, Gregory W. Henry<sup>6</sup>, Debra A. Fischer<sup>7</sup>, Kevin Apps<sup>8</sup>, Howard Isaacson<sup>3</sup>, Jason T. Wright<sup>9</sup>

Draft version May 26, 2018

## ABSTRACT

We report precise Doppler measurements of the nearby (d = 10.34 pc) M dwarf Gl 649 that reveal the presence of a planet with a minimum mass  $M_P \sin i = 0.328 \text{ M}_{\text{Jup}}$  in an eccentric (e = 0.30), 598.3 day orbit. Our photometric monitoring reveals Gl 649 to be a new variable star with brightness changes on both rotational and decadal timescales. However, neither of these timescales are consistent with the 600-day Doppler signal and so provide strong support for planetary reflex motion as the best interpretation of the observed radial velocity variations. Gl 649 b is only the seventh Doppler-detected giant planet around an M dwarf. The properties of the planet and host-star therefore contribute significant information to our knowledge of planet formation around low-mass stars. We revise and refine the occurrence rate of giant planets around M dwarfs based on the California Planet Survey sample of low-mass stars ( $M_{\star} < 0.6 \,\mathrm{M_{\odot}}$ ). We find that  $f = 3.4^{+2.2}_{-0.9}\%$  of stars with  $M_{\star} < 0.6 \,\mathrm{M_{\odot}}$  harbor planets with  $M_P \sin i > 0.3 \,\mathrm{M_{Jup}}$  and  $a < 2.5 \,\mathrm{AU}$ . When we restrict our analysis to metal-rich stars with [Fe/H] > +0.2 we find the occurrence rate is  $10.7^{+5.9}_{-4.2}\%$ .

Subject headings: techniques: radial velocities—planetary systems: formation—stars: individual (Gl 649)

#### 1. INTRODUCTION

Compared to the knowledge gleaned from the large sample of giant planets around Sun-like stars, little is known about the characteristics of Jovian planets around M dwarfs. This is due primarily to the empirical finding that the occurrence rate of detectable planets scales with stellar mass (Johnson et al. 2007); low-mass stars  $(M_{\star} < 0.6 \text{ M}_{\odot})$  simply do not harbor giant planets very frequently (Endl et al. 2003; Butler et al. 2006). The frequency of giant planets with  $M_P \sin i > 0.3 \,\mathrm{M}_{\mathrm{Jup}}$  around Sun-like stars is 8% within 2.5 AU (Cumming et al. 2008), and the occurrence of giant planets around M dwarfs is roughly a factor of 4 lower.

While the lower masses of M dwarfs decreases the likelihood of giant planet occurrence, a handful of Jovian planets have been discovered around low-mass stars. The sample of M dwarfs known to harbor at least one Doppler-detected giant planet  $(M_P \sin i > 0.2 \text{ M}_{Jup})$ is listed in Table 1 and shown in the H-R diagram in

Electronic address: johnjohn@astro.berkeley.edu

Based on observations obtained at the W.M. Keck Observatory, which is operated jointly by the University of California and the California Institute of Technology. Keck time has been granted by both NASA and the University of California.

<sup>2</sup> Department of Astrophysics, California Institute of Technol-ogy, MC 249-17, Pasadena, CA 91125

Department of Astronomy, University of California, Mail Code 3411, Berkeley, CA 94720 <sup>4</sup> Townes Fellow, Space Sciences Laboratory, University of Cali-

fornia, Berkeley, CA 94720-7450, USA

<sup>5</sup> Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822

<sup>6</sup> Center of Excellence in Information Systems, Tennessee State University, 3500 John A. Merritt Blvd., Box 9501, Nashville, TN

<sup>37209</sup> <sup>7</sup> Yale Astronomy Department, Box 208101, New Haven, CT 06520-8101, USA

<sup>8</sup> Cheyne Walk Observatory, Horley, Surrey, RH6 7LR, United Kingdom

The Pennsylvania State University, University Park, PA 16802

GI 649 HIP 79431 GI 849 GI 876 ≥́ e/H]= +0.2 K5 M5 10 3 4 5 6 V–K

FIG. 1.— Our Keck sample of low-mass stars plotted in the  $\{V K, M_K$  plane. The solid line is a fifth-order polynomial fit to the mean main sequence for stars within 10 pc, which Johnson & Apps (2009) identify as an isometallicity contoure with [Fe/H] equal to the mean value of the Solar neighborhood. The dashed line corresponds to [Fe/H] = +0.2 based on the calibration of Johnson & Apps. The five-point stars show the positions of all of the M dwarfs known to harbor at least one giant planet. The solid circle denotes the position of Gl 649.

Figure 1. Also given in that table are the stellar and planetary masses from the literature, and stellar metallicities from the broad-band photometric calibration of Johnson & Apps (2009).

These planets and their host stars demonstrate that stellar mass is not the only characteristic that correlates with the probability of a star harboring a planet. Stellar metallicity has been shown to be a strong predictor of planet occurrence around Sun-like stars (Fischer & Valenti 2005a), and the correlation between planet frequency and stellar metal content appears to hold for the M dwarfs, as well. Johnson & Apps (2009)



FIG. 2.— Our Keck/HIRES radial velocity time series for Gl 649. The dashed line shows the best-fitting Keplerian model. The rms scatter of the residuals (bottom panel) is 4.2 m s<sup>-1</sup>, and  $\sqrt{\chi_{\nu}^2} = 1.17$ .

found that M dwarfs with Jovian planets tend to be significantly metal-rich compared to a 10 pc, volume-limited sample of stars on the lower main sequence. For example, Gl 849 harbors a Jovian planet in a long-period orbit and is among the most metal-rich stars in the Solar neighborhood with [Fe/H] > +0.45.

If this preliminary trend proves to be real then it will provide valuable constraints for theoretical models of planet formation around a broad range of stellar characteristics. The effect of metallicity on planet occurrence will also inform the target selection of future Doppler and transit surveys targeting low-mass stars (e.g. Irwin et al. 2008), as well as the interpretation of results from direct-imaging, astrometric and microlensing surveys (e.g. Nielsen & Close 2009; Pravdo & Shaklan 2009; Dong et al. 2009).

As the time baselines, sample sizes and Doppler precision increase for the various Doppler surveys of low-mass stars, the relationships between the physical characteristics of stars and the properties of their planets will come into sharper focus. We are monitoring a sample of 147 late K and early M stars as part of the California Planet Survey at Keck Observatory with a current temporal baseline of  $\approx 12$  years and Doppler precision of 2–3 m s<sup>-1</sup> (Johnson et al. 2007; Howard et al. 2009b). In this contribution we announce the detection of a new Saturn-mass planet orbiting a nearby M dwarf. Gl 649 is only the eleventh M-type star known to harbor at least one Doppler-detected planet<sup>10</sup>, and it is

 $^{10}$  Several additional low–mass host stars have been discovered by gravitational microlensing surveys (Bond et al. 2004;

only the seventh low-mass star with a Doppler-detected giant planet (see also Bonfils et al. 2005; Maness et al. 2007; Forveille et al. 2009; Mayor et al. 2009, for examples of low-mass planet detections). In the following section we describe the stellar properties of Gl 649, and our spectroscopic observations and Doppler-shift measurements. In § 3 we test the validity of our interpretation of the observed radial velocity variations by measuring the false-alarm probability and by examining our photometric measurements. We conclude in § 4 with a summary and discussion of Gl 649 b, and we place this latest exoplanet in context with other giant planets discovered around M dwarfs.

### 2. OBSERVATIONS AND ANALYSIS

## 2.1. Stellar Properties

Gl 649 (= HIP 83043) is an M1.5 dwarf with a *Hipparcos* parallax-based distance of 10.34  $\pm$  0.15 parsecs (van Leeuwen 2007), apparent magnitude V = 9.7, and absolute magnitude  $M_V =$  9.627  $\pm$  0.053 (ESA 1997). We use the broadband metallicity calibration of Johnson & Apps (2009) to estimate [Fe/H] =+0.08\pm0.06, and we adopt the stellar mass estimate provided by the Delfosse et al. (2000) K<sub>s</sub>-band mass-luminosity relationship, which gives  $M_{\star} = 0.54 \pm 0.05 \,\mathrm{M}_{\odot}$ . Using the infrared flux method, Alonso et al. (1996) give an effective temperature  $T_{\rm eff} = 3700 \pm 60 \,\mathrm{K}$ . Wright et al. (2004) measured the emission in the CaII H emission line relative to the stellar photosphere on the Mt. Wilson scale (Duncan et al. 1991) and give a median

Gould et al. 2006; Beaulieu et al. 2006; Dong et al. 2009, e.g.).



FIG. 3.— Periodogram analysis of our RV time series, which reveals a strong peak at P = 592 days. The dashed lines show the analytic false-alarm probability of a peak arrising from noise sampled at our times of observation.

"grand S" value of 1.55. This S value places the chromospheric activity of Gl 649 among the top 20% of nearby early M-type stars, as shown by Rauscher & Marcy (2006) and by Gizis et al. (2002). Our spectra show Halpha to be in absorption, as was found for all Balmer lines observed in the spectrum of Gl 649 (Gizis et al. 2002). The stellar properties of Gl 649 are summarized in Table 2.

#### 2.2. Radial Velocities and Keplerian Fit

We began monitoring Gl 649 at Keck Observatory in 1999 October using the High-Resolution Echelle spectrometer (Vogt et al. 1994) in our standard Iodine cell setup with the B5 decker, giving a reciprocal resolution  $\lambda/\Delta\lambda = 55,000$  per ~ 4-pixel resolution element (Howard et al. 2009). We measured the Doppler shifts of the star from each star-times-iodine observation using the standard analysis procedure presented by Butler et al. (1996), with subsequent improvements over the years. For HIRES observations made prior to the 2004 CCD upgrade the measurement uncertainties range from 3.3–4.4 m s<sup>-1</sup>, and improve to 0.9–1.5 m s<sup>-1</sup> thereafter.

Our 44 radial velocities are presented in Table 3 (without jitter) and the time series is shown in Figure 2 (with jitter). The scatter in the measurements is larger than expected from the measurement errors, and a periodogram analysis of the data reveals strong power at periods near 592 days, with a corresponding analytic falsealarm probability < 0.0001 (Figure 3).

We used the partially-linearized Keplerian fitting code  $\texttt{RVLIN}^{11}$  described by Wright & Howard (2009) to search for a best-fitting orbital solution to the data. To ensure proper weighting of our measurements in the fitting procedure, we inflated the error bars to account for RV noise from astrophysical sources. This stellar "jitter" term is calculated based on the star's chromospheric activity, B-V color and absolute V-band magnitude using the formula of Wright (2005). We adopt a jitter estimate

3

of 3 m s<sup>-1</sup> for Gl 649, which we add in quadrature to the measurement errors.

We find that a single-planet Keplerian model with a period  $P = 598.3 \pm 4.2$  days, eccentricity  $e = 0.30 \pm 0.08$  and velocity semiamplitude  $K = 12.4 \pm 1.1$  m s<sup>-1</sup> results in a root-mean-squared (rms) scatter of 4.2 m s<sup>-1</sup> in the residuals and  $\sqrt{\chi_{\nu}^2} = 1.17$ , indicating an acceptable fit<sup>12</sup>. The resulting minimum planet mass is  $M_P \sin i = 0.328$  M<sub>Jup</sub>, and the semimajor axis is a = 1.135 AU. The best-fitting solution is shown in Figure 2, with the residuals to the fit shown in the lower panel. The orbital parameters are listed in Table 2.

The parameter uncertainties given above were estimated using a Markov Chain Monte Carlo (MCMC) algorithm with  $10^7$  links, in which a single randomlychosen parameter was perturbed at each link, with a perturbation size tuned such that 20–40% of the jumps were executed (see e.g. Ford 2005; Winn et al. 2008, and references therein). The resulting "chains" of parameters form the posterior probability distribution, from which we select the 15.9 and 84.2 percentile levels in the cumulative distributions (CDF) as the "one-sigma" confidence limits. In most cases the posterior probability distributions were approximately Gaussian.

## 3. TESTING THE NULL-HYPOTHESIS

#### 3.1. False-Alarm Probability

The Doppler semi-amplitude of our best-fitting model,  $K = 12.4 \text{ m s}^{-1}$ , is comparable to the measurement uncertainties and stellar jitter, which prompted us to test the null-hypothesis that the apparent periodicity arose by chance from larger-than-expected radial velocity fluctuations and sparse sampling. We tested this possibility calculating the false-alarm probability (FAP) based on the goodness of fit statistic  $\Delta \chi^2_{\nu}$  (Howard et al. 2009; Marcy et al. 2005; Cumming 2004), which is the difference between two values of  $\chi^2_{\nu}$ : one from the single-planet Keplerian fit and one from the fit of a linear trend to the data. Each trial is constructed by keeping the times of observation fixed and scrambling the measurements, with replacement. We record the  $\Delta \chi^2_{\nu}$  value after each trial and repeat this process for 10,000 trial data sets. For the ensemble set we compare the resulting distribution of  $\Delta \chi^2_{\nu}$  to the value from the fit to the original data.

We found that none of the  $10^4$  trials resulted in a higher value of  $\Delta \chi^2_{\nu}$ , which we interpret as a < 0.0001 probability that the 600-day periodicity is a spurious signal due to random fluctuations.

### 3.2. Photometric Variability

We note that our FAP value only addresses the existence of a periodicity in the radial velocities, but does not test its cause. As an additional test of the null hypothesis we acquired brightness measurements of Gl 649 in the Johnson V passband with the T3 0.4 m automatic photometric telescope (APT) at Fairborn Observatory. The APT observations cover five observing seasons between 2001 February and 2009 June and reveal

<sup>&</sup>lt;sup>11</sup> http://exoplanets.org/code/

<sup>&</sup>lt;sup>12</sup> We use  $\sqrt{\chi_{\nu}^2}$  to indicate the factor by which the observed scatter about the best-fitting model differs from our expectation based on the measurement errors. Thus, the scatter about our model is a factor of 1.17 larger than our average error bar.



FIG. 4.— Top Panel: Our 337 V - C photometric observations of Gl 649 in the Johnson V band acquired with the T3 0.4m APT at Fairborn Observatory. Second Panel: The K - C observations plotted with an identical scale as the top panel. Comparison of these two data sets shows that Gl 649 varies in brightness on night-to-night and year-to-year timescales. Third Panel: Observations from the second observing season plotted on an expanded x axis clearly show low-amplitude brightness variability in Gl 649. Fourth Panel: Frequency spectrum of the observations from season two gives a best period of 24.55 days. Bottom Panel: Plot of the data from season two phased with the 24.55-day period reveals coherent variability with a peak-to-peak brightness amplitude 0.012 mag.

photometric variability in Gl 649 on both rotational and decadal timescales. Details on APT operations, data acquisition and reduction procedures, and precision of the observations can be found in Henry et al. (1995b,a); Fekel et al. (2005); Eaton et al. (2003).

Our 337 Gl 649-minus-comparison (V-C) differential magnitudes are plotted against heliocentric Julian Date in the top panel of Figure 4. The comparison star is HD 152342 (V = 7.10, B - V = 0.35, F2V). Most obvious in this plot are the year-to-year changes in the mean magnitude of the observations. The mean magnitudes have a range of 0.0126 mag and suggest the possible existence of a spot (magnetic) cycle in Gl 649 with a length of at least several years (see, e.g., Henry 1999; Hall et al. 2009). The top panel also shows that the range in the V-C observations is ~0.02 mag within all five observing seasons. The standard deviations of the individual five seasons are all between 0.0057 mag and 0.0063 mag.

The check minus comparison (K-C) differential mag-

nitudes are plotted in the second panel of Figure 4 at the same magnitude scale as the top panel. The check star is HD 153897 (V = 6.57, B - V = 0.43, F4V). The yearly means of the K - C measurements have a range of only 0.0016 mag; their standard deviation is only 0.0007 mag. The standard deviations of the individual five seasons are all between 0.0034 mag and 0.0042 mag. Thus, the larger night-to-night scatter of the V - C measurements and the observed year-to-year change in the V - C mean magnitudes must both be intrinsic to Gl 649.

The observations from season 2 are replotted in the third panel of Figure 4, again on the same magnitude scale as in panels 1 and 2. Low-amplitude variability is clearly seen with a period of 20–30 days. Panel 4 shows our computed frequency spectrum for season 2, where the y axis plots the reduction factor in the variance of the observations for each trial frequency (Vaníček 1971). We find a clear period of 24.55 days; the observations are plotted phased with this period in the bottom panel of Figure 4. A least-squares sine fit to the phase curve gives a peak-to-peak amplitude of  $0.0117 \pm 0.0015$  mag. Similar analyses yield periods of 23.72, 27.92, 25.80, and 21.86 days for seasons one, three, four, and five, respectively. The mean of these five periods is  $24.8 \pm 1.0$  days, which we take to be the rotation period of Gl 649 revealed by rotational modulation in the visibility of cool starspots on the photosphere of Gl 649. This photometric variability is consistent with the level of chromospheric activity in the star, as mentioned above.

Henry et al. (1995b) show many examples of active stars with low-amplitude starspot variability. Queloz et al. (2001) and Paulson et al. (2004) show several examples of stellar spots masquerading as planets. In the case of Gl 649 described here, the photometric observations reveal variability timescales that are inconsistent with the 600-day radial velocity variations. Sinilarly, we examined the time-variability chromospheric emission from each of our spectroscopic observations. While the S-value, as measured from the Ca II H&K emission, has a variance of 0.14 dex, we observed no peridicities near the putative orbital period. The lack of photometric and chromospheric variability provide additional strong support for the interpretation of planetary reflex motion as the cause of the observed radial velocity variability in Gl 649.

#### 4. DISCUSSION

We have presented the discovery of a Saturn-mass planet ( $M_P \sin i = 0.328 \text{ M}_{\text{Jup}}$ ) orbiting the nearby, low-mass star Gl 649 ( $d = 10.34 \text{ pc}, 0.54 \text{ M}_{\odot}$ ). Gl 649 b resides in an eccentric (e = 0.30) orbit with a period of 598.3 days, corresponding to a semimajor axis a = 1.135 AU.

Gl 649 is only the seventh M dwarf known with a Doppler-detected giant planet, and the fifth detection from among the 147 low-mass stars we've monitored over the past decade at Keck Observatory (e.g. Johnson et al. 2007). The low Doppler amplitude of the planet ( $K = 12.4 \text{ m s}^{-1}$ ) highlights our need to attain high measurement precision to find low-mass planets, and to maintain that precision over long time baselines to detect planets at larger semimajor axes.

Johnson et al. (2007) recently analyzed the detection rate among our sample of low-mass stars and reported a

1.8% occurrence rate of planets with  $a\,<\,2.5$  AU. The increased time baseline and new detections of our sample suggest that a reanalysis of the frequency of planets around M dwarfs is warranted. Following Johnson et al. we first note that the  $\approx 10$  year time baseline of our survey, together with our radial velocity precision, provides us with sensitivity to planets with  $M_P \sin i \gtrsim 0.3 \, \mathrm{M_{Jup}}$  out to semimajor axes  $a \approx 2.5$  AU, assuming and average stellar mass  $M_{\star} =$  $0.5 M_{\odot}$ . Note that in the analysis that follows, we exclude the recently detected planets Gl 832 b, which was discovered by the Anglo-Australian Observatory planet search (Bailey et al. 2008), and HIP 79431 b, which was only recently added to the Keck survey as part of the metallicity-biased M-to-K program (Apps et al. 2010, submitted).

The probability density function (PDF) for the fraction of stars with planets given our number of detections k = 5 and total sample size N = 147 is given by the binomial distribution  $P(f|k, N) \propto f^5(1-f)^{147-5}$ . The overall occurrence rate from our sample is given by the maximum of the PDF, which we measure to be  $f = 3.4^{+2.2}_{-0.9}\%$ , where the upper and lower limits represent the 68.3% confidence interval measured from the cumulative distribution function.

The corresponding giant planet fraction around Sunlike stars was recently measured by Cumming et al. (2008, cf their Table 1), who report  $f = 7.6 \pm 1.3\%$ . In a similar study Bowler et al. (2009, submitted) measured the planet fraction around stars with  $M_{\star} > 1.5 \text{ M}_{\odot}$  to be  $26^{+9}_{-8}\%$ , albeit for minimum masses  $M_P \sin i \gtrsim 1 \text{ M}_{\text{Jup}}$ . Thus, the detection rate of giant planets around M dwarfs consistently lags behind that of higher mass stars, despite the enhanced detectability of planets around less massive stars since  $K \propto M_{\star}^{-2/3}$  for a fixed planet mass and period. The contrast between the measured planet fractions between M dwarfs and massive stars points to an even stronger correlation between stellar mass and planet occurrence than measured by Johnson et al. (2007).

The correlation between stellar mass and planet formation is an important piece of observational evidence in support of the core accretion model of planet formation. In this model, giant planets form in a bottomup process, starting with the collisions of small dust grains and proceeding up through the formation of large protoplanetary cores (see Ida & Lin 2004; Alibert et al. 2005, for reviews). Once these cores attain a critical mass of ~ 10  $M_{\oplus}$ , they can rapidly accrete gas from the surrounding disk. Given the limited lifetime of the gas disk, which dissipates on timescales less than 5 Myr (Hernández et al. 2008; Currie et al. 2009), the formation of gas giant planets is a race against time that is rarely won in the protoplanetary disks of low-mass stars. The low density of raw materials, low orbital frequencies  $(\Omega \propto 1/P \propto M_{\star}^{1/2}$  at fixed *a*), and unfavorable temperature profiles in the disks around M-type stars greatly inhibit the core growth, which results in a lower occurrence of giant planets (Laughlin et al. 2004; Ida & Lin 2005; Kennedy & Kenyon 2008; Dodson-Robinson et al. 2009).

Another important predictor of planet occurrence is stellar metallicity. Fischer & Valenti (2005b) showed the fraction of Sun-like stars with planets correlates strongly with [Fe/H], with an occurrence rate of ~ 3% for [Fe/H] < 0 and a rise to  $\approx 25\%$  for [Fe/H] > +0.3. Until recently, it was difficult to properly account for metallicity among the M dwarfs because the LTE spectral analysis tools used for more massive stars are not amenable to the complex spectra of low-mass stars (Maness et al. 2007). Because of the lack of knowledge about the metallicity distribution of M dwarfs in general, and low-mass stars with planets in particular, it was difficult to determine whether stellar mass or metallicity lay at the root cause of the puacity of planets around M dwarfs.

The mass/metallicity issue was recently addressed by Johnson & Apps (2009), who derived a revised broadband photometric metallicity calibration for M dwarfs. They examined a sample of M dwarfs with F, G and K wide binary companions. By anchoring the metallicity of the M dwarf to its earlier-type companion, Johnson & Apps observed that metal-rich M stars reside "above" the mean main sequence of the solar neighborhood when viewed in the  $\{V - K, M_K\}$  plane. Further, they noticed the majority of the 7 planetary systems (containing planets of all masses) that were known at the time contain metal-rich host stars.

Figure 1 shows that with the addition of 3 new planethost stars since the study of Johnson & Apps the planetmetallicity correlation among M dwarfs appears to persist. We can quantify this relationship by examining the fraction of stars in our Keck survey with  $[Fe/H] \ge 0$ that harbor giant planets. Including Gl 649, we find that all 4 of the stars that harbor at least one giant planet<sup>13</sup> fall within the subsamlple of 80 targets with  $[Fe/H] \ge 0$ . Based on this, we measure a planet fraction  $f = 5.5^{+2.7}_{-2.1}\%$  for [Fe/H] > 0. If we restrict our analysis to [Fe/H] > +0.2 (dashed line in Figure 1), 3 of these 33 "super-metal-rich" stars harbor planets, corresponding to  $f = 10.7^{+5.9}_{-4.2}\%$ .

The uncertainties in our measured planet fractions are large due to the small sample sizes involved. This underscores the need for extending the time baseline of our current survey and expanding the target list to include additional low-mass stars. Future surveys of nearby, low-mass stars such as the M2K planet search (Apps et al. 2010, submitted) and the *MEarth* transit survey (Irwin et al. 2009) will build upon the current sample and provide a clearer picture of the planet-metallicity relationship suggested from our analysis. A larger sample of planets detected around M dwarfs will also provide crucial leverage in understanding the relationship between stellar mass and planet properties, especially when compared to the growing sample of planets discovered around massive stars with  $M_{\star} > 1.5 \, \mathrm{M}_{\odot}$ .

We thank the many observers who contributed to the velocities reported here. We gratefully acknowledge the efforts and dedication of the Keck Observatory staff, especially Grant Hill and Scott Dahm for support of HIRES

<sup>&</sup>lt;sup>13</sup> We exclude Gl 317 in this analysis because it lacks a reliable parallax measurement and cannot be accurately placed in the  $\{V - K, M_K\}$  plane. The existing parallaxes given in the literature suggest that Gl 317 resides above the main sequence  $\Delta M_K = 0.08-0.37$ , indicating either a Solar composition or that the star is metal-rich  $(0.0 \leq [\text{Fe}/\text{H}] \leq +0.15)$ .

and Greg Wirth for support of remote observing. We are also grateful to the time assignment committees of NASA, NOAO, and the University of California for their generous allocations of observing time. We acknowledge R. Paul Butler and S. S. Vogt for many years of contributing to the data presented here. A. W. H. gratefully acknowledges support from a Townes Post-doctoral Fellowship at the U.C. Berkeley Space Sciences Laboratory. J.A.J. thanks the NSF Astronomy and Astrophysics Postdoctoral Fellowship program for support in the years leading to the completion of this work, and acknowl-

- Alibert, Y., et al. 2005, A&A, 434, 343
- Alonso, A., Arribas, S., & Martinez-Roger, C. 1996, A&AS, 117, 227
- Bailey, J., et al. 2008, arxiv:0809.0172
- -. 2009, ApJ, 690, 743
- Beaulieu, J.-P., et al. 2006, Nature, 439, 437
- Bond, I. A., et al. 2004, ApJ, 606, L155
- Bonfils, X., et al. 2005, A&A, 443, L15
- Butler, R. P., et al. 2006, PASP, 118, 1685
- Butler, R. P., et al. 1996, PASP, 108, 500
- Cumming, A. 2004, MNRAS, 354, 1165
- Cumming, A., et al. 2008, PASP, 120, 531
- Currie, T., et al. 2009, ApJ, 698, 1
- Delfosse, X., et al. 2000, A&A, 364, 217
- Dodson-Robinson, S. E., et al. 2009, ApJ, 707, 79
- Dong, S., et al. 2009, ApJ, 695, 970
- Duncan, D. K., et al. 1991, ApJS, 76, 383
- Eaton, J. A., Henry, G. W., & Fekel, F. C. 2003, The Future of Small Telescopes In The New Millennium. Volume II - The Telescopes We Use, 189
- Endl, M., et al. 2003, AJ, 126, 3099
- ESA, . 1997, VizieR Online Data Catalog, 1239, 0
- Fekel, F. C., Henry, G. W., & Lewis, C. 2005, AJ, 130, 794
- Fischer, D. A. & Valenti, J. 2005a, ApJ, 622, 1102
- -. 2005b, ApJ, 622, 1102
- Ford, E. B. 2005, AJ, 129, 1706
- Forveille, T., et al. 2009, A&A, 493, 645
- Gizis, J. E., Reid, I. N., & Hawley, S. L. 2002, AJ, 123, 3356
- Gould, A., et al. 2006, ApJ, 644, L37
- Hall, J. C., et al. 2009, AJ, 138, 312
- Henry, G. W. 1999, PASP, 111, 845
- Henry, G. W., et al. 1995a, ApJS, 97, 513
- Henry, G. W., Fekel, F. C., & Hall, D. S. 1995b, AJ, 110, 2926

edges support form NSF grant AST-0702821. G.W.M. acknowledges NASA grant NNX06AH52G. J.T.W. received support from NSF grant AST-0504874. G.W.H acknowledges support from NASA, NSF, Tennessee State University, and the State of Tennessee through its Centers of Excellence program. Finally, the authors wish to extend special thanks to those of Hawaiian ancestry on whose sacred mountain of Mauna Kea we are privileged to be guests. Without their generous hospitality, the Keck observations presented herein would not have been possible.

## REFERENCES

- Hernández, J., et al. 2008, ApJ, 686, 1195
- Howard, A. W., et al. 2009, ApJ, 696, 75
- Ida, S. & Lin, D. N. C. 2004, ApJ, 604, 388
- . 2005, ApJ, 626, 1045
- Irwin, J., et al. 2009, ApJ, 701, 1436
- Irwin, J., et al. 2008, arxiv:0807.1316
- Johnson, J. A. & Apps, K. 2009, ApJ, 699, 933
- Johnson, J. A., et al. 2007, ApJ, 670, 833
- Kennedy, G. M. & Kenyon, S. J. 2008, ApJ, 673, 502
- Laughlin, G., Bodenheimer, P., & Adams, F. C. 2004, ApJ, 612, L73
- Maness, H. L., et al. 2007, PASP, 119, 90
- Marcy, G., et al. 2005, Progress of Theoretical Physics Supplement, 158, 24
- Mayor, M., et al. 2009, ArXiv e-prints
- Nielsen, E. L. & Close, L. M. 2009, ArXiv:0909.4531
- Paulson, D. B., et al. 2004, AJ, 127, 1644
- Pravdo, S. H. & Shaklan, S. B. 2009, ApJ, 700, 623
- Queloz, D., et al. 2001, A&A, 379, 279
- Rauscher, E. & Marcy, G. W. 2006, PASP, 118, 617
- Rivera, E. J., et al. 2005, ApJ, 634, 625
- van Leeuwen, F. 2007, A&A, 474, 653 Vaníček, P. 1971, Ap&SS, 12, 10
- Vogt, S. S., et al. 1994, in Proc. SPIE Instrumentation in Astronomy VIII, David L. Crawford; Eric R. Craine; Eds. Volume 2198, p. 362, ed. D. L. Crawford & E. R. Craine, 362-+
- Winn, J. N., et al. 2008, ApJ, 683, 1076
- Wright, J. T. 2005, PASP, 117, 657
- Wright, J. T. & Howard, A. W. 2009, ApJS, 182, 205
- Wright, J. T., et al. 2004, ApJS, 152, 261

TABLE 1 Properties of M Dwarfs with Giant Planets

Gliese Number	<i>Hipparcos</i> Number	Spectral Type	Stellar Mass $(M_{\odot})$	[Fe/H]	$M_P \sin i ~(\mathrm{M_{Jup}})$	Semimajor Axis (AU)	Reference
876	113020	M4	0.32	+0.37	$0.6189,  1.9275^{\rm a}$	0.207,  0.130	Rivera et al. (2005)
849	109388	M3	0.45	+0.58	$0.83^{\mathrm{b}}$	2.35	Butler et al. (2006)
317		M4	0.24		1.17	0.95	Johnson et al. $(2007)$
832	106440	M2	0.45	-0.12	0.64	3.4	Bailey et al. (2009)
179	22627	M4	0.36	+0.30	0.9	2.42	Howard et al. $(2009)$
	79431	M3.5	0.50	+0.4	1.1	0.34	Apps et al. (2010, submitted)
649	83043	M1.5	0.54	+0.1	0.328	1.135	This work

<sup>a</sup> The Gl 876 planetary system contains two resonant Jovian planets and an inner "super-Earth" with  $M_P \sin i = 5.9 M_{\oplus}$ . <sup>b</sup> The orbit solution for Gl 849 includes a linear velocity trend  $dv/dt = -4.7 \text{ m s}^{-1} \text{ yr}^{-1}$ , which may correspond to a second

planet. <sup>c</sup> The orbit solution for Gl317 includes a linear velocity trend  $dv/dt = 7.6 \text{ m s}^{-1} \text{ yr}^{-1}$ , which may correspond to a second planet.

 TABLE 2

 Stellar Properties and Orbital

 Solution for GL 649

Parameter	Value
V	$9.70 \pm 0.04$
K	$5.62\pm0.02$
V - K	$4.08\pm0.05$
B - V	$1.52 \pm 0.04$
$M_V$	$9.63 \pm 0.05$
$M_K$	$5.55 \pm 0.02$
d (pc)	$10.34 \pm 0.15$
$M_{\star}  (\mathrm{M}_{\odot})$	$0.54 \pm 0.05$
$T_{eff}$ (K)	$3700 \pm 60$
[Fe/H]	$+0.08 \pm 0.06$
P (days)	$598.3 \pm 4.2$
P (years)	$1.638 \pm 0.011$
$K ({\rm ms^{-1}})$	$12.4 \pm 1.1$
e	$0.30\pm0.08$
$T_P$ (Julian Date-2400000)	$12876 \pm 22$
$\omega \text{ (degrees)}$	$352 \pm 15$
$M_P \sin i \ (M_{Jup})$	$0.328 \pm 0.032$
a (AU)	$1.135 \pm 0.035$
N <sub>obs</sub>	44
$rms (m s^{-1})$	4.2
$\sqrt{\chi_{\mu}^2}$	1.17
$\chi^2_{\mu}$	1.37

JD	RV	Uncertainty
-2440000	$(m \ s^{-1})$	$(m \ s^{-1})$
11409 824	-1 54	3 35
11705 913	25.38	3.80
12004.048	-8.26	3.73
12007.010	-10.05	4.01
12008.001	-11.88	3.92
12009.070	-2.72	3.65
12097.968	-8.16	3.60
12098.912	1.04	3.83
12099.844	4.90	3.63
12100.895	4.57	3.39
12127.911	0.00	3.72
12161.808	5.87	3.89
12189.748	2.16	4.36
12390.092	13.12	3.74
12486.759	-1.86	3.46
12514.750	-5.61	3.50
12030.722	-0.90	3.37
12074.090 19719.157	2.29	4.02
12712.137	-5.05 7.53	3.03
12804 920	13.40	3 75
12832 961	13.40 13.60	3.91
13179.964	-9.51	3.82
13195.812	0.17	3.38
13430.124	13.32	1.20
13547.904	12.84	0.99
13842.077	-3.95	0.97
13934.875	9.44	0.95
14640.044	14.79	1.16
14671.928	16.81	0.72
14673.897	25.43	0.58
14779.698	8.75	1.14
14964.087	4.24	0.96
14985.917	-3.91	0.96
15014.851	-2.75	0.98
15010.020	-7.25	0.97
15041.050	-2.00	1.01
15041.990	-9.03	1.00
15042.100	-9.45	1.00
15048.857	-3.67	1.28
15075.744	4.40	0.91
15111.736	-2.10	1.07

TABLE 3 RADIAL VELOCITIES FOR GL 649