## Technical aspects and dark matter searches

R Bernabei <sup>1,2</sup>, P Belli <sup>2</sup>, F Cappella <sup>3,4</sup>, R Cerulli <sup>5</sup>, C J Dai <sup>6</sup>, A d'Angelo <sup>3,4</sup>, H L He <sup>6</sup>, A Incicchitti <sup>4</sup>, H H Kuang <sup>6</sup>, X H Ma <sup>6</sup>, F Montecchia <sup>2,7</sup>, F Nozzoli <sup>1,2</sup>, D Prosperi<sup>3,4</sup>, X D Sheng<sup>6</sup> and Z P Ye<sup>6,8</sup>

- <sup>1</sup> Dip. di Fisica, Università di Roma "Tor Vergata", I-00133 Rome, Italy
- <sup>2</sup> INFN, sez. Roma "Tor Vergata", I-00133 Rome, Italy
- $^3$  Dip. di Fisica, Università di Roma "La Sapienza", I-00185 Rome, Italy
- <sup>4</sup> INFN, sez. Roma, I-00185 Rome, Italy
- <sup>5</sup> Laboratori Nazionali del Gran Sasso, I.N.F.N., Assergi, Italy
- $^6$  IHEP, Chinese Academy, P.O. Box 918/3, Beijing 100039, China
- <sup>7</sup> Laboratorio Sperimentale Policentrico di Ingegneria Medica, Università degli Studi di Roma "Tor Vergata"
- <sup>8</sup> University of Jing Gangshan, Jiangxi, China

E-mail: rita.bernabei@roma2.infn.it

Abstract. A variety of detectors has been proposed for dark matter direct detection, but most of them – by the fact – are still at R&D stage. In many cases, it is claimed that the lack of an adequate detectors' radio-purity might be compensated through heavy uses of MonteCarlo simulations, subtractions and handlings of the measured counting rates, in order to claim higher sensitivity (just for a particular scenario). The relevance of a correct evaluation of systematic effects in the use of MonteCarlo simulations at very low energy (which has always been safely discouraged in the field so far) and of multiple subtractions and handling procedures applied to the measured counting rate is shortly addressed here at some extent. Many other aspects would also deserve suitably deep investigations.

In this paper some arguments presented at the TAUP09 conference will be shortly summarized. More details, tables and figures can be found in the slides at the conference site[1].

Let us firstly comment the possibility of reliable evaluations of the background contributions at the keV energy region in the field of Dark Matter searches. As well known, it has been generally discouraged this procedure in the field of Dark Matter over more than twenty years. In fact, the estimation by a MonteCarlo simulation of the background component in the counting rate from the residual radioactivity requires a detailed knowledge of: i) the exact set-up geometry (detector or detectors' matrix, all materials, details of the assembling, of the shield layers, of the site, etc.); ii) the detector response function (e.g. energy resolution,  $\alpha/\beta$  ratio, channeling, etc.); iii) the nature, the position and the concentration of all the existing radioactive contaminants; iv) etc.. Unfortunately, apart from the geometrical layout of the set-up that are generally well known by people inside the experimental group, all the other quantities necessary in the MonteCarlo simulation require dedicated measurements. Moreover, there are some quantities (such as concentration of residual contaminants, etc.) that can be poorly known and just upper/lower limits are available; in some cases these quantities can be even totally unknown. As an example, the experimental energy resolution as a function of the energy and the energy scale

should be measured/verified down to the energy threshold, as done e.g. by the DAMA/LIBRA experiment where they are continuously measured by external/internal known sources from MeV down to the energy threshold [2]. On the contrary, in other experiments these quantities are instead extrapolated from calibrations at much higher energy (as done e.g. by liquid noble gas set-ups, where the energy threshold and the few keV energy scale are generally unproven, also because of position dependence, of non-uniform signal collection, etc.; see e.g. [3]). Regarding the presence of residual contaminants in the set-up, generally only limits on the contributions of the "standard" contaminants are given; these limits forbid any reliable estimation of the background (being unknown the exact values) and cannot be obviously exceeded (see also later). Moreover, possible presence of many non-standard contaminants should be also included. In addition, the MonteCarlo simulation also depends on the precise location of all the contaminants – that is generally unknown even for the "standard" ones - in complex set-ups. The situation is more complex for multi-detectors set-up and when the energy distribution refers to events where each detector has all the others in anticoincidence (single-hit events). Thus, it is trivial to conclude that a reliable precise simulation of the background counting rate – in particular at keV energy region – is not univocally determined and is a quite impossible task.

In addition, beyond the fore-mentioned arguments, we need to take into account that a MonteCarlo code cannot manage all the possible low energy atomic physical processes. This argument is still subject of improvements; in fact, as an example, non-negligible differences are also obtained by different versions of the same MonteCarlo code [4].

Some instructive examples are given by the trails in MonteCarlo simulations in ref. [5]. As shown there, these simulations noticeably differ from the measured energy distributions in the cases of XENON-10 and of ZEPLIN-III. In fact, they predict twice the measured rate for XENON-10 near 200 keV, and more than one order of magnitude the measured rate for ZEPLIN-III in the MeV range [5]. The same approach has been also pursued by the same authors [5] trying a MonteCarlo simulation of the background in the DAMA/LIBRA set-up. In particular, apart from errors in the details of the set-up geometry reconstruction and in the multiple-hit definition, many crude and arbitrary approximations in the nature and in the location of the residual contaminants have been arbitrarily assumed; in fact, e.g.: i) only standard contaminants, ii) only unbroken chains, iii) only uniform location of contaminants in the detectors, etc. have been taken into account. As a result of this rough, partial and arbitrary approach, the predicted rate has been estimated within a factor 10 lower than the measured one. Instead of refining the quality of the simulation or reasonably recognizing the impossibility of precise determination, the authors just pursued the exercise of arbitrarily increasing "by hand" the assumed values of the contaminants at levels much larger than the measured experimental limits [5]. This also implies an overestimate of the background in higher energy region with respect to the measured experimental rate. In conclusion, although the arbitrary and the erroneous adopted procedures, these authors do not succeed in reproducing either the low or the high energy spectra. Nevertheless, as a conclusion of this arbitrary exercise [5], this artificiallyboosted simulated spectrum has been subtracted by the measured one, attempting to obtain a limit for the unmodulated Dark Matter signal component. This example shows how subtraction procedures using MonteCarlo simulations in the few keV energy region can give rise to erroneous conclusions; thus, any constraint on Dark Matter signal on this basis would be an artefact.

Furthermore, let us also note that the measured spectra e.g. of the existing/past NaI(Tl) detectors (such as e.g. ANAIS, Frejus, NAIAD, ELEGANT, etc...) do not support even the shape presented in ref. [5]. In addition, well different counting rates at keV energy region are present even for detectors of the same experimental group, as e.g. the case of NAIAD in 1996 and in 2003 [1]. In conclusion, it does not exist an unique recipe for a precise and reliable MonteCarlo simulation of whatever set-up, and for NaI(Tl) in particular.

Let us finally remind that a safer approach has been presented in this conference by DAMA

collaboration; this shows that enough space is present in the measured counting rate of the DAMA/LIBRA keV energy spectrum for the unmodulated component of Dark Matter signal [1, 6].

In the second part of this contribution [1], problems related to the application of multiple subtraction procedures of the measured counting rate, as pursued by experiments trying to identify the presence of recoil nuclei in the measured energy spectrum, have been summarized. In fact, many of the existing Dark Matter candidates – also in the WIMP class – can give rise to signals that either have totally an electromagnetic nature (see e.g. [7, 8, 9]) or involve electromagnetic signals associated to nuclear recoils (see e.g. [10, 11]); obviously, approaches that are based on multiple subtraction procedure of the electromagnetic component of the counting rate are blind to similar scenarios. Moreover, well known side processes exist for recoils (such as recoils induced by neutrons, fission fragments, end-range alphas, surface electrons, etc.). This approach is generally pursued when the detectors suffer from a not-suitable radiopurity level in the sensitive target-material and in the surroundings.

Those activities generally apply a large number of cut procedures to the data; each one is affected by non-negligible systematic errors which are usually not suitable quantified. As an example, the XENON-10 experiment applies more than 10 different cuts to the data; the experiment collects  $\sim 10^4$  events but only 10 are claimed to survive to the cuts and handling procedures [12]. Very high reduction factors following applied multiple cuts are dangerous because of the difficult precise estimate of all the involved systematics. For example, it has been shown in ref. [13] that ZEPLIN-I has claimed a sensitivity 3 orders of magnitude larger than the one properly obtained when accounting for systematics. Thus, the robustness of some results appeared in the "race for the best exclusion-plot" (valid just in a single set of assumptions for a certain kind of WIMP) should be considered "cum grano salis". For example, for some liquid noble gas set-ups, apart from the robustness of the applied cuts themselves, the very low energy scale and the energy threshold are determined by extrapolating from calibrations at much higher energy and applying some kind of corrections for relevant non-uniformity of the detector's response; with a light yield of about 2.2 photoelectrons/keV and not specific calibrations even  $\approx 1.5-2$  keV electron equivalent is claimed as energy threshold [12, 1]. Considering the energy threshold dependence of the exclusion plots, up to several orders of magnitude differences can be present between claimed and realistic evaluation of the experimental sensitivity.

Another crucial aspect is the proper accounting for the existing experimental and theoretical uncertainties in the calculation and in the comparison of experiments using different target materials and approaches.

## References

- [1] F. Nozzoli, presentation given at this conference (TAUP 2009) http://taup2009.lngs.infn.it/slides/jul3/nozzoli.pdf or http://people.roma2.infn.it/~dama/pdf/nozzoli\_taup09.pdf.
- R. Bernabei et al., Nucl. Instr. & Meth. A 592 (2008) 297.
- [3] R. Bernabei et al., ISBN 978-88-95688-12-1, pages 1-53 (2009) Exorma ed. (arXiv:0806.0011[astro-ph]).
- [4] see e.g. R.M. Kippen, New Astr. Rev. 48 (2004) 221.
- [5] V. Kundryavtsev, presentation given at this conference (TAUP 2009); V. Kudryavtsev et al., arXiv:0912.2983v1.
- [6] R. Bernabei et al., Eur. Phys. J. C 56 (2008) 333.
- [7] R. Bernabei et al., Int. J. Mod. Phys. A 21 (2006) 1445.
- [8] R. Bernabei et al., Phys. Rev. D 77 (2008) 023506.
- [9] R. Bernabei et al., Mod. Phys. Lett. A 23 (2008) 2125.
- [10] R. Bernabei et al., Int. J. Mod. Phys. A 22 (2007) 3155.
- [11] R. Bernabei et al., Eur. Phys. J. C 53 (2008) 205.
- [12] J. Angle et al., Phys. Rev. Lett. 100 (2008) 021303.
- [13] A. Benoit et al., Phys. Lett. B 637 (2006) 156.