Noisy non-transitive quantum games

M. Ramzan, Salman Khan and M. Khalid Khan Department of Physics Quaid-i-Azam University Islamabad 45320, Pakistan (Dated: November 19, 2018)

We study the effect of quantum noise in 3×3 entangled quantum games. By considering different noisy quantum channels we analyze that how a two-player, three-strategy Rock-Scissor-Paper game is influenced by the quantum noise. We consider the winning non-transitive strategies R, S and P such as R beats S, S beats P, and P beats R. The game behaves as a noiseless game for maximum value of the quantum noise parameter. It is seen that Alice's payoff is heavily influenced by the depolarizing noise as compared to the amplitude damping noise. Depolarizing channel causes a monotonic decrease in players payoffs as we increase the amount of of quantum noise. In case of amplitude damping channel, the Alice's payoff function reaches its minimum for $\alpha = 0.5$ and is symmetrical. This means that larger values of quantum noise influence the game weakly. On the other hand, phase damping channel does not influence the game's payoff. Furthermore, the game's Nash equilibrium and non-transitive character of the game are not affected under the influence of quantum noise.

Keywords: Quantum noise; quantum channels; quantum games.

I. INTRODUCTION

In the recent past, rapid interest has been developed in the discipline of quantum information [1] that has led to the creation of quantum game theory [2-5]. During last few years, tremendous efforts have been made towards the development of quantum game theory [6-12]. In this connection, much work has been devoted to convert the classical games into quantum domain such as prisoners' dilemma game [13-15] and many other games [16-20] involving two and three players. Quantum games with 3×3 payoff matrices and larger have have been discussed by Wang et al. [21]. Recently,

Sousa et al. [22] have proposed that quantum games can be used to control the access of processes to the CPU in a quantum computer. More recently, Iqbal and Abbott [23] have reported that quantum games can directly be constructed from a system of Bell's inequalities using Arthur Fine's analysis.

In quantum information processing, the major problem is to faithfully transmit unknown quantum states through a noisy quantum channel. When quantum information is sent through a channel, the carriers of the information interact with the channel and get entangled with its many degrees of freedom. This gives rise to the phenomenon of decoherence on the state space of the information carriers. In real-world applications, the decoherence effects caused by the external environment are inevitable. Quantum channels [1] provide a natural theoretical framework for the study of decoherence in noisy quantum communication systems. Quantum games in the presence of decoherence have been discussed in the recent past by various authors [5, 24-26]. Recently, decoherence and correlated noise (memory) effects have been analyzed in different quantum games [8, 20, 27]. Quantum error correction [28-29] and entanglement purifications [30].can be employed to avoid the problem of decoherence.

In this paper, we study the effect of quantum noise in a 2-player 3-strategy entangled quantum game (*RSP* game). We consider different noisy channels parameterized by a quantum noise parameter α such that $\alpha \in [0, 1]$. The lower and upper limits of quantum noise parameter represent the fully coherent and fully decohered systems, respectively. It is seen that for maximum value of quantum noise parameter the game behaves as a noiseless game. The depolarizing channel influences the game's payoff more heavily as compared to the amplitude damping channel. The payoff function for amplitude damping channel, is symmetrical with its minimum at $\alpha = 0.5$. Furthermore, the phase damping channel does not influence the game.

II. NOISY QUANTUM RSP GAME

The rock, scissors and paper game is a game for two players typically played using the players' hands. It is a simple two-player, three-strategy game, the children's choosing game "rock (R), scissors (S) and paper (P)" denoted by RSP, in which rock beats scissors, scissors beats paper, and paper beats rock (R > S > P > R). The classical payoff matrix for this game is given in table 1. Since RSP game is a zero-sum game, therefore it has no pure strategy Nash equilibrium. However, the mixed strategy Nash equilibrium produces zero expected payoff for the classical game.

Quantum analog of the RSP game was formulated by Iqbal et al. [12] and Michael et al. [31].

In the quantum version of this game, the strategies R, S, and P are represented by three matrices $(U_1, U_2 \text{ and } U_3 \text{ equation 9 Ref. [31]})$. U(x, y), is the unitary matrix which is used to transform the initial qutrit state of the game. We assume that initially Alice and Bob share a maximally entangled two qutrit state of the form

$$|\Psi_{in}\rangle = \frac{1}{\sqrt{3}} \left(|00\rangle + |11\rangle + |22\rangle\right) \tag{1}$$

We can define the strategies of the players by the unitary operator U(x, y) of the form [31]

$$U(x,y) = \frac{1}{\sqrt{2}} \begin{bmatrix} e^{ix} \cos y & ie^{ix} \sin y & 0\\ i \sin y \cos x & \cos x \cos y & ie^{iy} \sin x\\ -\sin y \sin x & i \sin x \cos y & e^{iy} \cos x \end{bmatrix}$$
(2)

where $0 \le \{x, y\} \le \pi/2$ and the choice of x and y define a player's mixed strategy in a 3×3 game, just as the choice of θ does in the 2×2 game [3].

The interaction between the system and its environment introduces the decoherence to the system, which is a process of the undesired correlation between the system and the environment when the system evolves. The evolution of a state of a quantum system in a noisy environment can be described by the super-operator Φ in the Kraus operator representation [1] as

$$\rho_f = \Phi \rho_i = \sum_k E_k \rho_i E_k^{\dagger} \tag{3}$$

where the Kraus operators E_i satisfy the following completeness relation

$$\sum_{k} E_{k}^{\dagger} E_{k} = I \tag{4}$$

We have constructed the Kraus operators for the game from the single qutrit Kraus operators (as given in equations (6-9) below) by taking their tensor product over all n^2 combination of $\pi(i)$ indices

$$E_k = \underset{\pi}{\otimes} e_{\pi(i)} \tag{5}$$

where n is the number of Kraus operators for a single qutrit channel. The single qutrit Kraus operators for the amplitude damping channel are given by [32]

$$E_{0} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{1 - \alpha} & 0 \\ 0 & 0 & \sqrt{1 - \alpha} \end{pmatrix}, \quad E_{1} = \begin{pmatrix} 0 & \sqrt{\alpha} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E_{2} = \begin{pmatrix} 0 & 0 & \sqrt{\alpha} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
(6)

Similarly, the single qutrit Kraus operators for the phase damping channel are given as [32]

$$E_{0} = \sqrt{1 - \alpha} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad E_{1} = \sqrt{\alpha} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^{2} \end{pmatrix},$$
(7)

The single qutrit Kraus operators for the depolarizing channel are given by [33]

$$E_{0} = \sqrt{1 - \alpha}I, \ E_{1} = \sqrt{\frac{\alpha}{8}}Y, \ E_{2} = \sqrt{\frac{\alpha}{8}}Z, \ E_{3} = \sqrt{\frac{\alpha}{8}}Y^{2}, \ E_{4} = \sqrt{\frac{\alpha}{8}}YZ$$
$$E_{5} = \sqrt{\frac{\alpha}{8}}Y^{2}Z, \ E_{6} = \sqrt{\frac{\alpha}{8}}YZ^{2}, \ E_{7} = \sqrt{\frac{\alpha}{8}}Y^{2}Z^{2}, \ E_{8} = \sqrt{\frac{\alpha}{8}}Z^{2}$$
(8)

where

$$Y = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad Z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}$$
(9)

In the above equations α represents the quantum noise parameter and $\omega = e^{\frac{2\pi i}{3}}$. The final state of the game after the action of the channel can be written as

$$\rho_f = \Phi_\alpha(|\Psi_{in}\rangle \langle \Psi_{in}|) \tag{10}$$

where Φ_{α} is the super-operator realizing the quantum channel parametrized by the real number α (quantum noise parameter). After the action of the players unitary operations, the game's final state transforms to

$$\rho_f = (U_A(x,y) \otimes U_B(x,y))(\Phi_\alpha(|\Psi_{in}\rangle \langle \Psi_{in}|)(U_A^{\dagger}(x,y) \otimes U_B^{\dagger}(x,y))$$
(11)

The payoff operators for Alice and Bob can be written as

$$(P^{A,B})_{\text{Oper.}} = \$_{00}^{A,B} P_{00} + \$_{01}^{A,B} P_{01} + \$_{02}^{A,B} P_{02} + \$_{10}^{A,B} P_{10} + \$_{11}^{A,B} P_{11} + \$_{12}^{A,B} P_{12} + \$_{20}^{A,B} P_{20} + \$_{21}^{A,B} P_{21} + \$_{22}^{A,B} P_{22}$$
(12)

where $\$_{ij}^{A,B}$ are the elements of payoff matrix in the *i*th row and *j*th column of classical game as given in table 1 and

$$P_{00} = |00\rangle \langle 00|, P_{01} = |01\rangle \langle 01|, P_{02} = |02\rangle \langle 02|$$

$$P_{10} = |10\rangle \langle 10|, P_{11} = |11\rangle \langle 11|, P_{12} = |12\rangle \langle 12|$$

$$P_{20} = |20\rangle \langle 20|, P_{21} = |21\rangle \langle 21|, P_{22} = |22\rangle \langle 22|$$
(13)

The players payoffs can be calculated by using the relation

$$\$^{A,B}(x_i, y_i, \alpha) = \text{Tr}\{[(P^{A,B})_{\text{Oper.}}\rho_f]^{A,B}\}$$
(14)

where Tr represents the trace of the matrix.

A. Results using different noise models

By using equations (1-6, 11-13 and 17), the players payoffs for the amplitude damping channel can be obtained as

$$\begin{split} \$_{AD}^{4,B}(x_{i},y_{i},\alpha) &= \frac{1}{12} [(2 + \alpha^{2})\$_{00} + \{2 + \alpha(-4 + 5\alpha)\}\$_{00} \cos(2y_{1}) \cos(2y_{2}) \\ &+\alpha(2 + \alpha)\$_{00} (\cos(2y_{1}) + \cos(2y_{2})) + 2(-1 + \alpha) \times \\ [(-\$_{11} + \$_{12} + \$_{21} - \$_{22}) \cos(y_{1} + y_{2}) \\ &\sin(2x_{1}) \sin(2x_{2}) \{(-1 + \alpha) \cos(2y_{1}) \cos(2y_{2}) + \sin(y_{1}) \sin(y_{2})\} \\ &+\$_{00} \sin(2y_{1}) \sin(2y_{2})\} + \cos(x_{1})^{2} [-4(-1 + \alpha) \alpha\$_{20} \cos(y_{2})^{2} \\ &-\$_{10} \cos(2y_{1}) \{\alpha(2 + \alpha) + (2 + \alpha(-4 + 5\alpha)) \cos(2y_{2})\} \\ &+\$_{00} \{2 + \alpha^{2} + \alpha(2 + \alpha) \cos(2y_{2}) - 2(-1 + \alpha) \sin(2y_{1}) \sin(2y_{2})\}] \\ &+\cos(x_{2})^{2} [(2 + \alpha^{2})\$_{01} + \alpha(2 + \alpha)\$_{01} (\cos(2y_{1}) - \cos(2y_{2})) \\ &+ \{-2 + (4 - 5\alpha) \alpha\}\$_{01} \cos(2y_{1}) \cos(2y_{2}) - 2(-1 + \alpha)(2\alpha\$_{02} \cos(y_{1})^{2} \\ &+\$_{01} \sin(2y_{1}) \sin(2y_{2})) + \cos(x_{1})^{2} \{(2 + \alpha^{2})\$_{11} + 4(-1 + \alpha)^{2}\$_{22} \\ &+ (2 + \alpha(-4 + 5\alpha))\$_{11} \cos(2y_{1}) \cos(2y_{2}) - \alpha(2 + \alpha)\$_{11} (\cos(2y_{1}) + \cos(2y_{2})) \\ &-2(-1 + \alpha)(2\alpha\$_{12} \sin(y_{1})^{2} + 2\alpha\$_{21} \sin(y_{2})^{2} - \$_{11} \sin(2y_{1}) \sin(2y_{2}))\}] \\ &+\sin(x_{2})^{2} [-4(-1 + \alpha) \alpha\$_{01} \cos(y_{1})^{2} + \$_{02} \{2 + \alpha^{2} - \alpha(2 + \alpha) \cos(2y_{2}) \\ &+\cos(2y_{1}) [\alpha(2 + \alpha) + \{-2 + (4 - 5\alpha) \alpha\} \cos(2y_{2})] - 2(-1 + \alpha) \sin(2y_{1}) \sin(2y_{2})\} \\ &+\cos(x_{1})^{2} [(2 + \alpha^{2})\$_{12} + 4(-1 + \alpha)^{2}\$_{21} - \alpha(2 + \alpha)\$_{12} \cos(2y_{2}) \\ &+\cos(x_{1})^{2} [(2 + \alpha^{2})\$_{12} + 4(-1 + \alpha)^{2}\$_{21} - \alpha(2 + \alpha)\$_{12} \cos(2y_{2}) \\ &+\cos(x_{1})^{2} [(2 + \alpha^{2})\$_{12} + 4(-1 + \alpha)^{2}\$_{21} - \alpha(2 + \alpha)\$_{12} \cos(2y_{2}) \\ &+\sin(x_{1})^{2} [2 \$_{20} + \alpha^{2}\$_{20} + 4\alpha\$_{10} \cos(y_{2})^{2} - 4\alpha^{2}\$_{10} \cos(y_{2})^{2} \\ &-\alpha(2 + \alpha)\$_{20} \cos(2y_{1}) - \cos(2y_{2})) + \{-2 + (4 - 5\alpha) 3\$_{20} \cos(2y_{1}) \cos(2y_{2}) \\ &-\alpha(2 + \alpha)\$_{20} \sin(2y_{1}) \sin(2y_{2}) + 2\cos(x_{2})^{2} [2(-1 + \alpha)^{2}\$_{12} \\ &+2\sin(y_{1})^{2} [-(-1 + \alpha) \alpha(\$_{22} + \$_{21} \cos(2y_{2})^{2} - (-1 + \alpha)^{2}\$_{12} \\ &+2\sin(y_{1})^{2} (-(-1 + \alpha) \alpha(\$_{22} + \$_{21} \cos(2y_{2})^{2} + (-1 + \alpha)^{2}\$_{21} \sin(2y_{1}) \sin(2y_{2})] + \{\$_{21} + \alpha(2y_{1}) \sin(2y_{2})\} + \{\$_{21} + \alpha(2y_{1}) \sin(2y_{2})\} + \{\$_{21} + \alpha(2y_{1}) \sin(2y_{2})\} + \{\$_{22} + \alpha(2y_{21} - 2\alpha\$_{21} + \$_{22} + \alpha\$_{22} \\ &+\alpha(y_{1}) \sin(2y_{2})] + \{\$_{22} + \alpha(2)$$

$$+ (-1 + \alpha)^{2} (-1 + \alpha)^{2} (-1 +$$

By using equations (1-5, 8-13 and 17), the players payoffs for the depolarizing channel become

$$\begin{split} \$_{\text{Dep}}^{A,B}(x_{i},y_{i},\alpha) &= \frac{1}{3072} [(8-9\alpha)^{2}(\$_{11}-\$_{12}-\$_{21}+\$_{22})\cos(2(x_{1}+x_{2})) \\ &\times(5+3\cos(2(y_{1}+y_{2})))+2\{4\{64+3\alpha(-16+9\alpha)\}\$_{00} \\ &-3\alpha(-16+9\alpha)(2\$_{01}+2\$_{02}+2\$_{10}-\$_{11}-\$_{12}+2\$_{20} \\ &-\$_{21}-\$_{22})+64\{2\$_{01}+2\$_{02}+2\$_{10}+3\$_{11}+3\$_{12} \\ &+2\$_{20}+3(\$_{21}+\$_{22})\}+(8-9\alpha)^{2}\{(4\$_{00}-2\$_{01} \\ &-2\$_{02}-2\$_{10}+\$_{11}+\$_{12}-2\$_{20}+\$_{21}+\$_{22})\cos(2(y_{1}+y_{2})) \\ &+\{2(2\$_{10}-\$_{11}-\$_{12}-2\$_{20}+\$_{21}+\$_{22})\cos(2x_{1}) \\ &+(\$_{11}-\$_{12}-\$_{21}+\$_{22})\cos(2(x_{1}-x_{2}))+ \\ &2(2\$_{01}-2\$_{02}-\$_{11}+\$_{12}-\$_{21}+\$_{22})\cos(2x_{2})\}\sin(y_{1}+y_{2})^{2}\}] \end{split}$$

By using equations (1-5, 7, 10-13 and 17), the players payoffs for the phase damping channel are obtained as

$$\begin{split} \$_{\text{PD}}^{A,B}(x_{i},y_{i},\alpha) &= \frac{1}{192} \left[-8(2\$_{11}+2\$_{12}+2\$_{20}+\$_{21}+\$_{22})\cos(2x_{1}) \\ &+\cos(2x_{2})[8\{2\$_{01}-2\$_{02}-2\$_{11}+2\$_{12}+\$_{21}-\$_{22} \\ &+(2\$_{11}-2\$_{12}-\$_{21}+\$_{22})\cos(2x_{1}) \} -8\{2\$_{01}-2\$_{02} \\ &-\$_{21}+\$_{22}+(\$_{21}-\$_{22})\cos(2x_{1}) \}\cos(2y_{1})\cos(2y_{2}) \\ &+4\{2+3(-2+\alpha)\alpha\}\{2\$_{01}-2\$_{02}-\$_{21}+\$_{22} \\ &+(\$_{21}-\$_{22})\cos(2x_{1}) \}\sin(2y_{1})\sin(2y_{2}) \right] \\ &+16\cos(2x_{1})^{2}[2\$_{10}+\$_{11}+\$_{12}+2\$_{21}+2\$_{22} \\ &+(\$_{11}-\$_{12}-2\$_{21}+2\$_{22})\cos(2x_{2}) +\{-2\$_{10} \\ &+\$_{11}+\$_{12}+(\$_{11}-\$_{12})\cos(2x_{2})\}\cos(2y_{1})\cos(2y_{2}) \\ &+\frac{1}{2}\{2+3(-2+\alpha)\alpha\}\{2\$_{10}-\$_{11}-\$_{12} \\ &+(-\$_{11}+\$_{12})\cos(2x_{2})\}\sin(2y_{1})\sin(2y_{2}) \right] \\ &-8(\$_{11}-\$_{12}-\$_{21}+\$_{22})\sin(2x_{1})\sin(2x_{2})[2+3(-2+\alpha)\alpha \\ &+\{2+3(-2+\alpha)\alpha\}\cos(2(y_{1}+y_{2})]) +\sqrt{3}\alpha(-2+3\alpha)(\sin(2y_{1}) \\ &+\sin(2y_{2}))] +8[4\$_{00}+2\$_{01}+2\$_{02}+2\$_{11}+2\$_{12}+2\$_{20}+\$_{21} \\ &+\$_{22}+\{2\$_{00}-\$_{01}-\$_{02}+(-2\$_{20}+\$_{21}+\$_{22})\sin(2x_{1})^{2}] \\ &\times\{2\cos(2y_{1})\cos(2y_{2})+\{-2-3(-2+\alpha)\alpha\}\sin(2y_{1})\sin(2y_{2})] \end{split}$$

where the subscripts AD, Dep and PD in equations (18-20) represent the amplitude damping, depolarizing and phase damping channels respectively. It can be easily checked from equations (18-20) that by setting $\alpha = 1$, the game becomes a noiseless game.

III. DISCUSSIONS

In this work, we analyze the non-transitive two-player three-strategy entangled quantum game usually termed as RSP game under the influence of quantum noise. We consider different noisy quantum channels and show that that how the game's payoff is influenced by these channels. We consider the winning non-transitive strategies R, S and P such as R beats S, S beats P, and Pbeats R (R > S > P > R). It is seen that the game's payoff is differently influenced by different quantum channels.

In figure 1, we plot the Alice's payoff as a function of quantum noise parameter α for $x_1 = y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels. It is seen that Alice's payoff is heavily influenced by depolarizing noise as compared to the amplitude damping noise. It causes a monotonic decrease in players payoffs as the amount of quantum noise is increased. It is evident from the figure that in case of amplitude damping channel, the Alice's payoff reaches its minimum for $\alpha = 0.5$ and is symmetrical. This implies that the larger amount of quantum noise influences the game weakly. However, the phase damping channel has no influence on the payoffs of the players.

In figure 2, we plot the Alice's payoff as a function of her strategy x_1 for $y_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels. It can be seen from the figure that the phase damping channel does not influence the Alice's payoff. On the other hand, the Alice's payoff is strongly affected for amplitude damping and depolarizing channels.

In figure 3, we plot the Alice's payoff as a function of her strategy y_1 for $x_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels. It can be seen that the Alice's payoff is decreased due to the presence of quantum noise for both the amplitude damping and depolarizing channels. Whereas in case of phase damping channel, Alice's payoff remains unaffected in the presence of quantum noise. In addition, it can be easily checked that for maximum value of quantum noise parameter (i.e. at $\alpha = 1$) the game behaves as a noiseless game. For $\alpha = 1$, we obtain a single curve for all the three channels (the phase damping channel curve in figures 2 and 3). Hence, we can say that the game becomes a noiseless game for maximum value of quantum noise.

In figures 4-6, we present the 3D graphs of Alice's payoff as a function of α and her strategy x_1 for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping, depolarizing and phase damping channels respectively. One can easily see that the depolarizing channel influence the game more strongly as compared to the amplitude damping channel. From figures 4-6, it can also be seen that the Nash equilibrium of the game does not change under the influence of quantum noise. Furthermore, the non-transitive character of the classical game is not affected by the quantum noise.

IV. CONCLUSIONS

We study the influence of quantum noise on the Rock-Scissor-Paper (*RSP*) game under different noise models. We consider the winning non-transitive strategies R, S and P such as Rbeats S, S beats P, and P beats R. Our investigations show that for maximum value of quantum noise parameter the game behaves as a noiseless game. It is seen that game's payoff is strongly influenced by the depolarizing noise as compared to the amplitude damping noise. It is shown that under the influence of depolarizing channel the players payoffs decrease monotonically as a function of quantum noise. However, in case of amplitude damping channel, the payoff function reaches its minimum for $\alpha = 0.5$ and is symmetrical. Therefore, amplitude damping channel influences the game weakly for higher values of quantum noise. Furthermore, the phase damping channel does not influence the game's payoff. Therefore the game deserves a careful study during its implementation. It is also seen that the non-transitive character of the classical game and the game's Nash equilibrium are not affected by the quantum noise.

- Nielson M A and Chuang I L 2000 Quantum Computation and Quantum Information (Cambridge: Cambridge University Press)
- [2] Meyer D A 1999 Phys. Rev. Lett. 82 1052
- [3] Eisert J, Wilkens M and Lewenstein M 1999 Phy. Rev. Lett. 83 3077
- [4] Marinatto L and Weber T 2000 Phys. Lett. A 272 291
- [5] Flitney A P and Abbott D 2005 J. Phys. A 38 449
- [6] Cheon T and Iqbal A 2008 J. Phys. Soc. Japan 77 024801
- [7] Iqbal A and Abbott D 2009 J. Phy. Soc. Japan 78 014803
- [8] Ramzan M, Nawaz A, Toor A H and Khan M K 2008 J. Phys. A: Math. Theor. 41 055307
- [9] Iqbal A, Cheon T and Abbott D 2008 Phys. Lett. A 372 6564
- [10] Iqbal A and Toor A H 2001 Phys. Lett. A 280 249
- [11] Flitney A P and Abbott D 2002 Phys. Rev. A 65 062318
- [12] Iqbal A and Toor A H 2002 Phys. Rev. A 65 022036
- [13] Eisert J and Wilkens M 2000 J. Mod. Opt. 47 2543
- [14] Ramzan M and Khan M K 2008 J. Phys. A: Math. Theor. 41 435302
- [15] Ramzan M and Khan M K 2009 J. Phys. A: Math. Theor. 42 025301
- [16] Flitney A P, Ng J and Abbott D 2002 Physica A 314 35
- [17] Iqbal A and Toor A H 2002 Phys. Lett. A 293 103

- [18] Johnson N F 2001 Phys. Rev. A 63 020302(R)
- [19] D'Ariano G M, Gill R D, Keyl M, Kuemmerer B, Maassen H and Werner R F 2002 Quant. Inf. Comp. 2 355
- [20] Gawron P and Sladkowski J 2008 Int. J. Quant. Info. 6 667
- [21] Wang X B, Kwek L C and Oh C H 2000 Phys. Lett. A 278 44
- [22] de Sousa P B M and Ramos R V 2008 Quant. Inform. Process. 7 125
- [23] Iqbal A and Abbott D 2009 arXiv:quant-ph/0909.3369
- [24] Chen L K, Ang H, Kiang D Kwek L C and Lo C F 2003 Phys. Lett. A 316 317
- [25] Flitney A P and Hollenberg L C L 2007 Quantum Inf. Comput. 7 111
- [26] Salman K., Ramzan M and Khan M K 2010 Int. J. Theor. Phys. 49 31-41
- [27] Ramzan M and Khan M K 2010 to appear in Quant. Inform. Process.
- [28] Steane A 1996 Phys. Rev. Lett. 77 793
- [29] Preskill J 1998 Proc. R. Soc. Lond. A 454 385
- [30] Deutsch D, Ekert A, Josza R, Macchiavello C, Popescu S and Sanpera A 1996 Phys. Rev. Lett. 77 2818
- [31] Stohler M and Fischbach E 2005 Fizika B 14 235
- [32] Gregg jeager 2007 Journal of modern optics 54 1
- [33] Berg A 2002 IEEE Transaction on Information Theory 48 3096

Figures captions

Figure 1. Alice's payoff plotted as a function of quantum noise parameter α for $x_1 = y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.

Figure 2. Alice's payoff plotted as a function of her strategy x_1 for $y_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.

Figure 3. Alice's payoff plotted as a function of her strategy y_1 for $x_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.

Figure 4. Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping channel.

Figure 5. Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for depolarizing channel.

Figure 6. Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for phase damping channel.

Table Caption

Table 1. In the "rock, scissors, paper" game, a player can win regardless of the strategy chosen by an opponent. The first number in each entry corresponds to Alice's payoff and the second number corresponds to Bob. Winning strategies are non-transitive in that R > S > P > R. A payoff of +1 has been assigned to winning, -1 to losing and 0 for both in case of a tie.



FIG. 1: Alice's payoff plotted as a function of quantum noise parameter α for $x_1 = y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.



FIG. 2: Alice's payoff plotted as a function of her strategy x_1 for $y_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.



FIG. 3: Alice's payoff plotted as a function of her strategy y_1 for $x_1 = \pi/2$, $x_2 = y_2 = 0$ and $\alpha = 0.5$ for amplitude damping (solid line), depolarizing (dashed line) and phase damping (dotted line) channels.



FIG. 4: Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for amplitude damping channel.



FIG. 5: Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for depolarizing channel.



FIG. 6: Alice's payoff plotted as a function of her strategy x_1 and α for $y_1 = \pi/2$, $x_2 = y_2 = 0$ for phase damping channel.

TABLE I: In the "rock, scissors, paper" game, a player can win regardless of the strategy chosen by an opponent. The first number in each entry corresponds to Alice's payoff and the second number corresponds to Bob. Winning strategies are non-transitive in that R > S > P > R. A payoff of +1 has been assigned to winning, -1 to losing and 0 for both in case of a tie.

	Bob		
	R	\mathbf{S}	Р
R	(0, 0)	(1, -1)	(-1, 1)
Alice S	(-1, 1)	(0, 0)	(1, -1)
Р	(1, -1)	(-1, 1)	(0, 0)