The Nuclear Born Oppenheimer Method and Nuclear Rotations

Nouredine Zettili

Department of Physical & Earth Sciences Jacksonville State University Jacksonville, AL 36265

Abstract

We deal here with the application of the Nuclear Born Oppenheimer (NBO) method to the description of nuclear rotations. As an edifying illustration, we apply the NBO formalism to study the rotational motion of nuclei which are axially-symmetric and even, but whose shells are not closed. We focus, in particular, on the derivation of expressions for the rotational energy and for the moment of inertia. Additionally, we examine the connection between the NBO method and the self-consistent cranking (SCC) model. Finally, we compare the moment of inertia generated by the NBO method with the Thouless-Valantin formula and hence establish a connection between the NBO method and the large body of experimental data.

PACS numbers: 21.60.-n, 21.60.Ev, 21.10.Re

1 Introduction

Since nuclear and molecular rotation-vibration spectra present many striking analogies, and since the Born-Oppenheimer (BO) approximation[1] of molecular physics was shown to be very accurate¹ in describing molecular rotations and vibrations[2], it will be interesting to explore the possibility of using the BO approximation to describe nuclear collective rotations.

Exploiting the analogy between nuclear and molecular dynamics, Villars introduced a microscopic method[3], [4] to describe nuclear collective motion. This method, to be called the Nuclear Born-Oppenheimer (NBO) method, was developed along the lines of the molecular BO approximation by constructing a factorable trial function modeled after the BO ansatz.

Using an analytically solvable model[5], we have shown that the NBO method is very accurate for adiabatic collective motion[6]. Since the NBO method is a quantum mechanical prescription, we have shown that it offers a suitable framework for describing the zero-point fluctuations[7]; we have also shown that the method offers an accurate description of small-amplitude collective oscillations[8] and that it yields the random phase approximation (RPA) equations[9]. Additionally, we have applied the NBO method to study nuclear collective motion[10] and examined its connection with the collective model of Bohr[11].

So, having applied the NBO method to the study of small amplitude motion, we have yet to apply it to nuclear collective rotations. In this work we want to achieve just that aim; namely, we want to apply the general BO formalism outlined in Ref.[10] to the description of nuclear rotational states. As an illustration, we will apply the NBO formalism to study the rotations of nuclei that are axially-symmetric and even, but with non-closed shells. We will focus, in particular, on the derivation of expressions for the energy and for the moment of inertia. Additionally, we shall examine the connection of the NBO method with the successful self-consistent cranking (SCC) model.

In Sec. 2, we present a brief outline of the NBO formalism and how it applies to nuclear nuclear collective motion. We then devote Sec. 3 to the application of the NBO method to study the rotational states of axially symmetric nuclei; in

¹Using an elementary solvable model, Moshinksy and Kittel[2] have shown that the BO approximation is very accurate for both the molecular energy and wave function: $\frac{E_{\rm BO}}{E_{\rm exact}} = 1 - \frac{1}{4}\chi$ and $|\langle \psi_{\rm BO} | \psi_{\rm exact} \rangle|^2 = 1 - \frac{3}{128}\chi^2$, where χ is equal to the ratio of electronic to nuclear masses (i.e., $\chi = m_e/M \simeq 10^4$).

particular, we will derive an expression for the rotational energy. In Sec. 4, we present a discussion on the connection of the NBO method to the self-consistent cranking (SCC) model.

2 Synopsis of the Application of the NBO Method to Nuclear Collective Motion

To describe nuclear collective motion within the framework of the NBO method, we need to introduce a tensor operator $\hat{Q}_{\alpha\beta}$; that is, to be able to describe collective rotations and vibrations of nuclei, we need to introduce a set of operators that are the elements of a symmetric cartesian tensor operator $\hat{Q}_{\alpha\beta}$. In the rest of this work, we shall use Greek subscripts to refer to a space-fixed frame of reference; Latin subscripts will be used later to refer to a body-fixed frame. The operator $\hat{Q}_{\alpha\beta}$ is assumed to depend on the various nucleonic variables – positions \vec{x}_i , momenta, \vec{p}_i , and spins, \vec{s}_i . In addition, we assume that $\hat{Q}_{\alpha\beta}$ are one-body operators, symmetric, even under time reversal, have a continuous eigenvalue spectrum $(q_{\alpha\beta})$, and commute with any other component $\hat{Q}_{\gamma\delta}$ of \hat{Q} (*i.e.*, $[\hat{Q}_{\alpha\beta}, \hat{Q}_{\gamma\delta}] = 0$). Let $\hat{K}_{\alpha\beta}$ be the canonical conjugate of $\hat{Q}_{\alpha\beta}$:

$$[i\hat{K}_{\alpha\beta}, \ \hat{Q}_{\gamma\delta}] = \frac{1}{2} \left(\delta_{\alpha\gamma}\delta_{\beta\delta} + \delta_{\alpha\delta}\delta_{\beta\gamma} \right).$$
(2.1)

The NBO method consists of the following two essential steps[3], [4], [10]:

• First, we need to construct a suitable representation for the nucleus' Hamiltonian \hat{H} by decomposing it into a series

$$\hat{H} = \hat{H}_0 + \sum_{\alpha\beta} \hat{H}_{1\,\alpha\beta} \hat{K}_{\alpha\beta} + \frac{1}{2} \sum_{\alpha\beta} \sum_{\gamma\delta} \hat{H}_{2\,\alpha\beta\gamma\delta} \hat{K}_{\alpha\beta} \hat{K}_{\gamma\delta} + \cdots, \qquad (2.2)$$

where all coefficient operators \hat{H}_0 , $\hat{H}_{1\,\alpha\beta}$, $\hat{H}_{2\,\alpha\beta,\gamma\delta}$,... commute with $\hat{Q}_{\alpha\beta}$.

• Second, we make use a factorable trial function²

$$\langle x|\psi\rangle = \int \prod_{\alpha \le \beta} dq_{\alpha\beta} \langle x|\delta(q_{\alpha\beta} - \hat{Q}_{\alpha\beta})|\Phi(q)\rangle g(q) \quad , \tag{2.3}$$

where $\langle x | \Phi(q) \rangle$ is the intrinsic wave function, and g(q) is the collective amplitude.

²We will use x to abbreviate for the set of nucleonic variables – position \vec{x}_i , momentum \vec{p}_i , and spin, \vec{s}_i .

After constructing the Hamiltonian and the wave function, we can calculate the mean energy by a simple application of \hat{H} to $|\psi\rangle$:

$$\langle \psi | \hat{H} | \psi \rangle = \int \prod_{\alpha \leq \beta} dq_{\alpha\beta} \ g^*(q) \langle \Phi(q) | \delta(q_{\alpha\beta} - \hat{Q}_{\alpha\beta}) \{ \tilde{H}_0 | \Phi \rangle g(q) + \sum_{\alpha\beta} \tilde{H}_{1\,\alpha\beta} | \Phi \rangle k_{\alpha\beta} g(q)$$

$$+ \frac{1}{2} \sum_{\alpha\beta} \sum_{\gamma\delta} \tilde{H}_{2\,\alpha\beta,\gamma\delta} | \Phi \rangle k_{\alpha\beta} k_{\gamma\delta} g(q) + \cdots \},$$
 (2.4)

where the $k_{\alpha\beta}$ are operators that act on g(q); they obey commutation relations with the $q_{\alpha\beta}$ isomorphic with (2.1)

$$[ik_{\alpha\beta}, q_{\gamma\delta}] = \frac{1}{2} \left(\delta_{\alpha\gamma} \delta_{\beta\delta} + \delta_{\alpha\delta} \delta_{\beta\gamma} \right).$$
(2.5)

The few lowest expressions of \tilde{H}_K are given by

$$\tilde{H}_{0} = \hat{H} - \sum_{\alpha\beta} \left[\hat{H}, \ i\hat{Q}_{\alpha\beta} \right] \hat{G}_{\alpha\beta} + \frac{1}{2} \sum_{\alpha\beta} \sum_{\gamma\delta} \left[\left[\hat{H}, \ i\hat{Q}_{\alpha\beta} \right], \ i\hat{Q}_{\gamma\delta} \right] \hat{G}_{\alpha\beta} \hat{G}_{\gamma\delta} + \cdots,
\tilde{H}_{1\,\alpha\beta} = \left[\hat{H}, \ i\hat{Q}_{\alpha\beta} \right] - \sum_{\gamma\delta} \left[\left[\hat{H}, \ i\hat{Q}_{\alpha\beta} \right], \ i\hat{Q}_{\gamma\delta} \right] \hat{G}_{\gamma\delta} + \cdots,$$

$$\tilde{H}_{2\,\alpha\beta\,\gamma\delta} = \left[\left[\hat{H}, \ i\hat{Q}_{\alpha\beta} \right], \ i\hat{Q}_{\gamma\delta} \right] + \cdots,$$
(2.6)

where $\hat{G}_{\alpha\beta}$ is a one particle operator that acts on $|\phi\rangle$; it is defined by the action of $k_{\alpha\beta}$ on the parameter q in $|\phi\rangle$

$$k_{\alpha\beta}\langle x|\Phi(q)\rangle = \frac{1}{2i}(1+\delta_{\alpha\beta})\frac{\partial}{\partial q_{\alpha\beta}}\langle x|\Phi(q)\rangle = \langle x|\hat{G}_{\alpha\beta}|\Phi(q)\rangle.$$
(2.7)

We should note that the mean energy expression (2.4) was derived within a spacefixed or lab frame. However, in the description of permanently deformed (non spherical) nuclei, it is more convenient to employ a body-fixed frame of reference. Here, we take the axes of the body-fixed frame along the three principal axes of $q_{\alpha\beta}$ which are defined by the unit vectors \hat{e}_a (a = 1, 2, 3), and specify their orientation with respect to the space-fixed frame by three Euler angles[23] θ_s (i.e., θ, φ, ψ): $\hat{e}_a = \hat{e}_a(\theta_s)$ with $\hat{e}_a \cdot \hat{e}_b = \delta_{ab}$ and $\hat{e}_a \times \hat{e}_b = \mathcal{E}_{abc}\hat{e}_c$ where \mathcal{E}_{abc} is the antisymmetric tensor ($\mathcal{E}_{123} = 1 = -\mathcal{E}_{213}$ etc.). The collective degrees of freedom can be separated into rotational and vibrational terms by transforming $\hat{Q}_{\alpha\beta}$ to the body-fixed frame; that is, by means of the principal axes transformation of the tensor operator $\hat{Q}_{\alpha\beta}$:

$$q_{\alpha\beta} = \sum_{a=1}^{3} e_{\alpha a}(\theta) e_{\beta a}(\theta) q_a, \qquad (2.8)$$

where $e_{\alpha a} (\equiv \hat{e}_{\alpha} \cdot \hat{e}_{a})$, the α^{th} component of the unit vector \hat{e}_{a} , depends on the three Euler angles θ_{s} . In the transformation to the body-fixed frame, we have essentially replaced the six collective coordinates $q_{\alpha\beta}$ by the three q_{a} 's and the three Euler angles. The matrices $e_{\alpha\beta}$ obey the orthogonality relations: $\sum_{a} e_{\alpha a} e_{\beta a} = \delta_{\alpha\beta}$ and $\sum_{\alpha} e_{\alpha a} e_{\alpha b} = \delta_{ab}$.

We can now introduce rotation operators $\hat{\mathcal{L}}_a$ about the body-fixed axes. The Euler angles specifying the orientation of the intrinsic frame need to be viewed as dynamical variables; for instance, the unit vector \hat{e}_a satisfy the commutation rules of a vector operator

$$\left[\hat{\mathcal{L}}_{[ab]}, \ e_{\alpha c}\right] = i e_{\alpha a} \delta_{bc} - i e_{\alpha b} \delta_{ac}.$$
(2.9)

We can easily verify from (2.9) that these operators obey the commutation relations

$$\left[\hat{\mathcal{L}}_{a},\ \hat{\mathcal{L}}_{b}\right] = -i\mathcal{E}_{abc}\hat{\mathcal{L}}_{c},\tag{2.10}$$

which differ in sign from the commutation rules of ordinary angular momentum[12] because they refer to the moving axes and hence do not have the same commutation properties as angular-momentum components along space fixed axes. For instance, we have $[\hat{\mathcal{L}}_1, \hat{\mathcal{L}}_2] = -i\hat{\mathcal{L}}_3$. The space fixed components $\hat{\mathcal{L}}_{\alpha\beta}$ of $\vec{\mathcal{L}}$ can be obtained by rotation: $\hat{\mathcal{L}}_{\alpha\beta} = \sum_{ab} e_{\alpha a} e_{\beta b} \hat{\mathcal{L}}_{ab}$.

In conjunction with the replacement of $q_{\alpha\beta}$ by the variables q_a and θ_s , we seek an expression for the operator $k_{\alpha\beta}$ in terms of the $\hat{\mathcal{L}}_{\alpha\beta}$ and a set of three operators p_a conjugate to q_a , with $[p_a, q_b] = i\delta_{ab}$. We can verify[10] that $k_{\alpha\beta}$ transforms like an operator that acts on θ_s and q_a :

$$k_{\alpha\beta} = \frac{1}{2} \sum_{ab} \frac{e_{\alpha a} e_{\beta b}}{q_a - q_b} \hat{\mathcal{L}}_{[ab]} + \sum_{a} e_{\alpha a} e_{\beta a} p_a$$
$$= \sum_{ab} e_{\alpha a} e_{\beta b} \left[\frac{1}{2} \left(1 + \delta_{\alpha\beta} \right) \frac{\hat{\mathcal{L}}_{[ab]}}{q_a - q_b} + \delta_{ab} p_a \right].$$
(2.11)

Additionally, we can ascertain that $k_{\alpha\beta}$ is Hermitian with regard to the volume element $\prod_{\alpha < \beta} dq_{\alpha\beta}$, which can be shown to transform like:

$$\int d\tau = \int \prod_{\alpha \le \beta} dq_{\alpha\beta} = \int \prod_{a=1}^{3} dq_a (q_1 - q_2) (q_2 - q_3) (q_3 - q_1) d\Omega, \qquad (2.12)$$

where $d\Omega$ is the usual angular element[23] $d\Omega = \sin\theta d\theta d\varphi d\psi$.

Using the relations (2.11) and (2.12), we can now express (2.4) and (2.6) in the body-fixed frame. For this, note first that under the transformation (2.8) from

the Lab frame to the body-fixed system, the quantities $g(q_{\alpha\beta})$, $\langle x_{i\alpha}s_{i\alpha}|\Phi(q_{\alpha\beta})\rangle$, and $\prod_{\alpha\leq\beta}\delta(q_{\alpha\beta}-\hat{Q}_{\alpha\beta})$ become $f(q_a,\theta_s)$, $\langle x'_{ia}s'_{ia}|\phi(q_a)\rangle$, and $\prod_a\delta(q_a-\hat{Q}_{aa})\prod_{a\leq b}\delta(Q_{ab})$, respectively, which in turn will be abbreviated to $f(q,\theta)$, $\langle x'_is'_i|\phi(q)\rangle$, and $\delta(q-\hat{Q})$.

Next, we can show[10] that the action of the total angular momentum $\hat{J}_{\alpha\beta}$ on $|\psi\rangle$ can be expressed in terms of $\hat{\mathcal{L}}_{\alpha\beta}$ on the collective amplitude $f(q_a, \theta_s)$:

$$\langle x|\hat{J}_{\alpha\beta}|\psi\rangle = \int d\tau \langle x|\delta(q-\hat{Q})|\phi(q)\rangle \hat{\mathcal{L}}_{\alpha\beta}f(q,\theta).$$
(2.13)

In this new representation, the operator $\hat{G}_{\alpha\beta}$ of (2.6) is rotated into \hat{G}_{ab} :

$$\langle x_{ia}'s_{ia}'|\hat{G}_{ab}|\phi(q_a)\rangle = \frac{1}{2}(1-\delta_{ab})\frac{\langle x_{ia}'s_{ia}'|\hat{J}_{[ab]}|\phi(q_a)\rangle}{q_a-q_b} + \delta_{ab}\langle x_{ia}'s_{ia}'|\hat{G}_a|\phi(q_a)\rangle.$$
(2.14)

Finally, using Eq. (2.11), (2.13) and (2.14), we have shown in Ref.[10] that the mean energy (2.4) is given in the body-fixed frame of reference by³

$$\langle \psi | \hat{H} | \psi \rangle = \int d\tau f^*(q,\theta) \langle \phi(q) | \delta(q - \hat{Q}) \Big\{ \hat{H} - \sum_{a,b=1}^3 \dot{\hat{Y}}_{[ab]}(\hat{J}_{[ab]} - \hat{\mathcal{L}}_{[ab]})$$

$$- \sum_{a=1}^3 \dot{\hat{Q}}_a(\hat{G}_a - p_a) + \frac{1}{2} \sum_{abcd} \hat{B}_{[ab],[cd]}(\hat{J}_{[ab]} - \hat{\mathcal{L}}_{[ab]})(\hat{J}_{[cd]} - \hat{\mathcal{L}}_{[cd]})$$

$$+ \sum_{abc} \hat{B}_{[ab],c}(\hat{J}_{[ab]} - \hat{\mathcal{L}}_{[ab]})(\hat{G}_c - p_c)$$

$$+ \frac{1}{2} \sum_{ab} \hat{B}_{a,b}(\hat{G}_a - p_a)(\hat{G}_b - p_b) \Big\} |\phi(q)\rangle f(q,\theta),$$

$$(2.15)$$

where

$$\hat{Y}_{[ab]} = \frac{Q_{ab}}{q_a - q_b}, \qquad \dot{\hat{Y}}_{[ab]} = \left[i\hat{H}, \ \hat{Y}_{[ab]}\right], \quad \dot{\hat{Q}}_a = \left[i\hat{H}, \ \hat{Q}_a\right],$$
(2.16)

$$\hat{B}_{[ab],[cd]} = [\dot{\hat{Y}}_{[ab]}, \ i\hat{Y}_{[cd]}] , \quad \hat{B}_{[ab],c} = [\dot{\hat{Y}}_{[ab]}, \ i\hat{Q}_{c}] , \quad \hat{B}_{ab} = [\dot{\hat{Q}}_{a}, \ i\hat{Q}_{b}].$$
(2.17)

Note that, in deriving the mean energy (2.15), we have terminated the series (2.4) at the quadratic terms in $k_{\alpha\beta}$. This termination is justified by the validity of the adiabatic approximation in the present case, since we are dealing with nuclear dynamics for

³ Recall that $\hat{J}_{[ab]}$ and \hat{G}_a operate on the intrinsic state $|\phi(q_a)\rangle$, but $\hat{\mathcal{L}}_{[ab]}$ and p_a operate on the collective state $f(q_a, \theta_s)$ (i.e., $\hat{\mathcal{L}}_{[ab]}$ acts on θ_s and p_a on q_a).

which the time evolution of the collective variables is assumed to be slow on the scale of a single-particle (nucleonic) motion.

As we are going to see next, the rotational and vibrational degrees of freedom appear explicitly in the energy expression (2.15); we will also show how to derive expressions for the collective rotational energy and for the moment of inertia.

3 Description of Rotational States of Axially-Symmetric Nuclei

Consider a permanently deformed, non spherical nucleus. Since we are interested in rotational motion only, we assume the nucleus to be in its vibrational ground state. In this case, we assume that the collective amplitude $f(q_a, \Omega)$ of (2.15) separates into a vibrational part, $g_0(q_a - \bar{q}_a)$, and a rotational part, $D(\Omega)$:

$$f(q_a, \Omega) = g_0(q_a - \bar{q}_a)D(\Omega).$$
(3.18)

The vibrational collective amplitude $g_0(q_a - \bar{q}_a)$ represents here the zero-point oscillations about the equilibrium values, \bar{q}_a , of q_a (a = 1, 2, 3). Hence, the wave function $|\psi\rangle$ of the system becomes (*c.f.* Eq. (2.3)):

$$|\psi\rangle = \int d\tau d\Omega \,\,\delta(q - \hat{Q}) |\phi(q_a)\rangle g_0(q_a - \bar{q}_a) D(\Omega) \quad , \tag{3.19}$$

with $d\tau = (q_a - q_2)(q_2 - q_3)(q_3 - q_1)\prod_{a=1}^3 dq_a$ and $d\Omega = d\varphi d\psi \sin\theta d\theta$ (c.f. Eq. (2.12)), and where $\delta(q - \hat{Q})$ is used to abbreviate $\prod_{a=1}^3 \delta(q_a - \hat{Q}_{aa})\prod_{a < b} \delta(\hat{Q}_{ab})$. In this case, after expanding $|\phi(q_a)\rangle$ about $|\phi(\bar{q}_a)\rangle$, the mean energy (2.15) becomes:

$$\langle \psi | \hat{H} | \psi \rangle \simeq \int d\tau \int d\Omega g^*(q_a - \bar{q}_a) D^*(\Omega) \langle \phi(\bar{q}_a) | \delta(q - \hat{Q}) \Big\{ \hat{H} - \sum_{a=1}^3 \dot{Y}_a(\hat{J}_a - \hat{\mathcal{L}}_a) \\ + \frac{1}{2} \sum_{ab} B_{ab}(\hat{J}_a - \hat{\mathcal{L}}_a) (\hat{J}_b - \hat{\mathcal{L}}_b) \Big\} | \phi(\bar{q}_a) \rangle g(q_a - \bar{q}_a) D(\Omega) \\ + E_{\text{osc}}^0 + E_{\text{coupl}},$$

$$(3.20)$$

where we have used the notation $\hat{0}_c$ to abbreviate $\hat{0}_{ab}$ (a, b, c being cyclic permutations of the body-fixed axes 1,2,3). In this expression, $E_{\rm osc}^0$ is the energy of the zeropoint oscillations; $E_{\rm coupl}$ is the coupling-energy between the rotational and vibrational motions which we can neglect. As for $E_{\rm osc}^0$, we will drop it from all following meanenergy expressions, since it represents only a constant shift of the entire (rotational) energy spectrum. Note that, in the derivation of the mean-energy (3.20), we have approximated the operator \hat{B}_{ab} by its mean value $\langle \phi | \hat{B}_{ab} | \phi \rangle$ (i.e., $\langle \phi | \hat{B}_{ab} | \phi \rangle \equiv B_{ab}$). In what follows, the notation B_a will be used to abbreviate B_{aa}

For the sake of simplicity, we shall focus in this work only on deformed, even, and axially-symmetry nuclei. Consider the axis 3, of the body-fixed frame, to be the axis of symmetry for the system. As a consequence of the axial symmetry, we have: $\bar{q}_a = \bar{q}_2 \neq \bar{q}_3$ and $B_1 = B_2 \equiv B \neq B_3$.

Now, since \hat{H} , \hat{J}^2 and \hat{J}_z mutually commute⁴ commute, they possess joint eigenfunctions. The structure of our trial function allows it to be an exact eigenfunction of \hat{J}^2 and \hat{J}_2 , but provides only a variational approximation to the energy. In the case of axial symmetry, this trial function $|\psi\rangle$ can be obtained from (3.19) by expanding $|\phi(q_a)\rangle D(\Omega)$ in terms of the Wigner D-functions⁵:

$$|\psi_{IM}\rangle = \int d\tau d\Omega \ \delta(q - \hat{Q}) \sqrt{\frac{2I+1}{16\pi^2}} \sum_{K} \left[|\phi_K(q_a)\rangle \mathcal{D}^I_{MK}(\Omega) + (-1)^I |\phi_{-K}(q_a) \mathcal{D}^I_{M-K}(\Omega) \right] g_0(q_a - \bar{q}_a) , \qquad (3.21)$$

where $|\phi_K\rangle$ is an eigenfunction of \hat{J}_3 (*i.e.*, $\hat{J}_3|\phi_K\rangle = K|\phi_K\rangle$) and \mathcal{D}^I_{MK} is an eigenfunction to $\hat{\mathcal{L}}_3$ (*i.e.*, $\hat{\mathcal{L}}_3\mathcal{D}^I_{MK} = K\mathcal{D}^I_{MK}$), $\hat{\vec{\mathcal{L}}}^2$ and $\hat{\mathcal{L}}_z$. It then follows that

$$\vec{J}^{2}|\psi_{IM}\rangle = I(I+1)|\psi_{IM}\rangle \quad , \qquad \qquad J_{Z}|\psi_{IM}\rangle = M|\psi_{IM}\rangle \quad . \tag{3.22}$$

For the simpler case of the K=0 band, the wave function $|\psi_{IM}
angle$ is given by:

$$|\psi_{IM}\rangle = \int d\tau d\Omega \ \delta(q - \hat{Q}) \sqrt{\frac{2I+1}{8\pi^2}} |\phi_0(q_a)\rangle \mathcal{D}^I_{M0}(\Omega) g_0(q - \bar{q}_a) \quad . \tag{3.23}$$

Note that (as a consequence of axial symmetry) the following important relation holds for both forms, (3.21) and (3.23), of $|\phi\rangle D(\Omega)$:

$$(\hat{J}_3 - \hat{\mathcal{L}}_3) |\phi\rangle D(\Omega) = 0$$
 . (3.24)

In this case of axial symmetry, and after omitting $E_{\rm osc}^0 + E_{\rm coupl}$, we can see that the mean-energy (3.20) reduces to:

$$\langle \psi_{IM} | \hat{H} | \psi_{IM} \rangle = \int d\tau d\Omega g_0^* (q_a - \bar{q}_a) D^*(\Omega) \langle \phi(\bar{q}_a) | \delta(q - \hat{Q})$$
$$\times \left[\hat{H}_0 + \frac{B}{2} \sum_{a=1}^2 \hat{\mathcal{L}}_a^2 \right] | \phi(\bar{q}_a) \rangle g_0 D(\Omega)$$
$$+ \mathcal{E}_1 + \mathcal{E}_2 \quad ,$$
(3.25)

 $^{{}^{4}\}vec{J}$ is the total angular momentum and \hat{J}_{z} is its Z component with respect to the Lab frame. ⁵The definition of $\mathcal{D}_{M-K}^{I}(\Omega)$ used here is that of Bohr-Mottelson

with

$$\hat{H}_0 = \hat{H} - \sum_{a=1}^2 \dot{\hat{Y}} \hat{J}_a + \frac{1}{2} B \sum_{a=1}^2 \hat{J}_a^2 \quad , \qquad (3.26)$$

$$\mathcal{E}_{1} = \int d\tau d\Omega g_{0}^{*} D^{*} \langle \phi | \delta(q - \hat{Q}) \sum_{a=1}^{2} (\dot{\hat{Y}}_{a} - B\hat{J}_{a}) \hat{\mathcal{L}}_{a} | \phi \rangle g_{0}(q_{a} - \bar{q}_{a}) D(\Omega) ,$$

$$\mathcal{E}_{2} = (B_{12} + B_{21}) \int d\tau d\Omega g_{0}^{*} D^{*} \langle \phi | \delta(q - \hat{Q}) (\hat{J}_{1} - \hat{\mathcal{L}}_{1}) (\hat{J}_{2} - \hat{\mathcal{L}}_{2}) | \phi \rangle g_{0} D(\Omega) ,$$
(3.27)

where $|\phi(q_a)\rangle D(\Omega)$ is given by (3.21) or (3.23), depending on whether one is interested in the $K \neq 0$ band or the K = 0 band.

We should now specify the description of the intrinsic structure of the system. To this end, we assume that the intrinsic state $|\phi(\bar{q}_a)\rangle$ is given by a mean field approximation such that $\langle \phi(\bar{q}_a)|\hat{Q}_{11}|\phi(\bar{q}_a)\rangle = \bar{q}_1$ is equal to $\langle \phi|\hat{Q}_{22}|\phi\rangle = \bar{q}_2$ (*i.e.*, such that $\bar{q}_1 = \bar{q}_2$, the axial symmetry condition). This can be achieved by means of a constrained variational principle.

Let us now look at the determination of the collective tensor operator Q. We determine the particle-hole (ph) components of the tensor operator \hat{Q}_{ab} such that $\langle \phi | \dot{Y}_a - B \hat{J}_a | \phi \rangle$ is variationally stable, *i.e.*,

$$\delta\langle\phi|\hat{Y}_a - B\hat{J}_a|\phi\rangle = 0 \qquad (a = 1, 2) \quad . \tag{3.28}$$

This variational condition insures that the simple expression (3.21) for $|\psi\rangle$ is adequate to describe the rotational energy (term $\sim \hat{\mathcal{L}}^2$) correctly.

To determine the mean energy (3.25), we need to calculate \mathcal{E}_1 and \mathcal{E}_2 . In what follows, we are going to show that both \mathcal{E}_1 and \mathcal{E}_2 are identically zero. First, the term $\langle \phi | \delta(q - \hat{Q}) (\dot{Y}_a - B \hat{J}_a) | \phi \rangle$ in the integrand of \mathcal{E}_1 can be rewritten as⁶

$$\begin{aligned} \langle \phi | \delta(q - \hat{Q}) (\dot{\hat{Y}}_{a} - B\hat{J}_{a}) | \phi \rangle &\simeq \langle \phi | \delta(q - \hat{Q}) | \phi \rangle \langle \phi | \dot{\hat{Y}}_{a} - B\hat{J}_{a} | \phi \rangle \\ &+ \sum_{\sigma \mu} \langle \phi | \delta(q - \hat{Q}) | \phi_{\sigma \mu} \rangle \langle \phi_{\sigma \mu} | \dot{\hat{Y}}_{a} - B\hat{J}_{a} | \phi \rangle \end{aligned}$$

$$(3.29)$$

where σ, τ, \ldots refer to unoccupied (particle) states, while μ, λ, \ldots refer to occupied (hole) states. Using the condition (3.28), we see that the term $\langle \phi | \delta(q - \hat{Q}) (\dot{Y}_a - \dot{Q}) (\dot{Y}_a - \dot{Y}_a - \dot{Y}_a) (\dot$

⁶In this approximate expression, we have neglected the two-body part of the operator $\hat{Y}_a \equiv [\hat{H}, i\hat{Y}_a]$.

 $B\hat{J}_a)|\phi\rangle$ becomes equal to $\langle\phi|\delta(q-\hat{Q})|\phi\rangle\langle\phi|\hat{Y}_a-B\hat{J}_a|\phi\rangle$. Now, using the fact that $|\phi_k\rangle$ is an eigenfunction to \hat{J}_3 and that the action of both \dot{Y}_a and \hat{J}_a (a = 1, 2) on $|\phi_k\rangle$ generate $|\phi_{k\pm 1}\rangle$, we can ascertain that $\langle\phi|\hat{Y}_a-B\hat{J}_a|\phi\rangle$ is itself identically zero, and hence \mathcal{E}_1 is equal to zero. To see this, note that $(\hat{Y}_a-B\hat{J}_a)$ have non-zero matrix elements only between $|\phi_k\rangle$ and $\langle\phi_{K\pm 1}|$. So,

$$\langle \phi_k | \hat{Y}_a - B \hat{J}_a | \phi_k \rangle = 0 \qquad (a = 1, 2) \quad ,$$
 (3.30)

and also

$$\langle \phi_{-k} | \dot{\hat{Y}}_a - B \hat{J}_a | \phi_k \rangle = 0 \quad , \tag{3.31}$$

except for $K = \frac{1}{2}$ but, in our case, K is always an integer (because we are dealing with an even nucleus).

Second, \mathcal{E}_2 is identically zero, since both B_{12} and B_{21} can be shown to be equal to zero. To see this, using these expressions,

$$\left[\hat{J}_1, \ i\hat{Y}_1\right] = \frac{\hat{Q}_{33} - \hat{Q}_{22}}{q_3 - q_2} \quad , \qquad \left[\hat{J}_1, \ i\hat{Y}_2\right] = \frac{\hat{Q}_{12}}{q_1 - q_3} \quad , \qquad (3.32)$$

$$\left[\hat{J}_2, \ i\hat{Y}_2\right] = \frac{\hat{Q}_{33} - \hat{Q}_{11}}{q_3 - q_1} \quad , \qquad \left[\hat{J}_2, \ i\hat{Y}_1\right] = \frac{-\hat{Q}_{12}}{q_2 - q_3} \quad , \qquad (3.33)$$

we can easily show the following important relation:

$$\langle \phi | [\hat{J}_a, i\hat{Y}_b] | \phi \rangle = \delta_{ab} \qquad (a = 1, 2) \quad ,$$

$$(3.34)$$

since $\langle \hat{Q}_{ab} \rangle = \delta_{ab}$. Now, applying this relation to the variational principle (3.29), we can verify that $\langle \phi | [\dot{\hat{Y}}_{a}, \ \hat{Y}_{b}] | \phi \rangle$ is equal to $B \delta_{ab}$, *i.e.*,

$$B_{ab} = B\langle \phi | [\hat{J}_a, \ i\hat{Y}_b] | \phi \rangle = B\delta_{ab} \qquad (a = 1, 2) \quad . \tag{3.35}$$

Now, since both of \mathcal{E}_1 and \mathcal{E}_2 are zero, and using the relation $(\hat{J}_3 - \hat{\mathcal{L}}_3) |\phi\rangle D(\Omega) = 0$ of (3.24) and (3.28), we can show that the mean energy (3.25) reduces to

$$\langle \psi_{IM} | \hat{H} | \psi_{IM} \rangle = \int d\tau d\Omega g_0^*(\Omega) \langle \phi | \delta(q - \hat{Q}) \left[\hat{H} - \frac{1}{2} B(\hat{J}_1^2 + \hat{J}_2^2 + \hat{J}_3^2) | \phi \rangle g_0 D(\Omega) \right.$$

$$+ \frac{B}{2} \int d\tau d\Omega \langle \phi | \delta(q - \hat{Q}) | \phi \rangle g_0^* D^*(\Omega) \left[\hat{\mathcal{L}}_1^2 + \hat{\mathcal{L}}_2^2 + \hat{\mathcal{L}}_3^2 \right] g_0 D(\Omega) .$$

$$(3.36)$$

Using the approximation

$$\langle \phi | \delta(q - \hat{Q}) \left(\hat{H} - \frac{B}{2} \tilde{\vec{J}}^2 \right) | \phi \rangle \simeq \langle \phi | \delta(q - \hat{Q}) | \phi \rangle \langle \phi | \hat{H} - \frac{B}{2} \tilde{\vec{J}}^2 | \phi \rangle, \tag{3.37}$$

we can rewrite (3.36) in the following simpler form⁷

$$E_{I} = \frac{\langle \psi_{IM} | \hat{H} | \psi_{IM} \rangle}{\langle \psi_{IM} | \psi_{IM} \rangle} \approx \frac{1}{2} BI(I+1) + \langle \phi | \hat{H} - \frac{1}{2} B(\hat{J}_{1}^{2} + \hat{J}_{2}^{2} + \hat{J}_{3}^{2}) | \phi \rangle \quad ,$$
(3.38)

where we have used he fact that $\hat{\mathcal{L}}^2 \mathcal{D}^I_{MK} = I(I+1)\mathcal{D}^I_{MK}$. Note that the energy expression (3.38) has a term, $-\frac{1}{2}B\langle \vec{J}^2 \rangle$ which represents a substraction of a mean-rotational energy. This term is familiar from the standard Peierls-Yoccoz angular momentum projection method. We expect this approximate treatment of the δ -function in (3.38) to overestimate the mean-energy by a term of the order of half the zero-point vibration energy.

Moment of Inertia

Let us now look at the moment of inertia, which is given by B^{-1} . The inertial parameter B can be determined from eqs. (3.29) and (3.34); *i.e.*, it is given by the two equations

$$\langle \phi | \left[\hat{H}, \ i \hat{Z}_a \right] | \phi_{ph} \rangle = \langle \phi | \hat{J}_a | \phi_{ph} \rangle \quad , \qquad B^{-1} = \langle \phi | \left[\hat{Z}_a, \ i \hat{J}_a \right] | \phi \rangle \quad , \qquad (3.39)$$

where $\hat{Z}_a = \hat{Y}_a/B$. This expression for B^{-1} is of the well-known Thouless-Valantin form[13]. Note that if we neglect the residual two-body interactions from the Hamiltonian, expressions (3.39) would give rise to Inglis cranking formula[14]

$$\frac{1}{B} \approx 2 \sum_{\sigma\mu} \frac{|\langle \phi | \hat{J}_a | \phi_{\sigma\mu} \rangle|^2}{\mathcal{E}_{\sigma} - \mathcal{E}_{\mu}} \quad . \tag{3.40}$$

This approximate formula is well-known to overestimate the moment of inertia quite badly.

In what follows, we are going to examine the connection between the BO method and the large body of (rotational) data[15], *e.g.*, the moment of interia increases with angular momentum I. First, note that Eq. (3.38), which was derived for timereversal invariant $|\phi\rangle$, describe a rotational spectrum with constant moment of inertia, B^{-1} , in disagreement with data. To see this, consider the case K = 0 for which $\langle \phi_0 | \hat{J}_i | \phi_0 \rangle = 0$ (i = 1, 2, 3). Hence, the energy expression (9.2.20) becomes

$$E_I = \frac{1}{2}BI(I+1) + \langle \hat{H} \rangle - \Delta E_{f\ell} \quad , \tag{3.41}$$

⁷Recall that we have omitted the vibrational energy part, $E_{\rm osc}^0$.

with $\Delta E_{f\ell} = \frac{B}{2} \langle \Delta \hat{J}_1^2 + \Delta \hat{J}_2^2 \rangle$, the fluctuation energy which is generated by angular momentum fluctuations. In this case, therefore, the energy spectrum is that of a rigid rotor, since the moment of inertia B^{-1} , as given by (3.39), is constant. This contradicts, of course, the experimental facts.

Second, note that the failure of (3.38) to generate a moment of inertia, B^{-1} , which increases with angular momentum is due to a restrictive assumption on $|\phi\rangle$, the time-reversal invariance of $|\phi\rangle$ for K = 0. In what follows, we are going to show that the BO approach has a natural mechanism for introducing a moment of inertia which increases with I, provided the restrictive assumption on $|\phi\rangle$ is dropped. In addition, we will show that the energy we obtain for this case is lower than the energy, (3.41), obtained with a time-reversal invariant $|\phi\rangle$. To this end, let us consider a symmetry-violating $|\phi\rangle$ for which $\langle \hat{J}_1 \rangle$ is not zero but for which $\langle \hat{J}_2 \rangle$ and $\langle \hat{J}_3 \rangle$ are both zero. In this analysis, we will restrict ourselves to the simplest case: K = 0, and hence $|\phi_0\rangle$ is an eigenfunction to \hat{J}_3 with eigenvalue zero, $\hat{J}_3 |\phi_0\rangle \equiv 0$. In this case, the energy expression (3.38) reduces to

$$E_I = \frac{1}{2}BI(I+1) + \langle \hat{H} \rangle - \frac{B}{2} \langle \hat{J}_1 \rangle^2 - \Delta E_{f\ell} \quad . \tag{3.42}$$

Note that this energy is lower than the energy, (3.41), obtained with a time-reversal invariant $|\phi\rangle$ (provided the fluctuation energy is unchanged). In what follows, we shall neglect the angular momentum fluctuations, $\Delta E_{f\ell}$, in the determination of the mean field $|\phi\rangle$. The energy expression (3.42) then provides a basis for a variational determination of the symmetry violating $|\phi\rangle$:

$$\delta \langle \hat{H} - \omega \hat{J}_1 \rangle_{\omega} = 0 \quad , \qquad \omega = B \langle \hat{J}_1 \rangle_{\omega} \quad ,$$
 (3.43)

where the notation $\langle \hat{0} \rangle_{\omega}$ is used to abbreviate $\langle \phi(\omega) | \hat{0} | \phi(\omega) \rangle$. The parameter ω has, obviously, the significance of an angular velocity.

Now, we are in a position to show that $B(\omega)$ decreases when the angular momentum increases. To see this, using the relation $B(\omega) = \frac{\omega}{\langle \hat{J}_1 \rangle_{\omega}}$ of (3.43), we have

$$\frac{dB(\omega)}{d\omega} = \frac{1}{\langle \hat{J}_1 \rangle_{\omega}} \left[1 - \frac{1}{\langle \hat{J}_1 \rangle/\omega} \frac{d\langle \hat{J}_1 \rangle_{\omega}}{d\omega} \right] \quad . \tag{3.44}$$

Since $\langle \hat{J}_1 \rangle_{\omega}$ is well-known to increase with the angular velocity ω , and as shown in Fig. 1, the slope $\frac{d\langle \hat{J}_1 \rangle_{\omega}}{d\omega}$ is always larger than $\frac{\langle \hat{J}_1 \rangle_{\omega}}{\omega}$. Thus, the slope, $\frac{dB}{d\omega}$ of $B(\omega)$ is negative and, hence, $B(\omega)$ would behave as shown in Fig. 2: $B(\omega)$ decreases as the angular velocity ω increases.



Figure 1: $\langle \hat{J}_1 \rangle_{\omega}$ increases with the angular velocity ω .

Therefore, we conclude that the moment of inertia $B^{-1}(\omega)$ increases, indeed, with angular velocity ω , and hence with angular momentum also.

In what follows, we are going to show that there exists a non-zero value, ω_c , of ω at which the energy $E_I(\omega_c)$ of (3.42) is equal to its lowest value. To this end, let us write the energy expression (3.42) in the following form (from which we omit the fluctuation term, $\Delta E_{f\ell}$):

$$E_{I}(\omega) \simeq \langle \hat{H} - \omega \hat{J}_{1} \rangle_{\omega} + \frac{1}{2} B(\omega) \left[\langle \hat{J}_{1} \rangle_{\omega}^{2} + I(I+1) \right]$$

$$= \langle \hat{H} - \omega \hat{J}_{1} \rangle_{\omega} + \frac{1}{2} \left[\frac{\omega^{2}}{B(\omega)} + B(\omega) I(I+1) \right] \quad . \quad (3.45)$$

First, note that the derivative,

$$\frac{dE_I(\omega)}{d\omega} = \frac{1}{2} \left[I(I+1) - \langle \hat{J}_1 \rangle_{\omega}^2 \right] \frac{dB(\omega)}{d\omega} \quad , \tag{3.46}$$

of $E_I(\omega)$ vanishes at a value ω_c which is determined by $\langle \hat{J}_1 \rangle^2_{\omega_c} = I(I+1)$, i.e.,

$$\frac{dE_I(\omega)}{d\omega}\Big|_{\omega_c} = 0 \qquad \Longrightarrow \qquad \langle \hat{J}_1 \rangle_{\omega_c}^2 = I(I+1) \quad . \tag{3.47}$$

Second, we can easily show that the second derivative of $E_I(\omega)$,

$$\frac{d^2 E_I(\omega)}{d\omega^2}\Big|_{\omega=\omega_c} = \frac{\omega}{B^2(\omega)} \left[\frac{\omega}{B(\omega)} \left(\frac{dB(\omega)}{d\omega}\right)^2 - \frac{dB(\omega)}{d\omega}\right]\Big|_{\omega=\omega_c} \quad , \tag{3.48}$$



Figure 2: The inertial parameter $B(\omega)$ decreases as the angular velocity ω increases, since $dB/d\omega < 0$.

is positive, since, as shown above (c.f. Eq. (3.44)), $\frac{dB}{d\omega}$ is negative. Finally, we conclude that, using a trial function $|\phi(\omega)\rangle$ whose time-reversal symmetry is broken, one obtains, indeed, lower values for the inverse moment of inertia, $B(\omega)$, and for the energy than those calculated with a $T \cdot R$ invariant mean field.

Calculation of the energy difference: ΔE_I

Let us now calculate the energy difference, ΔE_I , between $E_I(\omega = 0)$ and $E_I(\omega_c)$. Using Eq. (3.46), we can show that

$$\Delta E_I = E_I(\omega_c) - E_I(0) = \int_0^{\omega_c} \frac{dE_I(\omega)}{d\omega} d\omega$$

= $\frac{1}{2} \left[B(\omega_c) - B(0) \right] I(I+1) + \frac{1}{2} \int_0^{\omega_c} \omega^2 \frac{d}{d\omega} \left(\frac{1}{B(\omega)} \right) d\omega$. (3.49)

This expression can, after a partial integration, be reduced to

$$\Delta E_{I} = -\frac{1}{2} B_{0} I(I+1) + \int_{0}^{\sqrt{I(I+1)}} \omega\left(\langle \hat{J}_{1} \rangle\right) d\langle \hat{J}_{1} \rangle$$
$$= -\int_{0}^{\sqrt{I(I+1)}} \omega\left(\langle \hat{J}_{1} \rangle\right) \left[\frac{B}{B(\omega)} - 1\right] d\langle \hat{J}_{1} \rangle \quad . \tag{3.50}$$

This expression shows that $E_I(\omega_c)$ is, indeed, lower than $E_I(0)$, since B_0 is larger than $B(\omega)$. So, if we know the dependence of the angular velocity ω on $\langle \hat{J}_1 \rangle_{\omega}$, we



Figure 3: Behavior of $E_I(\omega)$ as a function of the angular velocity ω , where $|\Delta E_I| = |E_I(0) - E_I(\omega_c)|$ is an increasing function of the angular momentum I

can easily calculate the energy difference between $E_I(0)$ and $E_I(\omega_c)$. Note that, the energy difference $|\Delta E_I|$ is an increasing function of the angular momentum I. The qualitative behavior of the energy $E_I(\omega)$, for various values of I, is plotted in Fig. 3.

4 Discussion and Conclusions

Let us summarize what we have achieved in this work. First, we have shown that the moment of inertia generated by the NBO method is identical to the Thouless-Valantin form. Second, the two relations (3.43) and (3.47) determine the intrinsic (symmetry-breaking) function $|\phi_{\omega}\rangle$ and the value, ω_c , of ω where $E_I(\omega_c)$ is equal to its lowest value, respectively. These two relations provide a bridge (connection) between the NBO method, which is a truly quantum mechanical description of collective motion, and the semi-classical approaches based on the idea of self-consistent cranking (SCC). Thus, we have established a connection between the NBO approach and the large body of experimental data, since the two relations (3.43) and (3.47) are known to provide reasonable descriptions of vast amounts of empirical data ranging from low-

lying rotational states to high angular momentum states.[16]–[21] So, the present (NBO) method appears to be well-equipped to describe low as well as high lying rotational states. Additionally, we should mention that work has been started to apply the NBO method to the description of the backbending phenomenon which was first observed by Johnson and his collaborators.[22]

In summary, we have studied here the rotational spectrum of even, axially-symmetric nuclei within the framework of the NBO method. We have made use of trial functions in which the intrinsic structure is described within a mean-field approximation. We have shown that the NBO formalism gives back the Thouless-Valantin moment of inertia. Then, we have established a connection between the NBO method and the SCC model, which has been successful in reproducing vast amounts of experimental data. Finally, we have shown that the introduction of a non time-reversal invariant intrinsic function both lowers the energy for a given I, and provides a moment of inertia that increases with the angular momentum I.

Acknowledgments

Supported in part by the Alabama Commission on Higher Education.

References

- [1] M. Born and J. R. Oppenheimer, Ann. Phys. 84 (1927) 457.
- [2] M. Moshinsky and C. Kittel, Proc. Nat. Acad. Sci. U.S.A., Vol. 60, no. 4, 1110 (Aug. 1968).
- [3] F. Villars, Nucl. Phys. A420 (1984) 61.
- [4] F. Villars, Nucl. Phys. A473 (1987) 539.
- [5] N. Zettili and F. Villars, Nucl. Phys. A469 (1987) 77.
- [6] N. Zettili and F. Villars, Nucl. Phys. A469 (1987) 93.
- [7] N. Zettili, Mod. Phys. Let. A, Vol. 6, No. 3, 183 (1991).
- [8] N. Zettili, Nucl. Phys. A576 (1994) 1.
- [9] N. Zettili, Phys. Rev. C., Vol. 51, No. 4, 1777 (1995).

- [10] N. Zettili and A. Boukahil, Int. Jour. Mod. Phys. E, Vol. 11, No. 3, (2002) 231.
- [11] A. Bohr, Mat. Fys. Medd. Dan. Vid. Selsk. 26, 14 (1952);
 A. Bohr and B. R. Mottelson, Mat. Fys. Medd. Dan. Vid. Selsk. 27, 16 (1953);
 A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol.1, (Benjamin: Reading, 1975).
- [12] J. P. Blaizot and G. Ripka, Quantum Theory of Finite Systems (MIT Press: Cambridge, 1986), p. 253;
 P. Ring and P. Schuck, The Nuclear Many-Body Problem (Springer Verlag, NY: 1980), p. 578.
- [13] D. J. Thouless and J. G. Valantin, Nucl. Phys. 31, 211 (1962).
- [14] D. R. Inglis, *Phys. Rev.* **96**, 1059 (1954); **97**, 701 (1955).
- [15] M. J. A. de Voigt, J. Dudek and Z. Szymanski, Rev. Mod. Phys. 55, 949 (1983).
- [16] J. L. Egido, et al., Nucl. Phys. A339, 390 (1980).
- [17] L. L. Riedinger, Nucl, Phys. A347, 141 (1980).
- [18] A. Faessler et al., Nucl. Phys. A256, 106 (1976).
- [19] I. Hamanoto, Nucl Phys. A271, 15 (1976).
- [20] A. L. Goodman, Nucl. Phys. A230, 466 (1974).
- [21] B. Banerjee, H. J. Mang and P. Ring, Nucl. Phys. A215, 366 (1973).
- [22] A. Johnson, H. Ryde and S. Hjorth, Nucl. Phys. A179, 753 (1972).
- [23] N. Zettili, Quantum Mechanics: Concepts and Applications, 2nd Ed., (John-Wiley: Chichester, 2009).