Multiple scattering of matter waves: an analytic model of the refractive index for atomic and molecular gases

Mikhail Lemeshko^{*} and Bretislav Friedrich[†]

Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

(Dated: September 27, 2018)

We present an analytic model of the refractive index for matter waves propagating through atomic or molecular gases. The model, which combines a WKB treatment of the long range attraction with the Fraunhofer model treatment of the short range repulsion, furnishes a refractive index in compelling agreement with recent experiments of Jacquey *et al.* [Phys. Rev. Lett. **98**, 240405 (2007)] on Li atom matter waves passing through dilute noble gases. We show that the diffractive contribution, which arises from scattering by a two dimensional "hard core" of the potential, is essential for obtaining a correct imaginary part of the refractive index.

PACS numbers: 03.75.Dg, 03.65.Nk, 34.50.-s

Keywords: Refraction of matter waves, Multiple scattering, Atom interferometry, Scattering theory, Models of atomic and molecular collisions, Diffraction.

I. INTRODUCTION

Multiple scattering of matter waves arises in a broad variety of contexts, ranging from gas transport [1] to the scattering of subatomic particles by atoms or molecules [2, 3] to cosmic ray showers [4]. Inherent to the concept of multiple matter wave scattering is the notion of a refractive index, n, which, by the optical theorem [5], is proportional to the forward scattering amplitude, f(0). While Im[f(0)] and thus Im[n] are related to the total integral scattering cross section and so can be measured in a standard beam scattering experiment, $\operatorname{Re}[f(0)]$ and thus $\operatorname{Re}[n]$ reflect the change of the wave's phase velocity and can only be accessed by matter wave interferometry. The first such interferometric measurement was carried out in 1995 by Schmiedmayer et al. [6, 7], who determined both the attenuation and the phase velocity change of sodium atom waves propagating through a number of gases. Subsequently, the Pritchard group was able to observe glory-type oscillations in the dependence of the refractive index on the sodium beam velocity [8-10]. In 2007, Jacquey et al. [11] implemented an improved, Mach-Zehner type of an atom interferometer and measured the index of refraction of lithium waves passing through noble gases. In 2008, Champenois et al. [12] and, independently, Hornberger and Vaccini [13], provided an analysis of how the motion of the scatterers affects the measured refractive index. This work showed that the formulae used to extract the refractive index from the experimental data [7, 14] and in the related theoretical treatments [15–18] lacked consistency with the Beer-Lambert law and thus were incorrect.

According to the scenario of Ref. [12], an atomic or ionic beam of "projectiles" p of mass m_p and the laboratory velocity \mathbf{v}_p propagates through a gas of density N_t made out of "target" particles t of mass m_t and laboratory velocity \mathbf{v}_t . For plane matter waves, the velocities $\mathbf{v}_{p,t}$ are related to the laboratory wave vectors $\mathbf{k}_{p,t}$ by $\hbar \mathbf{k}_{p,t} = m_{p,t} \mathbf{v}_{p,t}$. The relative motion of the target and projectile particles is described by the relative wave vector $\mathbf{k}_r = \mu \mathbf{v}_r / \hbar$, where $\mathbf{v}_r = \mathbf{v}_p - \mathbf{v}_t$ is the relative velocity, and $\mu = m_p m_t / (m_p + m_t)$ the reduced mass. Taking this scenario into account, Champenois *et al.* [12] derived a new formula for the refractive index of a matter wave propagating through a dilute gas

$$n = 1 + 2\pi N_t \frac{m_p + m_t}{m_t} \frac{\langle f(k_r, 0) \rangle}{k_p^2},$$
 (1)

where $f(k_r, 0) \equiv f(0)$ is the forward scattering amplitude pertaining to the scattering angle $\vartheta = 0$. The averaging in Eq. (1) is carried out over the relative wave vectors, k_r , corresponding to a normalized relative velocity distribution [12], see below.

Previous theoretical treatments of the refractive index of atomic waves, surveyed in Ref. [12], were based on WKB [7] or eikonal [10, 19] approximations, which become analytic only if the repulsive, short-range interaction, is neglected, cf. also refs. [7, 12]. In this contribution, we combine the WKB treatment of scattering for the long-range attraction with the Fraunhofer approximation for the short-range repulsion and thereby obtain an analytic model of the refractive index which accounts for both short- and long-range multiple matter-wave scattering. The resulting analytic refractive index provides an additional insight into multiple scattering and facilitates data analysis.

After briefly introducing the model, we apply it to the refractive index of Li atom waves propagating through Ar, Kr, and Xe gases and compare the results with the measurements of Jacquey *et al.* [11]. In addition, we exemplify the model's scope by examining the refraction of a Na⁺ ionic beam passing through a dilute N₂ gas.

^{*} mikhail.lemeshko@gmail.com

[†] brich@fhi-berlin.mpg.de

II. MODEL OF MATTER-WAVE REFRACTION

In the model introduced herein, we consider the scattering amplitude to consist of two parts:

$$f(k_r, 0) = f_{\text{short}}(k_r, 0) + f_{\text{long}}(k_r, 0)$$
 (2)

The short range part, $f_{\rm short}(k_r, 0)$, arises from the scattering by the "hard core," repulsive branch of the potential, and can be evaluated in closed form within the Fraunhofer model of matter wave scattering. The Fraunhofer model was introduced by Drozdov [20] and generalized by Blair [21] in the late 1950s to treat inelastic nuclear scattering, and adapted by Faubel [22] to account for rotationally inelastic thermal collisions between helium atoms and N₂ and CH₄ molecules. Recently, we generalized the Fraunhofer model to treat atom-molecule collisions in electric [23, 24], magnetic [25], and laser fields [26], and also to gain insight into the stereodynamics of molecular collisions [27, 28].

Within the Fraunhofer model, the hard core part of the potential amounts to a two-dimensional, impenetrable obstacle with sharp boundaries. The elastic scattering amplitude is given by the amplitude for the Fraunhofer diffraction of matter waves by such an obstacle [23],

$$f_{\rm short}(k_r,\vartheta) = ik_r R_0^2(k_r) \frac{J_1[k_r R_0(k_r)\vartheta]}{k_r R_0(k_r)\vartheta},\qquad(3)$$

where J_1 is the Bessel function [29] and R_0 is the radius of the interaction which, for an atom-atom potential, is given by the solution of the equation $V(R_0) = \hbar^2 k_r^2/(2\mu)$, with V(r) the interaction potential. For forward scattering, the amplitude of Eq. (3) becomes

$$f_{\rm short}(k_r, 0) = \frac{ik_r R_0^2(k_r)}{2}.$$
 (4)

Hence the Fraunhofer amplitude is purely imaginary and, therefore, contributes only to the imaginary part of the refractive index.

In order to account for the long-range part of the scattering amplitude, $f_{long}(k_r, 0)$, we make use of the WKB approximation, which is accurate for thermal collisions dominated by partial waves with large angular momenta l. We consider the general case of the inverse-power longrange potential,

$$V_{\rm long}(r) = -\frac{C_{\beta}}{r^{\beta}},\tag{5}$$

with $C_{\beta} > 0$ and $\beta > 3$. For an atom-atom interaction, $\beta = 6$, while $\beta = 4$ for an ion interacting with an atom or molecule. For the potential of Eq. (5), the WKB phase shift for the *l*-th partial wave is given by [5]

$$\delta_l^{(\beta)} = \frac{\mu C_\beta k_r^{\beta-2}}{2\hbar^2 l^{\beta-1}} \frac{\Gamma(\frac{1}{2})\Gamma(\frac{\beta-1}{2})}{\Gamma(\frac{\beta}{2})}.$$
 (6)

The WKB forward scattering amplitude [5],

$$f_{\text{WKB}}^{(\beta)}(k_r, 0) = \frac{1}{ik_r} \int_0^\infty l\left(e^{2i\delta_l^{(\beta)}} - 1\right) dl$$
(7)

then becomes, on substituting from Eq. (6):

$$f_{\text{long}}^{(\beta)}(k_r, 0) = k_r^{\frac{\beta-3}{\beta-1}} \left(\frac{\sqrt{\pi}}{2}\right)^{\frac{\beta+1}{\beta-1}} \left[\frac{\mu C_\beta}{\hbar^2} \frac{\Gamma(\frac{\beta-1}{2})}{\Gamma(\frac{\beta}{2})}\right]^{\frac{2}{\beta-1}} \times \left[\frac{\Gamma(\frac{3-\beta}{2-2\beta})}{\Gamma(\frac{1}{1-\beta})} + i\frac{\Gamma(\frac{\beta-2}{\beta-1})}{\Gamma(\frac{\beta+1}{2\beta-2})}\right] \quad (8)$$

III. REFRACTION OF ATOM MATTER WAVES

In the case of an atom beam passing through an atomic gas, $\beta = 6$ and the forward scattering amplitude becomes

$$f_{\text{atom}}(k_r, 0) = f_{\text{short}}(k_r, 0) + f_{\text{long}}^{(6)}(k_r, 0)$$
$$= \frac{ik_r R_0^2(k_r)}{2} + \left(\frac{ik_r}{2}\right)^{3/5} \Gamma\left(\frac{3}{5}\right) \left(\frac{3\pi}{16} \frac{\mu C_6}{\hbar^2}\right)^{2/5}$$
(9)

By substituting from Eq. (9) into Eq. (1) and making use of the distribution function given by Eq. (A3) of Ref. [12],

$$P(v_r) = \frac{2v_r}{\pi^{1/2}\alpha v_p} \exp\left[-\frac{v_p^2 + v_r^2}{\alpha^2}\right] \sinh\left[\frac{2v_p v_r}{\alpha^2}\right] \quad (10)$$

with $\alpha = \sqrt{2k_BT/m_t}$, k_B Boltzmann's constant, and T the temperature, we obtain the following expression for the reduced index of refraction,

$$\eta_{\text{atom}} \equiv \frac{n-1}{N_t}$$
$$= 2\pi \frac{\hbar}{m_p v_p^2} \left[i \langle v_r \rangle R_0^2 + \langle v_r^{3/5} \rangle \left(\frac{i}{2}\right)^{3/5} \Gamma\left(\frac{3}{5}\right) \left(\frac{3\pi}{16} \frac{C_6}{\hbar}\right)^{2/5} \right]$$
(11)

where the velocity averages are given by

$$\langle v_r \rangle = \int_0^\infty v_r P(v_r) dv_r, \qquad (12)$$

$$\langle v_r^{3/5} \rangle = \int_0^\infty v_r^{3/5} P(v_r) dv_r.$$
(13)

In general, the R_0 value depends on the relative velocity v_r , but for narrow velocity distributions, such as those implemented in the experiments of Jacquey *et al.* [11], the dependence is found to be negligible. We could thus simplify the resulting expression by fixing $R_0(v_r)$ to a constant value pertaining to the mean relative velocity, $R_0 \equiv R_0(\langle v_r \rangle)$.

While Table I compares the experimental values of η_{atom} obtained by Jacquey *et al.* [11] with our model results for Li matter waves propagating through Ar, Kr, and Xe gases, Figure 1 singles out the Li+Xe system and




FIG. 1. (Color online) Upper panel shows the real and imaginary parts of the reduced refractive index $\eta \equiv (n-1)/N_t$ $[m^3]$ multiplied by $10^{25}v_p^{7/5}$ for Li beam propagating through Xe gas, in dependence of the beam velocity v_p . Lower panel shows the corresponding ratio Re $(\eta)/\text{Im}(\eta)$. Model results (solid curves) are compared with experimental data of Ref. [11]. Dashed curves show the results of the model with the short-range potential excluded.

TABLE I. Reduced index of refraction, $\eta \equiv (n-1)/N_t$ [m³], for a Li beam propagating through different noble gases with a mean velocity $v_p = 1075$ m/s.

	Experiment	Model	Model	
	of Ref. [11]		$f_{\rm short}(k_r,0)\equiv 0$	
Ar				
20				
$10^{29} \mathrm{Re}(\eta)$	1.20 ± 0.11	0.97	0.97	
$10^{29} \mathrm{Im}(\eta)$	2.11 ± 0.06	1.65	1.33	
${\rm Re}(\eta)/{\rm Im}(\eta)$	0.56 ± 0.05	0.58	0.73	
Kr				
$10^{29} \mathrm{Re}(\eta)$	1.57 ± 0.10	1.25	1.25	
$10^{29} \mathrm{Im}(\eta)$	1.99 ± 0.07	2.04	1.71	
${\rm Re}(\eta)/{\rm Im}(\eta)$	0.78 ± 0.04	0.61	0.73	
Xe				
$10^{29} \mathrm{Re}(\eta)$	1.82 ± 0.07	1.52	1.52	
$10^{29} \mathrm{Im}(\eta)$	2.40 ± 0.06	2.42	2.08	
${\rm Re}(\eta)/{\rm Im}(\eta)$	0.70 ± 0.03	0.62	0.72	

be in a compelling agreement with the experiment. We note that the interference of the matter waves scattered by the short- and long-range parts of the interatomic potential causes the refractive index to exhibit glory-type oscillations. These are not correctly rendered by our analytic model, as the approximate scattering amplitudes, which were obtained by different methods, fail to "keep track" of one another.

Also presented in both Table I and Figure 1 are the values of the refractive index for the long range potential included but the short-range interaction excluded, i.e., for $f(k_r, 0) = f_{\text{long}}(k_r, 0)$, cf. Eq. (2). One can see that in such a case the imaginary part of the refractive index is substantially less than in the experiment.

In the limit of a cold atomic gas, i.e., for $T \to 0$, the velocity distribution (10) becomes a δ -function, which simplifies Eq. (11), since then $\langle v_r \rangle \to v_p$ and $\langle v_r^{3/5} \rangle \to v_p^{3/5}$. In this case, $\operatorname{Re}[\eta_{\operatorname{atom}}]$ is proportional to $v_p^{-7/5}$, while $\operatorname{Im}[\eta_{\operatorname{atom}}]$ is proportional to both v_p^{-1} and $v_p^{-7/5}$. However, since the v_p^{-1} term arises from the diffraction of matter waves from the hard core of the potential, it dominates $\operatorname{Im}[\eta_{\operatorname{atom}}]$ at large collision energies at the expense of the $v_p^{-7/5}$ term, which arises from the long-range attraction.

IV. REFRACTION OF ION MATTER WAVES

For an ion beam passing through an atomic or molecular gas, $\beta = 4$ and the scattering amplitude becomes

$$f_{\rm ion}(k_r,0) = f_{\rm short}(k_r,0) + f_{\rm long}^{(4)}(k_r,0) = \frac{ik_r R_0^2(k_r)}{2} + \left(\frac{ik_r}{2}\right)^{1/3} \Gamma\left(\frac{1}{3}\right) \left(\frac{\pi}{4} \frac{\mu C_4}{\hbar^2}\right)^{2/3}$$
(14)

leading to a reduced refractive index for an ionic beam:

$$\eta_{\rm ion} \equiv \frac{n-1}{N_t} = 2\pi \frac{\hbar}{m_p v_p^2} \left[i \langle v_r \rangle R_0^2 + \langle v_r^{1/3} \rangle \left(\frac{i}{2}\right)^{1/3} \Gamma\left(\frac{1}{3}\right) \left(\frac{\pi}{4} \frac{C_4}{\hbar}\right)^{2/3} \right],$$
(15)

with

$$\langle v_r^{1/3} \rangle = \int_0^\infty v_r^{1/3} P(v_r) dv_r \tag{16}$$

We note that in the case of an anisotropic interaction of, say, an ion or atom with a linear molecule given by a potential $V(r, \theta)$, the R_0 value is given by the spherical part of the hard core of the potential [23]. In order

TABLE II. Reduced index of refraction, $\eta \equiv (n-1)/N_t$ [m³], for a Na⁺ beam propagating through N₂ gas with a mean velocity $v_p = 5000$ m/s.

	Model	$\begin{array}{c} \text{Model} \\ f_{\text{short}}(k_r, 0) \equiv 0 \end{array}$
$\frac{10^{29} \text{Re}(\eta)}{10^{29} \text{Im}(\eta)}$ $\frac{\text{Re}(\eta)/\text{Im}(\eta)}{\text{Re}(\eta)}$	0.25 0.16 1.60	0.25 0.14 1.73

to extract R_0 , we first solve the equation $V(R(\theta), \theta) = \hbar^2 k_r^2/(2\mu)$ to obtain the hard core shape $R(\theta)$, which we then expand in a Legendre series,

$$R(\theta) = R_0 P_0(\cos \theta) + R_1 P_1(\cos \theta) + R_2 P_2(\cos \theta) + \dots,$$
(17)

from which we deduce the R_0 value.

The values of the refractive index for an ion beam passing through a molecular gas, as exemplified by the Na⁺-N₂ system, are listed in Table II at an ion projectile velocity $v_p = 5000$ m/s. The requisite value of R_0 was extracted from the potential energy surface of Ref. [31]. We see that the refractive index is less affected by switching off the repulsive part of the potential for the ion-molecule system than for the atom-atom system, cf. section III. This is due to the greater strength of the $-C_4r^{-4}$ interaction compared with that of the $-C_6r^{-6}$ potential and the more dominant role it thus plays in determining n.

We note that in the limit of zero temperature, $v_r \to v_p$,

- A. D. Cronin, J. Schmiedmayer, and D. E. Pritchard, Reviews of Modern Physics 81, 1051 (Jul 2009).
- [2] H. Rauch, in Advances in Quantum Phenomena, NATO Advanced Science Institutes, Series B. Physics, Vol. 347, edited by E. G. Beltrametti and J.-M. Lévy-Leblond (Plenum, New York, 1995) p. 113.
- [3] A. F. Borghesani, IEEE Transactions on Dielectrics and Electrical Insulations 13, 492 (2006).
- [4] L. Anchordoqui, M. T. Dova, A. Mariazzi, T. McCauley, T. Paul, S. Reucroft, and J. Swain, Annals of Physics **314**, 145 (Nov 2004).
- [5] L. D. Landau and E. M. Lifshitz, Quantum Mechanics, Non-Relativistic Theory (Oxford: Pergamon, 1977).
- [6] J. Schmiedmayer, C. R. Ekstrom, M. S. Chapman, T. R. Hammond, and D. E. Pritchard, in *Fundamentals of Quantum Optics III*, Lecture Notes In Physics No. 420, edited by F. Ehlotzky (Springer, Berlin, 1993) p. 21.
- [7] J. Schmiedmayer, M. S. Chapman, C. R. Ekstrom, T. D. Hammond, S. Wehinger, and D. E. Pritchard, Physical Review Letters 74, 1043 (Feb 1995).
- [8] J. Schmiedmayer, M. S. Chapman, C. R. Ekstrom, T. D. Hammond, D. A. Kokorowski, A. Lenef, R. A. Rubenstein, E. T. Smith, and D. E. Pritchard, in *Atom Inter-*

the real part of the refractive index (15) becomes proportional to $v_p^{-5/3}$. The imaginary part of the refractive index is determined by two terms, one of which is proportional to v_p^{-1} and the other to $v_p^{-5/3}$.

V. CONCLUSIONS

We presented an analytic model of the refraction of atom or ion matter waves passing through atomic or molecular gases. The values of the refractive index furnished by the model were found to be in good agreement with experiments of Jacquey *et al.* [11]. Our analysis has shown that in order to appraise the imaginary part of the refractive index correctly, we need to account for the diffraction by the repulsive hard core part of the interaction potential. This can be achieved by invoking the Fraunhofer model.

Within the Fraunhofer model of matter wave scattering, the elastic scattering amplitude depends solely on the spherical radius, R_0 , of the scatterer. As a result, an external field (e.g., magnetic), does not affect the elastic cross section furnished by the model. The corollary is that the field would not affect the index of refraction obtained from the analytic model presented herein either.

VI. ACKNOWLEDGEMENTS

We thank Jacque Vigué for discussions and to Gerard Meijer for encouragement and support.

ferometry, edited by P. R. Berman (Academic Press, New York, 1997) p. 1.

- [9] T. D. Hammond, M. S. Chapman, A. Lenef, J. Schmiedmayer, E. T. Smith, R. A. Rubenstein, D. A. Kokorowski, and D. E. Pritchard, Brazilian Journal of Physics 27, 193 (1997).
- [10] T. D. Roberts, A. D. Cronin, D. A. Kokorowski, and D. E. Pritchard, Physical Review Letters 89, 200406 (Oct 2002).
- [11] M. Jacquey, M. Büchner, G. Trénec, and J. Vigué, Physical Review Letters 98, 240405 (Jun 2007).
- [12] C. Champenois, M. Jacquey, S. Lepoutre, M. Büchner, G. Trénec, and J. Vigué, Physical Review A 77, 13621 (Jan 2008).
- [13] K. Hornberger and B. Vacchini, Physical Review A 77, 22112 (Feb 2008).
- [14] J. Vigué, Physical Review A 52, 3973 (Nov 1995).
- [15] R. C. Forrey, L. You, V. Kharchenko, and A. Dalgarno, Physical Review A 54, 2180 (Sep 1996).
- [16] C. Champenois, E. Audouard, P. Duplàa, and J. Vigué, Journal de Physique II 7, 523 (Apr 1997).
- [17] V. Kharchenko and A. Dalgarno, Physical Review A 63, 23615 (Feb 2001).

- [18] R. C. Forrey, V. Kharchenko, and A. Dalgarno, Journal of Physics B 35, L261 (Jun 2002).
- [19] S. Blanchard, D. Civello, and R. C. Forrey, Physical Review A 67, 13604 (Jan 2003).
- [20] S. I. Drozdov, Soviet Physics JETP 1, 588, 591 (Jan 1955).
- [21] J. S. Blair, in *Lectures in Theoretical Physics VIII C Nuclear Structure Physics*, edited by P. D. Kunz, D. A. Lind, and W. E. Brittin (The University of Colorado Press, Boulder, 1965) pp. 343–444.
- [22] M. Faubel, Journal of Chemical Physics 81, 5559 (Jan 1984).
- [23] M. Lemeshko and B. Friedrich, Journal of Chemical Physics **129**, 4301 (Jul 2008).
- [24] M. Lemeshko and B. Friedrich, Journal of Physical Chemistry A 113, 15055 (2009).

- [25] M. Lemeshko and B. Friedrich, Physical Review A 79, 012718 (Jan 2009).
- [26] M. Lemeshko and B. Friedrich, International Journal of Mass Spectrometry 280, 19 (Feb 2009).
- [27] M. Lemeshko and B. Friedrich, Physical Chemistry Chemical Physics, 1(Jan 2010).
- [28] M. Lemeshko, P. G. Jambrina, M. P. de Miranda, and B. Friedrich, Journal of Chemical Physics, submitted, arXiv: 1002.1572(2010).
- [29] G. N. Watson, *Theory of Bessel Functions* (Cambridge University Press, Cambridge, 1922).
- [30] J. Ahokas, T. Kiljunen, J. Eloranta, and H. Kunttu, Journal of Chemical Physics 112, 2420 (Feb 2000).
- [31] P. Soldán, V. Spirko, E. Lee, and T. Wright, Journal of Chemical Physics 111, 3420 (Jan 1999).