

# MPEMBA EFFECT, NEWTON COOLING LAW AND HEAT TRANSFER EQUATION

Vladan Panković, Darko V. Kapor  
 Department of Physics, Faculty of Sciences, 21000 Novi Sad,  
 Trg Dositeja Obradovića 4, Serbia,  
 vladan.pankovic@df.uns.ac.rs

PACS number: 05.70.-a

## Abstract

In this work we suggest a simple theoretical solution of the Mpemba effect in full agreement with known experimental data. This solution follows simply as an especial approximation (linearization) of the usual heat (transfer) equation, precisely linearization of the second derivation of the space part of the temperature function (as it is well-known Newton cooling law can be considered as the effective approximation of the heat (transfer) equation for constant space part of the temperature function).

In this work we shall suggest a simple theoretical solution of the Mpemba effect in full agreement with known experimental data [1]-[4]. This solution follows simply as an especial approximation (linearization) of the usual heat (transfer) equation, precisely linearization of the second derivation of the space part of the temperature function (as it is well-known Newton cooling law [5] can be considered as the effective approximation of the heat (transfer) equation for constant space part of the temperature function).

As it is well-known [5] usual Newton's law of cooling represents the following simple linear differential equation

$$\frac{dT}{dt} = -\alpha(T - T_e) \quad (1)$$

with simple solution

$$T = (T_0 - T_e) \exp[-\alpha t] + T_e. \quad (2)$$

Here  $T$  represents the temperature of a liquid with initial value  $T_0$ ,  $T_e$  - constant, smaller than  $T$ , temperature of a cooling wall (thermostat or environment) in the contact with liquid, while

$\alpha$  represents a cooling parameter that simplifiedly expresses thermodynamics of the cooling process. It is supposed implicitly that cooling parameter is independent of the initial temperature of the liquid as well as of the cooling wall temperature.

Consider now described by (1) or (2) thermodynamical interaction between liquid and cooling wall in two different cases, precisely for two different initial temperatures of the liquid, first one  $T_{10}$  and second one  $T_{20}$  so that  $T_{10}$  is larger than  $T_{20}$  , i.e.

$$T_{10} > T_{20}. \quad (3)$$

Then (2), (3) imply

$$\frac{T_1 - T_e}{T_2 - T_e} = \frac{T_{10} - T_e}{T_{20} - T_e} \quad (4)$$

where  $T_1$  and  $T_2$  represents the temperature of the liquid in the first case and second case in some (finite) later time moment  $t$ . It means that temperature of the liquid in the first case will be larger than the temperature of the liquid in the second case in any (finite) later time moment.

Previous conclusion implies unambiguously that usual Newton's cooling law (1) cannot model experimental data of the Mpemba effect [1]-[4] according to which, simply speaking, hot liquid freezes faster than cold liquid.

Consider, however, the standard (one-dimensional) heat transfer differential equation

$$\frac{\partial^2 T}{\partial x^2} = k \frac{\partial T}{\partial t}. \quad (5)$$

Here  $x$  represents the coordinate (along  $x$ -axis) between 0 and linear dimension of the vessel  $L > 0$ ,  $t$  - time moment between 0 and  $\infty$ , and  $k$  - corresponding (positive) constant, so that liquid temperature  $T$  represents the function of  $x$  and  $t$ . This equation, of course, represents mathematically partial differential equation of the second range.

Suppose additionally that  $T$  satisfies the following two, initial and "final" conditions,

$$T(x, 0) = T_0 \quad (6)$$

and

$$T(x, t) = T_e \quad (7)$$

for  $t \gg 0$  .

As it is well-known solution of heat equation (5) can be supposed in the following form

$$T(x, t) = X(x)\tau(t) + A \quad (8)$$

where  $X(x)$  represents the space part of  $T(x, t)$  depending of  $x$  only,  $\tau(t)$  - time part of  $T(x, t)$  depending of  $t$  only and  $A$  some constant.

Introduction of (8) in (5) yields

$$\tau \frac{d^2 X}{dx^2} = k X \frac{d\tau}{dt} \quad (9)$$

and further

$$\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{k}{\tau} \frac{d\tau}{dt} = -\beta \quad (10)$$

where  $\beta$  represents some (positive) constant. It, in fact, represents a system of two usual differential equations with simple solutions

$$\tau = B \exp\left[-\frac{\beta}{k}t\right] \quad (11)$$

$$X = C \cos[\beta^{\frac{1}{2}}x] + D \sin[\beta^{\frac{1}{2}}x] \quad (12)$$

where  $B$ ,  $C$  and  $D$  are some constants.

Then, according to (8), it follows

$$T = \exp\left[-\frac{\beta}{k}t\right](BC \cos[\beta^{\frac{1}{2}}x] + BD \sin[\beta^{\frac{1}{2}}x]) + A \quad (13)$$

which, according to initial and "final" conditions (6), (7) that imply

$$A = T_e \quad (14)$$

$$BC = BD = T_0 - T_e \quad (15)$$

, turns out in

$$T = (T_0 - T_e) \exp\left[-\frac{\beta}{k}t\right](\cos[\beta^{\frac{1}{2}}x] + \sin[\sin[\beta^{\frac{1}{2}}x]]) + T_e. \quad (16)$$

Obviously, in an approximation in which  $(\cos[\beta^{\frac{1}{2}}x] + \sin[\sin[\beta^{\frac{1}{2}}x]])$  can be approximately treated as a constant, i.e. average value of this expression close to 1, solution (16) can be effectively approximated by the solution of the Newton cooling law (2). Simply speaking, Newton cooling law can be considered as the effective approximation of the heat (transfer) equation for constant space part of the temperature function. But it implies that standard heat (transfer) equation, as well as Newton cooling law, cannot model experimental data of the Mpemba effect.

Consider, now, such situation in which temperature function changes very slowly by space coordinate change so that the following linearization of the temperature second space derivation can be approximately realized

$$\frac{\partial^2 T}{\partial x^2} \simeq \frac{T_0 - T_e}{L} \frac{\partial T}{\partial x}. \quad (17)$$

Such and similar kind of approximation is very often in different important domains of the physics, e.g. quasi-classical approximation of the quantum mechanics, i.e. WBK formalism, solid state physics, quantum field theory, inflationary cosmology etc. .

In such approximation standard heat (transfer) equation turns out in

$$\frac{T_0 - T_e}{L} \frac{\partial T}{\partial x} = k \frac{\partial T}{\partial t} \quad (18)$$

or, after application of (8), in

$$\frac{1}{X} \frac{dX}{dx} = \frac{kL}{T_0 - T_e} \frac{1}{\tau} \frac{d\tau}{dt} = -\beta. \quad (19)$$

It, in fact, represents a system of two usual differential equations with simple solutions

$$\tau = B \exp\left[-\beta \frac{T_0 - T_e}{kL} t\right] \quad (20)$$

$$X = C \exp[-\beta x] \quad (21)$$

where  $B, C$  are some constants.

Then, according to (8), it follows

$$T = BC \exp[-\beta x] \exp[-\beta \frac{T_0 - T_e}{kL} t] + A \quad (22)$$

which, according to initial and "final" conditions (6), (7) that imply

$$A = T_e \quad (23)$$

$$BC = T_0 - T_e \quad (24)$$

, turns out in

$$T = (T_0 - T_e) \exp[-\beta x] \exp[-\beta \frac{T_0 - T_e}{kL} t] + T_e. \quad (25)$$

Obviously here  $T$  decreases exponentially during time with exponent parameter proportional to  $T_0 - T_e$ . It means that, simply speaking, hot liquid can freeze faster than cold liquid in full agreement with experimental data of the Mpemba effect.

Precisely, consider two equivalent liquids in the same vessels so that

$$T_{20} > T_{10} \quad (26)$$

where  $T_{20}$  represents the larger initial temperature of the second liquid while  $T_{10}$  represents the smaller initial temperature of the first liquid. Suppose that the first liquid obtains some temperature  $T$  (25) in a time moment  $t_1$  while the second liquid obtains the same temperature  $T$  in some other time moment  $t_2$ . Then simple calculations, according suppositions and (25), yield

$$t_1 > \frac{T_{20} - T_e}{T_{10} - T_e} t_2 > t_2 \quad (27)$$

corresponding to Mpemba effect experimental data.

In this way it is shown that the suggested approximation of the heat transfer equation can successfully described the Mpemba effect experimental data. Moreover it follows that the Mpemba effect can appear only in situations when given approximation is satisfied, while in opposite situations Mpemba effect does not appear. Finally, according to (25), in situations when Mpemba effect appears this effect becomes sharply expressed when  $L$  is smaller in full agreement with experimental data on the Mpemba effect [1]-[4].

In conclusion it can be repeated and pointed out the following. In this work we suggest a simple theoretical solution of the Mpemba effect in full agreement with known experimental data. This solution follows simply as an especial approximation (linearization) of the usual heat (transfer) equation, precisely linearization of the second derivation of the space part of the temperature function (as it is well-known Newton cooling law can be considered as the effective approximation of the heat (transfer) equation for constant space part of the temperature function).

This work is dedicated to memory on Prof. Dr. Bratislav Tosic. He stated, we paraphrase, approximation is not only ground tool but holly spirit too of any physical theory.

## References

- [1 ] E. B. Mpemba, D. G. Osborne, Phys. Educ. **4** (1969) 172
- [2 ] D. Auerbach, A. J. Phys. **63** (1995) 882
- [3 ] M. Jeng, A. J. Phys. **74** (2006) 514
- [4 ] J. I. Katz, Am. J. Phys. **77** (2009) 27
- [5 ] C. T. O'Sullivan, Am. J. Phys. **58** (1990) 956