ON THE SYMMETRY INTEGRAL

GIOVANNI COPPOLA

to my Angel

ABSTRACT. We give a level one result for the "symmetry integral", say $I_f(N,h)$, of essentially bounded $f: \mathbb{N} \to \mathbb{R}$; i.e., we get a kind of "square-root cancellation" bound for the mean-square (in $N < x \leq 2N$) of the "symmetry sum" of, say, the arithmetic function $f := g * \mathbf{1}$, where $g: \mathbb{N} \to \mathbb{R}$ is such that $\forall \varepsilon > 0$ we have $g(n) \ll_{\varepsilon} n^{\varepsilon}$, and supported in [1, Q], with $Q \ll N$ (so, the exponent of Q relative to N, say the level $\lambda := (\log Q)/(\log N)$ is $\lambda < 1$), where the symmetry sum weights the f-values in (almost all, i.e. all but o(N) possible exceptions) the short intervals [x - h, x + h] (with positive/negative sign at the right/left of x), with mild restrictions on h (say, $h \to \infty$ and $h = o(\sqrt{N})$, as $N \to \infty$).

1. INTRODUCTION AND STATEMENT OF THE RESULTS.

We give upper bounds for the symmetry integral (compare [C-S], [C], [C1], [C2], [C3], [C5], [C6], esp.)

$$I_f(N,h) \stackrel{def}{=} \sum_{x \sim N} \Big| \sum_{|n-x| \leq h}' \operatorname{sgn}(n-x) f(n) \Big|^2,$$

with $x \sim N$ for $N < x \leq 2N$, $\operatorname{sgn}(r) := \frac{r}{|r|} \forall r \in \mathbb{R}, r \neq 0$, $\operatorname{sgn}(0) := 0$ and the dash means : the terms $n = x \pm h$ have to be halved (say, $\sum'_{|n-x| \leq h} \operatorname{sgn}(n-x)f(n) = \sum_{|n-x| \leq h} \operatorname{sgn}(n-x)f(n) - \frac{f(x+h)-f(x-h)}{2}$); this is one of the possible definitions : comparing the one(s) in the papers quoted above, the difference involved will be soon shown negligible, at least for the (large) class of essentially bounded arithmetic functions $f : \mathbb{N} \to \mathbb{R}$. Here we'll use the term "essentially" to leave (negligible) multiplicative factors bounded by arbitrary small powers of N. For example,

f is essentially bounded (abbrev. $f \ll 1$) $\stackrel{def}{\iff} \forall \varepsilon > 0 \ f(n) \ll_{\varepsilon} N^{\varepsilon}$

(here, $\forall n \leq 2N + h$, since we "don't see" f any further), with the usual Vinogradov notation (i.e., \ll_{ε} means "bounded in absolute value with a constant factor depending on ε "); and we'll abbreviate

$$F(N,h) \lll G(N,h) \quad \stackrel{def}{\Longleftrightarrow} \quad \forall \varepsilon > 0 \ |F(N,h)| \ll_{\varepsilon} N^{\varepsilon} G(N,h)$$

For example, to see that our new definition is "close" to the old one (an integral !), whenever $f \ll 1$,

$$\int_{N}^{2N} \Big| \sum_{|n-x| \le h} \operatorname{sgn}(n-x) f(n) \Big|^{2} dx \ll \sum_{N \le x < 2N} \Big| \sum_{|n-[x]| \le h}' \operatorname{sgn}(n-[x]) f(n) \Big|^{2} dx + N \ll I_{f}(N,h) + N + h^{2}$$

where $[x] \stackrel{def}{=}$ integer part of $x \in \mathbb{R}$, both the remainders $\ll N$ and $\ll h^2$ are clearly negligible : below the "diagonal" remainders, i.e. $\ll Nh$ (when $h \to \infty$ and h = o(N), for $N \to \infty$, as we'll assume henceforth).

The symmetry integral (now on, $I_f(N,h)$ defined above) measures, for f, the ALMOST-ALL (i.e., leave eventual o(N) exceptions) symmetry (around x) in the SHORT (since h = o(x)) INTERVALS [x - h, x + h].

For a motivation to study I_f , see esp. [C]. Our present arguments closely resemble [C7] ones.

Our main result is the following.

THEOREM. Let $N, h, Q \in \mathbb{N}$, with $Q \ll N$ and $h \to \infty$, $h = o(\sqrt{N})$ as $N \to \infty$. Assume $g : \mathbb{N} \to \mathbb{R}$ is independent of x, N, h, essentially bounded and supported in [1, Q]; set $f := g * \mathbf{1}$. Then

 $I_f(N,h) \ll Nh.$

Mathematics Subject Classification (2010): 11N37, 11N25.

We then get square-root cancellation for the symmetry integral of d_k (generating Dirichlet series ζ^k) : COROLLARY. Fix $k \ge 1$ integer. Let $N, h \in \mathbb{N}$ with $h = o(\sqrt{N})$. Then (the \ll -const.depends on k)

$$I_{d_k}(N,h) \ll Nh$$

Also, for the von Mangoldt function $\Lambda(n) := \log p \ \forall n = p^r, r \in \mathbb{N}, p \ prime, := 0 \ otherwise,$

$$I_{\Lambda}(N,h) \ll Nh.$$

We'll not prove the Corollary. (A kind of "DIRICHLET HYPERBOLA TRICK" for I_{d_k} gives $Q \ll N^{1-1/k}$, [C6].)

In passing, we note that an estimate of the kind in the Corollary, but for the Selberg integral of d_k would give, applying [C4] bounds, the proof of the Lindelöf Hypothesis ! (Also, we prove, with the Corollary, the almost-all symmetry of primes: for the corresponding Selberg integral, i.e. the classical Selberg integral of primes, this would give the Density Hypothesis !).

The paper is organized as follows:

- \triangle in section §2 we start proving the Theorem, then
- \triangle in next section, §3, we state and prove a small Lemma, in order to complete Theorem's proof;
- \bigtriangleup we conclude with some comments, in the final section.

The author trusts the possibility to treat the classic Selberg integral through the approach outlined in [C].

2. A weak majorant principle.

We start proving the Theorem, giving at next section the necessary Lemma. Here, we closely follow [C7]. PROOF. Assume now on $Q \ll N$, $g : \mathbb{N} \to \mathbb{R}$, with supp $(g) \subset [1, Q]$, $g * \mathbf{1} := f \ll 1$, $h \to \infty$ and h = o(N):

$$I_f(N,h) = \sum_{x \sim N} \Big| \sum_{q \leq Q, q \leq x+h} g(q) \chi_q(x) \Big|^2,$$

where we define, this time (compare [C-S], esp.), $\forall q \in \mathbb{N}$,

$$\chi_q(x) \stackrel{def}{=} - \sum_{\substack{|n-x| \le h \\ n \equiv 0 \pmod{q}}}' \operatorname{sgn}(n-x) = \sum_{\substack{\ell \mid q \\ \ell > 1}} \frac{\ell}{q} \sum_{j \le \frac{\ell}{2}} c_{j,\ell}^{\pm} \sin \frac{2\pi x j}{\ell},$$

and the FOURIER COEFFICIENTS ARE POSITIVE

$$c_{j,q}^{\pm} := \frac{4}{q} \cot \frac{\pi j}{q} \sin^2 \frac{\pi j h}{q} := \frac{1}{q} F_h^{\pm} \left(\frac{j}{q}\right) \ge 0 \quad \forall j \le \frac{q}{2}$$

(better, the finite Fourier expansion has non-negative coefficients); here (see [C-S], apply Parseval identity)

$$\sum_{j \le q}^{*} |c_{j,q}^{\pm}|^{2} \ll \sum_{0 < j \le q} |c_{j,q}^{\pm}|^{2} \ll \left\|\frac{h}{q}\right\| \ll \min\left(1, \frac{h}{q}\right).$$

As usual, $\|\alpha\| := \min_{n \in \mathbb{Z}} |\alpha - n|$ is the distance to the next integer, $\forall \alpha \in \mathbb{R}$.

The ORTHOGONALITY OF THE ADDITIVE CHARACTERS is the starting point (compare [C7]).

Here we'll not define explicitly the RAMANUJAN COEFFICIENTS, see [C7].

This time, not like in [C7], we use as a "majorant", for our f(n), the DIVISOR FUNCTION, d(n). In fact, this one has, at least for $h = o(\sqrt{N})$, see [C-S] (compare [C2]), square-root cancellation on average over $x \sim N$.

Then (see [C7], proof of the Proposition: by the way, misprints occur there, missing $\frac{1}{\ell^2}$ and $\frac{1}{\ell t}$)

$$I_f(N,h) = D_f^{\pm}(N,h) + 2\sum_{1<\ell,t\leq Q}\sum_{d\leq \frac{Q}{t}}\sum_{q\leq \frac{Q}{t}}\frac{g(\ell d)g(tq)}{\ell dtq} \sum_{\substack{j\leq \frac{\ell}{2}\\\delta:=\frac{j}{\ell}-\frac{\tau}{t}>0}}^* F_h^{\pm}\left(\frac{j}{\ell}\right)F_h^{\pm}\left(\frac{r}{t}\right)\sum_{\substack{x\sim N\\\dots}}\sin\frac{2\pi xj}{\ell}\sin\frac{2\pi xr}{t} = 0$$

$$=D_{f}^{\pm}(N,h)+\sum_{1<\ell,t\leq Q}\sum_{d\leq \frac{Q}{\ell}}\sum_{q\leq \frac{Q}{t}}\frac{g(\ell d)g(tq)}{\ell dtq}\sum_{\substack{j\leq \frac{\ell}{2}\\\delta:=\frac{j}{\ell}-\frac{r}{t}>0}}^{*}F_{h}^{\pm}\left(\frac{j}{\ell}\right)F_{h}^{\pm}\left(\frac{r}{t}\right)\left(\sum_{\substack{x\sim N\\\cdots}}\cos 2\pi\delta x-\sum_{\substack{x\sim N\\\cdots}}\cos 2\pi\sigma x\right),$$

where we abbreviate

$$\sum_{\substack{x \sim N \\ \dots}} := \sum_{\substack{x \sim N \\ x \ge \ell d - h \\ x \ge tq - h}},$$

indicating the present *diagonal* as

$$D_f^{\pm}(N,h) \stackrel{def}{=} \sum_{1 < \ell \le Q} \sum_{d_1 \le \frac{Q}{\ell}} \sum_{d_2 \le \frac{Q}{\ell}} \frac{g(\ell d_1)g(\ell d_2)}{\ell^2 d_1 d_2} \sum_{j \le \frac{\ell}{2}} F_h^{\pm} \left(\frac{j}{\ell}\right)^2 \sum_{\substack{x \ge N \\ x \ge \ell d - h \\ x \ge t q - h}} \sin^2 \frac{2\pi x j}{\ell},$$

with $\delta := j/\ell - r/t$ (> 0, see above) and

$$\sigma := \left\| \frac{j}{\ell} + \frac{r}{t} \right\| \in \left[0, \frac{1}{2} \right].$$

This time we appeal to the elementary Lemma of [C7] to treat the terms with well-spaced fractions (i.e., $\delta > 1/A$ or $\sigma > 1/A$, with 1/A = 1/NL, $L := \log N$, say), BUT we have A MINUS SIGN on the terms with $\sigma \leq 1/A$. These are treated exactly, when 1/A = o(1/N), in the next small Lemma.

3. Statement and proof of the lemma to complete the Theorem Proof.

We can state and show our

LEMMA. Let $A, N \in \mathbb{N}$ with $\frac{1}{A} = o(\frac{1}{N})$ when $N \to \infty$. Let $Q \ll N$ and R_{ℓ}, R_t, F_h^{\pm} be as above. Then

$$\delta := \frac{j}{\ell} - \frac{r}{t}, \ \sigma := \left\| \frac{j}{\ell} + \frac{r}{t} \right\| \ \Rightarrow \ \sum_{1 < \ell, t \le Q} R_{\ell}(f) R_{t}(f) \ \sum_{j \le \frac{\ell}{2} \atop \delta > 0, \sigma \le 1/A}^{*} F_{h}^{\pm} \left(\frac{j}{\ell} \right) F_{h}^{\pm} \left(\frac{r}{t} \right) \ \sum_{\substack{x \ge \lambda \\ x \ge tq - h}} \cos 2\pi \sigma x = 0.$$

PROOF. Abbreviating σ as above we have

$$\sigma \leq \frac{1}{A} \Rightarrow 0 < \frac{j}{\ell} + \frac{r}{t} \leq \frac{1}{A} \text{ or } 0 \leq 1 - \frac{j}{\ell} - \frac{r}{t} \leq \frac{1}{A};$$

first case gives an absurd: in particular $0 < j = o\left(\frac{\ell}{N}\right) = o\left(\frac{Q}{N}\right) = o(1), 0 < r = o\left(\frac{t}{N}\right) = o\left(\frac{Q}{N}\right) = o(1).$ Hence

$$0 \le \left(\frac{1}{2} - \frac{j}{\ell}\right) + \left(\frac{1}{2} - \frac{r}{t}\right) \le \frac{1}{A} \implies 0 \le \frac{1}{2} - \frac{j}{\ell} \le \frac{1}{A}, 0 \le \frac{1}{2} - \frac{r}{t} \le \frac{1}{A},$$

whence (use 1/A = o(1/N), here)

$$\frac{\ell}{2} - \frac{\ell}{A} \le j \le \frac{\ell}{2}, \quad \frac{t}{2} - \frac{t}{A} \le r \le \frac{t}{2}, \quad \Rightarrow \quad j = \left[\frac{\ell}{2}\right], r = \left[\frac{t}{2}\right]$$

which, together with previous ranges for $1 - j/\ell - r/t$, give, again from $\frac{1}{A} = o(\frac{1}{N})$,

$$0 \le \frac{1}{\ell} \left\{ \frac{\ell}{2} \right\} + \frac{1}{t} \left\{ \frac{t}{2} \right\} \le \frac{1}{A} \quad \Rightarrow \quad 2|\ell, 2|t.$$

Thus,

$$\frac{j}{\ell} = \frac{r}{t} = \frac{1}{2}$$

and the thesis, using $F_h^{\pm}\left(\frac{1}{2}\right) = 0$.

We may complete Theorem's proof. In fact, applying [C7] well-spaced Lemma (the conditions in ... give a small disturbance, since the same considerations of [C7] Proposition proof apply verbatim) with the same choice $A = NL = N \log N$, say, from present Lemma (which, together with the [C7] one, says terms with the x-sum of $\cos 2\pi\sigma x$ are negligible !)

$$I_f(N,h) \ll Nh +$$

$$+D_{d}^{\pm}(N,h) + \sum_{1 < \ell, t \le Q} \sum_{d \le \frac{2N+h}{\ell}} \sum_{q \le \frac{2N+h}{t}} \frac{1}{\ell dt q} \sum_{\substack{j \le \frac{\ell}{2} \\ \delta := \frac{j}{\ell} - \frac{r}{t} > 0}} F_{h}^{\pm} \left(\frac{j}{\ell}\right) F_{h}^{\pm} \left(\frac{r}{t}\right) \left(\sum_{\substack{x \sim N \\ \cdots}} \cos 2\pi \delta x - \sum_{\substack{x \sim N \\ \cdots}} \cos 2\pi \sigma x\right) \ll I_{d}(N,h) + Nh,$$

whence the thesis, since the symmetry integral of the function d(n), from [C-S] (requiring $h = o(\sqrt{N})$, here), is

 $\ll Nh.$ \Box

4. Some comments.

The method outlined in [C7] and here treats more in general second moments over "long intervals" of "short interval" sums (or averages). Both these approaches use the non-negativity of the Fourier coefficients (say, \tilde{F}_h there and F_h^{\pm} here) arising from the short interval inner average. The Selberg integral is another planet. I mean, no more free positivity there ! (No free meal...) However, there's still some hope ! First of all, the present approach seems weaker than [C7] one, whose proofs are easier; but not a problem the estimate in our present Lemma. The real complication comes when facing changing signs of Fourier coefficients for the Selberg integral. They require another approach and, also, will set some limits on the ranges of the divisors, say Q. Here we don't have any troubles, since we have a REAL (though approximate !) majorant principle, i.e. we use brute force and "bound any essentially bounded function with the divisor function". This is possible, since we KNOW the symmetry integral of d(n), not only, but it's also fantastically small (it's a square-root saving ! However, see my paper on Parma University Journal about square-free numbers: they've even better cancellation on their symmetry integral !).

The problem we have to face for both the original and the modified Selberg integral is the (already mentioned, see [C7] final comments) problem of the Wintner majorants that even here is represented (but immediately solved).

The majorating procedure is optimal (as we see now) in the case of the symmetry integral. Then, if you have a way to get non-trivial bounds from this (as Kaczorowski and Perelli do, as stated in [C]) for the Selberg integral, better to avoid a direct approach to $J_f(N, h)$.

However, the modified Selberg integral has its own interest; and the further difficulty to build good Wintner majorants to get non-trivial bounds is happily faced, in view of the non-trivial results obtained !

The same is of course true whenever a kind of majorant principle will be available for the Selberg integral, but (due to Fourier coefficients non-constant signs) with limitations, due to the technical analytic treatment.

I expect (and have already some scratch calculations) the analytic part to be fair (not impossible, not easy); then, the "*heavy*" part is building (non-trivial!) majorants. And next ... estimate Selberg's integral !

References

- [C] Coppola, G.- On the symmetry of divisor sums functions in almost all short intervals Integers 4 (2004), A2, 9 pp. (electronic). <u>MR 2005b:11153</u>
- [C1] Coppola, G.- On the Correlations, Selberg integral and symmetry of sieve functions in short intervals

 http://arxiv.org/abs/0709.3648v3 (to appear on: Journal of Combinatorics and Number Theory)
- [C2] Coppola, G.- On the Correlations, Selberg integral and symmetry of sieve functions in short intervals, II - Int. J. Pure Appl. Math. 58.3(2010), 281–298.
- [C3] Coppola, G.- On the Correlations, Selberg integral and symmetry of sieve functions in short intervals, III - http://arxiv.org/abs/1003.0302v1
- [C4] Coppola, G.- On the Selberg integral of the k-divisor function and the 2k-th moment of the Riemann zeta-function - http://arxiv.org/abs/0907.5561v1 - to appear on Publ. Inst. Math., Nouv. Sr.
- [C5] Coppola, G.- On the symmetry of arithmetical functions in almost all short intervals, V (electronic) http://arxiv.org/abs/0901.4738v2
- [C6] Coppola, G.- On some lower bounds of some symmetry integrals http://arxiv.org/abs/1003.4553v2
- [C7] Coppola, G.- On the modified Selberg integral http://arxiv.org/abs/1006.1229v1
- [C-S] Coppola, G. and Salerno, S.- On the symmetry of the divisor function in almost all short intervals -Acta Arith. 113 (2004), no.2, 189–201. <u>MR 2005a:11144</u>
 - [D] Davenport, H. Multiplicative Number Theory Third Edition, GTM 74, Springer, New York, 2000. <u>MR 2001f:11001</u>
 - [L] Linnik, Ju.V.- The Dispersion Method in Binary Additive Problems Translated by S. Schuur -American Mathematical Society, Providence, R.I. 1963. MR 29#5804
 - [T] Tenenbaum, G.- Introduction to Analytic and Probabilistic Number Theory Cambridge Studies in Advanced Mathematics, 46, Cambridge University Press, 1995. <u>MR 97e:11005b</u>
 - [V] Vinogradov, I.M.- The Method of Trigonometrical Sums in the Theory of Numbers Interscience Publishers LTD, London, 1954. MR 15,941b

Dr.Giovanni Coppola

DIIMA - Università degli Studi di Salerno

84084 Fisciano (SA) - ITALY

e-mail : gcoppola@diima.unisa.it

e-page : www.giovannicoppola.name