Introducing the Fission-Fusion Reaction Process: Using a Laser-Accelerated Th Beam to produce Neutron-Rich Nuclei towards the N=126 Waiting Point of the r-Process

D. Habs^{1,2}, P.G. Thirolf¹, M. Gross¹, K. Allinger¹, J. Bin¹, A. Henig¹, D. Kiefer¹, W. Ma², J. Schreiber²

¹ Fakultät für Physik, Ludwig-Maximilians Universität München, D-85748 Garching, Germany

Received: date / Revised version: date

Abstract We propose to produce neutron-rich nuclei in the range of the astrophysical r-process (the rapid neutron-capture process) around the waiting point N =126 [1,2,3] by fissioning a dense laser-accelerated thorium ion bunch in a thorium target (covered by a polyethylene layer, CH₂), where the light fission fragments of the beam fuse with the light fission fragments of the target. Via the 'hole-boring' (HB) mode of laser Radiation Pressure Acceleration (RPA) [4,5,6] using a high-intensity, short pulse laser, very efficiently bunches of ²³²Th with solid-state density can be generated from a Th layer (ca. 560 nm thick), placed beneath a deuterated polyethylene foil (CD₂ with ca. 520 nm), both forming the production target. Th ions laser-accelerated to about 7 MeV/u will pass through a thin CH₂ layer placed in front of a thicker second Th foil (both forming the reaction target) closely behind the production target and disintegrate into light and heavy fission fragments. In addition, light ions (d,C) from the CD₂ production target will be accelerated as well to about 7 MeV/u, inducing the fission process of ²³²Th also in the second Th layer. The laser-accelerated ion bunches with solid-state density, which are about 10¹⁴ times more dense than classically accelerated ion bunches, allow for a high probability that generated fission products can fuse again when the fragments from the thorium beam strike the Th layer of the reaction target.

In contrast to classical radioactive beam facilities, where intense but low-density radioactive beams of one ion species are merged with stable targets, the novel fission-fusion process draws on the fusion between neutron-rich, short-lived, light fission fragments both from beam and target. Moreover, the high ion beam density may lead to a strong collective modification of the stopping power in the target by 'snowplough-like' removal of target electrons, leading to significant range enhancement, thus allowing to use rather thick targets.

Using a high-intensity laser with two beams with a total energy of 300 J, 32 fs pulse length and 3 μ m focal diameter, as, e.g., envisaged for the ELI-Nuclear Physics project in Bucharest (ELI-NP) [7], order-of-magnitude estimates promise a fusion yield of about 10^3 ions per laser pulse in the mass range of A=180-190, thus enabling to approach the r-process waiting point at N=126. First studies on ion acceleration, collective modifications of the stopping behaviour and the production of neutron-rich nuclei can also be performed at the upcoming new laser facility CALA (Center for Advanced Laser Applications) in Garching.

1 Introduction

Elements like platinum, gold, thorium and uranium are produced via the rapid neutron capture process (r-process) at astrophysical sites like merging neutron star binaries or (core collapse) supernova type II explosions with outbursts of very high neutron density in the range of $10^{21}-10^{30}/cm^3$. We aim at improving our understanding of these nuclear processes by measuring the properties of heavy nuclei on (or near) the r-process path. According to a recent report by the US National Research Council of the National Academy of Science, the origin of the heaviest elements remains one of the 11 greatest unanswered questions of modern physics [8]. While the lower path of the r-process for the production of heavy elements is well explored, the nuclei around the N=126waiting point critically determine this element production mechanism. At present basically nothing is known about these nuclei.

Special ingredients of this proposal are: i) The very efficient Radiation Pressure Acceleration (RPA) mechanism for laser-based ion acceleration, especially exploiting the 'hole-boring' mode [4] producing pancake-like

 $^{^{2}}$ Max-Planck-Institut für Quantenoptik, D-85748 Garching, Germany




Fig. 1 Chart of the nuclides indicating various pathways for astrophysical nucleosynthesis: thermonuclear fusion reactions in stars (orange vector), s-process path (red vector) and the r-process generating heavy nuclei in the Universe (red pathway). The nuclei marked in black indicate stable nuclei. For the green nuclei some nuclear properties are known, while the yellow, yet unexplored regions extend to the neutron and proton drip lines. The blue line connects nuclei with the same neutron/proton ratio as for (almost) stable actinide nuclei. On this line the maximum yield of nuclei produced via fission-fusion (without neutron evaporation) will be located. The elliptical conture lines correspond to the expected maximum fission-fusion cross sections decreased to 50%, 10% and 0.1%, respectively, for primary 232 Th beams.

beam bunches of solid-state density. ii) The strongly reduced stopping power of these dense bunches in a second thick Th target, where the decomposition into fission fragments and the fusion of these fragments takes place. After the laser flash we want to extract rather long-lived isotopes (> 100 ms) in flight, separate them e.g. in a (gas-filled) recoil separator and study them via decay spectroscopy or lifetime and nuclear mass measurements.

In the following we outline the relevance of the project for nuclear astrophysics, describe the new laser acceleration scheme and in particular the new fission-fusion reaction method. Finally the planned ELI-Nuclear Physics facility will be briefly introduced, where the production of these nuclei and the experiments to measure their properties will be realized.

2 The Relevance of the N=126 Waiting Point for Nuclear Astrophysics

Fig. 1 shows the nuclidic chart marked with different nucleosynthesis pathways for the production of heavy elements in the Universe: the thermonuclear fusion processes in stars producing elements up to iron (orange arrow), the slow neutron capture process (s-process) along the valley of stability leading to about half of the heavier nuclei (red arrow) and the rapid neutron capture process (r-process) proceeding along pathways with neutron separation energies S_n in the range of 2–3 MeV. In this scenario, rather neutron-rich nuclei are populated in an intense neutron flux [9]. The r-process path ex-

hibits characteristic vertical regions for constant magic neutron numbers of 50, 82 and 126, where the r-process is slowed down due to low neutron capture cross sections when going beyond the magic neutron numbers. These decisive bottlenecks of the r-process flow are called waiting points [10].

The astrophysical site of the r-process nucleosynthesis is still under debate: it may be cataclysmic core collapse supernovae (II) explosions with neutrino winds [2, 3,11,12 or mergers of neutron-star binaries [13,14,15]. The r-process element abundances from galactic halo stars tell us that the r-process site for lighter and heavier neutron capture processes may occur under different astrophysical conditions [10]. For the heavier elements beyond barium, the isotopic abundancies are always very similar (called universality) and the process seems to be very robust. Perhaps also the recycling of fission fragments from the end of the r-process strengthens this stability. Presently, it seems more likely that a merger of neutron star binaries is the source for the heavier rprocess branch, while core collapsing supernova explosions contribute to the lighter elements below barium. The modern nuclear equations of state, neutrino interactions and recent supernova explosion simulations [3] lead to detailed discussions of the waiting point N=126. Here measured nuclear properties along the N=126 waiting point may help to clarify the sites of the r-process.

Fig. 2 shows the measured solar elemental abundances of the r-process nuclei together with a calculation, where masses from the Extended Thomas-Fermi plus Strutinski Integral (ETFSI) mass model [16] have been used to-




Fig. 2 Observed elemental solar abundances in the r-process mass range (black symbols with error bars) in comparison with calculated abundances (red line and symbols), normalized to silicon = 10^6 . The theoretical predictions show the elemental abundances for stable isotopes after α and β decay as obtained in the ETFSI-Q mass model [1,16] for a wide range of neutron densities n_n (in $1/cm^3$) and temperatures T_9 (in units of 10^9 K) and including shell quenching effects. Included with permission from [17].

gether with several neutron flux components, characterized by a temperature T_9 , neutron densities n_n and expansion time scales. A quenching of shell effects [18] was assumed in the nuclear mass calculations to achieve a better agreement between observed and calculated abundances.

The three pronounced peaks visible in the abundance distribution seem to be of different origin, which is also reflected in the theoretical calculations shown in Fig. 2. where contributions from different temperatures and neutron densities are superimposed to the observed data. We note the pronounced third peak in the abundance distribution around A = 180 - 200, corresponding to the group of elements around gold, platinum and osmium, where until now no experimental nuclear properties have been measured for r-process nuclei. Several astrophysical scenarios try to explain this third abundance peak. A detailed knowledge of nuclear lifetimes and binding energies in the region of the N=126 waiting point will narrow down the possible astrophysical sites. If, e.g., no shell quenching could be found in this mass range, the large dip existing for this case in front of the third abundance peak would have to be filled up by other processes like neutrino wind interactions. For cold decompressed neutron-rich matter, e.g., from neutron-star mergers, we find an equilibrium between (n, γ) and β decay. For the rather hot supernova-explosion scenario we find an equilibrium between (n,γ) and (γ,n) reactions. Considering the still rather large difficulties to identify convincing astrophysical sites for the third peak of the r-process with sufficiently occurrence rates, measurements of the nuclear properties around the N=126 waiting point will represent an important step forward in solving the difficult and yet confusing site selection of the third abundance peak of the r-process.

The key bottleneck nuclei of the N=126 waiting point around $Z \approx 70$ are about 15 neutrons away from presently known nuclei (see Fig. 1), with a typical drop of the production cross section for classical radioactive beam production schemes of about a factor of 10-20 for each additional neutron towards more neutron-rich isotopes. Thus presently nothing is known about these nuclei and even next-generation large-scale 'conventional' radioactive beam facilities like FAIR [19], SPIRAL II [20] or FRIB [21] will not be able to grant experimental access to the most important isotopes on the r-process path. The third peak in the abundance curve of r-process nuclei is due to the N=126 waiting point as visible in Fig. 1. These nuclei are expected to have rather long halflives of a few 100 ms. This waiting point represents the bottleneck for the nucleosynthesis of heavy elements up to the actinides. From the view point of astrophysics, it is the last region, where the r-process path gets close to the valley of stability and thus can be studied with the new isotopic production scheme discussed below. While the waiting point nuclei at N = 50 and N = 82 have been studied rather extensively [22, 23, 24, 25], nothing is known experimentally about the nuclear properties of waiting point nuclei at the N=126 magic number. Nuclear properties to be studied here are nuclear masses, lifetimes, β -delayed neutron emission probabilities $P_{\nu,n}$ and the underlying nuclear structure.

For the overall description of the r-process, the nuclear masses are typically taken from mass models like the macroscopic-microscopic Finite Range Droplet Model (FRDM). Alternatively, models more closely related to first principles like Hartree-Fock-Bogoliubov calculations are used [2,10]. Typically, somewhat less shell quenching is assumed for the heavier N=126 region compared to the N=82 region of the r-process.

Fig. 3 displays the difference between nuclear masses for the isotopic chain Z=55 (Cs) as calculated by various mass models and measured masses (data taken from AME95 [26]). The plot demonstrates the good agreement between measured and predicted masses in the mass range where experimental data are available, while drastic deviations occur outside these regions especially for the r-process region [25]. For the Cs isotopes with Z=55 as shown in Fig. 3 the mass measurements reach close to the r-process path. However, for the heavier elements relevant for the r-process waiting point N=126 around Z \approx 70, the known isotopes are about 15 neutrons away from the magic neutron number N=126 and Fig. 3 impressively illustrates the extremely large uncer-




Fig. 3 Differences between nuclear mass predictions from various theoretical mass models for Cs isotopes (Z=55) compared to measured masses taken from AME95 [26] as a function of the neutron number N (for Z=55). The figure is taken with permission from Ref. [25].

tainties presently expected for any theoretical prediction of nuclear masses, Q values, or β halflives which may partly be attributed to effects of nuclear deformation. This clearly points to the importance of direct measurements in this mass region, especially targeting nuclear masses.

Moreover, presently there exist difficulties to describe consistently the third abundance peak of the r-process, the 232 Th and 238 U cosmochronometers and the potential 'fission cycle' beyond A>260 [27].

From the viewpoint of nuclear structure theory, semimagic, heavy nuclei not too far away from stability are spherical and thus can be treated in general more successfully than heavy deformed nuclei. Shell model calculations with open proton- and neutron shells require extremely large dimensions of the configuration space. With realistic density functional theories, some extremely time consuming deformed RPA calculations with Skyrme or Gogny forces have been performed [28], but still spherical nuclei can be controlled much better. Also the subsystem of neutrons or protons allows to study systems with large isospin. These density functional calculations have to be fitted to experimental data of heavy nuclei, in order to enable reliable predictions for other nuclei [28]. If we improve our experimental understanding of this final bottleneck to the actinides at N=126, many new visions open up: (i) For many mass formulas (e.g. [29]), there is a branch of the r-process leading to extremely long-lived superheavy elements beyond Z=110 with lifetimes of about 10^9 years. If these predictions could be made more accurate, a search for these superheavy elements in nature would become more promising. (ii) At present the prediction for the formation of uranium and

thorium elements in the r-process is rather difficult, because there are no nearby magic numbers and those nuclei are formed during a fast passage of the nuclidic area between shells. Such predictions could be improved, if the bottleneck of actinide formation would be more reliably known. (iii) Also the question could be clarified if fission fragments are recycled in many r-process loops or if only a small fraction is reprocessed.

This description of our present understanding of the r-process underlines the importance of the present project for nuclear physics and, particularly, for astrophysics.

3 The Fission-Fusion Reaction Process

In the following section the various ingredients enabling the new fission-fusion reaction scenario are outlined. First the 'Radiation Pressure Acceleration' method of laser ion acceleration is described with special emphasis on the 'hole-boring' mode, which allows to generate ultradense ion beams. Consequently, collective effects are expected for the interaction of these ion beams with solid targets, leading to a significant reduction of the conventional electronic stopping power. Finally, the fission-fusion reaction process based on these ultra-dense laser-accelerated ion beams is described and an order-of-magnitude estimate for the achievable fusion yield will be presented.

3.1 Laser Ion Acceleration

Laser-accelerated energetic ion beams have been produced during the last few years from μm thick metallic

foils when irradiated by ultra-intense, short laser pulses [30,31,32]. In these experiments the high-energy electrons produced at the front of the target penetrated the target being opaque to the laser. At the rear side the electrons generate an electrostatic field, which ionizes and accelerates ions from the rear side. This acceleration mechanism was called 'Target Normal Sheath Acceleration (TNSA)'. It was explored in many experiments at various high-intensity laser laboratories [33,34,35,36]. A recent review [37] shows that the ion energy scales proportional to the square root of the laser intensity. Typical conversion efficiencies from laser energy to ion energy amount to less than 1%.

In the proposal of a new nuclear reaction scenario introduced in this work, we envisage to exploit instead the new Radiation Pressure Acceleration (RPA) mechanism for ion acceleration. It was first proposed theoretically [4,38,39,40,41]. Special emphasis has been given to RPA with circularly polarized laser pulses as this suppresses fast electron generation and leads to the interaction dominated by the radiation pressure [4,38]. It has been shown that RPA operates in two modes. In the first one, called 'hole-boring', the laser pulses interact with targets thick enough to allow to drive target material ahead of it as a piston, but without interacting with the target rear surface [4].

An alternative scenario, called 'light-sail' (LS) mode of RPA, occurs if the target is sufficiently thin for the laser pulse to punch through the target and accelerate part of the plasma as a single object [39,40]. Typically the 'hole-boring' mode leads to lower velocities of the accelerated ions, as envisaged for the present proposal. The first experimental observation of RPA in the 'hole-boring' regime was achieved only recently in experiments led by the Munich group [5,6].

The RPA laser ion acceleration mechanism in general provides a much larger efficiency for the conversion from laser energy to ion energy and allows for a generation of much larger ion energies in comparison to TNSA. Moreover, for circularly polarized laser light RPA holds promise of quasi-monoenergetic ion beams. Due to the circular polarization, electron heating is strongly suppressed. The electrons are compressed to a dense electron sheet in front of the laser pulse, which then via the Coulomb field accelerates the ions. This mechanism requires much thinner targets and ultra-high contrast laser pulses to avoid the pre-heating and expansion of the target before the interaction with the main laser pulse.

The RPA mechanism allows to produce ion bunches with solid-state density $(10^{22} - 10^{23}/\text{cm}^3)$, which thus are $\approx 10^{14}$ times more dense than ion bunches from classical accelerators. Correspondingly, the areal densities of these bunches are $\approx 10^7$ times larger. It is important to note that these ion bunches are accelerated as neutral ensembles together with the accompanying electrons and thus do not Coulomb explode.

For an estimate of the required laser intensities, focal spot area and target thickness, the 1-D RPA model as outlined in [4] is sufficient. It holds true for the relativistic 'hole-boring' regime of RPA. For the achievable ion energy E_i it yields the expression (circular polarized light)

$$E_i = E_u \cdot A = 2m_i c^2 \Xi / \left(1 + 2\sqrt{\Xi}\right), \tag{1}$$

where E_u is the energy per nucleon, A is the atomic mass number, m_i is the ion mass, c is the vacuum speed of light, and Ξ is the dimensionless pistoning parameter given by

$$\Xi = I_L/(m_i n_i c^3). \tag{2}$$

 I_L denotes the laser intensity and n_i the ion density. In the non-relativistic limit $\Xi << 1$, Eq. (1) reduces to $E_i = 2m_ic^2\Xi$, which together with Eq. (2) is equivalent to Macchi's Eq. (1) in [38]. The conversion efficiency of laser energy to ion energy, χ , follows from [4]

$$\chi = 2\sqrt{\Xi}/\left(1 + 2\sqrt{\Xi}\right). \tag{3}$$

The total number of ions, N_i , that can be accelerated results from the energy balance

$$N_i E_i = \chi W_L, \tag{4}$$

where W_L denotes the energy of the laser pulse.

The target arrangement we want to use is depicted in Fig. 4. It actually consists of two targets termed production target and reaction target. The first is composed of two spatially separated foils, one made from thorium and the other from deuterated polyethylene, CD_2 . They serve for the generation of a thorium ion beam and a beam containing carbon ions and deuterons. The reaction target has a sandwich structure. The first layer is made from CH₂ and causes fission of the accelerated thorium nuclei. The second layer is a pure thorium film. The accelerated carbon ions and deuterons lead to fission of these thorium nuclei. Fusion of the fragments created in both layers generates neutron-rich nuclei in a mass range towards the waiting point N=126. This reaction scheme works best when the thorium and carbon ions and the deuterons have each the energy of 7 MeV per nucleon (for details see Sect. 3.2 and the following sections). Accelerating 232 Th ions whose density $\rho_{\mathrm{Th}} = m_{\mathrm{Th}} n_{\mathrm{Th}}$ amounts to 11.7 g/cm³ to E_u =7 MeV per nucleon with laser light of 0.8 μ m wavelength needs, according to equations (1) and (2), an intensity of $1.2 \cdot 10^{23}$ W/cm².

$$a_L = \sqrt{\frac{f \cdot I_L[Wcm^{-2}] \cdot \lambda_L^2[\mu m^2]}{1.37 \cdot 10^{18}}}$$
 (5)

with f=1 for linear and f=1/2 for circular polarized light, respectively. Eq. (5) gives the value of 167 for $1.2 \cdot 10^{23}$ W/cm² and $\lambda_L=0.8 \mu m$, at circularly polarized light. The conversion efficiency, χ , reaches 11%

The dimensionless vector potential, a_L , follows from

 $(\Xi=3.8\cdot 10^{-3})$. Intensities of this level will be achievable with the APOLLON facility, which is under development at the ENSTA/Ecole Polytechnique in Palaiseau within the ILE project [49] and will form the backbone of the ELI Nuclear Physics project. The APOLLON single-beam pulses will provide $W_L=150~{\rm J}$ in $t_L=32~{\rm fs}$, corresponding to 4.7 PW. The sum of two of these beams is assumed to be available for the present estimate. Because of $W_L=I_L\cdot A_F\cdot t_L$ these values fix the focal spot area on the thorium production target, A_F , to 7.1 $\mu{\rm m}^2$ (3 $\mu{\rm m}$ diameter) and, from Eq. (4) the number of accelerated thorium ions, N_i , to $1.2\cdot 10^{11}$. The thickness of the thorium foil, $d_{\rm Th}$, follows from $N_i=A_F d_{\rm Th} n_{\rm Th}$ and amounts to 560 nm $(n_{\rm Th}=3\cdot 10^{22}/cm^3)$.

The data for the CD₂ case is obtained similarly. As shown in [42], the carbon ions and deuterons will experience the same energy per nucleon. The pistoning parameter and the conversion efficiency have hence the same values as before, $\Xi=3.8\cdot 10^{-3}$ and $\chi=0.11$. Eq. (2) then yields $1.0\cdot 10^{22}$ W/cm² ($a_L=48$) for the focal intensity, I_L , whereby for the polyethylene density, $\rho_{\rm PE}=m_cn_c+m_dn_d$, the value of 1 g/cm³ is taken. Assuming here again a focal spot diameter of of 3 μ m ($A_F=7.1$ μ m²), the required laser energy, $W_L=I_LA_Ft_L$, results in 23 J. The number of accelerated carbon ions and deuterons amounts to $1.4\cdot 10^{11}$ and $2.8\cdot 10^{11}$, respectively. The thickness of the polyethylene foil, $d_{\rm PE}$, is 520 nm.

Phase-stable acceleration [45,46] would yield monochromatic ion energy spectra. Whether this can be really achieved, in particular when several ion components with different charge-to-mass ratios are present, is hardly predictable on the basis of current experimental and theoretical knowledge on ion acceleration. So for a safe evaluation of the fusion process of the thorium fragments, the ion spectra are assumed to be broad (see Sect. 3.3). Predictions related to the important question of the beam stability based on 2-D or 3-D simulations show that plane foils heavily expand and break up due to the Rayleigh-Taylor instability [43]. Promising counter-measures include targets adequately modulated in density and shape [44,45].

3.2 Stopping Power for Dense Ion Bunches in a Solid Target

In nuclear physics the Bethe-Bloch formula [50] is used to calculate the atomic stopping of energetic individual ions

$$-\frac{dE}{dx} = 4\pi n_e \frac{Z_{eff}^2 e^4}{m_e v^2} \left[ln \left\{ \frac{2m_e v^2}{I_p (1 - (v/c)^2)} \right\} - \left(\frac{v}{c}\right)^2 \right],$$
(6)

where I_p denotes the ionization potential, n_e the electron density, m_e the mass of the electron, while v is the ion velocity and Z_{eff} is the effective charge of the ions.

For laser-accelerated ions the ion bunch densities reach solid-state density, which is about 14 orders of magnitude larger compared to beams from classical accelerators. In such a scenario collective effects become important. According to Ref. [51], the Bethe-Bloch equation can be decomposed into a first part describing binary collisions and a second term describing long-range collective contributions according to

$$-dE/dx = 4\pi n_e \frac{Z_{eff}^2 e^4}{m_e v^2} [ln(m_e v^2/e^2 k_D) + ln(k_D v/\omega_p)].$$
(7)

Here k_D is the Debye wave number and ω_p is the plasma frequency of the electrons. In Ref. [52] the mechanism of collective deceleration of a dense particle bunch in a thin plasma is discussed, where the particle bunch fits into a half of one plasma oscillation and is decelerated $10^5 - 10^6$ times stronger than predicted by the classical Bethe-Bloch equation [50] due to a strong collective wakefield. Now we discuss the opposite effect with a strongly reduced atomic stopping power that occurs when sending the energetic, solid-state density ion bunch into a solid carbon or thorium target. For this target the plasma wavelength ($\lambda_p \approx 5$ nm, driven by the ion bunch with a phase velocity corresponding to the thorium ion velocity) is much smaller than the ion bunch length ($\approx 560 \text{ nm}$) and collective acceleration and deceleration effects cancel. Only the binary collisions remain and contribute to the stopping power. Hence, we may consider the dense ion bunch as consisting of about 1750 atomic layers with a distance between the Th ions of about 3.2Å as obtained from the bulk density of metallic thorium (11.7 g/cm³). In this case the first layers of the ion bunch will attract the electrons from the target and like a snow plough will take up the decelerating electron momenta. Hence the predominant part of the ion bunch is screened from electrons and we expect a drastic reduction of the stopping power. The electron density n_e will be strongly reduced in the channel defined by the laser-accelerated ions, because many electrons are expelled by the ion bunch and the laser pulse. This effect requires detailed experimental investigations planned for the near future, aiming at verifying the perspective to use a significantly thicker reaction target. The classical ion range for e.g. 7 MeV/u thorium ions in carbon is 15 mg/cm^2 , corresponding to a range of 66 μ m, while this range amounts to only 40 μ m in a thorium target. However, if we aim at limiting the usable effective range to a thorium target thickness where the remaining projectile energy is still sufficient to induce fission, using the accelerated thorium ions directly to induce fission in the Th target would result in a usable target range of less than 10 μ m without invoking collective effects. However, the use of proton induced fission leads to a usable target thickness of about 50 μ m.

The expected reduced atomic stopping power will be supported by the strong laser heating of the electrons.

A reduction of the atomic stopping is essential to avoid a strong slowing down of the ions below the Coulomb barrier energies, where nuclear reactions are no longer possible. However, even without this reduced stopping power the basic properties of the novel reaction mechanism could still be studied, but with significantly reduced vields.

Taking collective effects into account by assuming a range enhancement by a factor of 100, we expect a usable thickness of several mm for a thorium target.

An optimized ion acceleration scheme will depend on measured stopping powers of the dense bunches in targets of different materials and thicknesses, including the ion beam energy as a further parameter to be optimized in preparatory studies.

3.3 The Fission-Fusion Process

The basic concept of the fission-fusion reaction scenario draws on the ultra-high density of laser-accelerated ion bunches. Choosing fissile isotopes as target material for a first target foil accelerated by an intense laser pulse will enable the interaction of a dense beam of fission fragments with a second target foil also consisting of fissile isotopes. So finally in a second step of the reaction process, fusion between (neutron-rich) beam-like and target-like fission products will become possible, generating extremely neutron-rich ion species.

For our discussion we choose 232 Th (the only component of chemically pure Th) as fissile target material, primarily because of its long halflife of $1.4\cdot10^{10}$ years, which avoids extensive radioprotection precautions during handling and operation. Moreover, metallic thorium targets are rather stable in a typical laser vacuum of 10^{-6} mbar, whereas e.g. metallic 238 U targets would quickly oxydize.

Nevertheless, in a later stage it may become advantageous to use also heavier actinide species in order to allow for the production of even more exotic fusion products.

In general, the fission process of the two heavy Th nuclei from beam and target will be preceded by the deep inelastic transfer of neutrons between the inducing and the fissioning nuclei. Here the magic neutron number in the superdeformed fissile nucleus with N=146 [53,54] may drive the process towards more neutron-rich fissioning nuclei, because the second potential minimum acts like a doorway state towards fission. Since in the subsequent fission process the heavy fission fragments keep their A and N values [55], these additional neutrons will show up in the light fission fragments and assist to reach more neutron-rich nuclei. This process will be of particular importance in the reaction scenario discussed in Sect. 3.3.2 for the case of collectively reduced stopping in the reaction target.

Fig. 4 shows a sketch of the proposed fission-fusion reaction scenario for two different situations, a) for the

case of normal electronic stopping as described by the Bethe-Bloch equation and b) for the case of reduced stopping due to collective effects in the target induced by the ultra-dense ion beam discussed earlier. The latter scenario will be discussed later.

As mentioned before, the accelerated thorium ions are fissioned in the CH_2 layer of the reaction target, whereas the carbon ions and deuterons generate thorium fragments in the thick thorium layer of the reaction target. This scenario is more efficient than the one where fission would be induced by the thorium ions only.

For practical reasons we propose to place the reaction target about 1 mm behind the production target, as indicated in Fig. 4.

3.3.1 Induced fission with normal electronic stopping In the scenario, where the earlier discussed collective effects in the target are not taken into account (marked with 'a)' in Fig. 4), the thorium layer of the reaction target would have a thickness of about 50 μ m.

Using a distance of 2.8 Å between atoms in solid layers of CH_2 , the accelerated light ion bunch $(1.4 \cdot 10^{11} \text{ ions})$ corresponds to 1860 atomic layers in case of a 520 nm thick CD₂ target. In order to allow for an optimized fission of the accelerated Th beam, the thicker Th layer of the reaction target, which is positioned behind the production target, is covered by about 70 μ m of polyethylene. This layer serves a twofold purpose: Primarily it is used to induce fission of the impinging Th ion beam, generating the beam-like fission fragments. Here polyethylene is advantageous compared to a pure carbon layer because of the increased number of atoms able to induce fission on the impinging Th ions. In addition, the thickness of this CH₂ layer has been chosen such that the produced fission fragments will be decelerated to a kinetic energy which is suitable for optimized fusion with the target-like fission fragments generated by the light accelerated ions in the Th layer of the reaction target, minimizing the amount of evaporated neutrons. After each laser shot, a new double-target has to be rotated into position.

In order to estimate the fission cross sections both of beam and target nuclei, we apply geometrical considerations based on the involved nuclear radii, which can be expressed for mass number A in the usual way as

$$R = 1.2 \cdot (A)^{1/3} fm. (8)$$

Neglecting the influence of surface diffuseness effects, the resulting fission cross section of the $^{232}\mathrm{Th}$ beam in the CH₂ layer of the reaction target amounts to $\sigma_{\mathrm{fis}}=\pi(R_1+R_2)^2=350~fm^2=3.5\cdot 10^{-28}~m^2~(3.5~\mathrm{b}).$ Correspondingly, the deuteron-induced fission in the Th reaction target occurs with a cross section of about 247 fm²= $2.47\cdot 10^{-28}\mathrm{m}^2~(2.47~\mathrm{b}).$ If we use the atomic distance of




Fig. 4 Sketch of the target arrangement envisaged for the fission-fusion reaction process based on laser ion acceleration. Part a) illustrates the situation in case no collective effects on the electronic stopping are taken into account. In this case the thickness of the CH₂ layer as well as the second thorium reaction target have to be limited to 70 μ m and 50 μ m, respectively, in order to enable fission of beam and target nuclei. This will allow for fusion between their light fragment as well as enable the fusion products to leave the second thorium reaction target. Part b) depicts an alternative scenario, where we consider also collective effects in the reaction target induced by the ultra-dense ion bunches. Here the first part of the thorium reaction target is used to decelerate the fission fragments from about 7 MeV/u to about 3 MeV/u, suitable for efficient fusion of neutron rich species. Due to the reduced electronic stopping a larger target thickness and thus increased fission and fusion yields can be expected. Further details are discussed in the text.

3.2 Å for thorium, we conclude a fission probability of about $4.1 \cdot 10^{-9}$ per atomic layer.

In order to estimate the required thickness of the CH₂ front layer of the reaction target, we have to take into account the range of the 7 MeV/u 232 Th ions, which is about 120 μ m. However, already after 70 μ m the kinetic energy of the Th ions has dropped to 3 MeV/u, which is about the energy required for the resulting fission fragments during the subsequent fusion step. Therefore, we estimate the thickness of the polyethylene layer to about 70 μ m, which corresponds to $\sim 2.5 \cdot 10^5$ atomic layers. Together with the above estimated fission probability per atomic layer and taking into account that from CH₂ three atoms will contribute to the fission process of the impinging Th beam, this results in a fission probability for the Th ion beam of about $3.1 \cdot 10^{-3}$ in the 70 μm CH₂ layer, thus generating about 3.7·10⁸ beam-like fission fragments per laser pulse.

The 99.7% of Th beam ions passing through the CH₂ layer will enter the Th layer of the reaction target with about 2.4 MeV/u, corresponding to a range of 21 μ m. In the first atomic layers a fraction of them will undergo fission before being slowed down too much, however, the resulting fragment energies will not be suitable for the

fusion step. A quantitative assessment of this component would require detailed simulations and will be finally addressed by experimental studies.

In general, the fission process proceeds asymmetric [55]. The heavy fission fragment for 232 Th is centred at A=139.5 (approximately at Z=54.5 and N=84) close to the magic numbers Z=50 and N=82. Accordingly, the light fission fragment mass is adjusted to the mass of the fixed heavy fission fragment, thus resulting for 232 Th in A_L =91 with $Z_L \approx 37.5$. During the fission process of ²³²Th for low excitation energies, typically 1.6 neutrons are emitted. However, for the discussion presented here we neglect this loss of neutrons, because 4 or 5 neutrons may be transferred to the fissioning nucleus in the preceding transfer step (particularly efficient and thus important in case of the Th-induced fission discussed in the following section). The width (FWHM) of the light fission fragment peak is typically $\Delta A_L = 14$ mass units, the 1/10 maximum width about 22 mass units [55].

So far we have considered the fission process of beamlike Th nuclei in the CH₂ layer of the reaction target. Similar arguments can be invoked for the deuteron- (and carbon) induced generation of (target-like) fission products in the subsequent thicker thorium layer of the reaction target, where deuteron- and carbon-induced fission will occur in the $^{232}{\rm Th}$ layer of the reaction target. Since we can consider the $2.8\cdot 10^{11}$ laser-accelerated deuterons (plus $1.4\cdot 10^{11}$ carbon ions) impinging on the second target per laser pulse as 1860 consecutive atomic layers, we conclude a corresponding fission probability in the Th layer of the reaction target of about $2.3\cdot 10^{-5},$ corresponding to $3.2\cdot 10^6$ target-like fission fragments per laser pulse. A thickness of the thorium layer of the reaction target of about $50~\mu{\rm m}$ could be exploited, where the kinetic proton energy would be above the Coulomb barrier to induce fission over the full target depth.

An essential effect to be taken into account is the widening of the fission fragment beam, because a kinetic energy of about 1 MeV/u is added to the fission fragments in arbitrary directions. However, the angular distribution of fission fragments from proton (or heavy ion) induced fission follows a $1/\sin(\Theta)$ distribution [55] (with Θ denoting the fragment angle with respect to the direction of the incoming beam inducing the fission process) and thus fragments are predominantly emitted in beam direction. Consequently, a fraction of a few percent will stay within the conical volume defined by the spot diameter of the laser focus on the production target.

Due to the additional kinetic energy of about 1 MeV/u of the fission fragments in the thick reaction target also the target-like fragment volume will expand. Here the very short bunch length of the fragment beam becomes important. The beam velocity is about 10% of the velocity of light and during the short fly-by time of the ions of only 1 fs the fission fragments of the target can only move a distance of 1 μ m, which is still small compared to the beam diameter of 3 μ m. Thus this enlargement of the target area is negligible.

In a second step of the fission-fusion scenario, we consider the fusion between the light fission fragments of beam and target to a compound nucleus with a central value of $A \approx 182$ and $Z \approx 75$.

Again we employ geometrical arguments for an order-of-magnitude estimate of the corresponding fusion cross section. For a typical light fission fragment with A=90, the nuclear radius can be estimated as 5.4 fm. Considering a thickness of 50 μ m for the Th layer of the reaction target that will be converted to fission fragments, equivalent to $1.6\cdot10^5$ atomic layers, this results in a fusion probability of about $1.8\cdot10^{-4}$.

With this estimate for the fusion cross section, we can finally derive an order-of-magnitude for the final yield of fusion products generated via the presented fission-fusion process of about 1-2 fusion products per laser shot. This estimate does not yet take into account any collective effects in the target that might result in much higher fission rates and accordingly increased fusion yields as discussed in the following section.

Besides the fusion of two light fission fragments other reactions may happen. The fusion of a light fission fragment and a heavy fission fragment would lead back to the original Th nuclei, with large fission probabilities, thus we can neglect these fusion cross sections. The fusion of two heavy fission fragments would lead to nuclei with $A \approx 278$, again nuclei with very high fission probability. Hence we have also neglected these rare fusion cross sections, although they may be of interest on their own. Thus we concentrate here only on the fusion of two light fission fragments. Besides studying nuclei close to the waiting point of the r-process with the magic neutron number N=126, we may investigate also neutron-rich isotopes with the magic proton number Z=82, which are of large interest in nuclear structure studies.

Very neutron-rich nuclei still have comparably small production cross sections, because weakly bound neutrons ($B_N \geq 3$ MeV) will be evaporated easily. The optimum range of beam energies for fusion reactions resulting in neutron-rich fusion products amounts to about 2.8 MeV/u according to PACE4 [56] calculations. So, e.g., the fusion of two neutron-rich $^{98}_{55}$ Br fission products with a kinetic energy of the beam-like fragment of 275 MeV leads with an excitation energy of about 60 MeV to a fusion cross section of 13 mb for $^{189}_{70}$ Yb₁₁₉, which is already 8 neutrons away from the last presently known Yb isotope.

One should note that the well-known hindrance of fusion for nearly symmetric systems (break-down of fusion) only sets in for projectile and target masses heavier than 100 amu [57,58]. Thus for the fusion of light fission fragments, we expect an unhindered fusion evaporation process.

In Fig. 1 the range of reachable fusion products from the fission-fusion process is indicated by the blue ellipses overlayed to the chart of nuclides. The proton to neutron ratio, which is approximately conserved during fission (indicated by the straight blue line connecting ¹H with $^{238}\mathrm{U}$) determines the slope of the inner blue elliptical contour in Fig. 1. Their eccentricity reflects the region of nuclei reachable within a range of 50% of the maximum fusion cross section, based on the large fluctuations of proton and neutron numbers of the participating fission fragments. So far the dimensions of the contour lines drawn in Fig. 1 and Fig. 5 have been estimated from usual fission distributions. Since the mass distributions of both light fission fragments exhibit a certain width (FWHM), the width of the resulting distribution after fusion will be about a factor of $\sqrt{2}$ larger. Those distributions will steepen when reaching further out to their tails. The other two elliptical contour lines correspond to the regions of fusion products expected to be reachable with 10% and 10^{-3} of the maximum cross sections, respectively.

Fig. 5 displays a closer view into the region of nuclides around the N=126 waiting point of the r-process, where nuclei on the r-process path are indicated by the green

colour, with dark green highlighting the key bottleneck r-process isotopes [59] at N=126 between Z=66 (Dy) and Z=70 (Yb). One should note that, e.g., for Yb the presently last known isotope is 15 neutrons away from the r-process path at N=126. The isotopes in light blue mark those nuclides, where recently β -halflives could be measured following projectile fragmentation and inflight separation at GSI [60]. Again the elliptical contour lines indicate the range of nuclei accessible with our new fission-fusion scenario on a level of 50%, 10% and 10^{-3} of the maximum fusion cross section between two neutron-rich light fission fragments in the energy range of about 2.8 MeV/u, respectively.

3.3.2 Fission-Fusion with collectively reduced electronic stopping So far we estimated the expected fission and fusion yield without refering to any collective effects in the reaction targets that may reduce the electronic stopping as discussed earlier. Now we extend this discussion by considering the expected reduction of the electronic stopping in the reaction target. This scenario would allow to extend the thickness of the Th production target to probably a few mm (situation 'b)' in Fig. 4).

While so far no experimental data or quantitative assessment on the amount of collective range enhancement is available, we assume for the discussion within this paragraph a factor of 100 and discuss the consequences.

In contrast to the previously discussed scenario without collective effects, we now propose to abandon the front carbon layer of the reaction target and use only a homogeneous, thick Th target as indicated in Fig. 4b). In this case we use the first part of the target primarily as stopping medium for the incoming energetic Th ions in order to decelerate them from initially about 7 MeV/u to about 3 MeV/u, which is suitable for the subsequent fusion step with target-like fragments from proton-induced fission. Since such a deceleration could be reached in about 16 μ m without collective effects, we estimate here about 0.2 mm from our 5 mm thick Th reaction target acting as stopper while producing fission fragments too fast for efficient fusion of extremely neutron-rich isotopes. This part of the reaction target is marked by the lighter red colour in Fig. 4b). However, this part amounts to only 4% of the reaction target and thus does not lead to a significant loss of usable fission yield. On the other hand, neutron transfer towards the deformed neutron shell closure at N=146 preceding fission will add 4 neutrons to the light fission fragment and thus significantly help to enhance fusion of very neutronrich isotopes. Compared to the situation of Fig. 4a), a thickness of the thorium target increased by a factor of 100 to about 5 mm due to the correspondingly reduced stopping would result in a full conversion of the Th beam into fission fragments (with $\sim 96\%$ in an energy range usable for the fusion step). Thus $1.2 \cdot 10^{11}$ beam-like light fission fragments would become available for the fusion stage of the reaction process.

Table 1 Compilation of relevant parameters determining the expected yield (per laser pulse) of the fission-fusion reaction process proposed in this work.

normal	reduced
stopping	stopping
560 nm	560 nm
520 nm	520 nm
$1.2 \cdot 10^{11}$	$1.2 \cdot 10^{11}$
$2.8 \cdot 10^{11}$	$2.8 \cdot 10^{11}$
$1.4 \cdot 10^{11}$	$1.4 \cdot 10^{11}$
$70~\mu\mathrm{m}$	_
$50~\mu\mathrm{m}$	5 mm
$3.7 \cdot 10^{8}$	$1.2 \cdot 10^{11}$
$2.3 \cdot 10^{-5}$	$2.3 \cdot 10^{-3}$
$3.2 \cdot 10^6$	$1.2 \cdot 10^{11}$
$1.8 \cdot 10^{-4}$	$1.8 \cdot 10^{-4}$
1.5	$4 \cdot 10^4$
	560 nm 520 nm $1.2 \cdot 10^{11}$ $2.8 \cdot 10^{11}$ $1.4 \cdot 10^{11}$ 70 μ m 50μ m $3.7 \cdot 10^{8}$ $2.3 \cdot 10^{-5}$ $3.2 \cdot 10^{6}$ $1.8 \cdot 10^{-4}$

Consequently the deuteron-induced fission yield in the reaction target would also rise by the same factor of 100 to a target fission probability of $2.3 \cdot 10^{-3}$. Here we use the assumption of a linear rise of the energy loss with target thickness, preserving a kinetic proton energy above the fission barrier to induce fission over the full target depth.

So we conclude that the expected collective stopping range enhancement will lead to a drastic increase of the fusion yield from about 1-2 fusion products per laser pulse to a value of about $4 \cdot 10^4$ exotic nuclides per pulse. Most likely only part of this estimated yield enhancement could finally be realized, so it may be adequate to finally quote the average between the two extremes, resulting in an estimate of about 10³ fusion products generated per laser pulse. However, it is obvious that collective effects from the ultra-dense ion bunches would significantly improve the experimental conditions towards the production of extremely neutron rich fusion products. Moreover, if we could use a layered production target instead of the presently separated arrangement, while still achieving quasi-monoenergetic thorium and CD_2 ion beams, then the primary fission fragment rates would each increase by a factor of 2, thus resulting in an increase of the fusion yield by a factor of 4.

The following table finally gives a quantitative overview of the two discussed experimental scenarios with and without collective stopping reduction, based on the parameters of the driver laser introduced earlier. All numbers refer to yields expected for one laser pulse.

While it will remain a challenge to directly study the key waiting point isotopes on the r-process path, it is on the other hand intriguing that a wide range of so far unknown isotopes will become accessible for experimental investigation.




Fig. 5 Chart of nuclides around the N=126 waiting point of the r-process path. The blue ellipses denote the expected range of isotopes accessible via the novel fission-fusion process. The indicated lines represent 0.5, 0.1 and 0.001 of the maximum fusion cross section after neutron evaporation. In green the N=126 nuclides relevant for the r-process are marked, with the dark green colour indicating the key bottleneck nuclei for the astrophysical r-process.

Presently the high-intensity APOLLON laser envisaged to be used for laser ion acceleration is designed to operate at a repetition rate of one laser pulse per minute. However, laser technology is progressing rapidly with large efforts presently devoted to the development of higher repetition rates, aiming of up to 10 Hz together with an increase of the laser pulse energy beyond 1 kJ. Moreover, since the yield of very neutron-rich fusion products grows strongly nonlinear with laser energy, a final use of several coincident APOLLON laser beams would be very advantageous.

Therefore, it is foreseeable that the above given estimate for the achievable rate of neutron-rich fusion products can be increased within the next years significantly by several orders of magnitude.

4 Experimental aspects

Exploring this 'terra incognita' of yet unknown isotopes towards the r-process waiting point at N= 126 certainly calls for a staged experimental approach. First studies should focus on the range and electronic stopping powers of dense laser-accelerated ion beams as discussed previously, followed by systematic optimizations of target properties in order to optimize the yield of fission fragments. Subsequently the A, Z and N distributions of the light thorium fission fragments should be characterized. Moreover, it is unclear in how far the first neutron transfer preceding fission will additionally broaden all these distributions. Also the yields for the fusion products should be measured in exploratory experiments, where

it will be crucial to optimize the kinetic energy of the beam-like fission products.

Fig. 6 shows a schematical view of the potential experimental setup of the presented reaction scenario. The high-intensity laser beam is tightly focussed onto the target assembly. This area will require heavy concrete shielding for radioprotection. The probably most essential and also most demanding experimental task will be the separation of the reaction products. Fusion products with about 2-3 MeV/u will have to be separated from faster beam-like fission fragments with about 7 MeV/u, or target-like fragments with about 1 MeV/u, which could be achieved with a velocity filter. However, the reaction products from various fusion channels varying in mass but not in velocity require a different separation scheme. Here one could use a recoil separator (as indicated in Fig. 6), where it may be advantageous to operate the separator in gas-filled mode. Alternatively, also a coarse magnetic dipole pre-separator followed by a gas stopping cell and an RFQ cooler/buncher could be used to inject the ions into an electrostatic mass separator like the 'Multi-Reflection Time-of-Flight' mass spectrometer [61], especially when aiming for fusion products with lifetimes significantly shorter than 100 ms. Such a spectrometer could be operated either as an isobar separator or directly for mass measurements with a mass accuracy of up to 10^{-7} .

In these first studies, a tape station could be used to transport the reaction products to a remote, well-shielded detector system, where the characterization of the implanted fusion products could be performed either via β -decay studies using, e.g., LaBr₃ scintillation detec-




Fig. 6 Schematical view of the experimental arrangement for fission-fusion studies. Measurements of fusion products will be performed in two stages, first aiming at an identification of the produced isotopes via decay spectroscopy using a transport system (e.g. tape), while later on precision mass measurements using a Penning trap system are envisaged.

tors or γ spectroscopy with high-resolution germanium detectors. This scenario has been labelled 'Phase 1' in Fig. 6. Since most of the fusion products have typical lifetimes of ≈ 100 ms, they will survive the transport to a secondary target and/or detector station.

In a later stage ('Phase 2'), the fusion products may be stopped in a buffer gas stopping cell [62], cooled and bunched in, e.g., a radiofrequency quadrupole ion guide before then being transferred to a Penning trap system for high-accuracy mass measurements. Such a setup would be similar to the SHIPTRAP facility at GSI [63] or ISOLTRAP at ISOLDE/CERN [64] for mass measurement with an accuracy of $\Delta m/m \sim 10^{-8}$, corresponding to about 10 keV/c^2 [65].

5 Conclusion

The exploration of nuclei far away from the valley of stability is a long-term endeavour of nuclear physics with strong relevance for astrophysical applications. In our present experimental proposal of a new nuclear reaction scheme, we address the heavy nuclei of the r-process nucleosynthesis path towards the waiting point at N=126, where our new production scheme holds promise to bring these extremely neutron-rich isotopes into reach of direct experimental studies with significantly higher yields than accessible with classical radioactive ion beam accelerator technology. With much more compact highpower, short-pulse laser systems we intend to develop an optimized production scheme for extremely neutronrich fusion products following induced fission from laseraccelerated ion beams. Exploiting the 'hole-boring' mode of the Radiation Pressure Acceleration mechanism will allow to generate ion beams of fissile species with solidstate density. A two-step production scheme of neutronrich nuclides ('fission-fusion') is proposed, where asymmetric fission preceded by a deep inelastic transfer reaction will be followed by fusion of the light fission fragments. Moreover, collective effects reducing the electronic stopping power in the target are expected for such ultradense ion bunches, allowing to use much thicker targets and thus increasing the fission yield significantly. The fusion of short-lived, neutron-rich fission fragment beams with short-lived, neutron-rich fission fragments in the target will result in very attractive production rates of extremely neutron-rich nuclides towards N=126 and Z>70. Order-of-magnitude estimates promise fusion rates of several 10³ fusion products per laser pulse, based on the laser parameters envisaged for the ELI-Nuclear Physics project in Bucharest (2 x 150 J, 32 fs). Whereas the present repetition rate of 1 laser pulse per minute limits the achievable fusion yield, ongoing development efforts for significantly higher repetition rates (aiming at up to 10 Hz) and increased laser energy (aiming at beyond 1 kJ) will open the perspective to increase the achievable yields within the fission-fusion reaction scheme by several orders of magnitude within the next years. In this way, high-power lasers used for laser ion acceleration can significantly contribute to access terra incognita in nuclear physics and astrophysical nucleosynthesis of heavy elements.

Acknowledgement

We acknowledge helpful discussions with M. Heil, K.L. Kratz, H.Th. Janka and P. Ring. The authors enjoyed the collaboration with V. Zamfir, who is heading the ELI-NP project, opening up many of the new perspectives presented here. We were supported by the DFG

Clusters of Excellence: 'Munich Centre for Advanced Photonics' (MAP) and 'Origin and Structure of the Universe' (UNIVERSE) and by the DFG Transregio TR18.

References

- K.L. Kratz, K. Farouqi, B. Pfeiffer, Nuclear physics far from stability and r-process nucleosynthesis, Prog. in Part. and Nucl. Phys. 59, 147 (2007).
- 2. M. Arnould, S. Goriely, K. Takahashi, The r-process of stellar nucleosynthesis: Astrophysics and nuclear physics achievements and mysteries, Phys. Rep. 450, 97 (2007).
- I.V Panov and H.-Th. Janka, On the dynamics of protoneutron star winds and r-process nucleosynthesis, Astr. Astroph. 494, 829 (2009).
- A.P.L. Robinson, P. Gibbon, M. Zepf, S. Kar, R.G. Evans and C. Bellei, Relativistically correct hole-boring and ion acceleration by circularly polarized laser pulses, Plasma Phys. Control. Fusion 51 (2009) 024004.
- A. Henig, S. Steinke, M. Schnürer, T. Sokollik, R. Hörlein, D. Kiefer, D. Jung, J. Schreiber, B.M. Hegelich, X.Q. Yan, J. Meyer-ter-Vehn, T. Tajima, P.V. Nickles, W. Sandner, and D. Habs, Radiation-Pressure Acceleration of Ion Beams Driven by Circularly Polarized Laser Pulses, Phys. Rev. Lett. 103, 245003 (2009).
- T. Tajima, D. Habs, X. Yan, Laser Acceleration of Ions for Radiation Therapy, Rev. Accel. Science and Technol. 2, 221 (2009).
- 7. http://www.eli-np.ro
- 8. E. Haseltine, http://discovermagazine.com2002febcover
- 9. C.E. Rolfs, *Cauldrons in the Cosmos*, Univ. of Chicago Press (1991).
- K.-L. Kratz, K. Farouqi, B. Pfeiffer, J.W. Truran, C. Sneden and J.J. Cowan, Explorations of the r-Processes: Comparisons between Calculations and Observations of Low-Metallicity Stars, Ap.J. 662, 39 (2007).
- H.-Th. Janka, K. Langanke, A. Marek, G. Martinez-Pinedo and B. Müller, *Theory of core-collapse super-novae*, Phys. Rep. 442, 38 2007.
- J.J. Cowan and F.-K. Thielemann, R-Process Nucleosynthesis in Supernovae, Physics Today 57 (2004) 47.
- C. Freiburghaus, S. Rosswog and F.-K. Thielemann, r-Process in Neutron Star Mergers, Ap. Jour. 525, L121 (1999).
- 14. R. Surman, G.C. McLaughlin, M. Ruffert, H.-Th. Janka and W.R. Hix, r-Process Nucleosynthesis in Hot Accretion Disk Flows from Black Hole-Neutron Star Mergers, Ap. Jour. 679, L117 (2008).
- 15. K. Farouqi, C. Freiburghaus, K.-L. Kratz, B. Pfeiffer, T. Rauscher and F.-K. Thielemann, Astrophysical conditions for an r-process in the high-entropy wind scenario of type II supernovae, Nucl. Phys. A758, 631c (2005).
- 16. J.M. Pearson, R. C. Nayak and S. Goriely, *Nuclear mass formula with Bogolyubov-enhanced shell-quenching: application to r-process*, Phys. Lett. **B387**, 455 (1996).
- 17. J.J. Cowan and C. Sneden, Heavy element synthesis in the oldest stars and the early Universe, Nature 440, 1151 (2006).
- 18. J. Dobaczewski, H. Flocard, J. Treiner, *Hartree-Fock-Bogolyubov description of nuclei near the neutron-drip line*, Nucl. Phys. **A 422**, 103 (1984).

- 19. FAIR, An International Accelerator Facility for Beams of Ions and Antiprotons, Baseline Technical Report, GSI (2006), http://www.gsi.defairreportsbtr.html.
- 20. SPIRAL II, Detailed Design Study APD Report, GANIL (2005), http:pro.ganil-spiral2.euspiral2what-is-spiral2apd
- 21. http://www.frib.msu.eduaboutmsu-frib-proposal
- 22. S. Baruah, G. Audi, K. Blaum, M. Dworschak, S. George, C. Guenaut, U. Hager, F. Herfurth, A. Herlert, A. Kellerbauer, H.-J. Kluge, D. Lunney, H. Schatz, L. Schweikhard and C. Yazidjian, Mass Measurements beyond the Major r-Process Waiting Point ⁸⁰Zn, Phys. Rev. Lett. 101, 262501 (2008).
- 23. M. Dworschak, G. Audi, K. Blaum, P. Delahaye, S. George, U. Hager, F. Herfurth, A. Herlert, A. Kellerbauer, H.-J. Kluge, D. Lunney, L. Schweikhard and C. Yazidjian, Restoration of the N=82 Shell Gap from Direct Mass Measurements of ^{132,134}Sn, Phys Rev. Lett. 100, 072501 (2008).
- 24. I. Dillmann, K.-L. Kratz, A. Wöhr, O. Arndt, B.A. Brown, P. Hoff, M. Hjorth-Jensen, U. Köster, A.N. Ostrowski, B. Pfeiffer, D. Seweryniak, J. Shergur and W. B. Walters, N=82 Shell Quenching of the Classical r-Process 'Waiting-Point' Nucleus ¹³⁰ Cd, Phys. Rev. Lett 91, 162503 (2003).
- K. Blaum, High-accuracy mass spectrometry with stored ions, Phys. Rep. 425, 1 (2006).
- G. Audi, A.H. Wapstra, The 1995 update to the atomic mass evaluation, Nucl. Phys. A 595, 409 (1995).
- I.V. Panov, E. Kolbe, B. Pfeiffer, T. Rauscher, K.-L. Kratz and F.-K. Thielemann, Calculations of fission rates for r-process nucleosynthesis, Nucl. Phys. A747 (2005) 633.
- 28. P. Ring, private communication (2010).
- 29. P. Möller, J.R. Nix, W.D. Myers and W.J. Swiatecki, *Nuclear Ground-State Masses and Deformations*, Atomic Data Nucl. Data Tabl. **59**, 185 (1995).
- B. M. Hegelich, B.J. Albright, J. Cobble, K. Flippo,
 S. Letzring, M. Paffett, H. Ruhl, J. Schreiber, R.K.
 Schulze and J.C. Fernandez, Laser acceleration of quasimonoenergetic MeV ion beams, Nature 439, 441 (2006).
- H. Schwörer, S. Pfotenhauer, O. Jäckel, K.-U. Amthor, B. Liesfeld, W. Ziegler, R. Sauerbrey, K.W.D. Ledingham and T. Esirkepov, Laser plasma acceleration of quasi-monoenergetic protons from microstructured targets, Nature 439, 445 (2006).
- S. Ter-Avetisyan, M. Schnürer, P.V. Nickles, M. Kalashnikov, E. Risse, T. Sokollik, W. Sandner, A. Andreev and V. Tikhonchuk, *Quasimonoenergetic Deuteron Bursts Produced by Ultraintense Laser Pulses*, Phys. Rev. Lett. 96, 145006 (2006).
- 33. R.A. Snavely, M.H. Key, S.P. Hatchett, T.E. Cowan, M. Roth, T.W. Phillips, M.A. Stoyer, E.A. Henry, T.C. Sangster, M.S. Singh, S.C. Wilks, A. MacKinnon, A. Offenberger, D.M. Pennington, K. Yasuike, A.B. Langdon, B.F. Lasinski, J. Johnson, M.D. Perry and E.M. Campbell, Intense High-Energy Proton Beams from Petawatt-Laser Irradiation of Solids, Phys. Rev. Lett. 85, 2945 (2000).
- 34. E. Clarke, K. Krushelnick, J.R. Davies, M. Zepf, M. Tatarakis, F.N. Beg, A. Machacek, P.A. Norreys, M.I.K. Santala, I. Watts, and A.E. Dangor, *Measurements of*

- Energetic Proton Transport through Magnetized Plasma from Intense Laser Interactions with Solids, Phys. Rev. Lett. 84, 670 (2000).
- 35. A. Maksimchuck, S. Gu, K. Flippo, D. Umstadter, V.Yu. Bychenkov, Forward Ion Acceleration in Thin Films Driven by a High-Intensity Laser, Phys. Rev. Lett. 84,4108 (2000).
- 36. S. P. Hatchett, C.G. Brown, T.E. Cowan, E.A. Henry, J.S. Johnson, M.H. Key, J.A. Koch, A.B. Langdon, B.F. Lasinski, R.W. Lee, A.J. Mackinnon, D.M. Pennington, M.D. Perry, T.W. Phillips, M. Roth, T.C. Sangster, M.S. Singh, R.A. Snavely, M.A. Stoyer, S.C. Wilks and K. Yasuike, Electron, photon, and ion beams from the relativistic interaction of Petawatt laser pulses with solid targets, Phys. Plasma 7, 2076 (2000).
- 37. L. Robson, P.T. Simpson, R.J. Clarke, K.W.D. Ledingham, F. Lindau, O. Lundh, T. McCanny, P. Mora, D. Neely, C.-G. Wahlström, M. Zepf, P. McKenna, Scaling of proton acceleration driven by petawatt-laserplasma interactions Nature Physics 3, 58 (2007) and refs. therein.
- 38. A. Macchi, F. Cattani, T.V. Liseykina and F. Cornolti, Laser Acceleration of Ion Bunches at the Front Surface of Overdense Plasmas, Phys. Rev. Lett. **94**, 165003 (2005).
- 39. O. Klimo, J. Psikal, J. Limpouch, V.T. Tikhonchuk, Monoenergetic ion beams from ultrathin foils irradiated by ultrahigh-contrast circularly polarized laser pulses, Phys. Rev. ST Accl. Beams 11, 031301 (2008).
- 40. A.P.L. Robinson, M. Zepf, S. Kar, R.G. Evans and C. Bellei, Radiation pressure acceleration of thin foils with circularly polarized laser pulses, New J. Phys. 10, 013021
- 41. S.G. Rykovanov, J. Schreiber, J. Meyer-ter-Vehn, C. Bellei, A. Henig, H.C. Wu and M. Geissler, Ion acceleration with ultra-thin foils using elliptically polarized laser pulses, New J. Phys. 10, 113005 (2008).
- 42. A.P.L. Robinson, D.-H. Kwon and K. Lancaster, Holeboring radiation pressure acceleration with two ion species, Plasma Phys. Control. Fusion 51 (2009) 095006.
- 43. T.-P. Yu, A. Pukhov, G. Shvets, M. Chen Stable Laser-Driven Proton Beam Acceleration from a Two-Ion-Species Ultrathin Foil Phys. Rev. Lett. 105, 065002 (2010).
- 44. M. Chen, A. Pukhov, T.P. Yu, Z.M. Sheng Enhanced Collimated GeV Monoenergetic Ion Acceleration from a Shaped Foil Target Irradiated by a Circularly Polarized Laser Pulse Phys. Rev. Lett. 103, 024801 (2009).
- 45. X.Q. Yan, C. Lin, Z.M. Sheng, Z.Y. Guo, B.C. Liu, Y.R. Lu, J.X. Fang, J.E. Chen Generating High-Current Monoenergetic Proton Beams by a Circularly Polarized Laser Pulse in the Phase-Stable $Acceleration\ Regime$ Phys. Rev. Lett. 100, 135003 (2008).
- 46. V.K. Tripathi, C.S. Liu, X. Shao, B. Eliasson, R.Z. Sagdeev Laser acceleration of monoenergetic protons in a selforganized double layer from thin foil Plasma Phys. Control. Fusion **51** (2009) 024014.
- 47. X.Q. Yan, H.C. Wu, Z.M. Sheng, J.E. Chen, J. Meyer-Self-Organizing GeV, Nanocoulomb, Collimated Proton

- Beam from Laser Foil Interaction at 7×10^{21} W/cm² Phys. Rev. Lett. 103, 135001 (2009).
- 48. X.Q. Yan, T. Tajima, M. Hegelich, L. Yin and D. Habs, Theory of laser ion acceleration from a foil target of nanometer thickness, Appl. Phys. B 98 711 (2010).
- 49. G. Mourou, The APOLLON/ILE 10 PW laser project, http://www.nipne.ro/eli_np_workshop/contributions.php
- 50. S. Segré, Nuclei and Particles, second edt., W.A. Benjamin, RA, London (1977)
- 51. S. Ichimaru, Basic Principles of Plasma Physics: A Statistical Approach, Benjamin, Reading, Ma (1973).
- 52. H.-C. Wu etal.,"Collective Deceleration"; arXiv:0909.1530v1[physics.plasm-ph], submitted to Phys. Rev. ST AB.
- 53. P.G. Thirolf and D. Habs, Spectroscopy in the Second and Third Minimum of Actinide Nuclei, Prog. in Part. and Nucl. Phys. 49, 325 (2002).
- 54. V. Metag., Nukleonica **20**, 789 (1975).
- 55. R. Vandenbosch and J.R. Huizenga, Nuclear fission, Academic Press, New York (1973)
- 56. A. Gavron, Statistical model calculations in heavy ion reactions, Phys. Rev. C21 (1980) 230; O.B. Tarasov and D. Bazin, Development of the program LISE: application to fusion-evaporation, Nucl. Instr. Meth. **B204** (2003) 174.
- 57. A.B. Quint, W. Reisdorf, K.-H. Schmidt, P. Armbruster, F-P. Heßberger, S. Hofmann, J. Keller, G. Münzenberg, H. Stelzer, H.-G. Clerc, W. Morawek, C.-C. Sahm, Investigation of the fusion of heavy nearly symmetric systems, Z. Phys. A 346, 119 (1993).
- 58. W. Morawek, D. Ackermann, T. Brohm, H.-G. Clerc, U. Gollerthan, E. Hanelt, M. Horz, W. Schwab, B. Voss, K.-H. Schmidt, F.-P. He??berger, Breakdown of the compound-nucleus model in the fusion-evaporation process for $^{110}Pd+^{110}Pd$, Z. Phys. **A 341**, 75 (1991).
- 59. NRC Rare Isotope Science Assessment Committee (RISAC) Report, 2007, National Academies Press, Washington/DC, USA.
- 60. T. Kurtukian-Nieto, J. Benlliure, K.H. Schmidt, A new analysis method to determine β -decay half-lives in experiments with complex background, Nucl. Instr. Meth. A589 (2008) 472
- 61. W.R. Plass, T. Dickel, U. Czok, H. Geissel, M. Petrick, K. Reinheimer, C. Scheidenberger, M.I.Yavor, Isobar separation by time-of-flight mass spectrometry for low-energy radioactive ion beam facilities, Nucl. Instr. Meth. B266 (2008) 4560.
- 62. J.B. Neumayr, L. Beck, D. Habs, S. Heinz, J. Szerypo, P.G. Thirolf, V. Varentsov, F. Voit, D. Ackermann, D. Beck, M. Block, A. Chaudhuri, Z. Di, S.A. Eliseev, H. Geissel, F. Herfurth, F.P. Heßberger, S. Hofmann, H.J. Kluge, M. Mukherjee, G. Münzenberg, M. Petrick, W. Quint, S. Rahaman, C. Rauth, D. Rodriguez, C. Scheidenberger, G. Sikler, Z. Wang, C. Weber, W.R. Plaß, M. Breitenfeldt, G. Marx, L. Schweikhard, A.F. Dodonov, Y. Novikov, M. Suhonen, The ion catcher device for SHIP-TRAP, Nucl. Inst. Meth. **B244**, 489 (2006). J.B. Neumayr, P.G. Thirolf, D. Habs, S. Heinz, V.S. Kolhinen, M. Sewtz, J. Szerypo, Performance of the MLL-
 - IonCatcher, Rev. Sci. Instr. 77, 065109 (2006).
- 63. M. Block, D. Ackermann, D. Beck, K. Blaum, M. Breitenfeld, A. Chauduri, A. Doemer, S. Eliseev, D. Habs,

- S. Heinz, F. Herfurth, F.P. Hessberger, S. Hofmann, H. Geissel, H.-J. Kluge, V. Kolhinen, G. Marx, J.B. Neumayr, M. Mukherjee, M. Petrick, W. Plass, W. Quint, S. Rahaman, C. Rauth, D. Rodriguez, C. Scheidenberger, L. Schweikhardt, M. Suhonen, P.G. Thirolf, Z. Wang, C. Weber, and the SHIPTRAP collaboration, *The ion trap facility SHIPTRAP*, Eur. Phys. Jour. A25 (suppl. 1) (2005) 49.
- 64. M. Mukherjee, D. Beck, K. Blaum, G. Bollen, J. Dilling, S. George, F. Herfurth, A. Herlert, A. Kellerbauer, H.-J. Kluge, S. Schwarz, L. Schweikhard, C. Yazidjian, ISOLTRAP: An on-line Penning trap for mass spectrometry on short-lived nuclides, Eur. Phys. J. A 35, 1-29 (2008).
- 65. M. Block, D. Ackermann, K. Blaum, C. Droese, M. Dworschak, S. Eliseev, T. Fleckenstein, E. Haettner, F. Herfurth, F.P. Heßberger, S. Hofmann, J. Ketelaer, J. Ketter, H.-J. Kluge, G. Marx, M. Mazzocco, Yu.N. Novikov, W. R. Plaß, A. Popeko, S. Rahaman, D. Rodriguez, C. Scheidenberger, L. Schweikhard, P.G. Thirolf, G.K. Vorobyev, C. Weber, Direct mass measurements above uranium bridge the gap to the island of stability, Nature 463, 785 (2010).