

Thermal quantum and classical correlations in two qubit XX model in a nonuniform external magnetic field.

Ali Saif M. Hassan,^{1,*} Behzad Lari,^{2,†} and Pramod S. Joag^{2,‡}

¹*Department of Physics, University of Amran, Amran, Yemen*

²*Department of Physics, University of Pune, Pune, India-411007.*

(Dated: May 4, 2021)

We investigate how thermal quantum discord (QD) and classical correlations (CC) of a two qubit one-dimensional XX Heisenberg chain in thermal equilibrium depend on temperature of the bath as well as on nonuniform external magnetic fields applied to two qubits and varied separately. We show that the behaviour of QD differs in many unexpected ways from thermal entanglement (EOF). For the nonuniform case, ($B_1 = -B_2$) we find that QD and CC are equal for all values of ($B_1 = -B_2$) and for different temperatures. We show that, in this case, the thermal states of the system belong to a class of mixed states and satisfy certain conditions under which QD and CC are equal. The specification of this class and the corresponding conditions are completely general and apply to any quantum system in a state in this class and satisfying these conditions. We further find that the relative contributions of QD and CC can be controlled easily by changing the relative magnitudes of B_1 and B_2 . Finally, we connect our results with the monogamy relations between the EOF, classical correlations and the quantum discord of two qubits and the environment.

PACS numbers: 03.65.Ud;75.10.Pq;05.30.-d

It is now quite well known that composite quantum systems can be in a class of states, called entangled states, in which the correlations between the constituents of the system cannot be achieved in a classical world [1, 2]. Although all pure entangled states possess such nonlocal quantum correlations, there are mixed entangled states which do not, in the sense of violating Bell inequalities [3]. The entanglement in quantum states and the resulting nonlocal quantum correlations form an area of intense research, due to their huge technological promise, especially in the areas of quantum communication and cryptography [4]. However, quantum correlations breaking Bell inequalities need not account for all quantum correlations in a composite quantum system in a given state. In order to account for the quantum correlation in a given state, we must find some means to divide the total correlation into a classical part and a purely quantum part. This is particularly important for mixed states, since their quantum correlations are many a time hidden by their classical correlations (CC). An answer to this requirement is given by quantum discord (QD), a measure of the quantumness of correlations introduced in Ref. [5]. Quantum discord is built on the fact that two classically equivalent ways of defining the mutual information turn out to be inequivalent in the quantum domain. In addition to its conceptual role, some recent results [6], suggest that quantum discord and not entanglement may be responsible for the efficiency of a mixed state based quantum computer. Present authors believe that QD will turn out to be a very useful tool to analyze mixed state quantum correlations and their consequences, as mixed state entanglement is very difficult and eluding to deal with [7]. To realize this hope we need a viable relation between mixed state entanglement and quantum discord

[8]. The pointers towards such a relation may be obtained by studying these properties for various quantum systems.

Motivated by these considerations, we present here the results of our investigation of the amount of QD and CC in a two qubit Heisenberg XX chain at finite temperature subjected to nonuniform external magnetic fields B_1 and B_2 acting separately on each qubit. We study two distinct cases namely $B_1 = -B_2$ (nonuniform field) and $B_1 = B_2$ (uniform field). In each case, we obtain the dependance of QD , CC and entanglement of formation (EOF) [17] in the system on the external magnetic field and temperature. Such a model is realized, for example, by a pair of qubits (spin 1/2) within a solid at finite temperature experiencing a spatially varying magnetic field. Such Heisenberg models can describe fairly well the magnetic properties of real solids [9] and are well adapted to the study of the interplay of disorder and entanglement as well as of entanglement and quantum phase transitions [10, 11]. The variation of entanglement [12] and QD [13, 14] in a two qubit Heisenberg XX chain with external magnetic field is already reported.

In order to quantify entanglement in a thermally mixed two qubit state, we use entanglement of formation derived using concurrence given by [15]

$$C = \max\{\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0\} \quad (1)$$

where $\lambda_i (i = 1, 2, 3, 4)$ are the square roots of the eigenvalues of the operator $\rho\tilde{\rho}$ in descending order

$$\tilde{\rho} = (\sigma_1^y \otimes \sigma_2^y)\rho^*(\sigma_1^y \otimes \sigma_2^y), \quad (2)$$

with $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4$, and ρ is the density matrix of the pair qubits; σ_1^y and σ_2^y are the normal Pauli operators. The entanglement of formation is related to the

concurrence by

$$EN = h \left(\frac{1 + \sqrt{1 - C^2}}{2} \right),$$

where $h(x) = -x \log_2 x - (1-x) \log_2 (1-x)$. Henceforth, in this paper, we denote the entanglement of formation (EOF) by EN . The concurrence $C = 0$ corresponds to an unentangled state and $C = 1$ corresponds to a maximally entangled state.

The thermalized Heisenberg system. The model Hamiltonian we study is given by

$$H = J(S_1^x S_2^x + S_1^y S_2^y) + B_1 S_1^z + B_2 S_2^z, \quad (3)$$

where $S^\alpha \equiv \sigma^\alpha/2$, ($\alpha = x, y, z$) are the spin 1/2 operators, σ^α are the Pauli operators and J is the strength of Heisenberg interaction. B_1 and B_2 are external magnetic fields. As stated in the introduction, by changing B_1 and B_2 separately, we want to study the effects of magnetic field on the thermal QD , CC and EN in a very general way. The eigenvalues and eigenvectors of H are

$$\begin{aligned} H|00\rangle &= -(B_1 + B_2)|00\rangle, \\ H|11\rangle &= (B_1 + B_2)|11\rangle, \\ H|\psi^\pm\rangle &= \pm D|\psi^\pm\rangle, \end{aligned} \quad (4)$$

where $D^2 = (B_1 - B_2)^2 + J^2$ and $|\psi^\pm\rangle = \frac{1}{\sqrt{N_\pm}}[|01\rangle + \frac{(B_1 - B_2) \pm D}{J}|10\rangle]$. We denote the eigenvalues corresponding to $|00\rangle, |11\rangle, |\psi^\pm\rangle$ by E_{00}, E_{11}, E_\pm respectively. In the standard basis, $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$, the density matrix $\rho(T)$ is given by

$$\rho(T) = \frac{1}{Z} \begin{bmatrix} u_1 & 0 & 0 & 0 \\ 0 & w_1 & v & 0 \\ 0 & v & w_2 & 0 \\ 0 & 0 & 0 & u_2 \end{bmatrix}, \quad (5)$$

where

$$\begin{aligned} u_1 &= e^{(B_1 + B_2)/kT}, \\ u_2 &= e^{-(B_1 + B_2)/kT}, \\ w_1 &= \cosh\left(\frac{D}{kT}\right) + \frac{(B_1 - B_2)}{D} \sinh\left(\frac{D}{kT}\right), \\ w_2 &= \cosh\left(\frac{D}{kT}\right) - \frac{(B_1 - B_2)}{D} \sinh\left(\frac{D}{kT}\right), \\ v &= -\frac{J \sinh\left(\frac{D}{kT}\right)}{D}, \end{aligned} \quad (6)$$

and $Z = Tr[\exp(\frac{-H}{kT})]$ is the partition function. In the following we select $|J|$ as the energy unit and set $k = 1$.

Quantum Discord. [5, 13] In classical information theory (CIT) the total correlation between two systems (two sets of random variables) A and B described by a joint distribution probability $p(A, B)$ is given by the mutual information (MI),

$$I(A, B) = H(A) + H(B) - H(A, B), \quad (7)$$

with the Shannon entropy $H(\cdot) = -\sum_j p_j \log_2 p_j$. Here p_j represents the probability of an event j associated to systems A, B , or to the joint system AB . Using Bayes's rule we may write MI as

$$I(A, B) = H(A) - H(A|B), \quad (8)$$

where $H(A|B)$ is the classical conditional entropy. In CIT these two expressions are equivalent but in the quantum domain this is no longer true [5, 16]. The first quantum extension of MI, denoted by $I(\rho)$, is obtained directly replacing the Shannon entropy in Eq.(7) with the von Neumann entropy, $S(\rho) = -Tr(\rho \log_2 \rho)$, with ρ , a density matrix, replacing probability distributions. To obtain a quantum version of Eq. (8) it is necessary to generalize the classical conditional entropy. This is done recognizing $H(A|B)$ as a measure of our ignorance about system A after we make a set of measurements on B . When B is a quantum system the choice of measurements determines the amount of information we can extract from it. We restrict ourselves to von Neumann measurements on B described by a complete set of orthogonal projectors, Π_j , corresponding to outcomes j .

After a measurement, the quantum state ρ changes to $\rho_j = [(I \otimes \Pi_j)\rho(I \otimes \Pi_j)]/p_j$, with I the identity operator for system A and $p_j = Tr[(I \otimes \Pi_j)\rho(I \otimes \Pi_j)]$. Thus, one defines the quantum analog of the conditional entropy as $S(\rho|\{\Pi_j\}) = \sum_j p_j S(\rho_j)$ and, consequently, the second quantum extension of the classical MI as [5]

$\mathcal{J}(\rho|\{\Pi_j\}) = S(\rho^A) - S(\rho|\{\Pi_j\})$. The value of $\mathcal{J}(\rho|\{\Pi_j\})$ depends on the choice of $\{\Pi_j\}$.

Henderson and Vedral [5] have shown that the maximum of $\mathcal{J}(\rho|\{\Pi_j\})$ with respect to $\{\Pi_j\}$ can be interpreted as a measure of classical correlations. Therefore, the difference between the total correlations $I(\rho)$ and the classical correlations $\mathcal{Q}(\rho) = \sup_{\{\Pi_j\}} \mathcal{J}(\rho|\{\Pi_j\})$ is defined as

$$D(\rho) = I(\rho) - \mathcal{Q}(\rho), \quad (9)$$

giving, finally, a measure of quantum correlations [5] called quantum discord (QD). For pure states QD reduces to entropy of entanglement [17], highlighting that in this case all correlations come from entanglement. However, it is possible to find separable (not-entangled) mixed states with nonzero QD [5, 16], meaning that entanglement does not cause all nonclassical correlations

contained in a composite quantum system. Also, QD can be operationally seen as the difference of work that can be extracted from a heat bath using a bipartite system acting either globally or only locally [18].

Results and discussion. Case I: $B_1 = -B_2$, and ($J > 0$).

In this case, $|\psi^-\rangle$ is the ground state with eigenvalue $E_- = -\sqrt{4B_1^2 + J^2}$. Other eigenvalues are $\{0, 0, \sqrt{4B_1^2 + J^2}\}$ for eigenvectors $\{|00\rangle, |11\rangle, |\psi^+\rangle\}$ respectively.

In this case the variation of QD , CC and EN with B_1 at different temperatures ($T=0.2, 0.9, 1.5$) is depicted in Figs. 1, 2, 3, respectively. We observe that QD and CC in the thermal state coincide for all values of B_1 for different temperatures ($T=0.2, 0.9, 1.5$).

In order to understand this observation, we take a close look at the thermal state. At temperature T , the thermal state is given by

$$\rho = \frac{1}{Z} [|00\rangle\langle 00| + |11\rangle\langle 11| + e^{\sqrt{4B_1^2 + J^2}/T} |\psi^-\rangle\langle\psi^-| + e^{-\sqrt{4B_1^2 + J^2}/T} |\psi^+\rangle\langle\psi^+|]. \quad (10)$$

This ρ has the Bloch representation [21]

$$\rho = \frac{1}{4} [I \otimes I + \sum_{i=1}^3 c_i \sigma_i \otimes \sigma_i], \quad (11)$$

where σ_i ($i = 1, 2, 3$) are the one-qubit Pauli operators. In the appendix to this paper we prove that, the class of mixed states as in Eq.(11) have equal classical and quantum correlations (like bipartite pure states [19]) provided

$$c_i = c_j > c_k$$

and

$$c_k = -c_i^2 \quad (12)$$

where $i \neq j \neq k \in \{1, 2, 3\}$. Here c_i, c_j, c_k are the diagonal elements of the correlation matrix defined by $c_{ij} = \text{Tr}(\rho \sigma_i \otimes \sigma_j)$. It is straightforward to check that the thermal state ρ in Eq.(10) which has form of Eq.(11) satisfies conditions (12). This explains the observations in Figs. 1, 2, 3, that the two qubit thermal state ρ for $B_1 = -B_2$ in Eq.(10) gives rise to equal QD and CC for all values of B_1 and temperature. In order to see why the common curve for QD and CC peaks at $B_1 = 0$ we can maximize the expression for $QD = CC$ with respect to B_1 and check that the maximum occurs at $B_1 = 0$.

From Figs. 1, 2, 3, we also see that EN as a function of B_1 has a peak at $B_1 = 0$ for $T = 0.2$, has a dip for $T = 0.9$ and goes to zero over an interval symmetric about $B_1 = 0$ for $T = 1.5$ [12]. From Eq. (10) we see that the concurrence of the thermal state is governed by




Figure 1 : QD and CC (dashed line) and EN (solid line) as a function of external magnetic field $B_1 = -B_2$ at $T = 0.2$




Figure 2 : QD and CC (dashed line) and EN (solid line) as a function of external magnetic field $B_1 = -B_2$ at $T = 0.9$




Figure 3 : QD and CC (dashed line) and EN (solid line) as a function of external magnetic field $B_1 = -B_2$ at $T = 1.5$

the admixture of the $|\psi^+\rangle$ and $|\psi^-\rangle$ states. We expect concurrence to fall as the state $|\psi^+\rangle$ classically mixes more and more with the ground state $|\psi^-\rangle$. For fixed $J = 1$ and a fixed temperature T , this happens for $B_1 = 0$. That is why we got a dip in the EN curve at $B_1 = 0$. The size of this dip increases with temperature. In fact the dip touches the B_1 axis when, at $B_1 = 0$ the concurrence is zero. To see this we note that the concurrence for ρ in

Eq.(10) is given by [20]

$$C = \frac{2}{Z} \max\{|v| - \sqrt{u_1 u_2}, 0\} \quad (13)$$

where v, u_1, u_2 are given in Eq. (6). Therefore, for $B_1 = 0$ and $J = 1$, $C \geq 0$ provided $\sinh \frac{1}{T} \geq 1$ or $T \leq 1.1346$. For $T = 1.5$ and $J = 1$, using the requirement $\sinh \frac{D}{T} = D$, we can find the range of B_1 around $B_1 = 0$ in which $C = 0$. This is $-1.1456 \leq B_1 \leq 1.1456$. Figs. 1,2,3 confirm the corresponding behaviour of EN .

Fig.4 shows the variation of EN, QD, CC with temperature at fixed values of $B_1 = -B_2$. As expected we have $QD = CC$ for all temperatures. Both EN and QD (CC) curves have plateau at low temperatures corresponding to their ground state values, as at these temperatures, the ground state is not thermally connected to other excited states. Other interesting observation is the vanishing of concurrence at a finite critical temperature T_c which increases with B_1 value, while QD and CC asymptotically go to zero with temperature. The increase in T_c with B_1 [12] can be understood from the thermal state Eq.(10) which says that higher temperatures are required to get a given admixture of $|\psi^-\rangle$ and $|\psi^+\rangle$ for higher B_1 values.




Figure 4 : QD and CC (dashed line) and EN (solid line) as a function of the absolute temperature T for (a) $B_1 = -B_2 = 1$ (b) $B_1 = -B_2 = 2$

We now deal with the case $B_2 = -aB_1$, $a \neq 1$ and positive. $B_1 = 0$ satisfies both, $B_2 = -aB_1$ and $B_2 = -B_1$, so that $QD = CC$ at $B_1 = 0$, for all temperatures T . For

a fixed temperature T , it turns out that $QD > CC$ for $B_1 \neq 0$ if $a > 1$ and $CC > QD$ for $B_1 \neq 0$ if $0 < a < 1$. This is depicted in Fig. 5, for $a = 2$ and $a = 1/2$ for $T = 1.5$. The dominance of QD over CC (or vice versa) varies continuously with a . This observation gives us the key to control the contributions of QD and CC to a two qubit thermal state in Heisenberg model via the continuous variation of the applied magnetic field. The behavior of concurrence in this case can be analyzed in a way similar to the case $B_2 = -B_1$ ($a = 1$). From Fig. 5, we see that for the same temperature, the range over which concurrence vanishes depends on a , this range decreases monotonically with a . Also, the peak position of concurrence (or, EN) on the B_1 axis shifts monotonically towards $B_1 = 0$ as a increases. Thus the main entanglement features can be controlled by varying external magnetic fields.




Figure 5 : QD (dashed line), CC (dash-dotted line) and EN (solid line) as a function of external magnetic field B_1 at $T = 1.5$ (a) $B_2 = -2B_1$ (b) $B_2 = -B_1/2$.

Case II : $B_1 = B_2$.

For uniform external magnetic field $B_1 = B_2$, Figs. 6, 7, 8, show the variation of QD, CC and EN with B_1 for temperatures $T = 0.2, 0.9, 1.5$, respectively. We see that all three quantities are symmetric about $B_1 = 0$ where they have their maxima. Further, $QD > CC$, except at $B_1 = 0$, where $QD = CC$. For higher temperatures, the qualitative behavior of QD and CC remains the same, while EN curve drops down below those of QD and CC . This can be qualitatively understood by looking at the

thermal state given by

$$\rho = \frac{1}{Z} [e^{2B_1/T} |00\rangle\langle 00| + e^{-2B_1/T} |11\rangle\langle 11| + e^{J/T} |\psi^-\rangle\langle \psi^-| + e^{-J/T} |\psi^+\rangle\langle \psi^+|]. \quad (14)$$

For small temperatures, the entanglement of the thermal state is largely dictated by that of $|\psi^-\rangle$ and becomes dominant. At higher temperatures, admixture due to other states reduces the entanglement, so that *QD* and *CC* dominate. Such a complementary behavior of entanglement and discord can serve as a pointer towards a possible connection between them.




Figure 6 : *QD* (dashed line), *CC* (dash-dotted line) and *EN* (solid line) as a function of external magnetic field B_1 where $B_1 = B_2$ at $T = 0.2$




Figure 7 : *QD* (dashed line), *CC* (dash-dotted line) and *EN* (solid line) as a function of external magnetic field B_1 where $B_1 = B_2$ at $T = 0.9$

Fig. 9 shows the variation of *QD*, *CC* and *EN* with temperature for $B_1 = B_2 = 1$ and $B_1 = B_2 = 2$. We see that for high temperatures, *QD* hugely dominates *EN*, showing the robustness of *QD* with temperature. As temperature becomes large *QD* and *CC* converge towards each other. For larger values of B_1 this happens at higher temperatures. As the temperature increases,




Figure 8 : *QD* (dashed line), *CC* (dash-dotted line) and *EN* (solid line) as a function of external magnetic field B_1 where $B_1 = B_2$ at $T = 1.5$

all the coefficients in the thermal mixture Eq.(14) tend to be equal and the thermal state approaches random mixture. Thus it seems that *QD* and *CC* approach each other as an arbitrary thermal state approaches a random mixture. Obviously, for random mixture $\rho = \frac{1}{4}(I \otimes I)$, $QD = CC = 0$. A quantitative analysis of the relative behaviors of *QD* and *CC* with temperature will be very interesting, but possibly have to wait for further developments in the theory.




Figure 9 : *QD* (dashed line), *CC* (dash-dotted line) and *EN* (solid line) as a function of the absolute temperature T for (a) $B_1 = B_2 = 1$ (b) $B_1 = B_2 = 2$




Figure 10 : EN_{AB} (solid line), QD_{AB}^{\leftarrow} (dashed line), EN_{AE} (dot line) and QD_{AE}^{\leftarrow} (dash-dotted line), as a function of external magnetic field $B_1 = -B_2$ at $T = 0.9$

It will be interesting to connect our results with the monogamy relations between the EN and the classical correlation [22] of two subsystems (qubits) and the environment

$$EN_{AB} + CC_{AE}^{\leftarrow} = S_A,$$

$$EN_{AE} + CC_{AB}^{\leftarrow} = S_A, \quad (15)$$

and the relation between EN and QD [23],

$$EN_{AB} + EN_{AE} = QD_{AB}^{\leftarrow} + QD_{AE}^{\leftarrow}, \quad (16)$$

showing us that EN and QD always exist in pairs. Here A, B label the qubits and E stands for the environment. We assume that environment (heat bath) comprises the universe minus the qubits A and B so that the state ρ_{ABE} is a pure state. Since the variation of all the quantities pertaining to the system AB with B_1 and T are obtained from the XX model, we can use Eq.(15,16) to find the corresponding dependence of EN_{AE} and QD_{AE}^{\leftarrow} on B_1 and T . Figs. 10 and 11, (for $B_1 = -B_2$) show the variation of EN_{AE} and QD_{AE}^{\leftarrow} with B_1 and T .

The monogamic relations also help us establish a necessary and sufficient condition for $QD_{AB}^{\leftarrow} = CC_{AB}^{\leftarrow}$ when the environment is present. This is : $QD_{AB}^{\leftarrow} = CC_{AB}^{\leftarrow}$ if and only if $\frac{1}{2}I_{AB} = EN_{AE} + EN_{AB} - QD_{AE}^{\leftarrow}$. To prove the necessity we note that when $QD_{AB}^{\leftarrow} = CC_{AB}^{\leftarrow}$, (that is, $QD_{AB}^{\leftarrow} = \frac{1}{2}I_{AB}$), Eq.(16) can be written as

$$\frac{1}{2}I_{AB} = EN_{AE} + EN_{AB} - QD_{AE}^{\leftarrow}. \quad (17)$$

Now suppose Eq.(17) is true. Then using Eq.(16) we have

$$QD_{AB}^{\leftarrow} = \frac{1}{2}I_{AB}$$

which implies $QD_{AB}^{\leftarrow} = CC_{AB}^{\leftarrow}$. Figs.12 and 13, show the variation of both sides of Eq.(17) with B_1 and T which establishes Eq.(17) for the XX model.




Figure 11 : EN_{AB} (solid line), QD_{AB}^{\leftarrow} (dashed line), EN_{AE} (dot line) and QD_{AE}^{\leftarrow} (dash-dotted line), as a function of the absolute temperature T for $B_1 = -B_2 = 1$.




Figure 12 : $\frac{1}{2}I_{AB}$ (solid line), $EN_{AE} + EN_{AB} - QD_{AE}^{\leftarrow}$ (dashed line), as a function of external magnetic field $B_2 = -B_1$ at $T = 1.5$




Figure 13 : $\frac{1}{2}I_{AB}$ (solid line), $EN_{AE} + EN_{AB} - QD_{AE}^{\leftarrow}$ (dashed line), as a function of the temperature T for $B_2 = -B_1 = -1$.

Summary. In this paper we have studied the variation of QD, CC and EN in two qubit XX Heisenberg chain

as functions of independently varied magnetic fields B_1 and B_2 on each qubit and also with temperature. We deal with two cases $B_1 = -B_2$ (nonuniform field) and $B_1 = B_2$ (uniform field). Our first observation is the complementary behavior of entanglement and QD/CC . For the nonuniform magnetic field, we get the interesting observation that QD and CC are equal for all $B_1 = -B_2$ values as well as all temperatures. Surprisingly, this observation is explained quite simply, using the symmetric form of the thermal state. A very interesting observation is that the relative contributions of QD and CC can be tunably controlled by varying the applied magnetic field. Another interesting finding is that the equality of QD and CC of the subsystem (qubits) imposes a constraint on the distribution of QD and EN over the subsystem and its environment. Further investigation of general Heisenberg models like XXZ along these lines may turn out to be interesting and fruitful.

Appendix. We prove the following statement.

If the quantum state has the Bloch representation [21]

$$\rho = \frac{1}{4} [I \otimes I + \sum_{i=1}^3 c_i \sigma_i \otimes \sigma_i], \quad (A1)$$

and $c_i = c_j > c_k$ and $c_k = -c_i^2$ where $i \neq j \neq k \in \{1, 2, 3\}$, then this state contains the same amount of quantum and classical correlation ($QD = CC$).

Proof: In Ref. [24] S. Luo evaluated analytically the quantum discord for a large family of two-qubit states, which have the maximally mixed marginal and their Bloch representation is

$$\rho = \frac{1}{4} [I \otimes I + \sum_{i=1}^3 c_i \sigma_i \otimes \sigma_i].$$

For this class of quantum states the quantum mutual information is

$$\mathcal{I}(\rho) = \frac{1}{4} [(1 - c_1 - c_2 - c_3) \log_2(1 - c_1 - c_2 - c_3) + (1 - c_1 + c_2 + c_3) \log_2(1 - c_1 + c_2 + c_3) + (1 + c_1 - c_2 + c_3) \log_2(1 + c_1 - c_2 + c_3) + (1 + c_1 + c_2 - c_3) \log_2(1 + c_1 + c_2 - c_3)].$$

We substitute the conditions above in the quantum mutual information. Putting $c = c_1 = c_2 > c_3$ and $c_3 = -c^2$, we get,

$$\mathcal{I}(\rho) = \frac{1}{4} [(1 - 2c + c^2) \log_2(1 - 2c + c^2) + (1 - c^2) \log_2(1 - c^2) + (1 + 2c + c^2) \log_2(1 + 2c + c^2)]$$

After some algebraic simplification, we get

$$\mathcal{I}(\rho) = (1 - c) \log_2(1 - c) + (1 + c) \log_2(1 + c)$$

which equals $2CC$ as in Ref. [24]. It is also easy to check that the above argument goes through when $c = c_1 = c_3 > c_2$ and $c_2 = -c^2$ and when $c = c_3 = c_2 > c_1$ and $c_1 = -c^2$, to get $\mathcal{I}(\rho) = 2CC$. Thus,

$$QD(\rho) = \mathcal{I}(\rho) - CC(\rho) = CC(\rho).$$

Acknowledgments:

ASMH thanks Pune University for hospitality during his visit when this work was carried out. PSJ thanks BCUD Pune University for financial support. We thank Guruprasad Kar and Prof. R. Simon for encouragement. We thank the anonymous referees whose comments have contributed towards the improvement of this paper.

* Electronic address: alisaif73@gmail.com

† Electronic address: behzadlari1979@yahoo.com

‡ Electronic address: pramod@physics.unipune.ac.in

- [1] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge Univ. Press, Cambridge, 2000).
- [2] R. F. Werner, Phys. Rev. A **40**, 4277 (1989); C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996).
- [3] J. S. Bell, Physics **1**, 195 (1964).
- [4] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991); C. H. Bennett *et al.* Phys. Rev. Lett. **70**, 1895 (1993); C. H. Bennett and D. P. DiVincenzo, Nature (London) **404**, 247 (2000).
- [5] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001); L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001).
- [6] E. Knill and R. Laflamme, Phys. Rev. Lett. **81**, 5672 (1998); S. L. Braunstein, C.M. Caves, R. Jozsa, N. Linden, S. Popescu, and R. Schack, Phys. Rev. Lett. **83**, 1054 (1999); D. A. Meyer, Phys. Rev. Lett. **85**, 2014 (2000); A. Datta, S. T. Flammia, and C. M. Caves, Phys. Rev. A **72**, 042316 (2005); A. Datta, and G. Vidal, Phys. Rev. A **75**, 042310 (2007); A. Datta, A. Shaji, and C. M. Caves, Phys. Rev. Lett. **100**, 050502 (2008); S. Luo Phys.Rev. A **77**, 022301 (2008).
- [7] A. S. M. Hassan and P. S. Joag, Quantum Inf. Comput. **8**, 773 (2007).
- [8] K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010).
- [9] P. R. Hammar *et al.*, Phys. Rev. B **59**, 1008 (1999).
- [10] S. Sachdev, Quantum Phase Transitions (Cambridge University Press, Cambridge, 2000); J. A. Hoyos and G. Rigolin, Phys. Rev. A **74**, 062324 (2006); L. F. Santos and G. Rigolin, *ibid.* **71**, 032321 (2005); T. R. de Oliveira *et al.*, *ibid.* **77**, 032325 (2008).
- [11] M. S. Sarandy Phys. Rev. A **80**, 022108 (2009).
- [12] Y. Sun, Y. Chen, and H. Chen, Phys. Rev. A **68**, 044301 (2003).
- [13] T. Werlang and G. Rigolin, Phys. Rev. A **81**, 044101 (2010).
- [14] J. Maziero, H. C. Guzman, L. C. Celeri, M. S. Sarandy, and R. M. Serra, Phys. Rev. A **82**, 012106 (2010).
- [15] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997); W. K. Wootters, *ibid.* **80**, 2245 (1998).
- [16] B. P. Lanyon, M. Barbieri, M. P. Almeida, and A. G. White, Phys. Rev. Lett. **101**, 200501 (2008).
- [17] C. H. Bennett *et al.*, Phys. Rev. A **53**, 2046 (1996).
- [18] W. H. Zurek, Phys. Rev. A **67**, 012320 (2003); R. Dillenschneider, Eurphys. Lett. **88**, 50003 (2009).
- [19] N. Li and S. Luo, Phys. Rev. A **76**, 032327 (2007).

- [20] M. A. O'Conner and W. K. Wootters, Phys. Rev. A **63**, 052302 (2001).
- [21] A. S. M. Hassan and P. S. Joag, Phys. Rev. A **77**, 062334 (2008).
- [22] M. Koashi, A. Winter, Phys. Rev. A **69**, 022309 (2004).
- [23] F. F. Fanchini, M. F. Cornelio, M. c. de Oliveira, A. O. Caldeira, e-print arXiv : 1006.2460v1.
- [24] S. Luo, Phys. Rev. A **77**, 042303 (2008).