

Two New LBV Candidates in the M 33 Galaxy

A. F. Valeev,¹ O. N. Sholukhova,¹ and S. N. Fabrika¹

¹*Special Astrophysical Observatory of the Russian AS, Nizhnij Arkhyz 369167, Russia*

We present two new luminous blue variable (LBV) candidate stars discovered in the M 33 galaxy. We identified these stars (Valeev et al. [1]) as massive star candidates at the final stages of evolution, presumably with a notable interstellar extinction. The candidates were selected from the Massey et al. catalog [2] based on the following criteria: emission in $H\alpha$, $V < 18^m5$ and $0^m35 < (B - V) < 1^m2$. The spectra of both stars reveal a broad and strong $H\alpha$ emission with extended wings (770 and 1000 km s⁻¹). Based on the spectra we estimated the main parameters of the stars. Object N 45901 has a bolometric luminosity $\log(L/L_\odot) = 6.0 - 6.2$ with the value of interstellar extinction $A_V = 2.3 \pm 0.1$. The temperature of the star's photosphere is estimated as $T_\star \sim 13000 - 15000$ K, its probable mass on the Zero Age Main Sequence is $M \sim 60-80 M_\odot$. The infrared excess in N 45901 corresponds to the emission of warm dust with the temperature $T_{\text{warm}} \sim 1000$ K, and amounts to 0.1 % of the bolometric luminosity. A comparison of stellar magnitude estimates from different catalogs points to the probable variability of the object N 45901. Bolometric luminosity of the second object, N 125093, is $\log(L/L_\odot) = 6.3 - 6.6$, the value of interstellar extinction is $A_V = 2.75 \pm 0.15$. We estimate its photosphere's temperature as $T_\star \sim 13000 - 16000$ K, the initial mass as $M \sim 90-120 M_\odot$. The infrared excess in N 125093 amounts to 5-6 % of the bolometric luminosity. Its spectral energy distribution reveals two thermal components with the temperatures $T_{\text{warm}} \sim 1000$ K and $T_{\text{cold}} \sim 480$ K. The [Ca II] $\lambda\lambda 7291, 7323$ lines, observed in LBV-like stars Var A and N 93351 in M 33 are also present in the spectrum of N 125093. These lines indicate relatively recent gas eruptions and dust activity linked with them. High bolometric luminosity of these stars and broad $H\alpha$ emissions allow classifying the studied objects as LBV candidates.

I. INTRODUCTION

Luminous blue variables are the most massive stars at one of the final stages of stellar evolution [3]. A small number of massive stars in the galaxies and a very short scale of the LBV phase ($10^4 - 10^5$ years) makes these objects unique. The study of LBV-type stars is essential for understanding the evolution of stars on the upper

part of the Main Sequence, the formation of WR stars, supernovae, relativistic stars (black holes), for the comprehension of the mechanisms of mass loss, and heavy-element enrichment in galaxies.

It is believed that the stars with more than $40 M_\odot$ undergo the LBV phase [4]. However, the relationship between the LBV stars and other types of massive stars at the final stages of evolution (red supergiants RSG, yellow supergiants

YSG, blue supergiants BSG, B[e]-supergiants, WR stars), and evolutionary transitions between these types of stars are still unclear [5]. LBV stars are closely linked with the late WR stars of the nitrogen sequence with hydrogen in their atmospheres (WNLh). Two cases of transition are known from observations: LBV \rightarrow WNLh (star V 532 in M 33, [6]) and WN3 \rightarrow WN11(LBV) \rightarrow WN4/5 (star HD5980 in the SMC galaxy, [7]).

Observational manifestations of LBV stars are very diverse [8], and the number of known objects of this type is little. It is hence not clear whether the LBV class of objects is homogeneous. For example, why have the S Dor variables [3, 8] not revealed giant eruptions, similar to the η Car and P Cyg stars. It is also difficult to draw specific conclusions about the place of the LBV phase in the evolution of massive stars, and even more so about the effects of binarity and rapid rotation at this stage of evolution.

It is obvious that the above problems can only be solved when a sufficiently large number of objects is thoroughly studied. Observations of more LBV-like stars increases our chances to witness the rare LBV \leftrightarrow WN-type transitions. This will allow testing the modern evolutionary sequences and the dependence of various stages of massive star evolution on galactic metallicity (Z).

Apparently, most of the LBV stars without notable interstellar extinction in our Galaxy are already discovered. However, it is probably still possible to discover a few dozen more such objects in the Galaxy using modern in-

frared surveys [9, 10]. In the M 33 galaxy almost all the LBV-like objects may be detected, as its fortunate orientation (its inclination to the line of sight is 56 deg. [11]), and a relatively close distance (950 Mpc, [12]) allow detailed spectroscopy of its bright stars.

The review [3] attributes five and 15 stars to the confirmed members of the LBV class in our Galaxy and in the galaxies of the Local Group, respectively. In particular, in the M 33 galaxy the authors identified four LBV stars. Clark et al. [13] mention 12 LBV stars and 23 LBV candidates in our Galaxy, while in the M 33 galaxy 37 LBV candidates are known to date [14].

According to the catalog [2], M 33 contains 2304 stellar objects with $V < 18^m5$. If we assume that the mean interstellar extinction of the brightest stars in the galaxy is $A_V \approx 1^m0$ (see, e.g., [15], where this value is estimated as $A_V \approx 0^m95 \pm 0^m05$), then with the distance modulus to M 33 of $(m-M)_0 = 24^m9$ [12], the stars with $V < 18^m5$ and $(B - V) < 0^m35$ will have the luminosity $M_V < -7^m4$ and color $(B - V)_0 \leq 0^m0$. In [1] we made photometry in the $H\alpha$ images of all the stars from the catalog [2] with the above restrictions on color and luminosity, and made a list of stars with an excess in $H\alpha$. These are bright supergiants of the Iab luminosity class and brighter, and the hottest Ib supergiants (with B0 spectra and earlier) [16]. We expected that all the potential LBV candidates will make it into our list. We hence isolated in [1] 185 blue emission objects ($V < 18^m5$ and

$(B - V) < 0^m35$), candidates for massive stars at the final stages of evolution.

Evidently, LBVs and similar objects may well have the extinction $A_V > 1^m0$, for which reason an *additional list* of stars with $V < 18^m5$ and $0^m35 < (B - V) < 1^m2$ with emission in $H\alpha$ was made in [1], containing 25 candidates.

We currently make spectroscopic observations of objects from both our lists [1]. In the first list we discovered an LBV-type star N 93351, which is located in the nuclear region of M 33 [17]. Two objects from our supplementary list (presumably with a notable own reddening) were earlier classified [14] as hot LBV candidates: N 6862 with $B - V = 0^m52$ and N 141751 with $B - V = 0^m77$.

We have recently obtained spectra of 15 candidate stars from the supplementary list, among them we discovered two new LBV candidates. In this paper we present the results of study of these two objects.

II. OBSERVATIONS AND DATA REDUCTION

A. Spectral Data

The spectra of two new LBV candidates, N 45901 and N 125093 were obtained on the 6-m BTA telescope of the Special Astrophysical Observatory of the Russian Academy of Sciences (SAO RAS) with the SCORPIO focal reducer. Table I presents the log of observations. During the observations we used the slit width of $1''$, its ori-

entation is shown in Fig. 1. The quality of the images during the observations was from $1''.2$ to $3''.0$.




Figure 1: Identification maps in the V-band for the objects N 45901 (top) and N 125093 (bottom), the arrows mark their positions. The slit orientation during the observations is shown. North is at the top, west on the left.

The data processing was performed in a standard way. After the spectra were cleaned for traces of cosmic particles, the following reduction procedures were completed: correction for the level of the electronic zero (bias), for flat field, wavelength calibration based on the spectrum of a NeAr lamp, correction for spectral sensitivity of the CCD, and reduction to fluxes in energy units.

Table I: The log of spectral observations

Object	Date	Range (resolution)	Exposure time
N 45901	2010.01.09	3700–7200 Å(12 Å)	2×1200^s
N 125093	2009.10.09	3700–7200 Å(12 Å)	1×600^s
	2009.10.23	3900–5700 Å(5 Å)	2×1200^s
	2009.11.09	5700–7400 Å(5 Å)	4×900^s

B. Photometric Data

We used the M33 images from the Spitzer Space Telescope archive (applicant: R.H. Gehrz), which were obtained on July 9 and August 22, 2004, and January 21 and June 10, 2005 in 4 filters (3.6, 4.5, 5.8 and 8.0 μm) with the infrared IRAC camera [18]. The camera has separate 256×256 pixel CCD chips for each band with a scale of $1''.2$ per pixel. For the photometry we used the images of the entire M33 galaxy, which are the result of co-adding of many small-sized original images into a single image of the galaxy with the simultaneous conversion of fluxes into energy units.

Other Spitzer telescope archive images we used were acquired on December 29, 2003 in the scanning mode with the MIPS instrument in the 24 μm filter [19]. A 128×128 pixel CCD chip with $2''.5$ per pixel is used in this range.

Aperture photometry of the Spitzer images was done using the MOPEX/APEX codes. The aperture correction was determined from single bright stars near the measured object. The fluxes

from the objects are listed in Table II, the flux errors do not exceed 2–3%.

The object N 125093 was not detectable in the MIPS camera images in 70 and 160 μm filters, but it is confidently measured in other bands.

The measurements in the 3.6 and 4.5 μm filters for the object N 45901 should be considered as upper limits because a neighbouring star is in $3''.5$. Its contribution may be significant since the mean FWHM across the field is about 1.8 pixels, and the star is located in less than 3 pixels. The PSF photometry may resolve the stars, but for such faint objects this technique will only allow making a rough estimate. In the 5.8 and 8.0 μm filters both the object and the nearby star get lost in the background noise. The object is not detectable in the 24 μm images.

Photometric measurements in the J (1.25 μm), H (1.65 μm) and K (2.17 μm) filters for N 125093 were adopted from the 2MASS catalog [20]. The images of the M33 galaxy in the survey were acquired in December 1997. To study the spectral energy distribution in N 125093, we used non-simultaneous observations, as we have no reliable information on the variability of this object. The 2MASS catalog has no data on N 45901.

In the Catalog of Variables in M33 [21] N 45901 (this catalog lists it as N 241989) was suspected in variability in one of the three MegaPrime/MegaCam filters of the CFHT telescope (in the SDSS i' band). In addition, it appears that N 45901 coincides with the $H\alpha$ emis-

sion object candidate from the list [22] (its number in this list is N 44). According to [22], the magnitude of this object is $V=18^m45$, whereas in the [2] catalog it is $V=17^m57$. The magnitude difference $\Delta V \approx 0^m9$ between two sets of observations, 1986–1987 [22], and 2000–2001 [2] can hardly be explained by calibration errors. This is why it is likely that the object N 45901 is indeed a variable star.

The magnitude of the object N 125093 in different catalogs (USNO-A2.0, USNO-B1.0, NOMAD Catalog, Guide Star Catalog versions 2.2 and 2.6) varies by 0^m65 in the V-band and by 2^m3 in the B-band. The differences in photometric and photographic systems can naturally lead to a conspicuous variation in the brightness estimates. Nevertheless, the differences are so remarkable that it requires additional observations to verify the possible variability.

III. SPECTRA

The main criterion which we used to relate these two new objects to the class of LBV candidates are their broad and strong emissions in $H\alpha$, their widths, taking into account the spectral resolution (FWHM), are 1000 and 770 km/s for N 125093 and N 45901, respectively. Their equivalent widths are 36\AA and 38\AA . In addition, $H\alpha$ emissions in these stars have significantly more extended wings. These lines are formed in the stellar wind. The second criterion is the stellar luminosity. Based on the spec-

tra of these stars (see below) we can suggest the temperature of the photosphere to be certainly higher than 10000 K. The observed colors of stars ($B - V = 0^m71$ and $B - V = 0^m85$ for N 45901 and N 125093, respectively), as well as the observed diffuse interstellar bands (DIBs) indicate strong interstellar reddening, i.e. high intrinsic luminosity of these candidates.

A. Candidate N 125093

The spectrum of N 125093 is demonstrated in Fig. 2 compared to the spectra of the known LBV stars in M 33, Var A, Var B, V 532 and a new quite firm LBV candidate in this galaxy, N 93351¹[17]. The He I emissions in the spectrum of this star are very weak, but noticeable, the brightest line He I $\lambda\lambda 5876$ has a low intensity. The red region of the spectrum reveals weak Fe II emission lines, and several forbidden [Fe II] lines. There are nebular lines, [O I] $\lambda\lambda 6300, 6360$, and weak [N II] $\lambda\lambda 6548, 6384$ lines in the wings of the broad $H\alpha$ emission.

The [Ca II] $\lambda\lambda 7291, 7323$ lines are quite interesting, they indicate a recent gas eruption and dust activity linked with it [17]. Var A was the brightest star in M 33 in 1950, and later [24, 25] revealed an infrared excess and bright [Ca II] lines. Typically, these lines are not ob-

¹ the spectrum of V 532 was kindly provided by Szeifert [23] and obtained with a resolution of 1.2 \AA . All the remaining spectra were obtained with the SCORPIO [17]. The spectra of Var A, Var B—with a spectral resolution of 12 \AA , while the spectrum of N 93351—with a spectral resolution of 5 \AA .

Table II: The observed parameters of the two new LBV candidates. The first two lines indicate the coordinates of stars. U, B, V, R, I values were adopted from [2], the J, H, K data is from [20], the last five lines list the results of our photometry of the Spitzer images. The errors of this photometry do not exceed 2–3%.

	N 45901	N 125093
Ra (J2000)	01:33:27.40	01:34:15.42
Dec (J2000)	+30:30:29.5	+30:28:16.4
U (mag)	18.509 ± 0.004	18.383 ± 0.004
B (mag)	18.279 ± 0.004	18.138 ± 0.004
V (mag)	17.572 ± 0.004	17.284 ± 0.004
R (mag)	17.193 ± 0.004	16.755 ± 0.004
I (mag)	16.807 ± 0.004	16.203 ± 0.005
J (mag)	–	15.385 ± 0.041
H (mag)	–	14.819 ± 0.062
K (mag)	–	14.120 ± 0.048
$3.6\mu m$ (mJy)	3.17×10^2 ^a	3.11×10^3
$4.5\mu m$ (mJy)	1.45×10^2 ^a	3.69×10^3
$5.8\mu m$ (mJy)	–	4.09×10^3
$8.0\mu m$ (mJy)	–	6.35×10^3
$24.0\mu m$ (mJy)	–	2.00×10^4

^awe consider these measurements as upper limits due to the presence of a nearby star

served in the classical LBV stars, they are visible in the cold hypergiants Var A in M 33 and IRC+10420 (but the star Var A is a typical LBV star based on a number of features, see [17]). Absorption the Fe II lines are visible in the blue region of the spectrum, some Ti II lines (like in the spectrum of N 93351) and two

Si II $\lambda\lambda 6347, 6371$ lines. However, the strongest metal line Fe II $\lambda 5169$ possesses emission components in the wings. In the spectrum of N 125093 diffuse interstellar extinction bands (DIBs) are clearly visible, the Na I $\lambda\lambda 5890, 5896$ doublet is as well sufficiently strong. The $H\beta$ emission in the spectrum of N 125093 is relatively weak and does not have broad wings at the given spectrum quality in the region.

We conducted the Gaussian analysis of the $H\alpha$ emission, and found that a narrow component of this line has FWHM= 5.9 \AA , which is not much different from our spectral resolution (5 \AA), but its broad component has FWHM= 22.5 \AA , which corresponds to the velocity dispersion in the outflowing wind of approximately 1000 km/s (after correcting for the spectral resolution). The narrow component of $H\beta$ line FWHM= 4.4 \AA .

Faint and extended H II regions were captured by the slit. Directly around N 125093 the H II emission is very weak.

To estimate the interstellar extinction, we used the $H\alpha/H\beta$ line flux ratio for the nebula, located near the star N 125093. For the gaseous nebulae this ratio amounts to approximately 2.87 in a wide range of temperatures and densities [26]. We used the law of interstellar reddening from the work of O’Donnell [27] at $R_V=3.07$. We obtained the value $A_V \lesssim 2.5$.

A comparison of the spectrum of N 125093 with the spectra of relatively hot stars Var B and V 532 [17] allows to conclude that the tempera-

ture of the photosphere of N 125093 is certainly lower than 20000 K. On the other hand, based on the presence of deep Si II absorptions we can construe that the temperature is definitely above 10000 K. The equivalent width of the Si II $\lambda 6347$ line is equal to $0.48 \pm 0.02 \text{ \AA}$. In the spectra of supergiants cooler than A0, the equivalent width of this line is equal to 0.7 \AA [28, 29], while in hotter supergiants the equivalent width becomes smaller.

From the value of the equivalent width of the Si II $\lambda 6347$ line, we can conclude that this is a B5–B7 star and its temperature is $> 10000 \text{ K}$. The spectrum of this star is very similar to the spectrum of N 93351, the temperature of which we earlier estimated [17] as 13000–16000 K. Hence we can make a rough estimate of the temperature of N 125093 from its spectral lines, $T \sim 12000 - 16000 \text{ K}$. We will later use these preliminary estimates of the temperature along with that of interstellar extinction in the spectral energy distribution fit of N 125093, where we will specify both the temperature and reddening.

B. Candidate N 45901

The spectrum of the star N 45901 is shown in Fig. 2. Unfortunately, due to weather conditions we were unable to obtain the spectrum of this star with a resolution of 5 \AA , and even the existing spectrum with a resolution of 12 \AA has a low signal-to-noise ratio. Nevertheless, there is

a bright and broad $H\alpha$ emission with extended wings in the spectrum of N 45901. Gaussian analysis revealed a two-component line profile: a bright narrow line and broad wings. The widths of the narrow and broad components, corrected for spectral resolution, are equal to 140 km/s and 770 km/s, respectively. There is a [N II] $\lambda 6584$ line in the broad $H\alpha$ wing. The second line of the [N II] doublet, which should be 3 times weaker than the first, is probably lost in the bright blue $H\alpha$ wing. We also found forbidden lines [O I] $\lambda\lambda 6300, 6360$, which are formed in the unresolved region around the star sized smaller than 4–5 pc (at the distance we adopted, the scale in M 33 is equal to $4.6 \text{ pc}''$). Despite the poor quality of the spectrum, especially in the blue region, the absorption lines Fe II and Ti II that are among the most intensive lines of these ions, are visible in this range. In the red range, we may even suspect the [Fe II] emissions. The region shorter than the $H\alpha$ line reveals signatures of two strongest Fe II emission lines.

Based on the presence of a broad and bright $H\alpha$ emission, as well as the likely occurrence of Fe II and Ti II lines we can make a rough estimate that the temperature of the photosphere of N 45901 is above 10000–12000 K. We did not detect any He I lines, hence its temperature is below 15000 K. Neither did we find any H II regions around the star N 45901, which might be due to the low quality of the spectrum. This is why we shall continue estimating the temperature of the star and the value of interstellar extinction from

Table III: SED modelling results. Two limiting versions of permissible values of interstellar reddening, and the corresponding bolometric luminosities and temperatures of the hot components (stars) in Kelvins are listed. The last two columns indicate the temperatures of warm and cold dust components, and their percentage contribution to the bolometric luminosity in the parentheses.

Object	A_V	$\log L/L_\odot$	T_*	T_{warm}	T_{cold}
N 45901	2.2	6.02	13200	1000(0.1)	
	2.4	6.21	15200	1000(0.1)	
N 125093	2.6	6.30	12700	1400(3.9)	480(3.2)
	2.9	6.60	16200	1400(2.0)	470(1.6)

its spectral energy distribution without preliminary evaluations of reddening.

IV. SPECTRAL ENERGY DISTRIBUTIONS AND STELLAR PARAMETERS

Before refining the temperatures and luminosities of these stars in their energy distributions, we estimate their interstellar reddening. It follows from the spectra that the temperatures of the photospheres of these stars are above 10000 K. For the supergiants this temperature corresponds to $(B - V)_0 = 0^m0$ at the bolometric correction $BC = -0^m3$ [30]. The observed colors of the stars are equal to $(B - V) = 0^m71$ for N 45901 and $(B - V) = 0^m85$ for N 125093. Therefore, their values of interstellar reddening are not below $A_V = 2.2$ and 2.6, respec-

tively. At the distance modulus to M 33 we adopted as equal to $(m - M)_0 = 24^m9$ [12], we find the lower limit of N 45901 luminosity as $M_V = -9^m5$, $M_{bol} = -9^m8$ and $M_V = -9^m9$, $M_{bol} = -10^m2$ for N 125093.

To construct the spectral energy distributions (SED), we used the data of optical and infrared photometry from Table II. We applied a black-body approximation for individual components: a hot star and one or two components of dust emission. The upper limit of the temperature of the warm dust component was set at 1500 K, since above this temperature the dust begins to evaporate.

Without the a priori information on temperature and reddening it is not possible to determine these two parameters independently. For the object N 125093, we have both a preliminary estimate of the stellar temperature, and the reddening (13000–16000 K, $A_V \lesssim 2.5$). For N 45901, we only have an evaluation of the stellar photosphere temperature (12000–15000 K). Indeed, the spectra of both our stars are very similar (Fig.2), hence, their temperatures should not differ greatly either.

The SEDs are demonstrated in Fig. 3. The radiation intensity drop in the U-band is associated with the presence of the Balmer jump in these stars. The magnitude of fading in the U-band is about the same for both stars. This independently confirms our conclusion that the temperatures of both stars are almost identical. We have not used the measurements obtained in the U-




Figure 2: The spectra of LBV candidates N 45901 and N 125093 along with the spectra of the known LBV in the M33 galaxy, Var A (an LBV or a red hypergiant) and N 93351 (a firm LBV candidate) (a, c), Var B and V 532 (b). The spectra of both new LBV candidates reveal broad $H\alpha$ emission lines. The main spectral lines detected in the spectra of candidates are marked.

band at the fits. We assume that the photometric measurements in the R, I (and, possibly, J-bands) may be distorted by an additional contribution of the free-free wind radiation [17]. However, the result of approximations shows that the contribution of this emission in the two stars is negligible.

The luminosity of each SED component was calculated as the area under the corresponding blackbody spectrum, and the bolometric luminosity of the object was deduced as the sum of luminosities of the components. The best solutions for these stars are listed in Table III. The photometric data (not corrected for reddening) are marked with crosses in Fig. 3. The measurements corrected for reddening ($A_V = 2.8$ for N 125093, and $A_V = 2.3$ for N 45901) are indicated by circles. The errors represent the accuracy of photometry. If the size of the circle is larger than the measurement error, the error bars are not visible. The solid lines represent individual blackbody components (three for N 125093, and two for N 45901), as well as the sum of all components.

For N 125093 we find the optimal value of the interstellar extinction as $A_V = 2.75 \pm 0.15$. It is close to the assessment we obtained from the spectra of the surrounding H II emission ($A_V \lesssim 2.5$). The stellar parameters obtained for the limiting values of $A_V=2.6$ and $A_V=2.9$ are listed in Table III. The temperature of the stellar photosphere is $T_\star \sim 13000 - 16000$ K, which is consistent with our estimate from the spec-

trum. The corresponding bolometric luminosity of N 125093 amounts to $\log(L/L_\odot) = 6.3 - 6.6$. We find two thermal components in the spectrum, a warm component with $T_{\text{warm}} \sim 1000$ K, and a cold component with $T_{\text{cold}} \sim 480$ K. The infrared excess in N 125093 is 5–6 % of the bolometric luminosity.

In the case of N 45901 we deduced the optimal estimate of interstellar extinction at $A_V = 2.3 \pm 0.1$. Accordingly, the temperature of the photosphere of N 45901 is estimated as $T_\star \sim 13000\text{--}15000$ K, and its bolometric luminosity amounts to $\log(L/L_\odot) = 6.0 - 6.2$. Infrared excess in N 45901 corresponds to the warm dust radiation with the temperature of $T_{\text{warm}} \sim 1000$ K and amounts to 0.1 % of the bolometric luminosity.

Figure 4 demonstrates the temperature–luminosity diagram for all the LBV stars known to date (Var B, Var C, Var 2, Var 83, V 532), as well as the stars Var A and N 93351 in the M 33 galaxy, the parameters of which were determined in our paper [17]. We show there our two new candidates N 45901 and N 125093, and the evolutionary tracks [31], computed for the stars with metallicity 0.007, corresponding to M 33. The temperatures and luminosities of these stars were deduced by a unified method of approximation of the SEDs engaging additional data on the temperatures of stars and interstellar extinction values, obtained from the spectra. For the majority of stars the boundary values of temperature are shown, which have met the approximation




Figure 3: SEDs of two new LBV candidates in M33. The results of photometry not corrected for reddening are marked with crosses. The corrected measurements (for N 45901 $A_V = 2.3$ and for N 125093 $A_V = 2.8$) are indicated by circles. The errors represent the accuracy of photometry. If the size of the circle is larger than the measurement error, the error bars are not visible. The solid lines represent individual blackbody components of radiation, as well as the sum of all components.

of spectral energy distributions. In the case of stars Var 83, Var A, and N 93351, the parameters of the two boundary values of interstellar extinction are indicated, hence the range of valid parameters (L,T) forms a closed region.

From this diagram, we find the mass estimates for the stars on the ZAMS. The masses of N 45901 and N 125093 are estimated as approximately $60 - 80M_{\odot}$, and $90 - 120M_{\odot}$, respectively.

V. CONCLUSION

Luminous blue variables are characterized by a strong and diverse brightness variability [8]. A large amplitude variability (over 1^m) is rare in the classical LBVs, such events may take place on the time scales of dozens of years. For example, an LBV prototype P Cyg, having demonstrated a giant eruption, possesses a relatively stable brightness over the past three centuries. In fact, any new object candidate for the LBV will remain only an “LBV candidate” until its brightness variability reaches a large amplitude (presumably not less than 0.3–0.5 magnitude, as the amplitudes of 0.1–0.2 magnitude is common for massive supergiants). Tens and hundreds of years may pass while we wait for a strong manifestation of brightness variability. Regardless of the spectrum, temperature, or luminosity, irrespective of which “typical LBV” features are found in one or another LBV candidate, such an object will remain only a candidate.

We hope that this state of physics of LBV objects will not persist long. There will be found more reliable criteria and classifications of the LBV class and this stage of massive star evolution in general. The chemical composition of the atmosphere, i.e. the relative hydrogen and helium abundance is likely to be such a reliable criterion. The measurements of abundances require spectral analysis based on the models of expanding atmospheres. Indeed, the LBV star well-known in our Galaxy, AG Car has the abun-




Figure 4: The temperature–luminosity diagram for all the LBV stars known to date (Var B, Var C, Var 2, Var 83, V 532), as well as the stars Var A and N 93351 from the M 33 galaxy (data adopted from [17]), with two new candidates N 45901 and N 125093 imposed. The evolutionary tracks, computed for metallicity $Z=0.007$ are adopted from [31].

dance $\text{He}/\text{H}=0.43$, which was deduced via detailed modelling of its spectrum [32]. A relation between the LBV stars and late WR stars of the nitrogen sequence with hydrogen in the atmospheres (WNLh, see the Introduction), which reveal a low abundance of hydrogen in the wind is well-known. It is obvious that the understanding of physics and evolutionary status of the LBV objects requires detecting large numbers of objects

bearing properties similar to LBV.

This paper presents a preliminary study of two new LBV candidates in the M 33 galaxy, which were discovered by us in the spectral observations of a list of emission object candidates with a presumably notable interstellar reddening [1]. The spectra of N 45901 and N 125093 are similar to the spectra of the stars at the LBV stage. They have strong and broad $H\alpha$, emis-

sions, forbidden [O I] and [N II] lines. The spectrum of the second star reveals [Ca II], [Fe II] and Fe II emission lines, Ti II and Fe II absorptions. The spectrum of the first star has an insufficient quality for a reliable identification, however the [O I], Fe II and Ti II lines are present in its spectrum. Additional spectra with a better resolution and signal-to-noise ratio are required to accurately examine the spectrum. Comparisons of brightness estimates from different catalogs indicate a probable variability of the object N 45901.

We found an infrared excess in both stars. The temperatures of warm and cold dust components in the star N 125093 (it possesses the [Ca II] emissions), amount to 1400 K and 470 K, respectively.

We estimate that the star N 45901 has a bolo-

metric luminosity of $\log(L/L_{\odot}) = 6.0 - 6.2$ and its probable mass on the initial main sequence is $M \sim 60 - 80 M_{\odot}$. The luminosity of N 125093 amounts to $\log(L/L_{\odot}) = 6.3 - 6.6$ and its initial mass is $M \sim 90 - 120 M_{\odot}$. All the above properties of N 45901 and N 12509 allow us to classify them as LBV candidates.

Acknowledgments

The authors thank T.A. Fatkhullin for help in observations, and E.L. Chentsov for the useful discussions of spectral classifications. This paper was supported by the grants of the Russian Foundation for Basic Research (№ 09-02-00163 and 10-02-00463), and the grant “Leading Scientific Schools of Russia” № 5473.2010.2.

-
- [1] A. F. Valeev, O. Sholukhova, and S. Fabrika, *Astrophysical Bulletin* **65**, 140 (2010).
- [2] P. Massey, K. A. G. Olsen, P. W. Hodge, et al., *Astronom. J.* **131**, 2478 (2006).
- [3] R. M. Humphreys and K. Davidson, *Publ. Astronom. Soc. Pacific* **106**, 1025 (1994).
- [4] G. Meynet, P. Eggenberger, and A. Maeder, *IAU Symposium* **241**, 13 (2006).
- [5] N. Smith and P. S. Conti, *Astrophys. J.* **679**, 1467 (2008).
- [6] S. Fabrika, O. Sholukhova, T. Becker, et al., *Astronom. and Astrophys.* **437**, 217 (2005).
- [7] G. Koenigsberger, L. Georgiev, D. J. Hillier, et al., *Astronom. J.* **139**, 2600 (2010).
- [8] A. M. van Genderen, *Astronom. and Astrophys.* **366**, 508 (2001).
- [9] V. V. Gvaramadze, A. Y. Kniazev, and S. Fabrika, *Monthly Notices Roy. Astronom. Soc.* **405**, 1047 (2010).
- [10] V. V. Gvaramadze, A. Y. Kniazev, S. Fabrika, et al., *Monthly Notices Roy. Astronom. Soc.* **405**, 520 (2010).
- [11] D. Zaritsky, R. Elston, and J. M. Hill, *Astronom. J.* **97**, 97 (1989).
- [12] A. Z. Bonanos, K. Z. Stanek, R. P. Kudritzki, et al., *Astrophys. J.* **652**, 313 (2006).
- [13] J. S. Clark, V. M. Larionov, and A. Arkharov, *Astronom. and Astrophys.* **435**, 239 (2005).
- [14] P. Massey, R. T. McNeill, K. A. G. Olsen, et al., *Astronom. J.* **134**, 2474 (2007).
- [15] S. Fabrika and O. Sholukhova, *Astrophys. J. Suppl.* **140**, 309 (1999).

- [16] K. U. Allen, *Astrofizicheskie velichini* (Mir, Moscow, 1977) [in Russian].
- [17] A. F. Valeev, O. Sholukhova, and S. Fabrika, Monthly Notices Roy. Astronom. Soc. **396**, L21 (2009).
- [18] G. G. Fazio, J. L. Hora, L. E. Allen, et al., Astrophys. J. Suppl. **154**, 10 (2004).
- [19] G. H. Rieke, E. T. Young, C. W. Engelbracht, et al., Astrophys. J. Suppl. **154**, 25 (2004).
- [20] R. M. Cutri, M. F. Skrutskie, S. van Dyk, et al., *2MASS All Sky Catalog of point sources*. (2003).
- [21] J. D. Hartman, D. Bersier, K. Z. Stanek, et al., Monthly Notices Roy. Astronom. Soc. **371**, 1405 (2006).
- [22] D. Calzetti, A. L. Kinney, H. Ford, et al., Astronom. J. **110**, 2739 (1995).
- [23] T. Szeifert, O. Stahl, B. Wolf, et al., *Astronomische Gesellschaft Abstract Series* **10**, 36 (1994).
- [24] R. M. Humphreys, T. J. Jones, and R. D. Gehrz, Astronom. J. **94**, 315 (1987).
- [25] R. M. Humphreys, T. J. Jones, E. Polomski, et al., Astronom. J. **131**, 2105 (2006).
- [26] D. E. Osterbrock and G. J. Ferland, *Astrophysics of gaseous nebulae and active galactic nuclei* (Univ. Sci. Books, Sausalite, 2006).
- [27] J. E. O'Donnell, Astrophys. J. **422**, 158 (1994).
- [28] E. L. Chentsov and A. N. Sarkisyan, Astrophysical Bulletin **62**, 257 (2007).
- [29] P. Didelon, Astrophys. J. Suppl. **50**, 199 (1982).
- [30] P. J. Flower, Astrophys. J. **469**, 355 (1996).
- [31] A. Claret, Astronom. and Astrophys. **453**, 769 (2006).
- [32] J. H. Groh, D. J. Hillier, A. Damineli, et al., Astrophys. J. **698**, 1698 (2009).