

On multidimensional analogs of Melvin's solution for classical series of Lie algebras

A. A. Golubtsova^{1,b} and V. D. Ivashchuk^{2,a,b}

- (a) *Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya Str., Moscow 119361, Russia*
- (b) *Institute of Gravitation and Cosmology, Peoples' Friendship University of Russia, 6 Miklukho-Maklaya Str., Moscow 117198, Russia*

Abstract

A multidimensional generalization of Melvin's solution for an arbitrary simple Lie algebra \mathcal{G} is presented. The gravitational model contains n 2-forms and $l \geq n$ scalar fields, where n is the rank of \mathcal{G} . The solution is governed by a set of n functions obeying n ordinary differential equations with certain boundary conditions. It was conjectured earlier that these functions should be polynomials (the so-called fluxbrane polynomials). A program (in Maple) for calculating of these polynomials for classical series of Lie algebras is suggested (see Appendix). The polynomials corresponding to the Lie algebra D_4 are obtained. It is conjectured that the polynomials for A_n -, B_n - and C_n -series may be obtained from polynomials for D_{n+1} -series by using certain reduction formulas.

1 Introduction

In this paper we deal with a special multidimensional generalization of the well-known Melvin solution [1]. This generalized solution is related a simple Lie algebra and is a special case of the so-called generalized fluxbrane solutions from [2]. For fluxbrane solutions see [2], [3]-[8] and references therein. For more general classes of solutions see also [9, 10].

We remind the reader that the generalized fluxbrane solutions are governed by functions $H_s(z) > 0$ defined on the interval $(0, +\infty)$ and obeying the non-linear differential equations

$$\frac{d}{dz} \left(\frac{z}{H_s} \frac{d}{dz} H_s \right) = P_s \prod_{s'=1}^n H_{s'}^{-A_{ss'}}, \quad (1.1)$$

with the following boundary conditions:

$$H_s(+0) = 1, \quad (1.2)$$

$s = 1, \dots, n$, where $P_s > 0$ for all s . Parameters P_s are proportional to brane charge density squared Q_s^2 and $z = \rho^2$, where ρ is a radial parameter. The boundary condition (1.2) guarantees the absence of singularity (in the metric) for $\rho = +0$.

Here we assume that $(A_{ss'})$ is a Cartan matrix for some finite dimensional simple Lie algebra \mathcal{G} of rank n ($A_{ss} = 2$ for all s). According to a conjecture suggested in [2], the solutions to Eqs.

¹siedhe@gmail.com

¹ivashchuk@mail.ru

(1.1), (1.2) governed by the Cartan matrix $(A_{ss'})$ are polynomials: $H_s(z) = 1 + \sum_{k=1}^{n_s} P_s^{(k)} z^k$, where $P_s^{(k)}$ are constants ($P_s^{(1)} = P_s$). Here $P_s^{(n_s)} \neq 0$ and $n_s = 2 \sum_{s'=1}^n A^{ss'}$ where $(A^{ss'}) = (A_{ss'})^{-1}$. Integers n_s are components of a twice dual Weyl vector in the basis of simple co-roots [11]. It was pointed in [2] that the conjecture on polynomial structure of H_s may be proved for A_n and C_n Lie algebras along a line as it was done for black-brane polynomials from [12] (see also [10]). It should be also noted that the set of polynomials H_s defines a special solution to open Toda chain equations [13, 14] corresponding to simple Lie algebra \mathcal{G} .

2 The solution

We consider a model governed by the action

$$S = \int d^D x \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \frac{1}{2} \sum_{s=1}^n \exp[2\lambda_s(\varphi)] (F^s)^2 \right\} \quad (2.1)$$

where $g = g_{MN}(x) dx^M \otimes dx^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbb{N}$), $F^s = dA^s = \frac{1}{2} F_{MN}^s dz^M \wedge dz^N$ is a 2-form, λ_s is a 1-form on \mathbb{R}^l : $\lambda_s(\varphi) = \lambda_{s\alpha} \varphi^\alpha$, $s = 1, \dots, n$; $\alpha = 1, \dots, l$. In (2.1), we denote $|g| = |\det(g_{MN})|$, $(F^s)^2 = F_{M_1 M_2}^s F_{N_1 N_2}^s g^{M_1 N_1} g^{M_2 N_2}$, $s = 1, \dots, n$.

Let us consider a family of exact solutions to the field equations corresponding to the action (2.1) and depending on one variable ρ . These solutions are defined on the manifold

$$M = (0, +\infty) \times M_1 \times M_2, \quad (2.2)$$

where M_1 is a one-dimensional manifold (say S^1 or \mathbb{R}) and M_2 is a (D-2)-dimensional Ricci-flat manifold. The solution reads

$$g = \left(\prod_{s=1}^n H_s^{2h_s/(D-2)} \right) \left\{ w d\rho \otimes d\rho + \left(\prod_{s=1}^n H_s^{-2h_s} \right) \rho^2 d\phi \otimes d\phi + g^2 \right\}, \quad (2.3)$$

$$\exp(\varphi^\alpha) = \prod_{s=1}^n H_s^{h_s \lambda_s^\alpha}, \quad (2.4)$$

$$F^s = -Q_s \left(\prod_{s'=1}^n H_{s'}^{-A_{ss'}} \right) \rho d\rho \wedge d\phi, \quad (2.5)$$

$s = 1, \dots, n$, where $w = \pm 1$, $g^1 = d\phi \otimes d\phi$ is a metric on M_1 and g^2 is a Ricci-flat metric on M_2 .

The functions $H_s(z) > 0$, $z = \rho^2$, obey the equations (1.1) with the boundary conditions (1.2) and

$$P_s = \frac{1}{4} K_s Q_s^2. \quad (2.6)$$

The parameters h_s satisfy the relations

$$h_s = K_s^{-1}, \quad K_s = B_{ss} > 0, \quad (2.7)$$

where

$$B_{ss'} \equiv 1 + \frac{1}{2-D} + \lambda_{s\alpha} \lambda_{s'\beta} h^{\alpha\beta}, \quad (2.8)$$

$s, s' = 1, \dots, n$, with $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$. Here $\lambda_s^\alpha = h^{\alpha\beta} \lambda_{s\beta}$ and

$$(A_{ss'}) = (2B_{ss'}/B_{s's'}) \quad (2.9)$$

is the Cartan matrix for a simple Lie algebra \mathcal{G} of rank n .

It may be shown that if the matrix $(h_{\alpha\beta})$ has an Euclidean signature and $l \geq n$, there exists a set of co-vectors $\lambda_1, \dots, \lambda_n$ obeying (2.9). Thus the solution is valid at least when $l \geq n$ and the matrix $(h_{\alpha\beta})$ is a positive-definite.

The solution under consideration may be verified just by substitution into the equations of motion corresponding to (2.1). It may be also obtained as a special case of the fluxbrane (for $w = +1$, $M_1 = S^1$) and S -brane ($w = -1$) solutions from [2] and [8], respectively.

If $w = +1$ and the (Ricci-flat) metric g_2 has a pseudo-Euclidean signature, we get a multidimensional generalization of the Melvin's solution [1]. Recall that the Melvin's solution corresponds to $n = 1$, $M_1 = S^1$ ($0 < \phi < 2\pi$), $M_2 = \mathbb{R}^2$, $g_2 = -dt \otimes dt + d\xi \otimes d\xi$ and $\mathcal{G} = A_1$. For $w = -1$ and g_2 of Euclidean signature we obtain a cosmological solution with a horizon (as $\rho = +0$) if $M_1 = \mathbb{R}$ ($-\infty < \phi < +\infty$).

3 Fluxbrane polynomials

Here we present polynomials corresponding to the Lie algebra $D_4 = \text{so}(8)$. These polynomials were obtained using a program written in Maple. The program is given in the Appendix (it was described in [15]).

For the Lie algebra D_4 we find the following set of polynomials

$$H_1 = 1 + P_1 z + \frac{1}{4} P_1 P_2 z^2 + \left(\frac{1}{36} P_1 P_2 P_3 + \frac{1}{36} P_1 P_2 P_4 \right) z^3 + \frac{1}{144} P_1 P_2 P_3 P_4 z^4 \quad (3.1)$$

$$+ \frac{1}{3600} P_1 P_2^2 P_3 P_4 z^5 + \frac{1}{129600} P_1^2 P_2^2 P_3 P_4 z^6,$$

$$H_2 = 1 + P_2 z + \left(\frac{1}{4} P_1 P_2 + \frac{1}{4} P_2 P_3 + \frac{1}{4} P_2 P_4 \right) z^2 + \left(\frac{1}{9} P_1 P_2 P_3 + \frac{1}{9} P_1 P_2 P_4 \right. \quad (3.2)$$

$$\left. + \frac{1}{9} P_2 P_3 P_4 \right) z^3 + \left(\frac{1}{144} P_1 P_2^2 P_3 + \frac{1}{144} P_1 P_2^2 P_4 + \frac{1}{144} P_2^2 P_3 P_4 + \frac{1}{16} P_1 P_2 P_3 P_4 \right) z^4$$

$$+ \frac{7}{600} P_1 P_2^2 P_3 P_4 z^5 + \left(\frac{1}{1600} P_1 P_2^3 P_3 P_4 + \frac{1}{5184} P_1 P_2^2 P_3^2 P_4 + \frac{1}{5184} P_1^2 P_2^2 P_3 P_4 \right.$$

$$\left. + \frac{1}{5184} P_1 P_2^2 P_3 P_4^2 \right) z^6 + \left(\frac{1}{32400} P_1^2 P_2^3 P_3 P_4 + \frac{1}{32400} P_1 P_2^3 P_3 P_4^2 + \frac{1}{32400} P_1 P_2^3 P_3^2 P_4 \right) z^7$$

$$+ \left(\frac{1}{518400} P_1^2 P_2^3 P_3 P_4^2 + \frac{1}{518400} P_1^2 P_2^3 P_3^2 P_4 + \frac{1}{518400} P_1 P_2^3 P_3^2 P_4^2 \right) z^8$$

$$+ \frac{1}{4665600} P_1^2 P_2^3 P_3^2 P_4^2 z^9 + \frac{1}{46656000} P_1^2 P_2^4 P_3^2 P_4^2 z^{10},$$

$$H_3 = 1 + P_3 z + \frac{1}{4} P_2 P_3 z^2 + \left(\frac{1}{36} P_1 P_2 P_3 + \frac{1}{36} P_2 P_3 P_4 \right) z^3 + \frac{1}{144} P_1 P_2 P_3 P_4 z^4 \quad (3.3)$$

$$+ \frac{1}{3600} P_1 P_2^2 P_3 P_4 z^5 + \frac{1}{129600} P_1 P_2^2 P_3^2 P_4 z^6,$$

$$H_4 = 1 + P_4 z + \frac{1}{4} P_2 P_4 z^2 + \left(\frac{1}{36} P_1 P_2 P_4 + \frac{1}{36} P_2 P_3 P_4 \right) z^3 + \frac{1}{144} P_1 P_2 P_3 P_4 z^4 \quad (3.4)$$

$$+ \frac{1}{3600} P_1 P_2^2 P_3 P_4 z^5 + \frac{1}{129600} P_1 P_2^2 P_3 P_4^2 z^6.$$

Setting $P_4 = +0$, we get a triple of polynomials (H_1, H_2, H_3) for the Lie algebras $A_3 = \mathfrak{sl}(4)$. For $P_3 = P_4$ we obtain a set of polynomials (H_1, H_2, H_3) for the Lie algebras $B_3 = \mathfrak{so}(7)$. The A_3 -polynomials for $P_1 = P_3$ give us a pair of $C_2 = \mathfrak{sp}(2)$ -polynomials (H_1, H_2) [8]. (For $P_1 = P_3 = P_4$ we obtain G_2 -polynomials from [8].)

This prescription could be generalized to higher ranks by using the following **Conjecture**: i) the set of A_n -polynomials is given by the first n polynomials for the Lie algebra D_{n+1} when $P_{n+1} = +0$; ii) the set of B_n -polynomials coincides with the first n polynomials for the Lie algebra D_{n+1} when $P_n = P_{n+1}$; iii) the set of C_{m+1} -polynomials coincides with the first $m+1$ polynomials for the Lie algebra A_{2m+1} when the following relations are imposed: $P_1 = P_{2m+1}$, $P_2 = P_{2m}$, ..., $P_m = P_{m+2}$. An analytical proof of this conjecture will be a subject of a separate work. At the moment, one can verify these relations just by using the program from the Appendix.

4 Conclusions

We have presented a multidimensional generalization of the Melvin's solution for an arbitrary simple Lie algebra \mathcal{G} . The solution is governed by a set of n fluxbrane polynomials. We have written a program for calculating of these polynomials for the classical series of Lie algebras (see the Appendix). The set of polynomials corresponding to the Lie algebra D_4 is obtained. The polynomials considered above define special solutions to open Toda chain equations corresponding to simple Lie algebras that may be of interest for certain applications of Toda chains.

We have conjectured (without proof) certain relations between polynomials belonging to different series of classical Lie algebras. These relations tells us that the most important is the calculation of D_{n+1} -polynomials, since all other polynomials (e.g. A_n -, B_n - and C_n -ones) may be obtained from the polynomials for D_{n+1} -series by using certain reduction formulas. A calculation of polynomials for exceptional Lie algebras (i.e. G_2 , F_4 , E_6 , E_7 , E_8) will be described in a separate publication.

As it was noted, the solution under consideration for $w = -1$ is a cosmological one. The study of this and/or some other similar cosmological solution governed by fluxbrane polynomials (e.g. in connection with the problem of acceleration) will be a subject of a forthcoming publications.

Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research grant Nr. 07 – 02 – 13624 – *ofits* and by a grant of People Friendship University (NPK MU).

Appendix

In this Appendix we present a program written for calculating fluxbrane polynomials for four classical series of simple Lie algebras: A_n , B_n , C_n and D_n . A description of this program is given in [15].

```
> with(LinearAlgebra):
> with(PolynomialTools):
> S:=3:
> A:=Matrix(S,S):
> AlgLie:=proc(algn,S,CartA:=Matrix(S,S))
  local i,mu,nu;
  i := 0; mu := 0; nu:=0;
  mu := S-1;
  nu:=S-2;
  for i to S do
```

```

    CartA[i, i] := 2
end do;
if (S>=1) and algn=an then
  for i from 1 to S-1 do
    CartA[i, i+1] := -1
  end do;
  for i from 1 to S-1 do
    CartA[i+1, i] := CartA[i, i+1]
  end do;
end if;
if (S >= 4) and algn = dn then
  for i from 1 to S-2 do
    CartA[i, i+1] := -1
  end do;
  for i from 1 to S-2 do
    CartA[i+1, i] := CartA[i, i+1]
  end do;
  CartA[S, nu] := CartA[mu, nu];
  CartA[nu, S] := CartA[S, nu]
end if;
if (S>=2) and algn=cn then
  for i from 1 to (S - 1) do
    CartA[i,i+1]:=-1
  end do;
  for i from 1 to (S - 2) do
    CartA[i+1,i]:=CartA[i,i+1]
  end do;
  CartA[S,mu]:=-2
end if;
if (S>=3) and algn=bn then
  for i from 1 to (S - 1) do
    CartA[i+1,i]:=-1
  end do;
  for i from 1 to (S - 2) do
    CartA[i,i+1]:=CartA[i+1,i]
  end do;
end if;
return CartA
end proc;
> AlgLie(algn, S, A);
> n := Vector[row](1 .. S):
> A1 := MatrixInverse(A);
> for i to S do n[i] := 2*add(A1[i, j], j = 1 .. S) end do:
> maxel := max(convert(n, list)[]):
> P := array(1 .. S, 1 .. maxel):
> H := Vector[row](1 .. S):
> for i to S do H[i] := 1+add(P[i, k]*z^k, k = 1 .. n[i]) end do:
> for i to S do for j to S do a[i, j] := A[i, j] end do end do:
> for i to S do h[i] := H[i] end do:
> for i to S do for v to S do c[i, v] := h[v]^(-a[i, v]) end do end do:
> equal := Vector[row](1 .. S):
> for i to S do
  equal[i]:= diff(z*(diff(H[i],z))/H[i],z)-P[i,1]*(product(c[i,m], m = 1..S))
end do:
> simequal := Vector[row](1 .. S):
> newequal := Vector[row](1 .. S):
> for i to S do simequal[i] := simplify(combine(value(equal[i]), power)) end do:
> for i to S do newequal[i] := numer(simequal[i]) end do:
> maxcoeff := Vector[row](1 .. S):
> for i to S do maxcoeff[i] := degree(newequal[i], z) end do:
> coefflist := table():
> for i to S do
  for c from 0 to maxcoeff[i] do
    coefflist[i, c] := coeff(newequal[i], z, c) = 0
  end do
end do:
> Sys := convert(coefflist, list):
> sol := solve(Sys):

```

```

> trans := {seq(seq(P[i,j] = P[i,j], i = 1..S), j = 1..maxel)}:
> sol := simplify(map2(subs, trans, sol)):
> P1 := map2(subs, sol, evalm(P)):
> for i to S do H[i]:= 1+add(P1[i,k]*z^k, k = 1..n[i]) end do;

```

References

- [1] M.A. Melvin, Pure magnetic and electric geons, *Phys. Lett.* **8**, 65 (1964).
- [2] V.D. Ivashchuk, Composite fluxbranes with general intersections, *Class. Quantum Grav.*, **19**, 3033-3048 (2002); hep-th/0202022.
- [3] D.V. Gal'tsov and O.A. Rytchkov, Generating branes via sigma models, *Phys. Rev. D* **58**, 122001 (1998); hep-th/9801180.
- [4] C.-M. Chen, D.V. Gal'tsov and S.A. Sharakin, Intersecting M -fluxbranes, *Grav. Cosmol.* **5**, No 1 (17), 45-48 (1999); hep-th/9908132.
- [5] M.S. Costa and M. Gutperle, The Kaluza-Klein Melvin solution in M-theory, *JHEP* **0103**, 027 (2001); hep-th/0012072.
- [6] M. Gutperle and A. Strominger, Fluxbranes in string theory, *JHEP* **0106**, 035 (2001); hep-th/0104136.
- [7] C.M. Chen, D.V. Gal'tsov and P.M. Saffin, Supergravity fluxbranes in various dimensions, *Phys. Rev. D* **65**, 084004 (2002); hep-th/0110164.
- [8] I.S. Goncharenko, V. D. Ivashchuk and V.N. Melnikov, Fluxbrane and S-brane solutions with polynomials related to rank-2 Lie algebras, *Grav. Cosmol.* **13**, No. 4 (52), 262-266 (2007); math-ph/0612079.
- [9] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.* **41**, 444-460 (2000); hep-th/9907019.
- [10] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.*, **18** R87-R152 (2001); hep-th/0110274.
- [11] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and Representations. A graduate course for physicists (Cambridge University Press, Cambridge, 1997).
- [12] V.D. Ivashchuk and V.N. Melnikov, P-brane black Holes for General Intersections, *Grav. Cosmol.*, **5**, No 4 (20), 313-318 (1999); gr-qc/0002085.
- [13] B. Kostant, *Adv. in Math.* **34**, 195 (1979).
- [14] M.A. Olshanetsky and A.M. Perelomov, *Invent. Math.*, **54**, 261 (1979).
- [15] A.A. Golubtsova and V.D. Ivashchuk, On calculation of fluxbrane polynomials corresponding to classical series of Lie algebras; arxiv: 0804.0757 [nlin.SI].