Bounds in 4D Conformal Field Theories with Global Symmetry

Riccardo Rattazzi^{*a*}, Slava Rychkov^{*b*}, Alessandro Vichi^{*a*}

^a Institut de Théorie des Phénomènes Physiques, EPFL, CH-1015 Lausanne, Switzerland

^b Laboratoire de Physique Théorique, École Normale Supérieure, and Faculté de Physique, Université Pierre et Marie Curie, France

Abstract

We explore the constraining power of OPE associativity in 4D Conformal Field Theory with a continuous global symmetry group. We give a general analysis of crossing symmetry constraints in the 4-point function $\langle \phi \phi \phi^{\dagger} \phi^{\dagger} \rangle$, where ϕ is a primary scalar operator in a given representation R. These constraints take the form of 'vectorial sum rules' for conformal blocks of operators whose representations appear in $R \otimes R$ and $R \otimes \overline{R}$. The coefficients in these sum rules are related to the Fierz transformation matrices for the $R \otimes R \otimes \overline{R} \otimes \overline{R}$ invariant tensors. We show that the number of equations is always equal to the number of symmetry channels to be constrained. We also analyze in detail two cases—the fundamental of SO(N) and the fundamental of SU(N). We derive the vectorial sum rules explicitly, and use them to study the dimension of the lowest singlet scalar in the $\phi \times \phi^{\dagger}$ OPE. We prove the existence of an upper bound on the dimension of this scalar. The bound depends on the conformal dimension of ϕ and approaches 2 in the limit $\dim(\phi) \to 1$. For several small groups, we compute the behavior of the bound at $\dim(\phi) > 1$. We discuss implications of our bound for the Conformal Technicolor scenario of electroweak symmetry breaking.

September 2010

1 Introduction

Conformal Field Theory was originally conceived in four and three dimensions, with applications to particle physics and critical phenomena in mind. However, it is in 2D that the most spectacular results and exact solutions have been obtained. In higher dimensions, there seems to be a general feeling that the constraining power of conformal symmetry by itself is insufficient to tell nontrivial things about dynamics. Hence the interest in various additional assumptions, like supersymmetry, or integrability for the planar $\mathcal{N} = 4$ super Yang-Mills, or the AdS/CFT duality. This is not fully satisfactory, since there are likely many 4D CFTs which do not fulfill any of these assumptions. For example, "conformal windows" of non-supersymmetric gauge theories.

And yet, in the early days of 4D CFT, it was hoped that the *OPE associativity* is such a strong constraint on the CFT data (the spectrum of operator dimensions and the 3-point function coefficients) that it could allow for a complete solution of the theory. Recently [1],[2],[3],[4] we have been taking a fresh look at this idea, originally proposed by Polyakov [5]. Our approach is not to try to *solve* the OPE associativity, but rather to try to deduce from it general bounds that any CFT must obey. We discovered that such general bounds do exist! The bounds found so far fall into two general classes:

• [1],[2] An upper bound on the gap in the operator spectrum: any unitary 4D CFT containing a scalar operator ϕ of dimension d_{ϕ} must also contain another scalar O appearing in the OPE $\phi \times \phi$ whose dimension d_O is bounded by a universal function of d_{ϕ} :

$$d_O \le f(d_\phi). \tag{1}$$

The function $f(d_{\phi})$ has been computed numerically by means of a well-defined algorithm.

• Upper bounds on the 3-point function coefficients $\lambda_{\phi\phi O}$ where ϕ is the same scalar as above and O is any primary operator in the OPE $\phi \times \phi$. They have the form:

$$|\lambda_{\phi\phi O}| \le g(d_{\phi}, d_O, l_O), \tag{2}$$

where g is a universal function of the dimensions and of the O's spin l_O . For $l_O = 0$ this function was evaluated in [3]. Very recently, the case $l_O = 2$, $d_O = 4$, corresponding to the stress tensor OPE coefficient, was considered in [6],[4]. In this case the upper bound (2) gives a lower bound on the central charge of the theory. In this paper, we will discuss a generalization of bounds of the first class to the case when CFT has a continuous global symmetry G (Abelian or non-Abelian), and an operator ϕ transforms in a nontrivial representation R of G. We will consider the OPE $\phi \times \phi^{\dagger}$ if R is complex, or $\phi \times \phi$ if R is real. We will be discussing how to show that this OPE necessarily contains a *singlet scalar* operator S whose dimension d_S is bounded by a universal function which depends only on ϕ 's dimension and transformation properties:

$$d_S \le f_S(d_\phi, R_G). \tag{3}$$

Thus the novelty with respect to [1], [2] is that we will be bounding the gap in a given global symmetry sector (singlet in this case).

It is useful to recall that the original motivation of [1] was to find a bound of precisely this type for the case G = SO(4) and ϕ in the fundamental. This in turn was needed in order to constrain the Conformal Technicolor scenario of electroweak symmetry breaking [7]. This connection was discussed extensively in [1], and we will come back to it in the discussion Section.

The paper is organized as follows. We first derive the "vectorial sum rules", which generalize the main equation encoding the OPE associativity—the sum rule of [1]—to the globally symmetric case. Not surprisingly, these new sum rules take a different form depending on G and R under consideration. To begin with, we treat three concrete examples: ϕ in the fundamental of SO(N); ϕ charged under a U(1); ϕ in the fundamental of SU(N). Here we provide explicit derivations, illustrating the necessary technical ingredients. We then consider the general case in some detail, and in particular show that the number of constraints from crossing symmetry is always equal to the number of unknown functions.

Then we discuss what our vectorial sum rules imply for the bounds on the singlet dimension. In principle, functions f_S can be computed by a straightforward generalization of the algorithm of [1]. However, numerical difficulties involved are much greater in the present case, because one is working in a linear space of a much bigger dimension. As a result, we cannot yet push our analysis to the point where it produces numerically significant bounds. Thus we follow a more modest strategy. First, we explain how one shows that a bound does exist for d_{ϕ} sufficiently close to 1. The main idea is to consider the case $d_{\phi} = 1$ and then to argue by continuity (which is possible since the vectorial sum rules are continuous in d_{ϕ}). This analysis also shows that f_S approaches 2 as $d_{\phi} \to 1$. In other words, just like in our previous work, the free field theory limit is approached continuously.

The next question is then to determine how fast f_S approaches this limit. Here we limit

ourselves to quoting several f_S values at $d_{\phi} > 1$, leaving the determination of a detailed shape of these bounds to future work.

We conclude by discussing consequences for phenomenology and promising research directions.

Note. As this work was being prepared for publication, a very interesting paper [6] appeared, which gave several important generalizations of our method and results. In particular, [6] generalized our method to the $\mathcal{N} = 1$ SCFT case, and used it to derive bounds on non-BPS quantities. They also derived lower bounds on the central charge and on the two-point functions of global symmetry currents, with or without supersymmetry. Finally, they presented a set of equations incorporating OPE associativity constraints in the case of a U(1) global symmetry. They have even derived a bound on the U(1) singlet dimension (in the supersymmetric case). We will be commenting on this partial overlap in more detail below.

2 Sum rules in CFTs with a global symmetry

2.1 Conventions

We begin with some preliminary comments and notational conventions. We will work in the Euclidean space. Just as in our previous work, availability of explicit expressions for 4D conformal blocks given by Dolan and Osborn [8] will play a crucial role. Consider a 4-point function $\langle \phi(x_1)\chi^{\dagger}(x_2)\chi(x_3)\phi^{\dagger}(x_4)\rangle$ where ϕ and χ are two primary operators, not necessarily Hermitean, assumed to have equal dimensions $d_{\phi} = d_{\chi} = d$. The OPE $\phi \times \chi^{\dagger}$ will contain a sequence of spin l, dimension Δ primary fields $O_{\Delta l}$:

$$\phi \times \chi^{\dagger} = \sum_{\Delta,l} \lambda_{\Delta,l} O_{\Delta,l} \,. \tag{4}$$

Here $\lambda_{\Delta,l}$ are the OPE coefficients, in general complex. We then normalize the conformal blocks via:

$$\left\langle \begin{array}{cc} \phi(x_1) \bullet & \bullet \phi^{\dagger}(x_4) \\ \chi^{\dagger}(x_2) \bullet & \bullet \chi(x_3) \end{array} \right\rangle = \sum_{\Delta,l} \frac{1}{x_{12}^{2d} x_{34}^{2d}} p_{\Delta,l} g_{\Delta,l}(u,v) , \qquad (5)$$

$$u \equiv x_{12}^2 x_{34}^2 / (x_{13}^2 x_{24}^2) = z\bar{z}, \quad v \equiv x_{14}^2 x_{23}^2 / (x_{13}^2 x_{24}^2) = (1-z)(1-\bar{z}), \tag{6}$$

$$g_{\Delta,l}(u,v) = +\frac{zz}{z-\bar{z}} [k_{\Delta+l}(z)k_{\Delta-l-2}(\bar{z}) - (z\leftrightarrow\bar{z})], \qquad (7)$$

$$k_{\beta}(x) \equiv x^{\beta/2} {}_{2}F_{1}\left(\beta/2, \beta/2, \beta; x\right) \,. \tag{8}$$

The points x_i are assumed to be near the vertices of a square, as the picture suggests. The ordering is important. Eq. (5) says that the exchanges of $O_{\Delta,l}$ and of its conformal descendants in the (12)(34) channel (\equiv s-channel) can be summed up in a 'conformal block' $g_{\Delta,l}(u,v)$. The coefficients $p_{\Delta,l}$ are given by

$$p_{\Delta,l} = |\lambda_{\Delta,l}|^2 > 0. \tag{9}$$

Compared to [8], and also to our previous work, we have dropped the $(-1/2)^l$ prefactor in the expression for $g_{\Delta,l}$. This normalization is more convenient for the following reason. In the new convention all conformal blocks are positive when operators are inserted at the vertices of a square in the shown order (this corresponds to $z = \bar{z} = 1/2$). This is just as it should be, because this configuration is reflection-positive in the Osterwalder-Shrader sense with respect to the vertical median line (notice that the fields in the two sides of the correlator are complex-conjugate of each other)¹. Thus any s-channel contribution to the correlator, even spin or odd, has to be positive. There is no disagreement with Doland and Osborn [8], because in their notation the extra minus sign would be offset by a change in the sign of the OPE coefficient in the RHS of the correlator.

The (14)(23) channel (\equiv t-channel) conformal block decomposition can be analyzed similarly. In this case we will need OPEs $\phi \times \phi^{\dagger}$ and $\chi \times \chi^{\dagger}$ and only fields appearing in both of these OPEs will give a nonzero contribution, proportional to the product of the two OPE coefficients.

In [1] we have analyzed the particular case when ϕ is Hermitean and $\chi = \phi$. In this case the s- and t-channels correspond to the same OPE ($\phi \times \phi$). In addition, only even spins contribute because of permutation symmetry $x_1 \leftrightarrow x_2$. Let us introduce the notation for the sum of all s-channel contributions:

$$G^{+} = \sum_{l \text{ even};\Delta} p_{\Delta,l} g_{\Delta,l}(u,v) , \qquad (10)$$

(+ means that we are summing over even spins only) and a tilde notation for a contribution of the same set of operators in the t-channel:

$$\widetilde{G}^{+} = G_{u\leftrightarrow v}^{+} = \sum p_{\Delta,l} g_{\Delta,l}(v, u) \,. \tag{11}$$

Here we used the fact that going from the s- to the t-channel, which means simply rotating the picture by 90° , interchanges u and v. In this notation the crossing symmetry constraint of [1] is written compactly as:

$$G^{+} = \left(\frac{u}{v}\right)^{d} \tilde{G}^{+} \,. \tag{12}$$

¹Actually, conformal blocks are positive on the whole interval $0 < z = \overline{z} < 1$. Configurations corresponding to such z, \overline{z} can be mapped onto a rectangle, which is reflection-positive.

The appearance of the $(u/v)^d$ factor in this relation is due to a nontrivial transformation of the prefactor $1/(x_{12}^{2d}x_{34}^{2d})$ in (5) under crossing.

Finally, an important technical remark. Unlike in [1], to extract full information from the 4-point function (5), we will have to consider not only the s- and t-channel OPEs, but also the u-channel ones (13)(24). Conformal blocks for such 'diagonal' OPEs are related to the nearest-neighbor conformal blocks discussed above by analytic continuation, which introduces spin-dependent signs into the crossing-symmetry constraints. A useful way to keep track of these signs is not to consider the u-channel OPE directly, but to instead apply the s- and t-channel decompositions to the 4-point function with the permuted insertion points:

$$\left\langle \begin{array}{cc} \phi(x_1)\bullet & \bullet\chi(x_4) \\ \chi^{\dagger}(x_2)\bullet & \bullet\phi^{\dagger}(x_3) \end{array} \right\rangle$$
(13)

Here, we transposed the fields in the right side of the correlator. Now in the t-channel we have the same OPE as we would have in the u-channel in (5). And in the s-channel we have the same OPE as in (5), except for the transposition. This transposition is taken into account by reversing the sign of the odd-spin contributions in the s-channel (and permuting the flavor indices accordingly, see below).

2.2 Fundamental of SO(N)

As a first example we will now consider the SO(N) global symmetry case, with a scalar primary operator ϕ_a transforming in the fundamental representation. We normalize the 2-point function of ϕ_a as $\langle \phi_a(x)\phi_b(0)\rangle = \delta_{ab} (x^2)^{-d}$, $d = d_{\phi}$. Consider the 4-point function

$$\left\langle \begin{array}{cc} \phi_a \bullet & \bullet \phi_d \\ \phi_b \bullet & \bullet \phi_c \end{array} \right\rangle \equiv \frac{1}{x_{12}^{2d} x_{34}^{2d}} \mathcal{G} \begin{bmatrix} a & d \\ b & c \end{array} \middle| u, v \right].$$
(14)

The operator insertion points are assumed numbered in the same order as in (5).

Operators appearing in the $\phi_a \times \phi_b$ OPE can transform under the global symmetry as singlets S, symmetric traceless tensors $T_{(ab)}$, or antisymmetric tensors $A_{[ab]}$:

$$\phi_a \times \phi_b = \delta_{ab} \mathbb{1}$$

$$+ \delta_{ab} S^{(\alpha)} \quad \text{(even spins)}$$

$$+ T^{(\alpha)}_{(ab)} \quad \text{(even spins)}$$

$$+ A^{(\alpha)}_{[ab]} \quad \text{(odd spins)}.$$

$$(15)$$

The index (α) shows that an arbitrary number of operators of each type may in general be present, of various dimensions Δ and spins l. However, permutation symmetry of the $\phi_a \phi_b$ state implies that the spins of the S's and T's will be even, while they will be odd for the A's.

We note in passing that the stress tensor will be an S of $\Delta = 4, l = 2$, while the conserved SO(N) current will be an A of $\Delta = 3, l = 1$. The OPE coefficients of these operators are related to the stress tensor and the current central charges by the Ward identities [11], which should allow to derive various bounds on these quantities by the method of [3]. The simplest cases of these bounds have already been explored in [4],[6]. In this paper we will not be making any assumptions about the central charges of the theory and will treat the stress tensor and the current on equal footing with all the other fields. (However, in future studies central charge information may be useful; see the discussion Section.)

On the other hand, it will be important for us that the unit operator 1 is always present in the $\phi_a \times \phi_b$ OPE, with a unit coefficient.

As mentioned in the Introduction, we are interested to learn something about the dimension of the lowest-dimension singlet scalar (an S of l = 0). This will require disentangling its contribution to the 4-point function from the possibly present low-dimension scalars of type T.

We will now see what the crossing symmetry says about the relative weights of various contributions in the $\phi \times \phi$ OPE. Applying the conformal block decomposition in the s-channel we get:

Here $G_{S,T,A}$ are defined as in (10), and sum up conformal blocks of all fields of a given symmetry. Remember that $G_{S,T}$ contain only even spins, while G_A only the odd ones. The unit operator contributes together with the singlets, and its conformal block is $\equiv 1$. To keep track of the index structure, we are using the graphical notation for tensors. Every line means that the corresponding indices are contracted with the δ tensor:

$$= \delta_{ab} \,\delta_{cd} \,, \,\, \text{etc.} \tag{17}$$

The index structure of the symmetric traceless and the antisymmetric tensor contributions in (16) is fixed by the symmetry (and by the tracelessness, in the case of G_T). The signs are fixed from the requirement that for $a = d \neq b = c$ all contributions have to be positive by reflection positivity, see Section 2.1. Apart from the sign and the index structure, we do not keep track of

the overall, positive, normalization of each term. In other words, we know that each G contains conformal blocks summed with positive coefficients, but we do not keep track of the normalization of these coefficients. This is sufficient for deriving constraints on the operator spectrum, which is the focus of this paper. On the other hand, normalization conventions will be important for any future studies of the OPE coefficients.

Next we apply the t-channel conformal block decomposition to the same 4-point function, and we get an alternative representation:

$$\mathcal{G}\begin{bmatrix} a & d \\ b & c \end{bmatrix} = \left(\frac{u}{v}\right)^d \left\{ \underbrace{\bullet \bullet \bullet}_{\bullet \bullet \bullet} \cdot (1 + \widetilde{G}_S) + \left(\underbrace{\bullet \bullet \bullet}_{\bullet \bullet} + \underbrace{\bullet \bullet \bullet}_{\bullet \bullet} - \frac{2}{N} \underbrace{\bullet \bullet \bullet}_{\bullet \bullet} \right) \cdot \widetilde{G}_T + \left(\underbrace{\bullet \bullet \bullet}_{\bullet \bullet} + \underbrace{\bullet \bullet \bullet}_{\bullet \bullet} \right) \cdot \widetilde{G}_A \right\}$$

Note that to get this equation requires only changing the index structure appropriately, permuting $u \leftrightarrow v$ (here we are using the tilde notation introduced in Section 2.1), and multiplying by $(u/v)^d$ to take into account how the $1/(x_{12}^{2d}x_{34}^{2d})$ transforms.

Now we equate the s- and t-channel representations and pick up coefficients before each of the 3 inequivalent tensor structures: λ , λ , λ , λ . We get 2 independent equations:

$$u^{-d}\left\{G_T - G_A\right\} = v^{-d}\left\{\widetilde{G}_T - \widetilde{G}_A\right\}, \qquad (18a)$$

$$u^{-d}\left\{1+G_S-\frac{2}{N}G_T\right\} = v^{-d}\left\{\widetilde{G}_T+\widetilde{G}_A\right\},\tag{18b}$$

and a third one which can be obtained from the second by $u \leftrightarrow v$:

$$v^{-d}\left\{1 + \tilde{G}_S - \frac{2}{N}\tilde{G}_T\right\} = u^{-d}\left\{G_T + G_A\right\},$$
(19)

Notice that for the SO(N) case using the u-channel OPE would not yield any new equation.

It will be convenient to rewrite the system (18a),(18b) in the following equivalent form:

$$F_T - F_A = 0, \qquad (20a)$$

$$F_S + \left(1 - \frac{2}{N}\right)F_T + F_A = 1, \qquad (20b)$$

$$H_S - \left(1 + \frac{2}{N}\right) H_T - H_A = -1,$$
 (20c)

where we introduced notation for (anti)symmetric linear combinations of G and G:

$$F(u,v) = \frac{u^{-d}G(u,v) - v^{-d}G(v,u)}{v^{-d} - u^{-d}},$$

$$H(u,v) = \frac{u^{-d}G(u,v) + v^{-d}G(v,u)}{u^{-d} + v^{-d}}.$$
(21)

Thus (20a) is obtained from (18a) just by grouping and dividing by $v^{-d} - u^{-d}$. Eq. (20b) is obtained by taking the difference of (18b) and (19) and moving the contribution of the unit operator to the RHS. Finally, Eq. (20c) follows by taking the sum of (18b) and (19), and again separating the unity contribution.

Note that the functions F(u, v) were already used in our previous work, while the appearance of H(u, v) is a new feature of the global symmetry analysis. Writing the equations in terms of these functions is convenient because they are highly symmetric with respect to the $z = \bar{z} = 1/2$ point (they have only even derivatives in $z + \bar{z}$ and $z - \bar{z}$ at this point).

The system (20a)-(20c) is then the main result of this Section. In an expanded notation, it can be written as a 'vectorial sum rule':

$$\sum p_{\Delta,l}^{S} \begin{pmatrix} 0\\F_{\Delta,l}\\H_{\Delta,l} \end{pmatrix} + \sum p_{\Delta,l}^{T} \begin{pmatrix} F_{\Delta,l}\\(1-\frac{2}{N})F_{\Delta,l}\\-(1+\frac{2}{N})H_{\Delta,l} \end{pmatrix} + \sum p_{\Delta,l}^{A} \begin{pmatrix} -F_{\Delta,l}\\F_{\Delta,l}\\-H_{\Delta,l} \end{pmatrix} = \begin{pmatrix} 0\\1\\-1 \end{pmatrix}.$$
 (22)

Here the functions $F_{\Delta,l}(u, v)$ and $H_{\Delta,l}(u, v)$ are related to the individual conformal blocks $g_{\Delta,l}$ by the same formulas as F and H are related to G. Their dependence on d is left implicit. In each sum we are summing vector-functions corresponding to the dimensions and spins present in this symmetry channel, with positive coefficients. The total must converge to the constant vector in the RHS.

Consequences of this new sum rule for the lowest singlet dimension will be discussed below. Let us do however a quick counting of degrees of freedom. In total we have three G-functions: G_S, G_T, G_A , each of which is restricted only to the odd or even spins. The vectorial sum rule gives three equations for their (anti)symmetric combinations F and H. This coincidence between the number of equations and unknowns is not accidental; see Section 2.5. One may hope that the constraining power is similar to the case without global symmetry, when we had one equation for only one function G^+ . We will see in Section 3 how this hope is realized.

2.3 U(1)

We next discuss the U(1) global symmetry, as a case intermediate between SO(N) and SU(N). On the one hand, we will be able to check that the U(1) constraints agree with the already considered SO(N) case for N = 2. On the other hand, the derivation will be similar to the SU(N) case which follows. In particular, we will be working with complex fields and will need the u-channel OPE. We want to derive constraints from crossing in the 4-point function of a charge 1 complex scalar ϕ . Charge normalization is unimportant. The nonvanishing correlators must have zero total charge, thus we are led to consider $\langle \phi \phi \phi^{\dagger} \phi^{\dagger} \rangle$. There are two basic OPEs:

Charge 0 sector: $\phi \times \phi^{\dagger} = 1 + \text{spins } 0, 1, 2...,$ (23)

Charge 2 sector:
$$\phi \times \phi = \text{even spins only}$$
. (24)

Let us begin by considering the configuration

$$\left\langle \begin{array}{cc} \phi \bullet \bullet \phi^{\dagger} \\ \phi^{\dagger \bullet} \bullet \phi \end{array} \right\rangle, \tag{25}$$

which is the same as in (5) for $\chi = \phi$. By doing the s- and t-channel conformal block decompositions and demanding that the answers agree we get a constraint:

$$u^{-d}\left\{1+G_0^++G_0^-\right\} = v^{-d}\left\{1+\widetilde{G}_0^++\widetilde{G}_0^-\right\}.$$
(26a)

Here the subscript 0 refers to the charge 0 fields appearing in the relevant $\phi \times \phi^{\dagger}$ OPE. As indicated in (23), this OPE contains both even and odd spin fields, whose contributions we separate in G_0^{\pm} . According to the discussion in Section 2.1, reflection positivity of (25) implies that even and odd spins contribute in (26a) with the same positive sign.

Next consider the configuration with the transposed right side of the correlator:

$$\left\langle \begin{array}{cc} \phi \bullet & \bullet \phi \\ \phi^{\dagger \bullet} & \bullet \phi^{\dagger} \end{array} \right\rangle \,.$$

Equating the s- and t-channel decompositions we get:

$$u^{-d} \left\{ 1 + G_0^+ - G_0^- \right\} = v^{-d} \widetilde{G}_2^+ \,. \tag{26b}$$

The LHS of this equation differs from the LHS of (26a) only by the reversed sign of the odd spin contribution (see Section 2.1). The t-channel decomposition appearing in the RHS is positive since the configuration is reflection-positive in this direction.

Eqs. (26a),(26b) solve the problem of expressing crossing constraints in a U(1) symmetric theory. Very recently, the same equations also appeared in [6]. The authors of [6] have noticed that they could get a bound on the lowest dimension singlet by using just Eq. (26a) (they only computed the bound in the supersymmetric case, but the general case must be similar). Dropping the other equation simplified the problem, but the downside was that they had to Taylor-expand up to a pretty high order (12) to extract the bound. Below we will show that if one uses all equations the second-order expansion is already sufficient to extract a bound.

Upon identification

$$G_S = G_0^+, \quad G_A = G_0^-, \quad G_T = \frac{1}{2}G_2^+$$
 (27)

the U(1) constraints become equivalent to the N = 2 case of the SO(N) constraints discussed above. The appearance of a positive factor 1/2 is consistent with the fact that we are keeping careful track of positivity but not of the normalization.

2.4 Fundamental of SU(N)

Our last example is the SU(N) case, with a primary scalar ϕ_i transforming in the fundamental. We have two basic OPEs:

$$\phi_i \times \phi_{\bar{\imath}}^{\dagger} = \delta_{i\bar{\imath}} \mathbb{1} + \delta_{i\bar{\imath}} \times \text{Singlets(spins 0,1,2...)} + \text{Adjoints(spins 0,1,2...)}, \qquad (28)$$

$$\phi_i \times \phi_j =$$
 's (even spins) + 's (odd spins). (29)

The representation content of the first OPE is $N \otimes \overline{N} = 1 + \text{Adj}$. Notice that, in general, there will be singlets and adjoints of any spin. The adjoint sector will contain the conserved current, but at present we are not using information about its coefficient. The second OPE contains symmetric and antisymmetric tensors, of even and odd spins respectively.

The constraints are now derived by a combination of what we did for SO(N) and U(1). First consider the following 4-point function configuration:

$$\left\langle \begin{array}{cc} \phi_{i\bullet} & {}_{\bullet}\phi_{\bar{j}}^{\dagger} \\ \phi_{\bar{i}}^{\dagger\bullet} & {}^{\bullet}\phi_{j} \end{array} \right\rangle \,.$$

The s- and t-channel conformal block decompositions are evaluated using the first OPE. Equating them, we get a constraint:

$$u^{-d} \left\{ \oint \left(1 + G_S^+ + G_S^- \right) + \left(\begin{array}{c} \bullet & \bullet \\ \bullet & \bullet \\ \bullet & \bullet \\ \end{array} - \frac{1}{N} \oint \left(0 \right) \left(G_{\mathrm{Adj}}^+ + G_{\mathrm{Adj}}^- \right) \right\} \\ = v^{-d} \left\{ \begin{array}{c} \bullet & \bullet \\ \bullet & \bullet \\ \end{array} \right. \left(1 + \widetilde{G}_S^+ + \widetilde{G}_S^- \right) + \left(\begin{array}{c} \bullet & \bullet \\ \bullet & \bullet \\ \end{array} \right) \left(- \frac{1}{N} \begin{array}{c} \bullet & \bullet \\ \bullet & \bullet \\ \end{array} \right) \left(\widetilde{G}_{\mathrm{Adj}}^+ + \widetilde{G}_{\mathrm{Adj}}^- \right) \right\}$$

Here lines denote SU(N)-invariant contractions of N (dots) and \overline{N} (circles) indices by $\delta_{i\overline{i}}$. The tensor structure of the Adj contributions is fixed by the tracelessness condition of the SU(N)

generators. The sign is fixed by the condition that for $i = \overline{j} \neq j = \overline{i}$ the s-channel contributions must be positive by reflection positivity.

Setting equal the coefficients before \int_{\bullet}^{\bullet} and \int_{\bullet}^{\bullet} we get two equations:

$$u^{-d}\left\{1 + G_S^+ + G_S^- - \frac{1}{N}(G_{\rm Adj}^+ + G_{\rm Adj}^-)\right\} = v^{-d}\left\{\widetilde{G}_{\rm Adj}^+ + \widetilde{G}_{\rm Adj}^-\right\},\tag{30a}$$

and a second one which is just the $u \leftrightarrow v$ version of the first.

Next we consider the transposed 4-point configuration:

$$\left\langle \begin{array}{cc} \phi_{i\bullet} & {}_{\bullet}\phi_{j} \\ \phi_{\overline{\imath}}^{\dagger\bullet} & {}^{\bullet}\phi_{\overline{\jmath}}^{\dagger} \end{array} \right\rangle$$

Equating the s- and t-channel decompositions, we get:

$$\begin{split} u^{-d} \left\{ \begin{bmatrix} \mathbf{I} & \mathbf{I} \\ \mathbf{I} & \mathbf{I} \end{bmatrix} (1 + G_S^+ - G_S^-) + \left(\begin{array}{c} \mathbf{I} & \mathbf{I} \\ \mathbf{I} & \mathbf{I} \end{bmatrix} \right) (G_{\mathrm{Adj}}^+ - G_{\mathrm{Adj}}^-) \right\} \\ &= v^{-d} \left\{ \left(\begin{bmatrix} \mathbf{I} & \mathbf{I} \\ \mathbf{I} \end{bmatrix} + \begin{array}{c} \mathbf{I} \\ \mathbf{I} \end{bmatrix} \right) \widetilde{G}_{\square} + \left(\begin{bmatrix} \mathbf{I} & \mathbf{I} \end{bmatrix} - \begin{array}{c} \mathbf{I} \\ \mathbf{I} \end{bmatrix} \right) \widetilde{G}_{\square} \right\} \end{split}$$

The s-channel decomposition is obtained from the previous case by transposing the index structure *and* flipping the sign of the odd-spin contributions. The t-channel decomposition is obtained by using the second OPE (29). The index structure is fixed by (anti)symmetry of the exchanged fields, while the signs are determined by demanding positive contributions for $i = \bar{i} \neq j = \bar{j}$ (which makes the configuration reflection-positive in the t-channel).

Collecting coefficients before the two inequivalent tensor structures, we get two more equations, which this time are independent:

$$u^{-d} \left\{ 1 + G_S^+ - G_S^- - \frac{1}{N} G_{Adj}^+ + \frac{1}{N} G_{Adj}^- \right\} = v^{-d} \left\{ \widetilde{G}_{\Box\Box} + \widetilde{G}_{\Box} \right\},$$
(30b)

$$u^{-d}\left\{G_{\mathrm{Adj}}^{+}-G_{\mathrm{Adj}}^{-}\right\}=v^{-d}\left\{\widetilde{G}_{\Box\Box}-\widetilde{G}_{\Box}\right\}.$$
(30c)

The system (30a)-(30c) solves the problem of expressing the crossing symmetry constraints. Like in the SO(N) case, we will find it convenient to rewrite it by separating the unit operator contributions and (anti)symmetrizing with respect to u and v. We end up with the following equivalent vectorial sum rule:

$$\begin{array}{rcl}
F_{S}^{+} & +F_{S}^{-} & +\left(1-\frac{1}{N}\right)F_{Adj}^{+} & +\left(1-\frac{1}{N}\right)F_{Adj}^{-} & = & 1\\
H_{S}^{+} & +H_{S}^{-} & -\left(1+\frac{1}{N}\right)H_{Adj}^{+} & -\left(1+\frac{1}{N}\right)H_{Adj}^{-} & = & -1\\
F_{S}^{+} & -F_{S}^{-} & -\frac{1}{N}F_{Adj}^{+} & +\frac{1}{N}F_{Adj}^{-} & +F_{\Box} & +F_{\Box} & = & 1\\
H_{S}^{+} & -H_{S}^{-} & -\frac{1}{N}H_{Adj}^{+} & +\frac{1}{N}H_{Adj}^{-} & -H_{\Box} & -H_{\Box} & = & -1\\
& & F_{Adj}^{+} & -F_{Adj}^{-} & +F_{\Box} & -F_{\Box} & = & 0\\
& & H_{Adj}^{+} & -H_{Adj}^{-} & -H_{\Box} & +H_{\Box} & = & 0.\\
\end{array}$$
(31)

Just like for SO(N), the number of components, six, is equal to the number of the OPE channels classified by representation×(spin parity): S^{\pm} , Adj^{\pm} , \Box , \Box .

2.5 General case

In this Section we will consider the case of an arbitrary global symmetry group G, with ϕ_{α} transforming in an irreducible representation R. We aim at a general analysis of crossing symmetry constraints. In particular, we would like to understand why the number of constraints came out equal to the number of unknown functions in the explicit SO(N) and SU(N) examples above.

We will assume that R is complex. The case of R real is analogous but simpler; necessary changes will be indicated below.

To understand the group theory aspect of the problem, we begin by counting the number of scalar invariants which can be made out of two ϕ 's and two ϕ^{\dagger} 's. These invariants can be constructed by decomposing the products $\phi_{\alpha} \times \phi_{\overline{\alpha}}^{\dagger}$ and $\phi_{\beta} \times \phi_{\overline{\beta}}^{\dagger}$ into irreducible representations and contracting those. The tensor product representation decomposes as:

$$R \otimes \bar{R} = \bigoplus_{i=1}^{n} r_i(+\bar{r}_i) , \qquad (32)$$

where $(+\bar{r}_i)$ indicates that the representations in the RHS must be either real or come in complex conjugate pairs. To simplify the discussion, assume for now that all r_i are real and different. In accord with the above decomposition, we have

$$\phi_{\alpha} \times \phi_{\bar{\alpha}}^{\dagger} = \sum_{i} \sum_{A_{i}} C_{\alpha \bar{\alpha} A_{i}}^{i} \Psi_{A_{i}}^{i} , \qquad (33)$$

where the objects Ψ_{A_i} transform in the r_i , and $C^i_{\alpha\bar{\alpha}A_i}$ are the Clebsch-Gordan coefficients $(A_i$ is the index in the r_i). Then we can construct exactly n invariant tensors by contracting two

Clebsch-Gordan coefficients:

$$T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}} = \sum_{A_{i}} C^{i}_{\alpha\bar{\alpha}A_{i}} C^{i}_{\beta\bar{\beta}A_{i}} , \qquad (34)$$

so that the product of two ϕ 's and two ϕ^{\dagger} 's can be decomposed into a sum of T's:

$$\phi_{\alpha}\phi^{\dagger}_{\bar{\alpha}}\phi_{\beta}\phi^{\dagger}_{\bar{\beta}} = \sum_{i} \xi_{i}T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}}$$
$$= \sum_{i} \tilde{\xi}_{i}T^{i}_{\alpha\bar{\beta}\beta\bar{\alpha}}, \qquad (35)$$

where in the second line we indicated that we can do the same construction in a crossed fashion, by starting with the $\phi_{\alpha} \times \phi_{\bar{\beta}}^{\dagger}$ product. The fact that both decompositions exist means that the invariant tensors satisfy a linear relation ('Fierz identity')

$$T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}} = \mathcal{F}^{i}_{\ i'}T^{i'}_{\alpha\bar{\beta}\beta\bar{\alpha}}.$$
(36)

The matrix \mathcal{F} is invertible and must satisfy $\mathcal{F}^2 = \mathbb{1}$, since crossing is a \mathbb{Z}_2 operation.

It is also possible to construct invariants by starting from $\phi_{\alpha} \times \phi_{\beta}$, which requires the tensor product

$$R \otimes R = \bigoplus_{j=1}^{n} \tilde{r}_j.$$
(37)

Assume for now that all \tilde{r} 's appearing in this product are also distinct (excluding as well the possibility for the same representation to occur both in the symmetric and antisymmetric part of the tensor product). Under this simplifying assumption, the number of \tilde{r} 's is the same as the number of r's. Indeed, we can construct invariant tensors

$$\widetilde{T}^{j}_{\alpha\beta\bar{\alpha}\bar{\beta}} = \sum_{A_{j}} C^{j}_{\alpha\beta A_{j}} C^{j}_{\bar{\alpha}\bar{\beta}A_{j}} , \qquad (38)$$

where $C^{j}_{\alpha\beta A_{j}}$ (resp. $C^{j}_{\bar{\alpha}\bar{\beta}A_{j}}$) are the Clebsch-Gordan coefficients for \tilde{r}_{j} in $R \times R$ (resp. $\bar{\tilde{r}}_{j}$ in $\bar{R} \times \bar{R}$). These must be related to T's by another Fierz identity

$$T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}} = \widetilde{\mathcal{F}}^{i}_{\ j}\widetilde{T}^{j}_{\alpha\beta\bar{\alpha}\bar{\beta}}\,,\tag{39}$$

where $\widetilde{\mathcal{F}}$ is again an invertible matrix. Notice however that $T \neq \widetilde{T}$ and thus $\widetilde{\mathcal{F}}^2 \neq \mathbb{1}$.

After this prelude, we come back to our problem of analyzing the crossing symmetry constraints of the CFT 4-point function.

Step 1. Let us compare the s- and t-channel conformal block decompositions:

$$\left\langle \begin{array}{c} \phi_{\alpha \bullet} & \bullet \phi_{\bar{\beta}}^{\dagger} \\ \phi_{\bar{\alpha}}^{\dagger \bullet} & \bullet \phi_{\beta} \end{array} \right\rangle = \sum_{i} \left\langle \begin{array}{c} \alpha \\ \bar{\alpha} \\ \bar{\alpha} \\ \bar{\alpha} \\ \bar{\alpha} \\ \bar{\alpha} \\ \beta \end{array} \right\rangle_{\beta} = \sum_{i} \left\langle \begin{array}{c} \alpha \\ \bar{\alpha} \\ \bar{\alpha} \\ \bar{\alpha} \\ \bar{\alpha} \\ \beta \end{array} \right\rangle_{\beta} . \tag{40}$$

Introduce functions G_i which sum up conformal blocks of operators in the representation r_i (which will in general occur in both even and odd spins). The tensor structure of these contributions will be given precisely by the invariant tensors T introduced above. The crossing symmetry constraint then takes the form:

$$\sum_{i} T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}}G_{i}(u,v) = \sum_{i} T^{i}_{\alpha\bar{\beta}\beta\bar{\alpha}}G_{i}(v,u).$$
(41)

Here we assume that the signs of T's have been chosen in agreement with reflection positivity. To simplify the notation we included the u^{-d} , v^{-d} prefactors in the definition of G_i . We also do not separate the unit operator explicitly.

Eq. (41) will be consistent with the first Fierz identity (36) if and only if

$$G_i(u,v) = \mathcal{F}_i^{i'} G_{i'}(v,u) \,. \tag{42}$$

Let us now define even and odd combinations:

$$^{(\pm)}G_i = G_i(u,v) \pm G_i(v,u).$$
(43)

These are the analogues of the F and H functions from Eq. (21). We put the index (\pm) on the left to stress that it has nothing to do with the spin parity index used in the previous Sections; these functions receive contributions from both even and odd spins. We have

$$\left(\mathbb{P}_{\pm}\right)_{i}^{i'}G_{i'}^{(\pm)} = 0\,,\tag{44}$$

where $\mathbb{P}_{\pm} = (1 \mp \mathcal{F})/2$ are projectors, $(\mathbb{P}_{\pm})^2 = \mathbb{P}_{\pm}$ by using $\mathcal{F}^2 = \mathbb{1}$. Going to the diagonal basis for \mathbb{P}_{\pm} , it is clear that Eq. (44) represents a total of *n* constraints.

Step 2. We next compare the s- and t-channel conformal block decompositions of the transposed 4-point function:

$$\left\langle \begin{array}{cc} \phi_{\alpha \bullet} & \bullet \phi_{\beta} \\ \phi_{\overline{\alpha}}^{\dagger \bullet} & \bullet \phi_{\overline{\beta}}^{\dagger} \end{array} \right\rangle = \sum_{i} \left\langle \begin{array}{c} \alpha \\ \overline{\alpha} \\ \overline{\alpha} \\ \overline{\beta} \end{array} \right\rangle \stackrel{\beta}{=} \sum_{j} \left\langle \begin{array}{c} \alpha \\ \overline{\beta} \\ \overline{\alpha} \\ \overline{\beta} \end{array} \right\rangle \stackrel{\beta}{=} \sum_{j} \left\langle \begin{array}{c} \alpha \\ \overline{\beta} \\ \overline{\beta} \\ \overline{\beta} \end{array} \right\rangle.$$
(45)

The crossing symmetry constraint can be written in terms of the invariant tensors introduced above as:

$$\sum_{i} T^{i}_{\alpha\bar{\alpha}\beta\bar{\beta}} [G^{+}_{i}(u,v) - G^{-}_{i}(u,v)] = \sum_{j} \widetilde{T}^{j}_{\alpha\beta\bar{\alpha}\bar{\beta}} G_{j}(v,u).$$

$$\tag{46}$$

Here we have shown explicitly that the odd spin parts G_i^- of the G_i flip signs compared to the above configuration (40). Note as well that each of the functions G_j will include even or odd spins only, depending if \tilde{r}_j occurs in the symmetric or antisymmetric part of $R \times R$.

For Eq. (46) to be consistent with the second Fierz identity (39), we must have

$$G_i^+(u,v) - G_i^-(u,v) = \widetilde{\mathcal{F}}_i^{\ j} G_j(v,u) \,. \tag{47}$$

Since the functions in the RHS and LHS now refer to completely different OPE channels (r_i in $\phi \times \phi^{\dagger}$ vs \tilde{r}_j in $\phi \times \phi$), this equation gives exactly 2n constraints when (anti)symmetrizing in u, v.

To summarize, we expect 3n constraints for 3n channels r_i^{\pm} , \tilde{r}_j . In particular, n = 3 for the fundamental of SU(N).

In case when R is a real representation, we only have one set of invariant tensors, whose Fierz dictionary matrix satisfies $\mathcal{F}^2 = 1$. In this case each of n representations in the $R \times R$ product will contribute with only even or odd spins. Only the first step of the above analysis is needed in this case. We will get n constraints for n channels. The fundamental of SO(N) corresponds to n = 3.

Generalizations. Let us now discuss how one can relax the assumptions on the content of $R \otimes \overline{R}$ and $R \otimes R$ taken in the above argument. In general, $R \otimes \overline{R}$ may contain repetitions of the same representation as well as conjugate pairs, while $R \otimes R$ may contain the same representation in both symmetric (s) and antisymmetric (a) part. As it will become clear below, these two things must happen simultaneously. A sufficiently representative example is $R = \overline{15}$ of G = SU(3) [9]:

$$15 \otimes \overline{15} = 1 + 64 + (8_1 + 8_2) + (27_1 + 27_2) + (10 + \overline{10}) + (35 + \overline{35}), \tag{48}$$

$$\overline{15} \otimes \overline{15} = 3_a + \overline{6}_s + 15'_s + 24_a + 42_a + 60_s + (15_s + 15_a) + (21_a + 24_s).$$
(49)

In $15 \otimes \overline{15}$ we have 8 and 27 appearing twice each, and also two conjugate pairs $(10 + \overline{10} \text{ and } 35 + \overline{35})$, while in $\overline{15} \otimes \overline{15}$, 15 and 24 appear both as s and a.

In cases like this, it is slightly more involved to count the quartic invariants. When counting in the $R \otimes \overline{R}$ channel, every conjugate pair $r + \overline{r}$ gives two invariants which for future purposes we (anti)symmetrize with respect to $(\alpha \bar{\alpha}) \leftrightarrow (\beta \bar{\beta})$:

$$\sum_{A} C^{r}_{\alpha\bar{\alpha}A} C^{\bar{r}}_{\beta\bar{\beta}A} \pm C^{\bar{r}}_{\alpha\bar{\alpha}A} C^{r}_{\beta\bar{\beta}A} \,. \tag{50}$$

In the same channel, a k-fold repetition of a real representation r gives rise to k^2 invariants:

$$\sum_{A} C^{r_i}_{\alpha \bar{\alpha} A} C^{r_j}_{\beta \bar{\beta} A} \qquad (i, j = 1 \dots k),$$
(51)

which can be (anti)symmetrized with respect to $(\alpha \bar{\alpha}) \leftrightarrow (\beta \bar{\beta})$, producing k(k+1)/2 symmetrics and k(k-1)/2 antisymmetrics.

When counting in the $R \otimes R$ channel, every representation r occurring both as s and a gives rise to 4 invariants

$$\sum_{A} C^{r_{s/a}}_{\alpha\beta A} C^{\bar{r}_{s/a}}_{\bar{\alpha}\bar{\beta}A} \,, \tag{52}$$

out of which two are symmetric and two antisymmetric in $(\alpha \bar{\alpha}) \leftrightarrow (\beta \bar{\beta})$.

The total number of invariants must of course be the same counted in $R \otimes \overline{R}$ and in $R \otimes R$ channel. This is indeed true in the above example, when both $15 \otimes \overline{15}$ and $\overline{15} \otimes \overline{15}$ give 14. The number of symmetric in $(\alpha \overline{\alpha}) \leftrightarrow (\beta \overline{\beta})$ invariants also agrees (10 in both channels). This is also true in general. An intuitive argument is as follows. The total number of invariants equals the number of independent coupling constants in the scalar potential $V(\phi_1, \phi_2^{\dagger}, \phi_3, \phi_4^{\dagger})$ where ϕ_i are four non-identical scalars transforming in R. This number should be the same whether you begin by contracting ϕ_1 with ϕ_2^{\dagger} or ϕ_3 . Analogously, the number of symmetric invariants is the number of quartic couplings if we identify $\phi_3 \equiv \phi_1, \phi_4 \equiv \phi_2$.

Each of the two Fierz identities (36) and (39) will now split into two, one for symmetric and one for antisymmetric invariants.

Let us now proceed to the crossing symmetry analysis of the 4-point function $\left\langle \phi_{\alpha} \phi_{\bar{\alpha}}^{\dagger} \phi_{\beta} \phi_{\bar{\beta}}^{\dagger} \right\rangle$. To begin with, out of all the invariant tensors discussed above, only the symmetric ones will appear as the coefficients in the conformal block expansions of this correlator². The $(\alpha \bar{\alpha}) \leftrightarrow (\beta \bar{\beta})$ symmetry is made manifest by applying a conformal transformation which maps a generic 4-point configuration in (40) onto a parallelogram. The 180° rotation symmetry of the parallelogram then acts on the indices as $(\alpha \bar{\alpha}) \leftrightarrow (\beta \bar{\beta})$, see Fig. 1.

 $^{^{2}}$ In a general Lorentz-invariant theory, the flavor structure of this correlator will involve both symmetric and antisymmetric tensors.



Figure 1: For any 4-point configuration, there exists a conformal transformation which maps it onto a parallelogram.

To see how this symmetry arises in the conformal block decomposition, consider the OPE

$$\phi_{\alpha} \times \phi_{\bar{\alpha}}^{\dagger} = \sum_{r \text{ real}} \left(\sum_{i=1}^{k_r} \lambda_O^i C_{\alpha \bar{\alpha} A}^{r_i} \right) O_A \tag{53}$$

$$+\sum_{r+\bar{r} \text{ pairs}} \lambda_O C^r_{\alpha\bar{\alpha}A} O_A + (-1)^l \lambda_O^* C^{\bar{r}}_{\alpha\bar{\alpha}A} O^{\dagger}_A.$$
(54)

Here in the first line we include all operators belonging to the real representations. If the representation is repeated k times in the $R \otimes \overline{R}$ product, there are k independent Clebsch-Gordan coefficients, and k independent real OPE coefficients λ_O^i . In the second line we have operators from the complex-conjugate pairs, whose OPE coefficients are always complex-conjugate, up to a spin-dependent minus sign.

By using this OPE in the s-channel conformal block decomposition of (40), we see that indeed only symmetric invariant tensors arise. Notice that the off-diagonal invariant tensors $(i \neq j)$ in the case of repeated representations will appear with coefficients $\lambda_O^i \lambda_O^j$ (×conformal block), which are not positive definite. We will discuss below what this means for the subsequent application of the derived constraints.

We then consider the t-channel conformal block decomposition of (40), and repeat the analysis of Step 1. The resulting number of constraints is equal to the number n_{sym} of symmetric invariants, while the number of representation×(spin parity) channels is $2n_{\text{sym}}$.

To generalize Step 2, we have to consider the t-channel decomposition of (45). In this channel, the OPE parity selection rules imply immediately that only symmetric tensor structures appear, in agreement with the above general result. If the s and a representations are not repeated, as in the $\overline{15} \otimes \overline{15}$ example, then only diagonal terms are present, and all conformal blocks enter with positive coefficients. Compared to Step 1, we will have n_{sym} new representation×(spin parity) channels and $2n_{\text{sym}}$ new constraints.

We are done: we have a total $3n_{\text{sym}}$ constraints for $3n_{\text{sym}}$ channels. Moreover, these constraints distinguish not only different representations appearing in the OPE, but also different copies of the same representation, and how they 'interfere' among each other.

Let us now come back to the fact that if repeated representations are present in $R \otimes \bar{R}$, the off-diagonal 'interference' channels have coefficients $\lambda_O^i \lambda_O^j$. To appreciate the difficulty that this creates, readers unfamiliar with our method of linear functionals are encouraged to read the rest of this Section after having read Section 3.

Consider then our abstract way (58) of representing the vectorial sum rule. It is crucial for us that when all coefficients p_{α} are allowed to vary subject to the positivity constraints $p_{\alpha} \geq 0$, linear combinations in the LHS fill a convex cone. In particular, this allows us to use the dual formulation of the problem in the form (75). Since the off-diagonal coefficients may be negative, the geometric interpretation in this case is not as obvious. Notice however that the off-diagonal coefficients cannot become arbitrarily negative since they are not independent of the diagonal ones. For a sharp formulation, consider a symmetric real matrix

$$P_{ij} = \sum_{O} \lambda_O^i \lambda_O^j \,, \tag{55}$$

where we allow for presence of more than one operator O with a given dimension, spin, and representation. The characterizing property of P is positive-definiteness:

$$P_{ij}s_is_j \ge 0 \qquad \forall s_i \in \mathbb{R} \,. \tag{56}$$

Now, as can be seen from this equation, the set of positive-definite matrices forms by itself a convex cone. It follows that the set of vectors in the LHS of the vectorial sum rule will remain a convex cone even if repeated representations are present. Constraints (56) replace the simple inequality $p_{\alpha} \geq 0$. In practical applications these constraints may have to be discretized by choosing a finite set of vectors s_i .

The dual formulation (75) is extended to the present case as follows. For the vectors \mathbf{x}_{ij} in the LHS of the sum rule corresponding to diagonal (i = j) and off-diagonal $(i \neq j)$ channels of the repeated representation, the simple condition $\Lambda[\mathbf{x}_{\alpha}] \geq 0$ must be replaced by the following condition on the matrix $\Lambda[\mathbf{x}_{ij}]$:

$$P_{ij}\Lambda[\mathbf{x}_{ij}] \ge 0 \quad \forall P \text{ positive-definite.}$$
 (57)

In other words, $\Lambda[\mathbf{x}_{ij}]$ must belong to the cone *dual* to the cone of positive-definite matrices. However, the latter cone is in fact self-dual, as can be easily inferred from the representation (55). Thus, $\Lambda[\mathbf{x}_{ij}]$ must be itself positive-definite.

In the above discussion, only real representations were allowed to repeat in $R \otimes R$. However, repetitions of complex pairs could be treated similarly; the only difference is that the corresponding P matrices will be positive-definite Hermitean rather than real.

3 Bounds on the lowest singlet scalar dimension

3.1 Generalities

The previous Section would be a futile exercise in group theory if our vectorial sum rules did not have any useful consequences. We will now discuss how they can be used to bound the gap in the singlet scalar sector. Consider the SO(N) case for definiteness. Given a CFT spectrum, the sum rule (22) can be viewed as an equation for the coefficients $p_{\Delta,l}^{S,T,A}$. If we start imposing restrictions on the spectrum, such as raising the singlet scalar gap, it is conceivable that this equation will not have any solution consistent with the positivity requirement $p_{\Delta,l} \ge 0$. This is in fact precisely what will happen.

An equivalent, geometric, way to view this is as follows. Let us rewrite the sum rule (22) schematically as an equation in a linear space V of functions from two variables u, v into \mathbb{R}^3 (vector space of vector-functions):

$$\sum p_{\alpha} \mathbf{x}_{\alpha} = \mathbf{y} \,. \tag{58}$$

Here vectors \mathbf{x}_{α} represent all vector-functions appearing in the LHS of (22), while the \mathbf{y} is the vector corresponding to the RHS.

For a fixed CFT spectrum and varying $p_{\alpha} \geq 0$, the vectors in the LHS of (58) fill a convex cone, so the question is whether the vector \mathbf{y} belongs to this cone. Imposing restrictions on the spectrum reduces the set of vectors \mathbf{x}_{α} generating the cone, and the cone shrinks. It may be that the new smaller cone no longer contains \mathbf{y} , see Fig. 2(a,b).

In this paper we will be dealing with two types of restrictions on the spectrum. First of all, we will always impose the unitarity bounds [10]

$$\Delta \ge 1 \quad (l=0), \qquad \Delta \ge l+2 \quad (l\ge 1). \tag{59}$$



Figure 2: Geometric interpretation of the sum rule: (a) the sum rule has a solution $\Leftrightarrow \mathbf{y}$ belongs to the cone; (b) the assumed spectrum is such that the sum rule does not allow for a solution \Leftrightarrow \mathbf{y} does not belong to the cone; (c) for $\Delta_S = \Delta_S^{cr}$, the \mathbf{y} belongs to the cone boundary.

These lower bounds on operator dimension are a completely general property of unitary 4D CFTs. Note that they depend only on spin and not, say, on the global symmetry representation in which the operator transforms.

Second, we will impose a lower bound on the dimension of scalar singlets:

$$\Delta \ge \Delta_S \quad (l = 0 \text{ singlets only}). \tag{60}$$

According to the above discussion, increasing Δ_S makes the cone shrink. Our goal will be to show that for Δ_S above a certain critical value Δ_S^{cr} , the **y** is not in the cone. This critical value will then be a theoretical upper bound on the dimension of the first singlet scalar, valid in an arbitrary unitary 4D CFT. This is the bound denoted by f_S in Eq. (3) of the introduction.

Note that the list of vectors generating the cone, and thus the cone itself, vary continuously with Δ_S . Since for $\Delta_S < \Delta_S^{cr}$ the **y** is inside the cone and for $\Delta_S > \Delta_S^{cr}$ it's outside, for $\Delta_S = \Delta_S^{cr}$ it must belong to the cone boundary, see Fig. 2(c).

Up to now we were keeping the external scalar dimension d_{ϕ} fixed. However, the vectors entering the sum rule depend on d_{ϕ} via Eq. (21). This dependence is continuous, and thus the cone will vary continuously with d_{ϕ} . It follows that the bound $f_S(d_{\phi})$, if it exists, will have a continuous dependence on d_{ϕ} .

The reader may be worried that the above discussion was not totally rigorous. Indeed, the vector space V is infinite-dimensional, and there may be subtleties of convergence. However, below we will always be considering a finite-dimensional subspace of V, by Taylor-expanding the sum rule up to a fixed finite order k. On the one hand, this means that the bounds that we will derive

will not be optimal (they will approach optimality in the $k \to \infty$ limit). On the other hand, finite-dimensional analysis is simpler both practically and from the point of view of mathematical rigor. In particular, the statement that f_S is a continuous function of d_{ϕ} is safe at finite k.

3.2 Best possible bound for $d_{\phi} = 1$

We will begin by analyzing carefully the case $d_{\phi} = 1$. The reader may wonder why this is necessary, since there is a theorem that a scalar saturating the unitarity bound is necessarily free. This theorem is easy to prove: starting from the 2-point function $\langle \phi(x)\phi(0)\rangle = x^{-2}$ we deduce $\langle \partial^2 \phi(x) \partial^2 \phi(0) \rangle = 0$ and thus $\partial^2 \phi = 0$ in the Hilbert space. In the SO(N) case, we will have N free real scalars ϕ_a . The $\phi_a \times \phi_b$ OPE will contain two scalar operators

$$S^{(0)} = :\phi_c \phi_c:, \quad T^{(0)}_{(ab)} = :\phi_a \phi_b - \frac{1}{N} \delta_{ab}(\phi_c \phi_c):, \tag{61}$$

the first a singlet, the second a symmetric traceless, both dimension 2. Bounding the singlet scalar dimension from above seems a waste of time: the exact solution shows that it must be exactly 2.

Yet it is not useless to give a derivation of such a result—that at $d_{\phi} = 1$ there must be a singlet scalar of dimension ≤ 2 —using our method based on the vectorial sum rule. The reason is very simple: unlike the derivation sketched above, our method is robust with respect to small variations in d_{ϕ} . If we show that it works at $d_{\phi} = 1$, it is guaranteed to work as well for d_{ϕ} sufficiently close to 1. Thus we will know that a bound *exists* for d_{ϕ} close to 1 and, since it is a continuous function of d_{ϕ} , that it approaches 2 as $d_{\phi} \to 1$. We believe that such an existence proof is conceptually important. It is also easier than actually computing the bound at $d_{\phi} > 1$. The latter problem will be discussed in the next Section.

Consider then the sum rule (22) for $d_{\phi} = 1$. This sum rule has at least one solution-the one corresponding to the theory of N free scalars. This solution is very special, in that among all the vectors appearing in the RHS, only those corresponding to twist $\Delta - l = 2$ fields will have nonzero coefficients. This is because in free theory no fields of other twists appear in the $\phi_a \times \phi_b$ OPE. Apart from the two $\Delta = 2$ scalars mentioned above, there are infinitely many twist 2 fields of the form

$$:\phi_a \overleftrightarrow{\partial}_{\mu_1} \dots \overleftrightarrow{\partial}_{\mu_l} \phi_b:, \qquad (62)$$

appropriately (anti)symmetrized in a,b to separate SO(N) representations. Expanding the free

field theory 4-point function

$$\langle \phi_a(x_1)\phi_b(x_2)\phi_c(x_3)\phi_d(x_4)\rangle = \frac{\delta_{ab}\delta_{cd}}{x_{12}^2x_{34}^2} + \text{crossings}$$
(63)

into twist 2 conformal blocks, one can find all the $p_{\Delta,l}$ coefficients. The *S* and *T* contributions can be disentangled since they have a different index structure, see Eq. (16). Even though we will not need the explicit expressions, we quote the result:

$$p_{l+2,l}^{T} = (l!)^{2}/(2l)!, \qquad p_{l+2,l}^{S} = (2/N)p_{l+2,l}^{T} \qquad (l \text{ even}),$$

$$p_{l+2,l}^{A} = (l!)^{2}/(2l)! \qquad (l \text{ odd}).$$
(64)

Notice that all $p_{\Delta,l} \ge 0$, consistent with unitarity.

We now proceed to showing that any solution of the sum rule at $d_{\phi} = 1$ must contain a singlet scalar of dimension ≤ 2 . In fact we will show an even stronger result—that any such solution must contain a singlet scalar of dimension exactly 2.

Let's group the twist 2 terms in (22) separately from the rest:

$$\sum_{\text{twist } 2} p_{\alpha} \mathbf{x}_{\alpha} + \sum_{\text{twist } \neq 2} p_{\beta} \mathbf{x}_{\beta} = \mathbf{y}$$
(65)

First we will show that in any solution all twist $\neq 2$ coefficients p_{β} must be zero. This is shown by exhibiting a linear functional $\Lambda_0: V \to \mathbb{R}$ such that

$$\Lambda_0[\mathbf{y}] = 0,$$

$$\Lambda_0[\mathbf{x}_{\alpha}] = 0 \quad \forall \text{ fields of twist } 2,$$

$$\Lambda_0[\mathbf{x}_{\beta}] > 0 \quad \forall \text{ fields of twist } \neq 2.$$
(66)

Applying Λ_0 to (65), and using the fact that $p_{\beta} \ge 0$, we conclude immediately that all $p_{\beta} = 0$.

The functional Λ_0 can be written explicitly by Taylor-expanding the vectors entering the sum rule around the point $z = \bar{z} = 1/2$. Let's introduce the coordinates a, b

$$z = \frac{1}{2} + a + b, \quad \bar{z} = \frac{1}{2} + a - b.$$
 (67)

Then the functions $F_{\Delta,l}$ and $H_{\Delta,l}$ are even with respect to both a and b, so that their derivatives $\partial_a^m \partial_b^n$ at a = b = 0 are nonzero only if both m and n are even. It turns out that the functional Λ_0 can be chosen as the following linear combination of second derivatives at the a = b = 0 point:

$$\Lambda_0 \left[\mathbf{x} \right] = A(\partial_a^2 x_1 - \partial_b^2 x_1) + B(\partial_a^2 x_2 - \partial_b^2 x_2) + C(\partial_a^2 x_3 - \partial_b^2 x_3) \,. \tag{68}$$

Here $x_{1,2,3}$ are the components of the vector-function: $\mathbf{x} = (x_1, x_2, x_3)^T$. There is a certain freedom in choosing the coefficients A, B, C; the following choice is one possibility which works for all $N \ge 2$:

$$A = 1, B = 2, C = 0.$$
(69)

The first relation (66) is trivially satisfied; the other two follow from the following curious property of the $F_{\Delta,l}$ second derivatives at the a = b = 0 point:

$$\partial_a^2 F_{\Delta,l} = \partial_b^2 F_{\Delta,l} \qquad \text{at twist } 2 ,$$

$$\partial_a^2 F_{\Delta,l} > \partial_b^2 F_{\Delta,l} \qquad \text{at twist } \neq 2 .$$
(70)

Since all $p_{\beta} = 0$, we are reduced to a simpler equation which involves only twist 2 fields:

$$\sum_{\text{twist } 2} p_{\alpha} \mathbf{x}_{\alpha} = \mathbf{y} \,. \tag{71}$$

We want to show that if we drop the scalar singlet from this equation, there are no solutions. This is shown by exhibiting a second linear functional Λ_1 with the following properties:

$$\Lambda_1[\mathbf{y}] < 0,$$

$$\Lambda_1[\mathbf{x}_{\alpha}] \ge 0 \text{ on all twist 2 fields except for the singlet scalar.}$$
(72)

Notice that any such Λ_1 must necessarily be negative on the singlet scalar, to allow at the very least the existence of one explicit solution (64).

This functional can be written again as a linear combination of derivatives at a = b = 0:

$$\Lambda_1[\mathbf{x}] = \sum_{i=1}^{3} \sum_{\substack{m,n \text{ even}\\0\le m+n\le k}} \frac{\lambda_{m,n}^i}{m!n!} \partial_a^m \partial_b^n x_i \,.$$
(73)

At present we can only find the coefficients $\lambda_{m,n}^i$ numerically. For $2 \leq N \leq 7$ it is enough to use derivatives up to the second order, just like in Λ_0 . For example, for N = 4 one can use the functional whose only nonzero coefficients are

$$\lambda_{0,0}^{1} = -8, \quad \lambda_{0,2}^{1} = 6,$$

$$\lambda_{0,0}^{2} = -11, \quad \lambda_{0,2}^{2} = 8,$$

$$\lambda_{0,0}^{3} = -10, \quad \lambda_{0,2}^{3} = 3.$$
(74)

Including derivatives up to the fourth order (k = 4) allows to find functionals in the range up to $N \leq 128$. While we have not checked higher N, we feel sufficiently confident that, adding

more and more derivatives, functional Λ_1 can be found for any N. With this small proviso, the demonstration that at $d_{\phi} = 1$ there must be a dimension 2 singlet scalar is complete.

It is perhaps useful to give a geometric representation of the given proof, see Fig. 3. The existence of Λ_0 means that there is a hyperplane (the zero set of Λ_0) such that all the twist 2 vectors as well as the vector **y** belong to it, while all twist $\neq 2$ vectors lie on one side of it. The existence of Λ_1 means that this hyperplane can be rotated so that the twist 2 singlet scalar and the rest of the twist 2 vectors lie on the opposite sides of the rotated hyperplane.



Figure 3: This figure gives a geometric interpretation of the proof that at $d_{\phi} = 1$ the sum rule has no solution unless the $\Delta = 2$ singlet scalar is included in the spectrum. The solid-contour plane represents the zero set of the functional Λ_0 . The vector \mathbf{y} and all the twist 2 vectors (black dots) lie in the $\Lambda_0 = 0$ plane. On the other hand, all twist $\neq 2$ vectors, which for varying Δ trace separate curves labeled by spin and representation, lie on one side of this plane ($\Lambda_0 > 0$). The $\Lambda_0 = 0$ plane can be slightly rotated so that the \mathbf{y} vector and the twist 2 singlet scalar lie on one side of the rotated plane, while the rest of the twist 2 vectors lie on the opposite side. The rotated plane (dashed contour) can be described by an equation $\Lambda_0 + \varepsilon \Lambda_1 = 0$ for a small ε .

Let us now discuss the SU(N) case. The statement is the same: at $d_{\phi} = 1$ the OPE $\phi \times \phi^{\dagger}$ must contain a dimension 2 singlet scalar. This is derived from the SU(N) vectorial sum rule (31) by using the same method of Λ_0 and Λ_1 . Just like for SO(N), the functional Λ_0 can be given by using only second derivatives. For Λ_1 , one can use derivatives up to the second order for N = 2, 3, while derivatives up to the fourth order work for at least all $N \leq 50$.

Finally, the reader may want to compare the above discussion with Section 5.4 of [1], where

the $d_{\phi} = 1$ case was analyzed analogously, and somewhat more explicitly, for CFTs without global symmetry.

3.3 Some numerical results at $d_{\phi} > 1$

As explained in Section 3.1, the bound f_S is a continuous function of d_{ϕ} . The previous Section has established that $f_S = 2$ at $d_{\phi} = 1$. The next question is to understand how fast the bound deviates from 2 at we increase d_{ϕ} above 1. To do this we need to be able to compute the bound at any given d_{ϕ} . This is done by the method of linear functionals [1], already used above to analyze the $d_{\phi} = 1$ case. At any fixed d_{ϕ} , we will be looking for a linear functional Λ such that

$$\Lambda[\mathbf{y}] < 0,$$

$$\Lambda[\mathbf{x}_{\alpha}] \ge 0 \qquad \forall \text{ scalar singlets with } \Delta \ge \Delta_S \text{ and} \qquad (75)$$

$$\forall \text{ other fields (subject to the unitarity bounds)}.$$

If such a functional exists, then the spectrum without any singlet scalars below Δ_S cannot solve the vectorial sum rule and cannot be realized in any CFT. Geometrically, Eqs. (75) says that there is a hyperplane separating the **y** vector from the cone generated by such a spectrum. The bound f_S is computed as the smallest Δ_S for which a functional satisfying (75) can be found.

Assuming that Λ is of the form (73), Eqs. (75) define a linear programming problem for the coefficients $\lambda_{m,n}^i$. However, the number of constraints is formally infinite, which requires careful discretizations and truncations. We refer to [1],[2], and also to [6], for a detailed description of how these numerical difficulties are overcome.

There is no difference of principle between the system (75) and the analogous system analyzed in the case without global symmetry in our previous work; it is just bigger. There are several families of vectors \mathbf{x}_{α} , labeled by representation and spin parity, and each vector now represents a *vector*-function. As a result, for the same number of derivatives (parameter k in (73)), the globally symmetric case involves Q times more constraints and Q times more functional coefficients, where Q = 3; 6 for SO(N); SU(N). This makes computations much more time-consuming as well as increases numerical instabilities. In the case without global symmetry, the initial analysis of [1] used k = 6, and subsequently we were able to push k up to 18 [2], producing a very strong bound. For the globally symmetric case, we have so far not been able to go beyond k = 4 for SO(N) and k = 2 for SU(N).

Here is an account of these exploratory calculations:

- In every case that we looked at, we found a bound for d_{ϕ} near 1 which approached 2 continuously in the $d_{\phi} \rightarrow 1$ limit. Thus we have checked the existence theorem from the previous Section.
- We have seen that the bound is a monotonically increasing function of d_{ϕ} at $d_{\phi} > 1$.
- Typically, we have only computed the bound in the interval $1 \le d_{\phi} \le d_*$ where $f_S(d_*) \approx 4$. This is because knowing whether the singlet scalar is relevant or irrelevant is a particularly interesting question. It is also sufficient to get an idea about how strong the bound is, for a given k. In the following table, we give the d_* for the cases that we considered. At the current level of accuracy, our f_S interpolates almost linearly between 2 and 4 as d_{ϕ} increases from 1 to d_* .

G	$U(1) \equiv SO(2)$	SO(3)	SO(4)	SU(2)	SU(3)
d_*	1.063 $(k=2)$	$1.032 \ (k=2)$	$1.017 \ (k=2)$	1.016	1.003
	$1.12 \ (k=4)$	$1.08 \ (k=4)$	$1.06 \ (k=4)$	(k=2)	(k=2)

Table 1: First line: the global symmetry groups considered. The external scalar was assumed to transform in the fundamental. Second line: the value of external scalar's dimension d_{ϕ} for which the bound f_S on the singlet scalar dimension (monotonically increasing from 2 for $d_{\phi} = 1$) was seen to cross 4.

As we already mentioned in Section 2.3, a U(1) bound on the singlet scalar dimension was recently published in [6]. It is not possible to compare our and their results directly: on the one hand, they assume supersymmetry and expand to a very high order (k = 12) which should make their bound stronger, but on the other hand they use but one scalar component of the full vectorial sum rule, which makes their bound weaker. It seems that these two effects compensate each other, so that their bound is roughly comparable to our U(1), k = 4 bound. Notice however that they could not see any bound for $d \ge 1.16$, while we checked by using the full vectorial sum rule that a bound continues to exist even for larger d.

4 Discussion

In this paper we extended the constraints from OPE associativity and crossing first derived in Ref. [1] to CFTs with global symmetry. Focussing on the scalar 4-point function we derived a set of sum rules that constrains the operator content in all the possible channels with given symmetry and parity $(-1)^{\ell}$ quantum numbers. More precisely, by explicit examples and by a general argument, we have shown that the number of sum rules equals the number of possible channels. This results suggest that, in principle, one could obtain (correlated) constraints on the operators appearing in each channel. In analogy with previous studies these constraints could involve the dimension and the fusion coefficients of the lowest lying operators in a given channel.

As a first exploration we have studied the possibility of obtaining an upper bound on the dimension of the scalar singlet of lowest dimension appearing in the OPE of $\phi \times \phi$ and $\phi \times \phi^{\dagger}$. That question is also relevant to asses the viability of the so called Conformal Technicolor (CT) scenario [7]. The goal of CT is to achieve a natural separation between the electroweak scale and the scale of Flavor. That is motivated by the experimental success of the Cabibbo-Kobayashi-Maskawa pattern of flavor violation. The way CT works is that the electroweak symmetry breaking sector above the weak scale flows to a strongly coupled CFT. The role of the Higgs field is then played by a composite operator H, while electroweak gauge interactions arise from the weak gauging of a $SU(2) \times U(1)$ subgroup of the global symmetry group G of the CFT. In that situation, suppression of flavor violation is the more robust the closer to 1 is the dimension d_H of H. On the other hand, a natural separation of mass scale requires all total singlet scalars to be only marginally relevant or irrelevant. Indicating by S the lowest dimension singlet $\subset H \times H^{\dagger}$, we thus need d_S slightly below or above 4 ³. A major constraint on d_H is imposed by the top coupling, that runs like $\lambda_t(\mu) = \lambda_t(\mu_{EW})(\mu/\mu_{EW})^{d_H-1}$ and quickly becomes strong for $d_H - 1 = O(1)$. Notice that one major difficulty to achieve the dream of CT is that for $d_H = 1$ we must have $d_S = 2$ and the hierarchy problem in all its splendor.

To be more quantitative about the needed pattern of field dimensions, assumptions on the physics of flavor must be made. Making the optimistic, although plausible, assumption that flavor violation in the light families is either suppressed by their mixing to the third family of by their Yukawa couplings ⁴, the range $d_H \leq 1.7$, $d_S \geq 3.5 \div 4$ is sufficient. In that situation the scale where the top Yukawa becomes strong can be as low as ~ 100 TeV, so that the window where CT is active is not very big. On the other hand, the more conservative, but robust, assumption that all flavor violating operators are equally important at the Flavor scale requires the more constrained

³Of course if d_S is strictly > 4 we need some other marginally relevant coupling, or a strongly relevant coupling which can be taken small because of an extra symmetry, to generate the weak scale by dimensional transmutation.

⁴For instance for the $\Delta S = 2$ operators contributing to $K\bar{K}$ -mixing, this amounts to assuming respectively an extra suppression factor $\sim (V_{st}V_{dt})^2$ or $y_s y_d/y_t^2$ compared to operators involving just the third family quarks.

pattern $d_H - 1 \leq 0.2$, $d_S \geq 4$. That second situation corresponds to a flavor scale around 10^5 TeV, with the CFT describing physics in a sizeable window of scales. In view of the above, it would be interesting to derive, if it exists, an upper bound on d_S as a function of d_H which was one goal of the present paper. Indeed in [1] a first step toward an answer was given by deriving an upper bound on min (d_S, d_T) , where d_T is the dimension of the triplet scalar $\subset H \times H^{\dagger}$. The method of [1] was however "symmetry blind" in that it could not resolving the singlet and triplet channels. In [2] a further refinement of the bound was obtained by working up to 18 derivatives in function space. The bound is a monotonically growing function of d_H crossing 4 at about $d_H = 1.6$. That result is compatible with the flavor-optimistic CT scenario. Given the clear signs of convergence of the symmetry blind bound [2], and also given the remarkable success of the method in 2D CFTs where the bound basically tracks minimal models, we are tempted to conclude that the flavor-optimistic CT scenario is plausible. However the bound of [2], if interpreted as a bound on d_S (that is if $d_S < d_T$), would thoroughly rule out the flavor-robust CT.

In the present paper we have instead shown that it is possible to obtain an independent bound on the singlet. We have rigorously shown that the bound on d_S exists and goes smoothly to 2 as $d_{\phi} \to 1$ for ϕ a fundamental in SO(N) (SU(N)) and $N \leq 128$ ($N \leq 50$). We have further worked it out numerically for a few small groups, and in particular for SO(4), which is the smallest group of phenomenological relevance. The results are not yet very strong, as seen in Table 1. In particular, in SO(4), d_S crosses 4 already for $d_{\phi} = 1.06$, way within the interesting region of flavor-robust CT. One reason for the weakness of the bound is that our numerical method based on the Linear Programming algorithm does not converge fast enough when the function space is truncated beyond k = 4 derivatives. So our best bound just corresponds to working up to k = 4. One reason of the extra difficulty with respect to the symmetry blind case is that we are now dealing with a triple sum rule, rather than with a single one, and the complexity grows $3^2 = 9$ times faster with k. Notice that even in the symmetry blind case the bounds at low k are not very strong. Indeed one has the progression $d_* \approx 1.12$ (k = 2) [1], $d_* \approx 1.18$ (k = 4) (unpublished), $d_* \approx 1.35$ (k = 6) [1], eventually increased to $d_* \approx 1.6$ for k = 18 [2]. Assuming a similar rate of improvement for the globally symmetric bound, and assuming optimistically that we could push the analysis to similarly high values of k (which would likely require new ideas in algorithm implementation), we could expect to get to $d_* \approx 1.1 \div 1.2$ for the SO(4) case. This is more or less at the edge of interest of the flavor-robust CT.

It is important to understand why the globally symmetric bounds are so weak, and why they are getting even weaker at larger N, as shown in Table 1. One could imagine two alternative

explanations. One, boring, possibility is that crossing symmetry for the 4-point function of just one operator ϕ_{α} is simply not an efficient constraint in presence of global symmetry. It would be more interesting if, perhaps, our result is telling us something physical, namely that for larger global symmetry, the role of the singlet S in maintaining consistency of the theory is indeed getting smaller, so that it can be allowed to decouple. A very partial hint of that could be the fact, already emphasized in [1], that in the O(N) model in $4 - \epsilon$ dimensions the anomalous dimension of the singlet is $O(\epsilon)$ while that of the symmetric traceless is $O(\epsilon/N)$. One way to test which of the two possibilities is true is to derive a twin bound on the symmetric traceless T (without imposing any constraints on the singlet S). If also that bound were found to be weak, the first possibility would be favored. If, on the other hand, the symmetric traceless bound would turn up much stronger than the singlet one, and perhaps comparable in strength to the general bound of [1], [2] on the min (d_T, d_S) , then this would be an indication that tensor is much more important than singlet in maintaining the crossing symmetry. Note that, as far as we know, there's no simple reason why the bounds get weaker with larger N. At this stage this is just an experimental fact.

There is yet another piece of information which, if taken into account, could change the picture qualitatively. In our numerical study we did not make any assumption about the stress tensor and symmetry current central charges. In principle our bound could become stronger under the condition that these central charges are bounded from above, corresponding to a perhaps more reasonable theory, that is one that does not contain too many degrees of freedom. Indeed, in connection to our main phenomenological application, the common wisdom is that constraints from the S parameter point towards a small EWSB sector, so that the central charges should be small. This is especially true of the current central charge, because the current-current spectral density enters directly into the spectral representation of the S parameter. Since conformal symmetry is broken in the IR, only the high energy tail of this density will be controlled by the CFT central charge. This does not allow to make this connection precise, but still large central charge seems to be disfavored.

What one could do then is to study a lower bound on the central charge as a function of the gap in the singlet scalar sector. For the stress tensor central charge in the case without global symmetry, precisely such a study was performed in [4]. These studies are made possible by the existence of O(1) universal bounds on the coefficients $p_{\Delta,l}$ in the conformal block decomposition [3], and that the coefficients $p_{4,2}$ and $p_{3,1}$ can be related to the inverse central charges by the Ward identities [11]. In [4], we found that higher gap sometimes requires a significant increase in the stress tensor central charge. If the current central charge is shown to have an even stronger

dependence on the singlet scalar gap, this may indicate a potential difficulty with the S parameter for CT.

Notice that central charge studies are also interesting in their own right, without connection to CT. For example, can one show that an SO(N) theory with a fundamental necessarily has central charges larger than free theory of N scalars? For the stress tensor central charge and the case without global symmetry this was shown in [4],[6] (in a range of d_{ϕ} near 1).

In passing, one could imagine one day bounds like those discussed in this paper would make contact with the studies of IR fixed points of gauge theories performed on the lattice. Since the very existence of an IR fixed point implies that these theories cannot contain a singlet scalar with dimension below 4, our bounds could provide rigorous theoretical constraints on the lattice measurements of the fermion bilinear operator dimension [12].

The framework laid out in this paper will likely lead to many applications beyond those mentioned above. Be aware that the next crucial steps are an algorithm improvement and/or finding if there exists a set of questions for which the convergence is faster, so that interesting bounds can be obtained already at small k. This is a new field to explore!

Acknowledgements

We are grateful to Erik Tonni for collaboration at the early stages of this project. The work of R.R. and A.V. is supported by the Swiss National Science Foundation under contract No. 200020-126941. The work of S.R. was supported in part by the European Programme "Unification in the LHC Era", contract PITN-GA-2009-237920 (UNILHC).

References

- R. Rattazzi, V. S. Rychkov, E. Tonni and A. Vichi, "Bounding scalar operator dimensions in 4D CFT," JHEP 0812, 031 (2008) arXiv:0807.0004.
- [2] V. S. Rychkov and A. Vichi, "Universal Constraints on Conformal Operator Dimensions," Phys. Rev. D 80, 045006 (2009) arXiv:0905.2211.
- [3] F. Caracciolo and V. S. Rychkov, "Rigorous Limits on the Interaction Strength in Quantum Field Theory," Phys. Rev. D 81, 085037 (2010) arXiv:0912.2726.

- [4] R. Rattazzi, S. Rychkov and A. Vichi, "Central Charge Bounds in 4D Conformal Field Theory," arXiv:1009.2725 [hep-th].
- [5] A. M. Polyakov, "Nonhamiltonian approach to conformal quantum field theory," Zh. Eksp. Teor. Fiz. 66, 23 (1974).
- [6] D. Poland and D. Simmons-Duffin, "Bounds on 4D Conformal and Superconformal Field Theories," arXiv:1009.2087 [hep-th].
- [7] M. A. Luty and T. Okui, "Conformal technicolor," JHEP 0609, 070 (2006) arXiv:hep-ph/0409274. M. A. Luty, "Strong Conformal Dynamics at the LHC and on the Lattice," arXiv:0806.1235. J. Galloway, J. A. Evans, M. A. Luty and R. A. Tacchi, "Minimal Conformal Technicolor and Precision Electroweak Tests," arXiv:1001.1361 [hep-ph].
- [8] F. A. Dolan and H. Osborn, "Conformal four point functions and the operator product expansion," Nucl. Phys. B 599, 459 (2001) arXiv:hep-th/0011040. "Conformal partial waves and the operator product expansion," Nucl. Phys. B 678, 491 (2004) arXiv:hep-th/0309180.
- [9] R. Slansky, Phys. Rept. **79**, 1 (1981).
- [10] S. Ferrara, R. Gatto and A. F. Grillo, "Positivity Restrictions On Anomalous Dimensions," Phys. Rev. D 9, 3564 (1974); G. Mack, "All Unitary Ray Representations Of The Conformal Group SU(2,2) With Positive Energy," Commun. Math. Phys. 55, 1 (1977).
- [11] H. Osborn and A. C. Petkou, "Implications of Conformal Invariance in Field Theories for General Dimensions," Annals Phys. 231, 311 (1994) arXiv:hep-th/9307010.
- [12] F. Bursa, L. Del Debbio, L. Keegan, C. Pica and T. Pickup, "Mass anomalous dimension in SU(2) with two adjoint fermions," Phys. Rev. D 81, 014505 (2010) arXiv:0910.4535 [hep-ph]. L. Del Debbio, B. Lucini, A. Patella, C. Pica and A. Rago, "The infrared dynamics of Minimal Walking Technicolor," Phys. Rev. D 82, 014510 (2010) arXiv:1004.3206 [hep-lat]. T. De-Grand, Y. Shamir and B. Svetitsky, "Running coupling and mass anomalous dimension of SU(3) gauge theory with two flavors of symmetric-representation fermions," arXiv:1006.0707 [hep-lat]. F. Bursa, L. Del Debbio, L. Keegan, C. Pica and T. Pickup, "Mass anomalous dimension in SU(2) with six fundamental fermions," arXiv:1007.3067 [hep-ph].