

On Quadratic Twists of Hyperelliptic Curves

Mohammad Sadek

Abstract

Let C be a hyperelliptic curve of good reduction defined over a discrete valuation field K . Given $d \in K^* \setminus K^{*2}$, we find the minimal regular model of the quadratic twist of C by d . Then we prove that there exists an infinite family of hyperelliptic curves of genus 2 defined over \mathbb{Q} violating the Hasse principle.

1 Introduction

We assume that C is a hyperelliptic curve defined over \mathbb{Q} by the equation $y^2 = f(x) \in \mathbb{Z}[x]$, where $\deg(f) \geq 3$. The genus of C will be called g . If $d > 1$ is a square free integer, then we write C_d for the quadratic twist of C by d . In particular, C_d is defined by $dy^2 = f(x)$.

We try to find an explicit description of the minimal regular model of C_d/\mathbb{Q}_p in terms of the minimal regular model of C/\mathbb{Q}_p itself, when C has good reduction.

Let Δ denote the minimal discriminant of C/\mathbb{Q}_p , see §2 of [6]. For every prime $p \nmid \Delta$, C/\mathbb{Q}_p has good reduction. Hence the minimal regular model of C/\mathbb{Q}_p is smooth. The minimal regular model of C_d/\mathbb{Q}_p , $p \mid d$, is obtained as the minimal desingularisation of a quotient of a smooth scheme by a twisted action of some finite group, see §3. In fact, we prefer to handle the problem over the maximal unramified extension of \mathbb{Q}_p to avoid any complications which might appear because the residue field \mathbb{F}_p is not algebraically closed.

As we investigate minimal regular models of quadratic twists of a hyperelliptic curve C over \mathbb{Q}_p , we can not see how to do the cases C_d , where one of the prime divisors p of d is a bad prime of C and f has no simple root when reduced modulo p . The difficulty lies in the wide range of possibilities of the structure of the minimal regular model of C/\mathbb{Q}_p when C has bad reduction. Furthermore, we are not aware of any reference which discusses the desingularization of quotient singularities of models of algebraic curves when these singularities are not ordinary double points.

A curve C/\mathbb{Q} is said to violate the Hasse principle if $C(\mathbb{Q}_p) \neq \emptyset$ for every prime $p \leq \infty$ while $C(\mathbb{Q}) = \emptyset$. In [1], P. Clark finds the first natural family of curves violating the Hasse Principle over \mathbb{Q} . By natural I mean the family is of prior arithmetic-geometric interest. The precise statement of his result is as follows: If $131 < N \neq 163$ is a square free integer and $C(N, p)$ is the quadratic twist of the modular curve $X_0(N)$ by the Atkin-Lehner involution and the quadratic extension $\mathbb{Q}(\sqrt{p})/\mathbb{Q}$, then the set of primes $p \equiv 1 \pmod{4}$ such that $C(N, p)$ violates the Hasse principle over \mathbb{Q} has positive density. In §4 of [2], for every $g \geq 2$ a hyperelliptic curve of genus g violating the Hasse principle is constructed.

For a hyperelliptic curve C/\mathbb{Q} , it is not difficult to construct quadratic twists of C which are everywhere locally soluble. We introduce a strategy to look for some of these everywhere locally soluble quadratic twists.

Finally we give an explicit example of an infinite family of genus 2 hyperelliptic curves defined over \mathbb{Q} which violate the Hasse principle. More precisely, we introduce an infinite set of square free integers S , then for every $d \in S$ we construct an infinite family $\{C_\lambda\}_\lambda$ of genus 2 curves such that the quadratic twist of each C_λ by d violates the Hasse principle.

2 Hyperelliptic Curves

We assume K is a perfect field with $\text{char } K \neq 2$ and algebraic closure \overline{K} . Two hyperelliptic equations with coefficients in K

$$y^2 = f(x) \text{ and } z^2 = f'(u)$$

represent the same curve with genus $g \geq 1$ if

$$x = \frac{au + b}{cu + d}, \quad y = \frac{ez}{(cu + d)^{g+1}} \text{ where } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(K), \quad e \in K^*.$$

We associate a discriminant Δ to a hyperelliptic equation $y^2 = f(x)$ as in §2 of [6]. This equation defines a smooth curve if and only if $\Delta \neq 0$.

By a *hyperelliptic curve* C over K we mean a smooth curve of genus $g \geq 2$ endowed with a separable morphism $C \rightarrow \mathbb{P}_K^1$ of degree 2. There exists a hyperelliptic equation $y^2 = f(x) \in K[x]$ describing C with $\deg f \in \{2g + 1, 2g + 2\}$. The fact that C is smooth implies $f(x)$ is separable over \overline{K} . The equation $y^2 = f(x)$ has one singularity at infinity. If $\deg f = 2g + 1$, the singularity at infinity corresponds to one point ∞ on the

hyperelliptic equation. If $\deg f = 2g + 2$, the singularity corresponds to two points ∞^+ and ∞^- on the hyperelliptic equation, and these can be distinguished by the value of the rational function y/x^{g+1} . If $\deg f = 2g + 2$, then there exists a hyperelliptic equation $z^2 = f'(u)$ describing C over \overline{K} with $\deg f' = 2g + 1$, and it describes C/K if and only if $f(x)$ has a zero in K .

Let C be a smooth hyperelliptic curve of genus $g \geq 2$ defined over K by the equation $y^2 = f(x)$. We will denote the hyperelliptic involution on C by $\iota : C \rightarrow C$. Let $K' = K(\sqrt{d})$ be a separable quadratic extension of K where $d \in K \setminus K^{*2}$. By a quadratic twist C_d of C we mean the hyperelliptic curve obtained from the curve C/K' by twisting the curve C/K by the cohomology class corresponding to K' in $H^1(K, \langle \iota \rangle)$.

This means that if σ generates $\text{Gal}(K'/K)$, then the twisted action of $\text{Gal}(K'/K)$ on $C(K')$ is given by $Q \rightarrow \iota(\sigma Q)$. To produce an explicit equation describing C_d , we consider the quadratic character $\chi : \text{Gal}(K'/K) \rightarrow \{\pm 1\}$ associated with K'/K , i.e., $\chi(\sigma) = \sqrt{d}^\sigma / \sqrt{d}$. Then we define a cocycle in

$$H^1(K, \langle \iota \rangle) \cong H^1(K, \mathbb{Z}/2\mathbb{Z}) = \text{Hom}(\text{Gal}(\overline{K}/K), \mathbb{Z}/2\mathbb{Z}) = \text{Hom}(\text{Gal}(\overline{K}/K), \pm 1)$$

by

$$\xi : \text{Gal}(\overline{K}/K) \rightarrow \{\pm 1\}; \quad \xi_\tau = [\chi(\tau)].$$

Now, $\overline{K}(C) = \overline{K}(x, y)$ and $\overline{K}(C_d) = \overline{K}(x, y)_\xi$ (the twist of the function field by the cocycle ξ). Since $\iota(x, y) = (x, -y)$, the action of σ on $\overline{K}(x, y)_\xi$ is described by

$$\sqrt{d}^\sigma = \chi(\sigma)\sqrt{d}, \quad x^\sigma = x, \quad y^\sigma = \chi(\sigma)y.$$

Thus the functions which are fixed by $\text{Gal}(K'/K)$ in $\overline{K}(x, y)_\xi$ are $x' = x$, $y' = y/\sqrt{d}$, hence they are in $K(C_d)$. They satisfy the equation

$$dy'^2 = f(x').$$

The curves C and C_d are isomorphic over K' via $(x', y') \mapsto (x', y'\sqrt{d})$.

3 Minimal regular models of hyperelliptic curves

We assume that K is a complete discrete valuation field with ring of integers \mathcal{O}_K , valuation ν , uniformiser t and residue field k with $\text{char } k \neq 2$. Set $S = \text{Spec } \mathcal{O}_K$.

Let C be a hyperelliptic curve defined over K . In [6], Liu associates to C a projective model W , a *Weierstrass model* of C , defined over \mathcal{O}_K , arising from a hyperelliptic equation of C with integral coefficients. The discriminant Δ_W of W is the discriminant

of this hyperelliptic equation. The model W is said to be *minimal* if Δ_W is minimal, i.e., $\nu(\Delta_W)$ is the least possible valuation among the valuations of the discriminants of the hyperelliptic equations related to our equation via the transformations of the form given in §2. If $C(K) \neq \emptyset$, then W being minimal implies that the minimal regular model of C is the minimal desingularisation of W , see Corollaire 5 of [6].

Let W be a minimal Weierstrass model of C . The curve C has good reduction if $\nu(\Delta_W) = 0$, see §3 of [6]. In this case, the minimal regular model \mathcal{C} of C over S is smooth. Moreover, \mathcal{C} is the unique smooth model of C over S , ([7], Proposition 10.1.21 (b)).

Lemma 3.1. *Assume that C has good reduction over k . Let K'/K be a finite extension with residue field k'/k . Then $C \times_K K'$ has good reduction over k' .*

PROOF: Let \mathcal{C} be a minimal Weierstrass model of C . Let ν' be the valuation corresponding to K' . Since $\nu' = e_{K'/K} \times \nu$, where $e_{K'/K}$ is the ramification index of K'/K , one has $\nu'(\Delta_{\mathcal{C}}) = e_{K'/K} \times \nu(\Delta_{\mathcal{C}}) = 0$. \square

We state the following lemma which help us see whether a hyperelliptic curve has a rational point over a discrete valuation field or not.

Lemma 3.2 ([7], Corollary 9.1.32). *Let C be an algebraic curve of genus $g \geq 1$ defined over K . Let $\mathcal{C} \rightarrow S$ be the minimal regular model of C . Assume that $C(K) \neq \emptyset$. Then a point $P \in C(K)$ is reduced to a point $\tilde{P} \in \mathcal{C}_k(k)$, and \mathcal{C}_k is smooth at \tilde{P} . In particular, \tilde{P} belongs to a single irreducible component of multiplicity 1 in \mathcal{C}_k .*

Lemma 3.3. *Let C be a smooth hyperelliptic curve over K . Assume that C has good reduction over k . Let K'/K be a quadratic extension with residue field k'/k . Let \mathcal{C}' be the minimal regular model of $C \times_K K'$. Then \mathcal{C}' is smooth. Moreover, $\mathcal{C}'_{k'}$ consists of one irreducible component of multiplicity 1.*

PROOF: Since C has good reduction over k , then it extends to a smooth relative curve $\mathcal{C}/\mathcal{O}_K$. This relative curve is the minimal regular model of C over \mathcal{O}_K . Hence it consists of one irreducible component Γ . If $C(K) \neq \emptyset$, then $\text{mult}(\Gamma) = 1$, see Lemma 3.2, otherwise $\text{mult}(\Gamma) = 2$. The reason for the latter statement is as follows: Let D be the image of Γ in \mathbb{P}_S^1 under the morphism $g : \mathcal{C} \rightarrow \mathbb{P}_S^1$. We will denote the generic points of Γ and D by ξ and ξ' respectively. The morphism g restricts to $\mathcal{O}_{\Gamma, \xi} \rightarrow \mathcal{O}_{D, \xi'}$. The valuations ν_{Γ} and ν_D are the corresponding normalised valuations to Γ and D respectively. Remember that $\nu_D(t) = 1$ because D is reduced. One has $[\text{Frac}(\mathcal{O}_{\Gamma, \xi}) : \text{Frac}(\mathcal{O}_{D, \xi'})] = 2$. Since Γ is not reduced, we deduce that t ramifies in $\text{Frac}(\mathcal{O}_{\Gamma, \xi})$. Thus $\nu_{\Gamma}(t) = 2$.

Lemma 3.1 implies that the minimal regular model of $C \times K'$ is smooth. Again it consists of one irreducible component Γ' of multiplicity 1. This is clear if $\text{mult}(\Gamma) = 1$. If $\text{mult}(\Gamma) = 2$, then $\text{mult}(\Gamma') = \text{mult}(\Gamma)/[K' : K]$, see for example §2.4 of [8]. It is true that the mentioned reference gives result when $\bar{k} = k$, but if we take a base change over the maximal unramified extension of K , then the multiplicity of components will not change. \square

In what follows we assume k is algebraically closed. Hence $K' = K(\sqrt{t})$ is the unique totally and tamely ramified quadratic extension of K . Furthermore, K'/K is Galois. Let $G := \text{Gal}(K'/K) = \langle \sigma \rangle$.

Again C/K is a hyperelliptic curve. We assume that C/K has good reduction. We are concerned with the twisted action of σ on $C(K')$ given by $Q \rightarrow \iota(\sigma Q)$, where $\iota : C \rightarrow C$ is the hyperelliptic involution on C and $\sigma(Q)$ is the usual Galois action of σ on $C(K')$. Now the automorphism $\sigma : C \rightarrow C$ extends to the minimal regular models \mathcal{C} and \mathcal{C}' of C and $C \times K'$ respectively. We will denote the extended automorphisms by σ again.

Let C_t be the quadratic twist of C by t . In what follows we obtain the minimal regular model of C_t/K as the minimal desingularisation of the quotient scheme $\mathcal{C}'/\langle \sigma \rangle$ by the twisted action of σ . In other words, we construct the minimal regular model of C_t/K from the minimal regular model of C/K .

The first step is to find the fixed points of the twisted action of σ on \mathcal{C}' . This is because the singular points of $\mathcal{C}'/\langle \sigma \rangle$ lie among the images of the points of \mathcal{C}' fixed by σ . Let $y^2 = f(x)$ be a minimal hyperelliptic equation describing C . Then the assumption that C has good reduction implies that $f(x)$ has no repeated roots over k . If $P \in C \times K'$, we denote its reduction by \tilde{P} .

Proposition 3.4. *Let $P \in C \times K'$. The following are equivalent:*

- (i) P is fixed under the twisted action of σ .
- (ii) $\tilde{P} = (\tilde{x}, \tilde{y})$ is fixed under the twisted action of σ .
- (iii) $f(\tilde{x}) = 0$ over k .

PROOF: The twisted action of σ on P is given by $\iota(\sigma P)$. Let $P = (x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})$, where $x_0, x_1, y_0, y_1 \in K$ is fixed if and only if $(x_0 - x_1\sqrt{t}, -y_0 + y_1\sqrt{t}) = (x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})$. Hence $x_1 = y_0 = 0$. Whence P is fixed if and only if $P = (x_0, y_1\sqrt{t})$ where $x_0, y_1 \in K$. The latter is equivalent to $\tilde{P} = (\tilde{x}_0, 0)$, or equivalently $f(x_0) = 0$ in k . Now if $(x, y) \in \mathcal{C}_k$, then it is σ -fixed if and only if $(x, y) = (x, -y)$, i.e., $2y = 0$ but $2 \in k^*$, hence $y = 0$. So (i) \Leftrightarrow (ii) holds. \square

Recall that C and C_t are isomorphic over K' . Therefore both $C \times K'$ and $C_t \times K'$ have the same minimal regular model $\mathcal{C}' \rightarrow \text{Spec } \mathcal{O}_{K'}$. The model \mathcal{C}' is smooth. Since \mathcal{C}' is projective, the S -quotient scheme $Z := \mathcal{C}' / \langle \sigma \rangle$ is constructed by glueing together the rings of invariants of $\langle \sigma \rangle$ -invariant affine open sets of \mathcal{C}' . Moreover, Z is a normal scheme and hence its singular points are closed points of the special fiber.

Proposition 3.5. *Let $\sigma_k : \mathcal{C}'_k \rightarrow \mathcal{C}'_k$ and $\sigma_k^{\text{red}} : \mathcal{C}'_k^{\text{red}} \rightarrow \mathcal{C}'_k^{\text{red}}$ be the natural morphisms induced by σ . Then the natural map*

$$\mathcal{C}'_k^{\text{red}} / \langle \sigma_k^{\text{red}} \rangle \rightarrow Z_k^{\text{red}} := (\mathcal{C}' / \langle \sigma \rangle)_k^{\text{red}}$$

is an isomorphism over k .

PROOF: See ([10], Facts II, III). □

In fact if $\alpha : \mathcal{C}' \rightarrow Z$ is the quotient map, then α induces a natural map $\mathcal{C}'_k \rightarrow Z_k^{\text{red}}$ which factors as follows:

$$\mathcal{C}'_k \rightarrow \mathcal{C}'_k / \langle \sigma_k \rangle \rightarrow Z_k^{\text{red}}$$

where the second map is the normalisation map of Z_k^{red} , see ([9], p.21).

Proposition 3.6. *The generic fiber of $Z := \mathcal{C}' / \langle \sigma \rangle$ is isomorphic to C_t / K .*

PROOF: The generic fiber of Z is given by

$$Z_K = Z \times_{\mathcal{O}_K} K = (\mathcal{C}' / \langle \sigma \rangle) \times_{\mathcal{O}_K} K = (\mathcal{C}' \times_{\mathcal{O}_K} K) / \langle \sigma \rangle.$$

But one has

$$\mathcal{C}' \times_{\mathcal{O}_K} K = \mathcal{C}' \times_{\mathcal{O}_{K'}} \mathcal{O}_{K'} \times_{\mathcal{O}_K} K = \mathcal{C}' \times_{\mathcal{O}_{K'}} K' = C \times K'.$$

Since we consider the twisted action of σ , we have $Z_K = C_t$. □

Now we aim to prove that the minimal desingularisation \tilde{Z} of the S -quotient scheme $Z = \mathcal{C}' / \langle \sigma \rangle$ is the minimal regular model of C_t / K . Moreover, we will show that this model consists of one irreducible component of multiplicity 2, the image of the special fiber \mathcal{C}'_k of \mathcal{C}' in Z , and a finite number of multiplicity-1 irreducible components each of which corresponds to a singular point of Z . In particular, if $\rho : \mathcal{C}' \rightarrow Z$ is the quotient map, we will prove that since \mathcal{C}' is smooth then the only singular points of Z are the images of the σ -fixed points of \mathcal{C}' under ρ .


Consider the morphism

$$\mathcal{C}'_k \xrightarrow{\beta} \mathcal{C}'_k / \langle \sigma_k \rangle.$$

Let $y^2 = f(x)$ be a minimal hyperelliptic equation defining C . Then this equation still defines the minimal regular models \mathcal{C} and \mathcal{C}' which are smooth because C has good reduction. The ramification points of β are the zeros of $f(x)$ over k plus the point at infinity when $\deg f = 2g + 1$. Since $f(x)$ splits completely into linear factors over k , because $k = \bar{k}$, the number of the ramification points of β is $2g + 2$.

Theorem 3.7. *Let C/K be a hyperelliptic curve of genus g . Assume that C has good reduction over k . Let \mathcal{C}' be the minimal regular model of $C \times K'$. Again $\text{Gal}(K'/K) = \langle \sigma \rangle$. Let Z denote the quotient of \mathcal{C}' by the twisted action of σ by the hyperelliptic involution. Then Z is singular exactly at the images x_1, \dots, x_m , $m = 2g + 2$, of the ramification points of the morphism $\beta : \mathcal{C}'_k \rightarrow \mathcal{C}'_k / \langle \sigma_k \rangle$ in Z .*

Let $\tilde{Z} \rightarrow Z$ be the minimal desingularisation of Z . Then \tilde{Z} is the minimal regular model of the quadratic twist C_t of C over K' . Moreover, \tilde{Z}_k consists of an irreducible component of multiplicity 2 and components $\Gamma_1, \dots, \Gamma_m$ of multiplicity 1, each of which corresponds to blowing-up one of the x_i 's, see the following figure.



PROOF: We have seen that the generic fiber Z_K is the curve C_t/K , Proposition 3.6. Furthermore Z is a normal scheme. As we know that \mathcal{C}'_k consists of one irreducible component of multiplicity 1, Lemma 3.3, the image of this irreducible component in Z_k has multiplicity $[K' : K] = 2$, see Fact IV of [10].

To obtain the minimal desingularisation of Z we only need to blow-up the singularities of Z . We will show first that x_1, \dots, x_m are exactly the singular points of Z . If the points x_1, \dots, x_m were regular, then the morphism $\alpha : \mathcal{C}' \rightarrow Z$ would be flat above x_1, \dots, x_m . Hence by Zariski's purity Theorem, the branch locus $\alpha(\mathcal{C}' \setminus U)$, where U is the largest open subscheme of \mathcal{C}' such that $\alpha|_U : U \rightarrow Z$ is étale, is of codimension 1, see ([7], Exercise 8.2.15), a contradiction.

Now we blow-up Z at each x_i , $i = 1, \dots, m$ to construct the minimal desingularisation $\tilde{Z} \rightarrow Z$. Since x_i is a singular point of Z , it is known that if v_1, v_2 are local parameters of the local ring \mathcal{O}_{Z, x_i} , then the twisted action of σ_k on v_1, v_2 is described as follows:

$$\sigma_k(v_1) = -v_1, \sigma_k(v_2) = -v_2,$$

see ([17], Lemma 2.2) or ([14], Lemma 2). Let $T = \text{Spec } \hat{\mathcal{O}}_{Z, x_i}$ and $T^G = \text{Spec } \hat{\mathcal{O}}_{Z, x_i}^G$. Because x_i is non-regular then T^G is a non-regular local scheme. The monomials $s_1 = v_1^2, s_2 = v_1 v_2, s_3 =$

v_2^2 are invariant under the action of σ_k and any σ_k -invariant polynomial in v_1, v_2 is a polynomial in these. Therefore desingularising T^G is equivalent to desingularising $\text{Spec } k[s_1, s_2, s_3]/(s_2^2 - s_1 s_3)$. It is known that we need only one blow-up to desingularise $k[s_1, s_2, s_3]/(s_2^2 - s_1 s_3)$ at its only singular point corresponding to the maximal ideal $\mathfrak{m} = (s_1, s_2, s_3)$. In fact, the exceptional curve of the blowing-up is one irreducible component Γ_i isomorphic to \mathbb{P}_k^1 , see for example ([7], Example 8.1.5). Since σ_k acts trivially on x_i , the multiplicity of the irreducible component Γ_i is 1, see ([3], Proposition 7.3 (ii)).

Now each Γ_i has self-intersection -2 , whereas the multiplicity-2 component has self-intersection $-g - 1 \leq -3$. Therefore according to Castelnuovo's criterion \tilde{Z} contains no exceptional divisors, and hence \tilde{Z} is the minimal regular model of C_t/K . \square

Remark 3.8. In the above theorem, we see that the arithmetic genus of each of the components Γ_i , $1 \leq i \leq 2g + 2$, is 0, because $\Gamma_i \cong \mathbb{P}_k^1$. We can apply Hurwitz's formula, see ([7], Remark 10.4.8), to find the arithmetic genus of the multiplicity-2 irreducible component Θ . In fact, we have

$$\begin{aligned} 2p_a(C'_k) - 2 &= 2(2p_a(\Theta) - 2) + \sum_{1 \leq i \leq m} 1 \\ &= 4p_a(\Theta) + 2g - 2. \end{aligned}$$

Therefore we deduce $p_a(\Theta) = 0$ and $\Theta \cong \mathbb{P}_k^1$.

In Theorem 3.7 we proved that the quadratic twist C_t/K of a hyperelliptic curve C/K of genus $g \geq 2$ and with good reduction has a minimal regular model consisting of a component $\Theta \cong \mathbb{P}_k^1$ of multiplicity 2, and $2g + 2$ irreducible components $\Gamma_1, \dots, \Gamma_{2g+2}$ each is isomorphic to \mathbb{P}_k^1 and of multiplicity 1. The intersection numbers are given by $\Theta \cdot \Gamma_i = 1$ for every i and $\Gamma_i \cdot \Gamma_j = 0$ for every $i \neq j$. In case $g = 1$, this is reduction type I_0^* , see ([16], Chapter IV, §9). While for $g = 2$, this is reduction type $[I_{0-0-0}^*]$ given for example in p.155 of [13].

4 Quadratic twists which are not everywhere locally soluble

We start this subsection with the following result on irreducible polynomials over \mathbb{Q} .

Lemma 4.1. *Let $f(x) \in \mathbb{Z}[x]$ be an irreducible polynomial over \mathbb{Q} . Then there exists an infinite set of primes S_f in \mathbb{Q} such that $f(x)$ has no linear factors over \mathbb{F}_p for every $p \in S_f$.*

PROOF: See ([11], Remark 8.40(d)). \square

Chebotarev Density Theorem implies that the density $\delta(S_f)$ of S_f exists and satisfies $\delta(S_f) > 0$, see for example Exercise 11.3.7 of [12].

Let C be a hyperelliptic curve over \mathbb{Q} of genus $g \geq 1$. Then C has good reduction over all but finitely many finite places of \mathbb{Q} , see ([7], Proposition 10.1.21 (a)). The finite set of primes in \mathbb{Q} of bad reduction of C will be denoted by S_Δ .

Let C_d be a quadratic twist of C with $d > 1$ a square free integer. Since C and C_d are isomorphic over $K = \mathbb{Q}(\sqrt{d})$, it follows that they have the same minimal regular model over the ring of integers of any completion of K at one of its finite places. If $p \notin S_\Delta$, then this implies that for every prime $\mathfrak{p} \in K$ lying above p , both $C \times K_{\mathfrak{p}}$ and $C_d \times K_{\mathfrak{p}}$ have good reduction, see Lemma 3.1, where $K_{\mathfrak{p}}$ denotes the completion of K at \mathfrak{p} . In particular, the minimal regular model of $C_d \times K_{\mathfrak{p}}$ is smooth.

We will denote the maximal unramified extension of the p -adic field \mathbb{Q}_p by \mathbb{Q}_p^{un} . The residue field of \mathbb{Q}_p^{un} is $\overline{\mathbb{F}}_p$. We write \mathbb{Z}_p^{un} for the ring of integers of \mathbb{Q}_p^{un} . The following fact follows directly from Theorem 3.7.

Corollary 4.2. *Let $C : y^2 = f(x) \in \mathbb{Z}[x]$ be a hyperelliptic curve of genus $g \geq 1$ over \mathbb{Q} . Assume moreover that $f(x)$ is of even degree and irreducible over \mathbb{Q} . Let C_d be a quadratic twist of C with $d > 1$ a square free integer. If d has an odd prime factor $p \in S_f \setminus S_\Delta$, then $C_d(\mathbb{Q}_p) = \emptyset$. Hence $C_d(\mathbb{Q}) = \emptyset$.*

In particular, there are infinitely many quadratic twists of C which have no rational points over \mathbb{Q} because they are not everywhere locally soluble. In addition, there is a positive density family of prime quadratic twists of C which are not everywhere locally soluble.

PROOF: Note that $K := \mathbb{Q}_p^{un}(\sqrt{d}) = \mathbb{Q}_p^{un}(\sqrt{p})$ because the residue field $\overline{\mathbb{F}}_p$ is algebraically closed. Since $p \notin S_\Delta$, one has $C \times K$ is smooth and $f(x)$ factors completely into linear factors over $\overline{\mathbb{F}}_p$.

Let $\mathcal{C} \rightarrow \text{Spec } \mathcal{O}_K$ be the minimal regular model of $C \times K$ and $\mathcal{C}_d \rightarrow \text{Spec } \mathbb{Z}_p^{un}$ the minimal regular model of C_d/\mathbb{Q}_p^{un} . According to Theorem 3.7, the special fiber $(\mathcal{C}_d)_{\overline{\mathbb{F}}_p}$ consists of an irreducible component of multiplicity 2 and multiplicity-1 irreducible components $\Gamma_1, \dots, \Gamma_m$, $m = 2g+2$, corresponding to blowing-up the singular points of $\mathcal{C}/\langle \sigma \rangle$, where $\text{Gal}(K/\mathbb{Q}_p^{un}) = \langle \sigma \rangle$ and the action of σ on \mathcal{C} is the twisted action by the hyperelliptic involution introduced in §3. These singular points correspond to the simple roots of $f(x)$ over $\overline{\mathbb{F}}_p$ (plus the point at infinity if $\deg f = 2g + 1$). But since $f(x)$ has no simple root over $\overline{\mathbb{F}}_p$ as $p \in S_f$, and $\deg f = 2g + 2$, it follows that each Γ_i is defined over $\overline{\mathbb{F}}_p$ and none of the Γ_i 's is defined over \mathbb{F}_p . Indeed, $\mathcal{C}_d \rightarrow \text{Spec } \mathbb{Z}_p$ consists only of the multiplicity-2 component. According to Lemma 3.2, $C_d(\mathbb{Q}_p) = \emptyset$. \square

5 A family of genus 2 curves violating the Hasse principle

In the previous section we treated quadratic twists C_d of a hyperelliptic curve $C : y^2 = f(x) \in \mathbb{Z}[x]$, where f is irreducible over \mathbb{Q} , when f does not have a linear factor modulo at least one of the prime divisors of d . In case $\delta(S_f) = 1$, we see that the set of prime quadratic twists of C which might violate the Hasse principle has 0-density.

We give a strategy to look for everywhere locally soluble quadratic twists. Among these we can look for twists violating the Hasse principle, i.e., ones with no \mathbb{Q} -rational points.

Let \mathcal{P} denote the set of primes in \mathbb{Q} . Given an irreducible polynomial $f(x) \in \mathbb{Z}[x]$ over \mathbb{Q} , we write $\mathcal{P} = S_f \cup S_f^{lin}$.

Let C/\mathbb{Q} be an everywhere locally soluble hyperelliptic curve described by the hyperelliptic equation $y^2 = f(x)$ where $f(x) \in \mathbb{Z}[x]$ is irreducible of even degree. Let $d > 1$ be a square free integer such that if $q \mid d$ is prime, then $q \in S_f^{lin}$. The following steps ensures that $C_d(\mathbb{Q}_p) \neq \emptyset$ for every $p \leq \infty$.

- (i) If $(x_0, y_0) \in C(\mathbb{R})$, then $f(x_0)$ is a square in \mathbb{R} . Since $d > 0$, one has $(x_0, y_0/\sqrt{d}) \in C_d(\mathbb{R})$.
- (ii) Let $p \in S_f^{lin}$. Then $f(x)$ has a linear factor over \mathbb{F}_p , hence a linear factor over \mathbb{Q}_p by Hensel's Lemma. Therefore $C_d(\mathbb{Q}_p) \neq \emptyset$.
- (iii) Let $p \in S_f \setminus S_\Delta$. Since $p \nmid d$, one has that the completion of $\mathbb{Q}(\sqrt{d})$ at any of the primes lying above p is an unramified extension of \mathbb{Q}_p . Therefore $C \cong_{\mathbb{Q}_p^{un}} C_d$. Remembering that $p \notin S_\Delta$, our assumption that $C(\mathbb{Q}_p) \neq \emptyset$ implies that the special fiber of C over \mathbb{F}_p and hence the special fiber of C_d over \mathbb{F}_p consists of one smooth irreducible component of multiplicity 1. This is because the minimal regular model is stable under unramified base changes. Thus using Hensel's Lemma, one lifts a smooth \mathbb{F}_p -point to a point in $C_d(\mathbb{Q}_p)$.
- (iv) Let $p \in S_\Delta \cap S_f$. To overcome the various possible structures of the special fiber of the minimal regular model \mathcal{C} of C/\mathbb{Q}_p , we will assume that the special fiber of \mathcal{C} consists only of multiplicity-1 irreducible components. For example when $g = 2$, one can assume that \mathcal{C} has one of the reduction types [VII], [VIII – 1], [IX – 2], $[I_0 - I_0 - m]$, $m > 0$, see [13]. Now again since $p \nmid d$, then $C \cong_{\mathbb{Q}_p^{un}} C_d$ and so $\mathcal{C} \rightarrow \mathbb{Z}_p^{un}$ is the minimal regular model of $C_d \times \mathbb{Q}_p^{un}$. Whence the special fiber of the minimal regular model of C_d/\mathbb{Q}_p consists only of multiplicity-1 irreducible components. Therefore Hensel's Lemma implies that $C_d(\mathbb{Q}_p) \neq \emptyset$.

As we have seen when $f(x)$ is irreducible, the set S_f^{lin} might be finite. Then the above strategy will only yield a finite number of everywhere locally soluble quadratic twists. So it seems reasonable to dispense with the irreducibility of $f(x)$ and allow ourselves to do reducible polynomials $f(x) \in \mathbb{Z}[x]$ with no linear factors over \mathbb{Q} .

It is conjectured that for any elliptic curve E over \mathbb{Q} , asymptotically, there are at least half

of the quadratic twists E_d of E which have rank 0. A lot of work has been done to investigate this conjecture for special elliptic curves, yet the general conjecture is still open. In fact, we are going to use elliptic curves that we know they have positive proportion of quadratic twists with rank 0 to construct infinite families of hyperelliptic curves of genus 2 violating the Hasse principle.

Now let $E : y^2 = x(x^2 + ax + b)$, $a, b \in \mathbb{Q}$, be an elliptic curve defined over \mathbb{Q} with a rational 2-torsion point. E_d is the quadratic twist of E by a square free integer d . If there exists a constant $X_0 > 0$ such that for every $X > X_0$, one has

$$\#M_E^0(X) \gg X \text{ where } M_E^0(X) := \{d : |d| \leq X, \text{rank}(E_d(\mathbb{Q})) = 0\},$$

then we will say that E satisfies condition (*). Condition (*) is equivalent to saying that a positive proportion of the quadratic twists of E have Mordell-Weil rank 0.

We put $x = s^2 + \lambda$, $\lambda \in \mathbb{Z}^\times$, to obtain the following family of hyperelliptic curves $\{C_\lambda\}_\lambda$ of genus 2

$$C_\lambda : y^2 = (s^2 + \lambda)((s^2 + \lambda)^2 + a(s^2 + \lambda) + b).$$

We denote the involution of C_λ given by $(s, y) \mapsto (-s, y)$ by ι . The map $(s, y) \mapsto (s^2 + \lambda, y)$ is the quotient map $C_\lambda \rightarrow C_\lambda / \langle \iota \rangle \cong E$.

In the following theorem we generate an infinite family of genus 2 curves violating the Hasse principle. Let E be an elliptic curve satisfying condition (*). The set $S(E)$ is the infinite set of square free integers d such that $\text{rank}(E_d(\mathbb{Q})) = 0$.

Theorem 5.1. *Let $E : y^2 = x(x^2 + ax + b)$ be an elliptic curve over \mathbb{Q} satisfying condition (*). For all but finitely many $d \in S(E)$, there exists an infinite number of integers λ such that the quadratic twist by d of each C_λ/\mathbb{Q} , where*

$$C_\lambda : y^2 = (s^2 + \lambda)((s^2 + \lambda)^2 + a(s^2 + \lambda) + b),$$

violates the Hasse principle.

PROOF: Firstly we will prove that for each integer λ and for all but finitely many $d \in S(E)$, one has $C_{\lambda,d}(\mathbb{Q}) = \emptyset$.

Let E_d be the quadratic twist of E by d . Since E satisfies condition (*), a positive proportion of the quadratic twists of E have Mordell-Weil rank 0. For $d \in S(E)$, one has

$$\text{rank } E(\mathbb{Q}(\sqrt{d})) = \text{rank } E(\mathbb{Q}) + \text{rank } E_d(\mathbb{Q}) = \text{rank } E(\mathbb{Q}),$$

see for example ([15], Exercise 10.16). It is known that the torsion subgroup of $E(\mathbb{Q}(\sqrt{d}))$ is the same as the torsion subgroup of $E(\mathbb{Q})$ for all but finitely many d 's, see for example [5].

Let $C_{\lambda,d}/\mathbb{Q}$ be the quadratic twist of C_λ by d . Consider the morphism

$$C_{\lambda,d}(\mathbb{Q}) \hookrightarrow C_\lambda(\mathbb{Q}(\sqrt{d})) \rightarrow E(\mathbb{Q}(\sqrt{d})),$$

where the inclusion is given by $(s, y) \mapsto (s, y/\sqrt{d})$, and the second morphism is the quotient morphism $C_\lambda \rightarrow C_\lambda/\langle \iota \rangle \cong E$ given by $(s, y) \mapsto (s^2 + \lambda, y)$. Since $E(\mathbb{Q}) = E(\mathbb{Q}(\sqrt{d}))$ for all but finitely many $d \in S(E)$, it follows that $C_{\lambda,d}(\mathbb{Q}) = \emptyset$ for every such d .

In what follows, we fix d to be one of the square free integers such that $E(\mathbb{Q}) = E(\mathbb{Q}(\sqrt{d}))$. We will choose $\lambda \in \mathbb{Z}^\times$ such that $C_{\lambda,d}$ has points everywhere locally, and hence $C_{\lambda,d}$ violates the Hasse principle. The hyperelliptic curve $C_{\lambda,d}$ is given by

$$dy^2 = f_\lambda(s) := (s^2 + \lambda)((s^2 + \lambda)^2 + a(s^2 + \lambda) + b).$$

For every $\lambda \in \mathbb{Z}^\times$, there is an M_1 such that for all primes $q > M_1$, the hyperelliptic curve $C_{\lambda,d}$ has good reduction modulo q . In particular, $C_{\lambda,d}/\mathbb{F}_q$ consists of one smooth irreducible component. By Weil bounds, there exists an integer M_2 such that for every prime $q > M_2$, $C_{\lambda,d}/\mathbb{F}_q$ has an \mathbb{F}_q -rational point. Thus for every prime q such that $q > M := \max(M_1, M_2)$, Hensel's Lemma implies that $C_{\lambda,d}(\mathbb{Q}_q) \neq \emptyset$.

If $d > 0$, then pick $s \in \mathbb{R}$ big enough such that $f_\lambda(s) > 0$, whence $C_{\lambda,d}(\mathbb{R}) \neq \emptyset$. If $d < 0$, then for any $\lambda < 0$, find $s \in \mathbb{R}$ which makes $f_\lambda(s) < 0$, again $C_{\lambda,d}(\mathbb{R}) \neq \emptyset$.

Now we choose λ to satisfy $C_{\lambda,d}(\mathbb{Q}_2) \neq \emptyset$. Since $(s, y) \mapsto (s^2 + \lambda, y)$ is the quotient map $C_\lambda \rightarrow E$ and $(0, 0) \in E(\mathbb{Q}_2)$, we choose λ such that $-\lambda \in \mathbb{Z} \cap \mathbb{Q}_2^{*2} \setminus \mathbb{Z}^2$. Hence $(\sqrt{-\lambda}, 0) \in C_{\lambda,d}(\mathbb{Q}_2)$.

Finally we choose λ to treat the primes $q \leq M$, $q \mid d$ and $q \neq 2$. In fact, we choose λ such that $-\lambda$ is a quadratic residue modulo each of these primes. Therefore the factor $s^2 + \lambda$ factorises into two distinct linear factors modulo each such q . Thus Hensel's Lemma yields $\sqrt{-\lambda} \in \mathbb{Q}_q$. Whence $(\sqrt{-\lambda}, 0) \in C_{\lambda,d}(\mathbb{Q}_q)$. \square

Corollary 5.2. *There is an infinite set S of square free integers such that: for each $d \in S$, there exists an infinite family of hyperelliptic curves $\{C_\lambda\}_\lambda$, where*

$$C_\lambda : y^2 = (s^2 + \lambda - 1)(s^2 + \lambda)(s^2 + \lambda + 1),$$

whose quadratic twists by d violate the Hasse principle.

PROOF: The elliptic curve $E : y^2 = x^3 - x$ satisfies condition (*), see [4]. Hence the statement of the corollary follows as a direct consequence of Theorem 5.1. \square

Corollary 5.3. *Let $b \geq 2$ be a non perfect square integer admitting a solution $(u, v) \in \mathbb{Z}^2$ for the equation $u^2 - bv^2 = -4$. Let $a = bv$. There is an infinite set S of square free integers such that: for each $d \in S$, there exists an infinite family of hyperelliptic curves $\{C_\lambda\}_\lambda$, where*

$$C_\lambda : y^2 = (s^2 + \lambda)((s^2 + \lambda)^2 + a(s^2 + \lambda) + b),$$

whose quadratic twists by d violate the Hasse principle.

PROOF: According to Theorem 1.1 of [18], the elliptic curve E/\mathbb{Q} given by $y^2 = x(x^2 + ax + b)$, where a, b satisfy the conditions stated above, satisfies condition (*). Now we conclude by Theorem 5.1. \square

References

- [1] P. Clark. An "anti-Hasse principle" for prime twists. *International Journal of Number Theory*, 4:627–637, 2008.
- [2] D. Coray and C. Manoil. On large Picard groups and the hasse principle for curves and K3 surfaces. *Acta Arithmetica*, 76(2):165–189, 1996.
- [3] L. Halle. Stable reduction of curves and tame ramification. *Mathematische Zeitschrift*, 265(3):529–550, 2009.
- [4] D.R. Heath-Brown. The size of Selmer groups for the congruent number problem I. *Invent. Math.*, 111(1):171–195, 1993.
- [5] S. Kwon. Torsion subgroups of elliptic curves over quadratic extensions. *Journal of Number Theory*, 62:144–162, 1997.
- [6] Q. Liu. Modèles entiers des courbes hyperelliptiques sur un corps de valuations discrète. *Trans. Amer. Math. Soc.*, 348(11):4577–4610, November 1996.
- [7] Q. Liu. *Algebraic Geometry and Arithmetic Curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002.
- [8] D. Lorenzini. Models of curves and wild ramification. To appear in *Pure and Applied Math. Quarterly*.
- [9] D. Lorenzini. Wild quotient singularities of surfaces, preprint.
- [10] D. Lorenzini. Dual graphs of degenerating curves. *Math. Ann.*, 287:135–150, 1990.
- [11] J.S. Milne. *Algebraic Number Theory (v3.01)*. 2008. Available at www.jmilne.org/math/.
- [12] Maruti Ram Murty and Jody Esmonde. *Problems in algebraic number theory*, volume 190 of *Graduate Texts in Mathematics*. Springer-Verlag, 2nd edition, 2005.
- [13] Y. Namikawa and K. Ueno. The complete classification of fibres in pencils of curves of genus two. *Manuscripta Math.*, 9:143–186, 1973.
- [14] A.N. Paršin. Minimal models of curves of genus 2 and homomorphisms of abelian varieties defined over a field of finite characteristic. *Math. USSR Izv.*, 6(1):65–108, 1972.
- [15] J. Silverman. *The arithmetic of elliptic curves*. GTM 106. Springer-Verlag, New York, 1986.
- [16] J. Silverman. *Advanced topics in the arithmetic of elliptic curves*. GTM 151. Springer-Verlag, 1995.
- [17] H. Xue. Minimal resolution of Atkin-Lehner quotients of $X_0(N)$. *Journal of Number Theory*, 129:2072–2092, 2009.
- [18] G. YU. Rank 0 quadratic twists of a family of elliptic curves. *Compositio Mathematica*, 135:331–356, 2003.