

# FUNCTIONAL EQUATIONS FOR ZETA FUNCTIONS OF $\mathbb{F}_1$ -SCHEMES

OLIVER LORSCHIED

ABSTRACT. For a scheme  $X$  whose  $\mathbb{F}_q$ -rational points are counted by a polynomial  $N(q) = \sum a_i q^i$ , the  $\mathbb{F}_1$ -zeta function is defined as  $\zeta_{\mathcal{X}}(s) = \prod (s - i)^{-a_i}$ . Define  $\chi = N(1)$ . In this paper we show that if  $X$  is a smooth projective scheme, then its  $\mathbb{F}_1$ -zeta function satisfies the functional equation  $\zeta_{\mathcal{X}}(n - s) = (-1)^\chi \zeta_{\mathcal{X}}(s)$ . We further show that the  $\mathbb{F}_1$ -zeta function  $\zeta_G(s)$  of a split reductive group scheme  $G$  of rank  $r$  with  $N$  positive roots satisfies the functional equation  $\zeta_G(r + N - s) = (-1)^\chi (\zeta_G(s))^{(-1)^r}$ .

## 1. INTRODUCTION

In recent years around a dozen different suggestion of what a scheme over  $\mathbb{F}_1$  should be appeared in literature (cf. [6]). The common motivation for all these approaches is to provide a framework in which Deligne's proof of the Weyl conjectures can be transferred to characteristic 0 in order to proof the Riemann hypothesis. Roughly speaking,  $\mathbb{F}_1$  should be thought of as a field of coefficients for  $\mathbb{Z}$ , and  $\mathbb{F}_1$ -schemes  $\mathcal{X}$  should have a base extension  $\mathcal{X}_{\mathbb{Z}}$  to  $\mathbb{Z}$  which is a scheme in the usual sense.

Though it is not clear yet whether one of the existing  $\mathbb{F}_1$ -geometries comes close to this goal, and thus in particular it is not clear what the appropriate notion of an  $\mathbb{F}_1$ -scheme should be, the zeta function  $\zeta_{\mathcal{X}}(s)$  of such an elusive  $\mathbb{F}_1$ -scheme  $\mathcal{X}$  is determined by the scheme  $X = \mathcal{X}_{\mathbb{Z}}$ .

Namely, let  $X$  be a variety of dimension  $n$  over  $\mathbb{Z}$ , i.e. a scheme such that  $X_k$  is an variety of dimension  $n$  for any field  $k$ . Assume further that  $X$  has a counting polynomial

$$N(q) = \sum_{i=0}^n a_i q^i \in \mathbb{Z}[q],$$

i.e. the number of  $\mathbb{F}_q$ -rational points is counted by  $\#X(\mathbb{F}_q) = N(q)$  for every prime power  $q$ . If  $X$  descends to an  $\mathbb{F}_1$ -scheme  $\mathcal{X}$ , i.e.  $\mathcal{X}_{\mathbb{Z}} \simeq X$ , then  $\mathcal{X}$  has the zeta function

$$\zeta_{\mathcal{X}}(s) = \lim_{q \rightarrow 1} (q - 1)^\chi \zeta_X(q, s)$$

where  $\zeta_X(q, s) = \exp\left(\sum_{r \geq 1} N(q^r) q^{-sr} / r\right)$  is the zeta function of  $X \otimes \mathbb{F}_q$  if  $q$  is a prime power and  $\chi = N(1)$  is the order the pole of  $\zeta_X(q, s)$  in  $q = 1$  (cf. [9]). This expression comes down to

$$\zeta_{\mathcal{X}}(s) = \prod_{i=0}^n (s - i)^{-a_i}$$

([9, Lemme 1]).

From this it is clear that  $\zeta_{\mathcal{X}}(s)$  is a rational function in  $s$  and that its zeros (resp. poles) are at  $s = i$  of order  $-a_i$  for  $i = 0, \dots, n$ . The only statement from the Weyl conjectures which is not obvious for zeta functions of  $\mathbb{F}_1$ -schemes is the functional equation.

---

I like to thank Takashi Ono for drawing my attention to the symmetries occurring in the counting polynomials of split reductive group schemes, and I like to thank Markus Reineke for his explanations on the comparison theorem for liftable smooth varieties.

## 2. THE FUNCTIONAL EQUATION FOR SMOOTH PROJECTIVE $\mathbb{F}_1$ -SCHEMES

Let  $X$  be an (irreducible) smooth projective variety of dimension  $n$  with a counting polynomial  $N(q)$ . Let  $b_0, \dots, b_{2n}$  be the Betti numbers of  $X$ , i.e. the dimensions of the singular homology groups  $H_0(X_{\mathbb{C}}), \dots, H_{2n}(X_{\mathbb{C}})$ . By Poincaré duality, we know that  $b_{2n-i} = b_i$ . As a consequence of the comparison theorem for smooth liftable varieties and Deligne's proof of the Weil conjectures, we know that the counting polynomial is of the form

$$N(q) = \sum_{i=0}^n b_{2i} q^i$$

and that  $b_i = 0$  if  $i$  is odd (cf. [2] and [8]). Thus  $\chi = \sum_{i=0}^n b_{2i}$  is the Euler characteristic of  $X_{\mathbb{C}}$  in this case (cf. [4]).

Suppose  $X$  has an elusive model  $\mathcal{X}$  over  $\mathbb{F}_1$ . Then  $\mathcal{X}$  has the zeta function  $\zeta_{\mathcal{X}}(s) = \prod_{i=0}^n (s-i)^{-b_{2i}}$ .

**Theorem 1.** *The zeta function  $\zeta_{\mathcal{X}}(s)$  satisfies the functional equation*

$$\zeta_{\mathcal{X}}(n-s) = (-1)^{\chi} \zeta_{\mathcal{X}}(s)$$

*and the factor equals  $-1$  if and only if  $n$  is even and  $b_n$  is odd.*

*Proof.* We calculate

$$\begin{aligned} \zeta_{\mathcal{X}}(n-s) &= \prod_{i=0}^n ((n-s)-i)^{-b_{2i}} \\ &= \prod_{i=0}^n (-1)^{b_{2i}} (s-(n-i))^{-b_{2i}} \\ &= (-1)^{\chi} \prod_{i=0}^d (s-(n-i))^{-b_{2n-2i}} \end{aligned}$$

where we used  $b_{2n-2i} = b_{2i}$  in the last equation. If we now substitute  $i$  by  $n-i$  in this expression, we obtain

$$\zeta_{\mathcal{X}}(n-s) = (-1)^{\chi} \prod_{i=0}^n (s-i)^{-b_{2i}} = (-1)^{\chi} \zeta_{\mathcal{X}}(s).$$

If  $n$  is odd, then there is an even number of non-trivial Betti numbers and  $\chi = 2b_0 + 2b_2 + \dots + 2b_{n-1}$  is even. If  $n$  is even, then  $\chi = 2b_0 + 2b_2 + \dots + 2b_{n-2} + b_n$  has the same parity as  $b_n$ . Thus the additional statement.  $\square$

**Remark.** Note the similarity with the functional equation for motivic zeta functions as in [3, Thm. 1]. Amongst other factors, also  $(-1)^{\chi(M)}$  appears in the functional equation of the zeta function of a motive  $M$  where  $\chi(M)$  is the (positive part of the) Euler characteristic of  $M$ .

## 3. THE FUNCTIONAL EQUATION FOR REDUCTIVE GROUPS OVER $\mathbb{F}_1$

The above observations imply further a functional equation for reductive group schemes over  $\mathbb{F}_1$ . Note that Soule's and Connes and Consani's approaches towards  $\mathbb{F}_1$ -geometry indeed succeeded in descending split reductive group schemes from  $\mathbb{Z}$  to  $\mathbb{F}_1$  (cf. [1], [5], [7]).

Let  $G$  be a split reductive group scheme of rank  $r$  with Borel group  $B$  and maximal split torus  $T \subset B$ . Let  $N$  be the normalizer of  $T$  in  $G$  and  $W = N(\mathbb{Z})/T(\mathbb{Z})$  be the Weyl group. The Bruhat decomposition of  $G$  (with respect to  $T$  and  $B$ ) is the morphism

$$\coprod_{w \in W} BwB \longrightarrow G,$$

induced by the subscheme inclusions  $BwB \rightarrow G$ , which has the property that it induces a bijection between the  $k$ -rational points for every field  $k$ . We have  $B \simeq \mathbb{G}_m^r \times \mathbb{A}^N$  as schemes where  $N$  is the number of positive roots of  $G$ , and  $BwB \simeq \mathbb{G}_m^r \times \mathbb{A}^{N+\lambda(w)}$  where  $\lambda(w)$  is the length of  $w \in W$ . With this we can calculate the counting polynomial of  $G$  as

$$N(q) = \# \coprod_{w \in W} BwB(\mathbb{F}_q) = (q-1)^r q^N \sum_{w \in W} q^{\lambda(w)}.$$

The quotient variety  $G/B$  is a smooth projective scheme of dimension  $N$  with counting function  $N_{G/B}(q) = ((q-1)^r q^N)^{-1} N(q) = \sum_{w \in W} q^{\lambda(w)}$ . Let  $b_0, \dots, b_{2N}$  be the Betti numbers of  $G/B$ , then we know from the previous section that  $N_{G/B}(q) = \sum_{l=0}^N b_{2l} q^l$  and that  $b_{2N-2l} = b_{2l}$ .

Thus we obtain for the counting polynomial of  $G$  that

$$\begin{aligned} N(q) &= q^N \left( \sum_{k=0}^r (-1)^{r-k} \binom{r}{k} q^k \right) \cdot \left( \sum_{l=0}^N b_{2l} q^l \right) \\ &= \sum_{i=0}^d \left( \sum_{k+l=i-N} (-1)^{r-k} \binom{r}{k} b_{2l} \right) q^i \end{aligned}$$

where  $d = r + 2N$  is the dimension of  $G$  and with the convention that  $\binom{r}{k} = 0$  if  $k < 0$  or  $k > r$ . Denote by  $a_i = \sum_{k+l=i-N} (-1)^{r-k} \binom{r}{k} b_{2l}$  the coefficients of  $N(q)$ .

**Lemma.** *We have  $a_0 = \dots = a_{N-1} = 0$  and  $a_{d-i} = (-1)^r a_{i+N}$ .*

*Proof.* The first statement follows from the fact that  $N(q)$  is divisible by  $q^N$ . For the second statement we use the symmetries  $\binom{r}{k} = \binom{r}{r-k}$  and  $b_{2N-2l} = b_{2l}$  to calculate

$$\begin{aligned} a_{d-i} &= \sum_{k+l=d-i-N} (-1)^{r-k} \binom{r}{k} b_{2l} \\ &= \sum_{k+l=d-i-N} (-1)^r (-1)^k \binom{r}{r-k} b_{2N-2l}. \end{aligned}$$

When we substitute  $k$  by  $r-k$  and  $l$  by  $N-l$  in this equation and use  $d = r + 2N$ , then we obtain

$$a_{d-i} = (-1)^r \sum_{k+l=(i+N)-N} (-1)^{r-k} \binom{r}{k} b_{2l},$$

which is the same as  $(-1)^r a_{i+N}$ .  $\square$

Suppose  $G$  has an elusive model  $\mathcal{G}$  over  $\mathbb{F}_1$ . Then  $\mathcal{G}$  has the zeta function  $\zeta_{\mathcal{G}}(s) = \prod_{i=0}^n (s-i)^{-a_i}$ . Let  $\chi = N(1) = \sum_{i=0}^d a_i$ .

**Theorem 2.** *The zeta function  $\zeta_{\mathcal{G}}(s)$  satisfies the functional equation*

$$\zeta_{\mathcal{G}}(r+N-s) = (-1)^\chi \left( \zeta_{\mathcal{G}}(s) \right)^{(-1)^r}.$$

*Proof.* We use of the previous lemma and  $r + N = d - N$  to calculate that

$$\begin{aligned}\zeta_{\mathcal{G}}(r + N - s) &= \prod_{i=0}^n (r + N - s - i)^{-a_i} \\ &= \prod_{i=0}^n (d - N - s - i)^{-(-1)^r a_{d-N-i}}.\end{aligned}$$

After substituting  $i$  by  $d - N - i$ , we find that

$$\begin{aligned}\zeta_{\mathcal{G}}(d - N - s) &= \prod_{i=0}^n (i - s)^{-(-1)^r a_i} \\ &= (-1)^{\sum a_i} \left( \prod_{i=0}^n (s - i)^{-a_i} \right)^{(-1)^r} \\ &= (-1)^{\chi} \left( \zeta_{\mathcal{G}}(s) \right)^{(-1)^r}.\end{aligned}\quad \square$$

**Remark.** Kurokawa calculates the  $\mathbb{F}_1$ -zeta functions of  $\mathbb{P}^n$ ,  $\mathrm{GL}(n)$  and  $\mathrm{SL}(n)$  in [4]. One can verify the functional equation for these examples immediately.

#### REFERENCES

- [1] A. Connes and C. Consani. *On the notion of geometry over  $\mathbb{F}_1$* . Preprint, arXiv:0809.2926, 2008.
- [2] A. Deitmar. *Remarks on zeta functions and K-theory over  $\mathbb{F}_1$* . Proc. Japan Acad. Ser. A Math. Sci., vol. 82, 2006.
- [3] B. Kahn. *Motivic zeta functions of motives*. Pure and Applied Mathematics Quarterly 5, 1 (2009) 507–570
- [4] N. Kurokawa. *Zeta functions over  $\mathbb{F}_1$* . Proc. Japan Acad. Ser. A Math. Sci. 81 (2005), no. 10, 180–184.
- [5] J. López Peña and O. Lorscheid. *Torified varieties and their geometries over  $\mathbb{F}_1$* . Online first at Mathematische Zeitschrift, 2009.
- [6] J. López Peña and O. Lorscheid. *Mapping  $\mathbb{F}_1$ -land: an overview over geometries over the field with one element*. To be published in the proceeding of the conferences on  $\mathbb{F}_1$ , arXiv:0909.0069, 2009.
- [7] O. Lorscheid. *Algebraic groups over the field with one element*. arXiv:0907.3824, 2009.
- [8] M. Reineke. *Moduli of representations of quivers*. Trends in representation theory of algebras and related topics, EMS Ser. Congr. Rep. (2008), 589–637.
- [9] C. Soulé. *Les variétés sur le corps à un élément*. Mosc. Math. J. 4 (2004), 217–244.

CITY COLLEGE OF CUNY, DEPT. MATH., 160 CONVENT AVE., NYC, NY 10031, U.S.  
E-mail address: olorscheid@ccny.cuny.edu