

Corrections to “Unified Laguerre polynomial-series-based distribution of small-scale fading envelopes”

Yin Sun*, *Student Member, IEEE*, Árpád Baricz, and Shidong Zhou*, *Member, IEEE*.

Abstract—In this correspondence, we point out two typographical errors in Chai and Tjhung’s paper and we offer the correct formula of the unified Laguerre polynomial-series-based cumulative distribution function (cdf) for small-scale fading distributions. A Laguerre polynomial-series-based cdf formula for non-central chi-square distribution is also provided as a special case of our unified cdf result.

Index Terms—Generalized Laguerre polynomial series expansion; non-central chi-square distribution; small-scale fading; unified distribution.

I. TWO TYPOGRAPHICAL ERRORS IN CHAI AND TJHUNG’S PAPER [1]

In an interesting paper, Chai and Tjhung [1] proposed new unified probability density function (pdf) and cumulative distribution function (cdf) formulas based on the generalized Laguerre polynomial series expansion that cover a wide range of small-scale fading distributions in wireless communications. Many known Laguerre polynomial-series-based small-scale fading distributions are special cases of their results, which include the multiple-waves-plus-diffuse-power (MWDP) fading, $\kappa - \mu$ (noncentral chi), Nakagami- m , Rician (Nakagami- n), Nakagami- q (Hoyt), Rayleigh, Weibull, and $\alpha - \mu$ (Stacy) distributions.

First we present two typographical errors of [1]:

1. Chai and Tjhung stated that the orthogonality relation of generalized Laguerre polynomial is expressed as [1, Eq. (2)]

$$\int_0^{\infty} e^{-x} L_l^{\beta}(x) L_n^{\beta}(x) dx = \begin{cases} \Gamma(1 + \beta) \binom{n+\beta}{n}, & \text{if } n = l, \\ 0, & \text{if } n \neq l, \end{cases}$$

Copyright (c) 2010 IEEE. Personal use of this material is permitted. However, permission to use this material for any other purposes must be obtained from the IEEE by sending a request to pubs-permissions@ieee.org.

*Corresponding authors.

The work of Y. Sun and S. Zhou was supported by the National Basic Research Program of China under Grant 2007CB310608, National Natural Science Foundation of China under Grant 60832008 and Lab project from Tsinghua National Lab on Information Science and Technology (sub-project): Key technique for new distributed wireless communications system.

The work of Á. Baricz was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and by the Romanian National Authority for Scientific Research CNCSIS-UEFISCSU, project number PN-II-RU-PD 388/2010.

Y. Sun and S. Zhou are with the State Key Laboratory on Microwave and Digital Communications, Tsinghua National Laboratory for Information Science and Technology and Department of Electronic Engineering, Tsinghua University, Beijing, China. Address: Room 4-407, FIT Building, Tsinghua University, Beijing 100084, People’s Republic of China. e-mail: sunyin02@gmail.com, zhousd@tsinghua.edu.cn. Á. Baricz is with the Department of Economics, Babeş-Bolyai University, RO-400591, Cluj-Napoca, Romania. e-mail: bariczocsi@yahoo.com.

where L_n^{μ} is the generalized Laguerre polynomial, defined as [2, Eq. (22.3.9)]

$$L_n^{\mu}(x) = \frac{e^x x^{-\mu}}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\mu}) = \sum_{k=0}^n \binom{n+\mu}{n-k} \frac{(-x)^k}{k!},$$

$$\mu > -1, n = 0, 1, 2, \dots$$

However, there is a typographical error in [1, Eq. (2)]. The correct formula is

$$\int_0^{\infty} x^{\beta} e^{-x} L_l^{\beta}(x) L_n^{\beta}(x) dx = \begin{cases} \Gamma(1 + \beta) \binom{n+\beta}{n}, & \text{if } n = l, \\ 0, & \text{if } n \neq l. \end{cases}$$

2. The Laguerre polynomial expansion coefficients C_n for the pdf $f_X(x)$ of random variable X was determined by [1, Eq. (10)]. The fading envelop random variable R is assumed to satisfy

$$x = r^{\alpha}/b,$$

where α, b are parameters. Chai and Tjhung claimed that [1, Line -9 in p. 3990]

$$E_X [x^k] = E_R [r^{\alpha k}] / b,$$

which contains a small typographical error. The correct expression is given by

$$E_X [x^k] = E_R [r^{\alpha k}] / b^k.$$

Using this and [1, Eq. (10)], the coefficients C_n can be expressed by the αk th moment of the fading envelope R .

II. THE CORRECT UNIFIED CDF FORMULA

Chai and Tjhung presented an unified cdf formula for the fading envelope R in [1, Eq. (13)], i.e.,

$$\begin{aligned} F_R(R) &= \int_0^R f_R(r) dr \\ &= \frac{\alpha}{b^{\beta+1}} \sum_{n=0}^{\infty} C_n \frac{n!}{\Gamma(n + \beta + 1)} \\ &\quad \times \int_0^R \exp\left(-\frac{r^{\alpha}}{b}\right) r^{\alpha(\beta+1)-1} L_n^{\beta}\left(\frac{r^{\alpha}}{b}\right) dr \\ &= \frac{1}{n} \left(\frac{R^{\alpha}}{b}\right)^{\beta+1} \exp\left(-\frac{R^{\alpha}}{b}\right) \\ &\quad \times \sum_{n=0}^{\infty} C_n \frac{n!}{\Gamma(n + \beta + 1)} L_{n-1}^{\beta+1}\left(\frac{R^{\alpha}}{b}\right), \quad (1) \end{aligned}$$

where α, β, b are some parameters depending on the type of fading distribution and are presented in [1, Section V] for some special cases.

Unfortunately, the third line of the formula (1) is not valid, because $\frac{1}{n}$ is outside of the summation and the generalized Laguerre polynomial $L_{n-1}^{\beta+1}$ is not defined for $n = 0$. The correct formula of the unified cdf is

$$F_R(R) = \left(\frac{R^\alpha}{b}\right)^{\beta+1} \exp\left(-\frac{R^\alpha}{b}\right) \times \sum_{n=1}^{\infty} C_n \frac{\Gamma(n)}{\Gamma(n+\beta+1)} L_{n-1}^{\beta+1}\left(\frac{R^\alpha}{b}\right) + \frac{1}{\Gamma(\beta+1)} \gamma\left(\beta+1, \frac{R^\alpha}{b}\right), \quad (2)$$

where we used that $L_0^\mu(x) = 1$ and $C_0 = 1$. Here $\gamma(\cdot, \cdot)$ is the lower incomplete gamma function, defined by [2, Eq. (6.5.2)]

$$\gamma(a, x) = \int_0^x t^{a-1} e^{-t} dt.$$

Therefore, the unified Laguerre polynomial-series-based cdf contains one term expressed with the aid of the lower incomplete gamma function.

III. ONE SPECIAL CASE OF THE UNIFIED CDF FORMULA EQ. (2)

We now provide a Laguerre polynomial-series-based cdf formula for non-central chi-square distribution as a special case of Eq. (2). The pdf of non-central chi-square distribution $\chi_{\nu, \lambda}^2$ with ν degrees of freedom and non-centrality parameter λ is given by [3, Eq. (29.4)]

$$f_{\chi_{\nu, \lambda}^2}(r) = \frac{e^{-(r+\lambda)/2}}{2} \left(\frac{r}{\lambda}\right)^{\nu/4-1/2} I_{\nu/2-1}(\sqrt{\lambda r}),$$

which is equivalent to the pdf of $\kappa - \mu$ power distribution defined by [4, Eq. (2)]. Tiku proposed the Laguerre polynomial expansion of $f_{\chi_{\nu, \lambda}^2}(r)$ in 1965 [5], which is given by (see also [3, Eq. (29.11)])

$$f_{\chi_{\nu, \lambda}^2}(r) = \frac{e^{-r/2}}{2} \left(\frac{1}{2}r\right)^{\left(\frac{\nu}{2}-1\right)} \sum_{j=0}^{\infty} \frac{\left(-\frac{\lambda}{2}\right)^j}{\Gamma\left(\frac{1}{2}\nu + j\right)} L_j^{\frac{\nu}{2}-1}\left(\frac{1}{2}r\right), \quad \nu > 0, \lambda \geq 0. \quad (3)$$

Gideon and Gurland obtained a Laguerre polynomial expansion of the cdf of non-central chi-square distribution in 1977 [6] (see also [3, Eq. (29.13)] with the parameter set of L_0), i.e.,

$$F_{\chi_{\nu, \lambda}^2}(R) = \int_0^R f_{\chi_{\nu, \lambda}^2}(r) dr = \left(\frac{R}{2}\right)^{\frac{\nu}{2}} \exp\left(-\frac{R}{2}\right) \sum_{n=1}^{\infty} \frac{\left(-\frac{\lambda}{2}\right)^n}{\Gamma(n+\nu/2)n} L_{n-1}^{\frac{\nu}{2}}\left(\frac{R}{2}\right) + \frac{1}{\Gamma(\nu/2)} \gamma\left(\frac{\nu}{2}, \frac{R}{2}\right), \quad \nu > 0, \lambda \geq 0. \quad (4)$$

Some other Laguerre polynomial expansions of $F_{\chi_{\nu, \lambda}^2}(R)$ are available in [6] and [3, Eq. (29.13)]. We note that the pdf and cdf formulas in Eq. (3) and Eq. (4) can be derived by choosing $\alpha = 1$, $b = 2$, $\beta = \nu/2 - 1$ and $C_n = (-\lambda/2)^n/n!$ in [1, Eq. (11)] and Eq. (2) in previous section.

REFERENCES

- [1] C. C. Chai, T. T. Tjhung, "Unified Laguerre polynomial-series-based distribution of small-scale fading envelopes," *IEEE Trans. Veh. Technol.*, vol. 58, no. 8, pp. 3988-3999, Oct. 2009.
- [2] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 9th ed. New York, NY: Dover Press, 1972.
- [3] N. L. Johnson, S. Kotz, N. Balakrishnan, *Continuous Univariate Distributions*, vol. 2, second ed., John Wiley & Sons, Inc., New York, 1995.
- [4] M. D. Yacoub, "The $\kappa - \mu$ distribution and the $\eta - \mu$ distribution," *IEEE Antennas Propag. Mag.*, vol. 49, no. 1, pp. 68-81, Feb. 2007.
- [5] M. L. Tiku, "Laguerre series forms of non-central χ^2 and F distributions," *Biometrika*, vol. 52, no. 2-3, pp. 415-426, 1965.
- [6] R. A. Gideon and J. Gurland, "Some alternative expansions for the distribution function of a noncentral chi-square variable," *SIAM J. Math. Anal.*, vol. 8, pp. 100-110, 1977.