

**A STUDY ON THE  $q$ -EULER NUMBERS AND THE  
FERMIONIC  $q$ -INTEGRAL OF THE PRODUCT OF  
SEVERAL TYPE  $q$ -BERNSTEIN POLYNOMIALS ON  $\mathbb{Z}_p$**

TAEKYUN KIM

ABSTRACT. In this paper, we investigate some properties between  $q$ -Bernstein polynomials and  $q$ -Euler numbers by using fermionic  $p$ -adic  $q$ -integrals on  $\mathbb{Z}_p$ . From these properties, we derive some interesting identities related to the  $q$ -Euler numbers.

**§1. Introduction**

Let  $p$  be a fixed odd prime number. Throughout this paper  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , and  $\mathbb{C}_p$  will denote the ring of  $p$ -adic rational integers, the fields of  $p$ -adic rational numbers, and the completion of algebraic closure of  $\mathbb{Q}_p$ , respectively. Let  $v_p$  be the normalized exponential valuation of  $\mathbb{C}_p$  with  $|p|_p = p^{-v_p(p)} = \frac{1}{p}$ . When one talks of  $q$ -extension,  $q$  is variously considered as an indeterminate, a complex number  $q \in \mathbb{C}$  or  $p$ -adic number  $q \in \mathbb{C}_p$ . If  $q \in \mathbb{C}$ , one normally assumes  $|q| < 1$ , and if  $q \in \mathbb{C}_p$ , one normally assumes  $|1 - q|_p < 1$ . We use the notations of  $q$ -numbers as  $[x]_q = \frac{1-q^x}{1-q}$  and  $[x]_{-q} = \frac{1-(-q)^x}{1+q}$ . Let  $C[0, 1]$  denote the set of continuous function on  $[0, 1]$ . For  $f \in C[0, 1]$ ,  $q$ -Bernstein operator is introduced in [2] as follows:

$$(1) \quad \mathbb{B}_{n,q}(f|x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} [x]_q^k [1-x]_{\frac{1}{q}}^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x, q).$$

Here  $\mathbb{B}_{n,q}(f|x)$  is called  $q$ -Bernstein operator of order  $n$  for  $f$ . For  $k, n \in \mathbb{Z}_+(= \mathbb{N} \cup \{0\})$ , the  $q$ -Bernstein polynomials of degree  $n$  is defined by

$$(2) \quad B_{k,n}(x, q) = \binom{n}{k} [x]_q^k [1-x]_{\frac{1}{q}}^{n-k}, \text{ (see [2]).}$$

---

*Key words and phrases.*  $q$ -Bernstein polynomials,  $q$ -Euler numbers,  $q$ -Stirling numbers, fermionic  $p$ -adic integrals.

2000 AMS Subject Classification: 11B68, 11S80, 60C05, 05A30

The present Research has been conducted by the research Grant of Kwangwoon University in 2010

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

Let  $UD(\mathbb{Z}_p)$  be the space of uniformly differentiable functions on  $\mathbb{Z}_p$ . For  $f \in UD(\mathbb{Z}_p)$ , the fermionic  $p$ -adic  $q$ -integrals on  $\mathbb{Z}_p$  is defined by

$$(3) \quad I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \text{ (see [3, 4, 5]).}$$

As well known definition, Euler polynomials are defined by

$$(4) \quad \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \text{ (see [1-11]).}$$

In the special case,  $x = 0$ ,  $E_n(0) = E_n$  are called the  $n$ -th Euler numbers. By (4), we see that the recurrence formula of Euler numbers is given by

$$(5) \quad E_0 = 1, \text{ and } (E + 1)^n + E_n = 0, \text{ if } n > 0, \text{ (see [2, 4, 5]),}$$

with the usual convention of replacing  $E^n$  by  $E_n$ . When one talks of  $q$ -analogue,  $q$  is variously considered as an indeterminate, a complex number  $q \in \mathbb{C}$ , or a  $p$ -adic number  $q \in \mathbb{C}_p$ . If  $q \in \mathbb{C}$ , we normally assume  $|q| < 1$ . If  $q \in \mathbb{C}_p$ , we normally always assume that  $|1 - q|_p < 1$ . As the  $q$ -extension of (5), the  $q$ -Euler numbers are defined by

$$(6) \quad \xi_{0,q} = 1, \text{ and } q(q\xi + 1)^n + \xi_{n,q} = 0, \text{ if } n > 0, \text{ (see [5]),}$$

with the usual convention of replacing  $\xi^n$  by  $\xi_{n,q}$ . The  $q$ -Euler polynomials are also defined by

$$(7) \quad \xi_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} [x]_q^{n-l} q^{lx} \xi_{l,q}, \text{ for } n \in \mathbb{Z}_+, \text{ (see [5]).}$$

By (6) and (7), we easily get

$$(8) \quad q\xi_{n,q}(1) + \xi_{n,q} = 1, \text{ if } n > 0, \text{ (see [4, 5]).}$$

In [5], Witt's formula for the  $q$ -Euler polynomial is given by

$$(9) \quad \xi_{n,q}(x) = \int_{\mathbb{Z}_p} [y + x]_q^n d\mu_{-1}(y) = \frac{[2]_q}{(1 - q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{q^{lx}}{1 + q^l}, \text{ for } n \geq 0.$$

By (9), we easily see that

$$\xi_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} q^{lx} \xi_{l,q} = (q^x \xi + 1)^n.$$

In this paper, we investigate some properties between  $q$ -Bernstein polynomials and  $q$ -Euler numbers by using fermionic  $p$ -adic  $q$ -integrals on  $\mathbb{Z}_p$ . From these properties, we derive some interesting identities related to the  $q$ -Euler numbers.

## §2. $q$ -Euler numbers and $q$ -Bernstein Polynomials

In this section we assume that  $q \in \mathbb{C}_p$  with  $|1 - q|_p < 1$ . From (6), (7) and (9), we note that

$$\begin{aligned}
(10) \quad q^2 \xi_{k,q}(2) - q - q^2 &= q^2 (q(q\xi + 1) + 1)^k - q - q^2 \\
&= q^2 \left( 1 + \sum_{l=1}^k \binom{k}{l} q^l (q\xi + 1)^l \right) - q - q^2 \\
&= -q \sum_{l=1}^k \binom{k}{l} q^l \xi_{l,q} - q = -q(q\xi + 1)^k = \xi_{k,q}, \text{ if } k > 0.
\end{aligned}$$

Therefore, we obtain the following theorem.

**Theorem 1.** *For  $n \in \mathbb{N}$ , we have*

$$\xi_{n,q}(2) = 1 + \frac{1}{q} + \frac{1}{q^2} \xi_{n,q}.$$

By (9), we easily get

$$\int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) = \xi_{n,q}(0) = \xi_{n,q} = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^l}, \text{ for } n \geq 0,$$

and

$$(11) \quad \xi_{n,q}(x) = \int_{\mathbb{Z}_p} [x+y]_q^n d\mu_{-q}(y) = \sum_{l=0}^n \binom{n}{l} q^{lx} \int_{\mathbb{Z}_p} [y]_q^l d\mu_{-q}(y) = \sum_{l=0}^n \binom{n}{l} [x]_q^{n-l} q^{lx} \xi_{l,q}.$$

It is easy to show that  $[1-x]_q^n = (-1)^n q^n [x-1]_q^n$ . Thus, we have

$$(12) \quad \int_{\mathbb{Z}_p} [1-x]_q^n d\mu_{-q}(x) = (-1)^n q^n \int_{\mathbb{Z}_p} [x-1]_q^n d\mu_{-q}(x) = (-1)^n q^n \xi_{n,q}(-1).$$

By (11) and (12), we obtain the following proposition.

**Proposition 2.** For  $n \in \mathbb{Z}_+$ , we have

$$\int_{\mathbb{Z}_p} [1-x]_{\frac{1}{q}}^n d\mu_{-q}(x) = \sum_{l=0}^n \binom{n}{l} (-1)^l \xi_{l,q} = (-1)^n q^n \xi_{n,q}(-1).$$

From (3), we can derive the following equation (13).

$$(13) \quad \int_{\mathbb{Z}_p} [1-x+x_1]_{\frac{1}{q}}^n d\mu_{-\frac{1}{q}}(x_1) = (-1)^n q^n \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{q^{lx}}{1+q^{l+1}}.$$

By (9) and (13), we obtain the following proposition.

**Proposition 3.** For  $n \in \mathbb{Z}_+$ , we have

$$(14) \quad \int_{\mathbb{Z}_p} [1-x+x_1]_{\frac{1}{q}}^n d\mu_{-\frac{1}{q}}(x_1) = (-1)^n q^n \int_{\mathbb{Z}_p} [x+x_1]_q d\mu_{-q}(x_1).$$

By (12) and (14), we get

$$(15) \quad q^n (-1)^n \xi_{n,q}(-1) = (-1)^n q^n \int_{\mathbb{Z}_p} [x_1-1]_q^n d\mu_{-q}(x_1) = \int_{\mathbb{Z}_p} [2+x_1]_{\frac{1}{q}}^n d\mu_{-\frac{1}{q}}(x_1) = \xi_{n,\frac{1}{q}}(2).$$

Therefore, by (15), we obtain the following corollary.

**Corollary 4.** For  $n \geq 0$ , we have

$$\xi_{n,\frac{1}{q}}(2) = (-1)^n q^n \xi_{n,q}(-1).$$

From (12) and Corollary 4, we have

$$(16) \quad \int_{\mathbb{Z}_p} [1-x]_{\frac{1}{q}}^n d\mu_{-q}(x) = (-1)^n q^n \xi_{n,q}(-1) = \xi_{n,\frac{1}{q}}(2).$$

By (12), (16) and Theorem 1, we obtain the following theorem.

**Theorem 5.** For  $n \in \mathbb{N}$ , we have

$$\int_{\mathbb{Z}_p} [1-x]_{\frac{1}{q}}^n d\mu_{-q}(x) = q^2 \int_{\mathbb{Z}_p} [x]_{\frac{1}{q}}^n d\mu_{-\frac{1}{q}}(x) + q + 1 = [2]_q + q^2 \xi_{n,\frac{1}{q}}.$$

Now we consider the  $p$ -adic analogue of (1) and (2) on  $\mathbb{Z}_p$ . For  $x \in \mathbb{Z}_p$  and  $f \in UD(\mathbb{Z}_p)$ , we define  $p$ -adic  $q$ -Bernstein operator of order  $n$  for  $f$  as follows:

$$\mathbb{B}_{n,q}(f|x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} [x]_q^k [1-x]_{\frac{1}{q}}^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x, q).$$

For  $n, k \in \mathbb{Z}_+$ , the  $p$ -adic  $q$ -Bernstein polynomials of degree  $n$  is defined by

$$(17) \quad B_{k,n}(x, q) = \binom{n}{k} [x]_q^k [1-x]_{\frac{1}{q}}^{n-k}, \text{ where } x \in \mathbb{Z}_p.$$

Taking the fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  for one  $q$ -Bernstein polynomials in (17), we get

$$(18) \quad \begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) d\mu_{-q}(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} [x]_q^k [1-x]_{\frac{1}{q}}^{n-k} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_q^{k+l} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \xi_{k+l, q}. \end{aligned}$$

From (17), we note that

$$(19) \quad B_{k,n}(x, q) = B_{n-k, n}(1-x, \frac{1}{q}), \text{ where } n, k \in \mathbb{Z}_+ \text{ and } x \in \mathbb{Z}_p.$$

By (19), we get

$$(20) \quad \begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) d\mu_{-q}(x) &= \int_{\mathbb{Z}_p} B_{n-k, n}(1-x, \frac{1}{q}) d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} [1-x]_{\frac{1}{q}}^{n-l} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \xi_{n-l, \frac{1}{q}}(2). \end{aligned}$$

By comparing the coefficients on the both sides of (18) and (20), we obtain the following theorem.

**Theorem 6.** For  $n, k \in \mathbb{Z}_+$  with  $n \geq k$ , we have

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \xi_{k+l, q} = \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \xi_{n-l, \frac{1}{q}}(2).$$

By Theorem 1 and Theorem 6, we obtain the following corollary.

**Corollary 7.** Let  $n, k \in \mathbb{Z}_+$  with  $n > k$ . Then we have

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \xi_{k+l, q} = q^2 \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \xi_{n-l, \frac{1}{q}}, \text{ if } k > 0.$$

Moreover,

$$\sum_{l=0}^n \binom{n}{l} (-1)^l \xi_{l, q} = [2]_q + q^2 \xi_{n, \frac{1}{q}}.$$

For  $m, n, k \in \mathbb{Z}_+$  with  $m, n > k$ , we consider the fermionic  $p$ -adic  $q$ -integral for multiplication of two  $q$ -Bernstein polynomials on  $\mathbb{Z}_p$ .

$$\begin{aligned} & \int_{\mathbb{Z}_p} B_{k,n}(x, q) B_{k,m}(x, q) d\mu_{-q}(x) = \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} [x]_q^{2k} [1-x]_{\frac{1}{q}}^{n+m-2k} d\mu_{-q}(x) \\ (21) \quad & = \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \int_{\mathbb{Z}_p} [1-x]_{\frac{1}{q}}^{n+m-l} d\mu_{-q}(x) \\ & = \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \xi_{n+m-l, \frac{1}{q}}(2). \end{aligned}$$

From (21) and Theorem 1, we can derive the following equation (22).

$$\begin{aligned} (22) \quad & \int_{\mathbb{Z}_p} B_{k,n}(x, q) B_{k,m}(x, q) d\mu_{-q}(x) = \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \xi_{n+m-l, \frac{1}{q}}(2) \\ & = \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} ([2]_q + q^2 \xi_{n+m-l, \frac{1}{q}}). \end{aligned}$$

For  $n, m, k \in \mathbb{Z}_+$ , it is easy to show that

$$(23) \quad \int_{\mathbb{Z}_p} B_{k,n}(x, q) B_{k,m}(x, q) d\mu_{-q}(x) = \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \xi_{l+2k, q}.$$

Therefore, by (22) and (23), we obtain the following theorem.

**Theorem 8.** Let  $m, n, k \in \mathbb{Z}_+$  with  $m+n > 2k$ . Then we have

$$\sum_{l=0}^{m+n-2k} \binom{n+m-2k}{l} (-1)^l \xi_{l+2k, q} = q^2 \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \xi_{n+m-l, \frac{1}{q}}, \text{ if } k > 0.$$

Moreover,

$$\sum_{l=0}^{m+n} \binom{n+m}{l} (-1)^l \xi_{l, q} = q^2 \xi_{n+m, \frac{1}{q}} + [2]_q.$$

By induction hypothesis, we obtain the following theorem.

**Theorem 9.** Let  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  ( $s \in \mathbb{N}$ ) with  $n_1 + n_2 + \dots + n_s > sk$ . Then we have

$$\begin{aligned} & \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \xi_{l+sk, q} \\ &= q^2 \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk-l} \xi_{n_1+\dots+n_s-l, \frac{1}{q}}, \text{ if } k > 0. \end{aligned}$$

Moreover,

$$\sum_{l=0}^{n_1+\dots+n_s} \binom{n_1+\dots+n_s}{l} (-1)^l \xi_{l, q} = [2]_q + \xi_{n_1+\dots+n_s, \frac{1}{q}}.$$

## REFERENCES

- [1] M. Acikgoz, S. Araci, *A study on the integral of the product of several type Bernstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, (2010).
- [2] T. Kim, *A note on  $q$ -Bernstein polynomials*, Russ. J. Math. Phys( accepted).
- [3] T. Kim, *Barnes type multiple  $q$ -zeta function and  $q$ -Euler polynomials*, J. Phys. A: Math. Theor. **43** (2010), 255201, 11pp.
- [4] T. Kim, *New approach to  $q$ -Euler polynomials of higher order*, Russ. J. Math. Phys. **17** (2010), 218-225.
- [5] T. Kim,  *$q$ -Euler numbers and polynomials associated with  $p$ -adic  $q$ -integral*, J. Nonlinear Math. Phys. **14** (2007), 15-27.
- [6] V. Gupta, T. Kim, J. Choi, Y.-H. Kim, *Generating function for  $q$ -Bernstein,  $q$ -Meyer-König-Zeller and  $q$ -Beta basis*, Automation Computers Applied Mathematics **19** (2010), 7-11.
- [7] L.C. Jang, W.-J. Kim, Y. Simsek, *A study on the  $p$ -adic integral representation on  $\mathbb{Z}_p$  associated with Bernstein and Bernoulli polynomials*, Advances in Difference Equations **2010** (2010), Article ID 163217, 6pp.
- [8] Y. Simsek, M. Acikgoz, *A new generating function of  $q$ -Bernstein-type polynomials and their interpolation function*, Abstract and Applied Analysis **2010** (2010), Article ID 769095, 12pp.
- [9] T. Kim, L. C. Jang, H. Yi, *A note on the modified  $q$ -Bernstein polynomials*, Discrete Dynamics in Nature and society **2010** (2010), Article ID 706483, 13pp.
- [10] B. A. Kupershmidt, *Reflection symmetries of  $q$ -Bernoulli polynomials*, J. Nonlinear Math. Phys. **12** (2005), 412-422.
- [11] T. Kim, *Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -stirling numbers by the fermionic  $p$ -adic integrals on  $\mathbb{Z}_p$* , Russ. J. Math. Phys. **16** (2009), 484-491.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University, Seoul  
139-701, S. Korea e-mail: tkkim@kw.ac.kr