

The Shape Parameter in the Shifted Surface Spline

Lin-Tian Luh

Department of Mathematics, Providence University

Shalu, Taichung

email:ltluh@pu.edu.tw

phone:(04)26328001 ext. 15126

fax:(04)26324653

October 28, 2018

Abstract. The purpose of this article is to explore the optimal choice of shape parameter which is an important and longstanding problem in the theory of radial basis functions(RBF). We already handled it for multiquadric and Gaussian in [9, 10, 11, 12, 13]. Here we focus on shifted surface spline and present concrete criteria for the choice of shape parameter.

key words: radial basis function, shifted surface spline, shape parameter.

AMS subject classification: 41A05, 41A15, 41A25, 41A30, 41A63, 65D10

1 Introduction

The so-called shifted surface spline is defined by

$$h(x) := (-1)^m (|x|^2 + c^2)^{\frac{\lambda}{2}} \log(|x|^2 + c^2)^{\frac{1}{2}}, \quad \lambda \in Z_+, \quad m = 1 + \frac{\lambda}{2}, \quad c > 0, \quad x \in R^n, \quad \lambda, \quad n \text{ even} \quad (1)$$

, and

$$h(x) := (-1)^{\lceil \lambda - \frac{n}{2} \rceil} (|x|^2 + c^2)^{\lambda - \frac{n}{2}}, \quad \lambda \in Z_+ = \{1, 2, 3, \dots\}, \quad n \text{ odd and } \lambda > \frac{n}{2} \quad (2)$$

, where $|x|$ is the Euclidean norm of x , and λ, c are constants. The number $\lceil \lambda - \frac{n}{2} \rceil$ denotes the smallest integer greater than or equal to $\lambda - \frac{n}{2}$.

This function was introduced by Dyn, Levin and Rippa in [4] for $x \in R^2$. Then it's extended to R^n for $n \geq 1$. For further details we refer the reader to [2, 3, 5, 18, 19, 21, 22, 23, 24].

Note that (2) is just multiquadric and will not be explored here. In this paper we only deal with (1). As is well known, $h(x)$ is conditionally positive definite of order $m = 1 + \frac{\lambda}{2}$ where λ is of our choice. The constant c is just the so-called shape parameter whose optimal choice is of our primary concern.

Since $h(x)$ is c.p.d.(conditionally positive definite) of order m , there exists a unique function $s(x)$ of the form

$$s(x) = p(x) + \sum_{j=1}^N c_j h(x - x_j) \quad (3)$$

which interpolates any scattered set of data points $(x_1, y_1), \dots, (x_N, y_N)$ as long as $\{x_1, \dots, x_N\}$ is polynomial-nondegenerate in R^n . Here $p(x) \in P_{m-1}$ is a polynomial of degree $\leq m-1$ and c'_j s are chosen so that

$$\sum_{j=1}^N c_j q(x_j) = 0$$

for all polynomials q in P_{m-1} .

We will show that whenever x and the data points are fixed, the upper bound of $|f(x) - s(x)|$ will be made minimal by a good choice of c . Here $f(x)$ denotes the underlying function of the data points.

1.1 Basic Ingredients

Definition 1.1 For $n = 1, 2, \dots$, the integers γ_n are defined by $\gamma_1 = 2$ and $\gamma_n = 2n(1 + \gamma_{n-1})$ for $n > 1$.

The numbers γ_n will appear in our criteria.

In this paper all approximated functions belong to a semi-Hilbert space $\mathcal{C}_{h,m}(R^n)$, abbreviated as $\mathcal{C}_{h,m}$, where m denotes the order of conditional positive definiteness of the function h . There is a semi-norm on $\mathcal{C}_{h,m}$, denoted by $\|\cdot\|_h$. The definition and characterization of this space can be found in [7, 8, 15, 16]. We will not repeat them. A subspace of $\mathcal{C}_{h,m}$, denoted by B_σ , is of our special concern and is defined as follows.

Definition 1.2 For any $\sigma > 0$,

$$B_\sigma := \{f \in L^2(R^n) : \hat{f}(\xi) = 0 \text{ for all } |\xi| > \sigma\}$$

is the set of band-limited functions, where \hat{f} denotes the Fourier transform of f .

We need some constants as follows.

Definition 1.3 Let n, λ and m be as in (1). The constants ρ and Δ_0 are defined according to the following three cases.

(a) Suppose $n - \lambda > 3$. Let $s = \lceil \frac{n-\lambda-3}{2} \rceil$. Then

$$\rho = 1 + \frac{s}{2m+3} \text{ and } \Delta_0 = \frac{(2m+2+s)(2m+1+s) \cdots (2m+3)}{\rho^{2m+2}}.$$

(b) Suppose $n - \lambda \leq 1$. Let $s = -\lceil \frac{n-\lambda-3}{2} \rceil$. Then

$$\rho = 1 \text{ and } \Delta_0 = \frac{1}{(2m+2)(2m+1) \cdots (2m-s+3)}.$$

(c) Suppose $1 < n - \lambda \leq 3$. Then

$$\rho = 1 \text{ and } \Delta_0 = 1.$$

Finally, the Fourier transform [6] of (1) should also be introduced. It's of the form

$$\hat{h}(\xi) = l(\lambda, n) |\xi|^{-\lambda-n} \tilde{\mathcal{K}}_{\frac{n+\lambda}{2}}(c|\xi|) \quad (4)$$

where $l(\lambda, n) = (2\pi)^{-\frac{n}{2}} \cdot 2^{\frac{\lambda}{2}} \cdot (\frac{\lambda}{2}!)$ [16] and $\tilde{\mathcal{K}}_\nu(t) = t^\nu \mathcal{K}_\nu(t)$, $\mathcal{K}_\nu(t)$ being the modified Bessel function of the second kind [1].

2 Fundamental Theory

The cornerstone of our theory is the exponential-type error bound raised by the author in [14]. We cite it directly with a slight modification to make it easier to understand.

In the following theorem

$$d = d(E, X) := \sup_{y \in E} \inf_{x \in X} |y - x|$$

denotes fill distance.

Theorem 2.1 *Suppose h is defined as in (1). Then, given any positive number b_0 , there are positive constants d_0 and ω , $0 < \omega < 1$, which depend on b_0 , for which the following is: If $f \in \mathcal{C}_{h,m}$ and s is the h spline defined in (3) that interpolates f on a subset X of R^n , then*

$$|f(x) - s(x)| \leq \sqrt{l(\lambda, n)} \cdot (2\pi)^{\frac{1}{4}} \cdot \sqrt{n\alpha_n} \cdot \sqrt{\Delta_0} \cdot c^{\frac{\lambda}{2}} \cdot \omega^{\frac{1}{d}} \cdot \|f\|_h \quad (5)$$

holds for all x in a cube $E \subseteq \Omega$, where Ω is a set which can be expressed as the union of rotations and translations of a fixed cube of side b_0 , provided that (a) E has side $b \geq b_0$ and (b) $0 < d \leq d_0$. Here, α_n denotes the volume of the unit ball in R^n , c was defined in (1) and Δ_0 was defined in Definition 1.3.

The numbers d_0 and ω can be expressed specifically as

$$d_0 = \frac{1}{6C\gamma_n(m+1)}, \quad \omega = \left(\frac{2}{3}\right)^{\frac{1}{6c\gamma_n}}$$

where

$$C = \max \left\{ 2\rho' \sqrt{n} e^{2n\gamma_n}, \frac{2}{3b_0} \right\}, \quad \rho' = \frac{\rho}{c}.$$

The numbers γ_n and ρ were defined in Definition 1.1 and 1.3, respectively, and $m = 1 + \frac{\lambda}{2}$ as in (1).

Remark: (a)The seemingly complicated theorem is in fact not difficult to understand. We expressed each constant specifically for later use. (b)The set X of centers is usually contained in the cube E . However it's harmless to extend it to a more general form as a subset of R^n . (c) $\|f\|_h$ is the semi-norm of f mentioned in subsection 1.1.

On the right-hand side of the inequality (5), the only things dependent of the shape parameter c are $c^{\frac{\lambda}{2}}$, $\omega^{\frac{1}{d}}$ and $\|f\|_h$. Also, d_0 is dependent of c . It's not hard to imagine a good choice of c will minimize the error bound (5). However it's nontrivial to express this error bound by an explicit function of c , and clarify the relation between d_0 and c . Further treatment of (5) is necessary.

Lemma 2.2 *Let $\sigma > 0$ and h be as in (1). For any $f \in B_\sigma$,*

$$\|f\|_h \leq C_0(m, n) \cdot \left(\frac{2}{\pi}\right)^{\frac{1}{4}} \cdot \sigma^{\frac{1+n+\lambda}{4}} \cdot c^{\frac{1-n-\lambda}{4}} \cdot e^{\frac{c\sigma}{2}} \cdot \|f\|_{L^2} \quad (6)$$

, where

$$C_0(m, n) = \frac{(2\pi)^{-n} \sqrt{m!}}{\sqrt{l(\lambda, n)}}.$$

Proof. By Corollary3.3 of [16],

$$\begin{aligned}
\|f\|_h &= \left\{ \sum_{|\alpha|=m} \frac{m!}{\alpha!} \|(D^\alpha f)^\wedge\|_{L^2(\rho)}^2 \right\}^{1/2} \\
&= \left\{ \sum_{|\alpha|=m} \frac{m!}{\alpha!} \int |(D^\alpha f)^\wedge(\xi)|^2 \cdot \frac{1}{(2\pi)^{2n} |\xi|^{2m} \hat{h}(\xi)} d\xi \right\}^{1/2} \\
&= (2\pi)^{-n} \left\{ \sum_{|\alpha|=m} \frac{m!}{\alpha!} \int \frac{\xi^{2\alpha} |\hat{f}(\xi)|^2}{|\xi|^{2m} l(\lambda, n) |\xi|^{-\lambda-n} \tilde{\mathcal{K}}_{\frac{n+\lambda}{2}}(c|\xi|)} d\xi \right\}^{1/2} \\
&\leq \frac{(2\pi)^{-n} \sqrt{m!}}{\sqrt{l(\lambda, n)}} \cdot \left\{ \int \frac{|\hat{f}(\xi)|^2}{|\xi|^{-\lambda-n} \tilde{\mathcal{K}}_{\frac{n+\lambda}{2}}(c|\xi|)} d\xi \right\}^{1/2} \\
&= \frac{(2\pi)^{-n} \sqrt{m!}}{\sqrt{l(\lambda, n)}} \cdot \left\{ \int \frac{|\hat{f}(\xi)|^2}{|\xi|^{-\lambda-n} (c|\xi|)^{\frac{n+\lambda}{2}} \mathcal{K}_{\frac{n+\lambda}{2}}(c|\xi|)} d\xi \right\}^{1/2} \\
&\leq C_0(m, n) c^{-\frac{n-\lambda}{4}} \left\{ \int |\hat{f}(\xi)|^2 |\xi|^{\frac{n+\lambda}{2}} \cdot \frac{1}{\sqrt{\frac{\pi}{2}} \cdot \frac{e^{-c|\xi|}}{\sqrt{c|\xi|}}} d\xi \right\}^{1/2} \quad \text{by [20] where} \\
C_0(m, n) &:= \frac{(2\pi)^{-n} \sqrt{m!}}{\sqrt{l(\lambda, n)}} \\
&= C_0(m, n) \left(\frac{2}{\pi} \right)^{\frac{1}{4}} c^{\frac{1-n-\lambda}{4}} \left\{ \int |\hat{f}(\xi)|^2 |\xi|^{\frac{1+n+\lambda}{2}} e^{c|\xi|} d\xi \right\}^{1/2}.
\end{aligned}$$

If $f \in B_\sigma$, then

$$\|f\|_h \leq C_0(m, n) \left(\frac{2}{\pi} \right)^{\frac{1}{4}} \sigma^{\frac{1+n+\lambda}{4}} c^{\frac{1-n-\lambda}{4}} e^{\frac{c\sigma}{2}} \|f\|_{L^2}.$$

‡

Corollary 2.3 Let $\sigma > 0$. If $f \in B_\sigma$, then (5) can be transformed into

$$|f(x) - s(x)| \leq \sqrt{m!} (2\pi)^{-n} \sqrt{2n\alpha_n} \sigma^{\frac{1+n+\lambda}{4}} \sqrt{\Delta_0} c^{\frac{1-n-\lambda}{4}} e^{\frac{c\sigma}{2}} \omega^{\frac{1}{d}} \|f\|_{L^2}. \quad (7)$$

Proof. $B_\sigma \subseteq \mathcal{C}_{h,m}$ is a simple result of Corollary3.3 of [16]. Now (7) is just a combination of (5) and (6). ‡

On the right-hand side of (7), the only things dependent of c are $c^{\frac{1-n+\lambda}{4}}$, $e^{\frac{c\sigma}{2}}$ and $\omega^{\frac{1}{d}}$, where d denotes fill distance. It seems that one can abstract a function of c from the long expression of (7) after every thing independent of c is fixed, including d .

However as c changes, the upper bound d_0 of d changes also. As required by Theorem2.1, $d \leq d_0$ where

$$d_0 = \frac{1}{6C\gamma_n(m+1)}.$$

The number $C \rightarrow \infty$, and hence $d_0 \rightarrow 0$, as $c \rightarrow 0^+$. In order to keep $d \leq d_0$, the minimal possible choice of c is

$$c_0 := 12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n(m+1)d \quad (8)$$

where d is fixed. Therefore, we can choose c from the interval $[c_0, \infty)$ only.

There is a logical problem about d , d_0 , c and b_0 . Before c and b_0 are given, d_0 is unknown and we do not know whether d is eligible. This question is not difficult to resolve. For any $b_0 > 0$, we require $d < \frac{1}{6C\gamma_n(m+1)}$ where we let $C = \frac{2}{3b_0}$ temporarily. Then $d < \frac{b_0}{4\gamma_n(m+1)}$. Note that $2\rho'\sqrt{n}e^{2n\gamma_n} = \frac{2}{3b_0}$ iff $c = 3b_0\rho\sqrt{n}e^{2n\gamma_n}$. With the same b_0 and $c \geq 3b_0\rho\sqrt{n}e^{2n\gamma_n}$, we have

$$C = \max \left\{ 2\rho'\sqrt{n}e^{2n\gamma_n}, \frac{2}{3b_0} \right\} = \frac{2}{3b_0}$$

and

$$d_0 = \frac{b_0}{4\gamma_n(m+1)}.$$

Then $d < d_0$ is satisfied.

With the same b_0 and $c < 3b_0\rho\sqrt{n}e^{2n\gamma_n}$, we have

$$C = \max \left\{ 2\rho'\sqrt{n}e^{2n\gamma_n}, \frac{2}{3b_0} \right\} = 2\rho'\sqrt{n}e^{2n\gamma_n}$$

and

$$d_0 = \frac{1}{6C\gamma_n(m+1)} = \frac{c}{12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n(m+1)}.$$

In order to keep $d < d_0$, we require

$$c \geq c_0 := 12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n(m+1)d$$

where d was given above satisfying $d < \frac{b_0}{4\gamma_n(m+1)}$. Therefore, once the cube side b_0 is given, we first choose $d < \frac{b_0}{4\gamma_n(m+1)}$. Then put the restriction $c \in [c_0, \infty)$ where $c_0 = 12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n(m+1)d$. The condition $d < d_0$ will always be satisfied.

Theoretically d , and hence c_0 , can be arbitrarily small. However practically the problem of ill-conditioning when constructing $s(x)$ has to be considered. In this paper we explore theoretically the optimal choice of c for $c_0 \leq c < \infty$.

3 Criteria of Choosing c

In the long expression of (7) the only things dependent of c are $c^{\frac{1-n+\lambda}{4}}$, $e^{\frac{c\sigma}{2}}$ and $\omega^{\frac{1}{d}}$. Let's define

$$MN(c) := c^{\frac{1-n+\lambda}{4}} \cdot e^{\frac{c\sigma}{2}} \cdot \omega^{\frac{1}{d}} \quad (9)$$

and call it an MN function. The optimal choice of c will then be the number minimizing $MN(c)$, when every thing independent of c is fixed.

The value of ω highly depends on the cube side b_0 . Theoretically, $\omega \rightarrow 0$ and is very influential as $c \rightarrow \infty$ and $b_0 \rightarrow \infty$ simultaneously. However, a lot of time b_0 is fixed and cannot approach ∞ . Therefore we divide our criteria into two classes.

3.1 b_0 fixed

Recall that $C = \max\{2\rho'\sqrt{n}e^{2n\gamma_n}, \frac{2}{3b_0}\}$ and $\rho' = \frac{\rho}{c}$. The two values $2\rho'\sqrt{n}e^{2n\gamma_n}$ and $\frac{2}{3b_0}$ are equal if and only if

$$c = 3b_0\rho\sqrt{n}e^{2n\gamma_n} =: c_1 \quad (10)$$

Then, for $c \in [c_0, c_1)$, $C = 2\rho'\sqrt{n}e^{2n\gamma_n}$. For $c \in [c_1, \infty)$, $C = \frac{2}{3b_0}$. So we have

$$\begin{aligned} \omega^{\frac{1}{d}} &= \left(\frac{2}{3}\right)^{\frac{1}{6C\gamma_n d}} \\ &= \begin{cases} \left(\frac{2}{3}\right)^{\frac{c}{12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n d}} & \text{if } c \in [c_0, c_1), \\ \left(\frac{2}{3}\right)^{\frac{b_0}{4\gamma_n d}} & \text{if } c \in [c_1, \infty) \end{cases} \\ &= \begin{cases} e^{\frac{c \cdot \ln \frac{2}{3}}{12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n d}} & \text{if } c \in [c_0, c_1), \\ e^{\frac{b_0 \ln \frac{2}{3}}{4\gamma_n d}} & \text{if } c \in [c_1, \infty). \end{cases} \end{aligned}$$

Putting this result into $MN(c)$, we thus have

$$MN(c) = \begin{cases} c^{\frac{1-n+\lambda}{4}} \cdot e^{c\left(\frac{\sigma}{2} + \frac{\ln \frac{2}{3}}{12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n d}\right)} & \text{if } c \in [c_0, c_1), \\ c^{\frac{1-n+\lambda}{4}} \cdot e^{\frac{c\sigma}{2}} \cdot e^{\frac{b_0 \ln \frac{2}{3}}{4\gamma_n d}} & \text{if } c \in [c_1, \infty) \end{cases} \quad (11)$$

which is a continuous function. Our goal is to find c which minimizes $MN(c)$.

As for the fill distance d , we require $d < \frac{b_0}{4\gamma_n(m+1)}$ once the cube side b_0 is given, as mentioned in the end of section 2.

We have the following cases where n , λ , m , σ , γ_n , ρ and d were defined in (1), Definition 1.1, 1.2, 1.3 and Theorem 2.1. Moreover we introduce a constant

$$k := \frac{\sigma}{2} + \frac{\ln \frac{2}{3}}{12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n d}.$$

Case1. $\boxed{1-n+\lambda > 0}$ and $\boxed{k \geq 0}$ Let $f \in B_\sigma$ be the approximated function, $b_0 > 0$ be the cube side as in Theorem 2.1, and $d < \frac{b_0}{4\gamma_n(m+1)}$ be the fill distance. If $1-n+\lambda > 0$ and $k \geq 0$, the optimal choice of c for $c \in [c_0, \infty)$ is $\boxed{c = 12\rho\sqrt{n}e^{2n\gamma_n}\gamma_n(m+1)d}$.

Reason: In this case $MN(c)$ is an increasing function in its domain $[c_0, \infty)$. We therefore choose $c = c_0$, defined in (8), which minimizes $MN(c)$ in (11) and the error bound in (7).

Case2. $\boxed{1-n+\lambda > 0}$ and $\boxed{k < 0}$ Let $f \in B_\sigma$ be the approximated function, $b_0 > 0$ be the cube side as in Theorem 2.1, and $d < \frac{b_0}{4\gamma_n(m+1)}$ be the fill distance. If $1-n+\lambda > 0$ and $k < 0$, the optimal choice of c for $c \in [c_0, \infty)$ is the value minimizing $MN(c)$ over the interval $[c_0, c_1]$ where c_1 was defined in (10).

Reason. In this case $MN(c)$ is increasing on $[c_1, \infty)$. Therefore the minimum value of $MN(c)$ in $[c_0, \infty)$ must happen in $[c_0, c_1]$.

Remark:(a) This case rarely happens because k is usually positive. (b) In fact, the optimal c in Case2 can be obtained exactly. Let $g(c) := c^{\frac{1-n+\lambda}{4}} e^{ck}$ which is just $MN(c)$ on $[c_0, c_1]$. Then $g(c)$ is increasing on $(0, \frac{1-n+\lambda}{-4k}]$ and decreasing on $[\frac{1-n+\lambda}{-4k}, \infty)$. So the optimal c is (i) c_0 if $c_1 \leq \frac{1-n+\lambda}{-4k}$, (ii) c_1 if $\frac{1-n+\lambda}{-4k} \leq c_0$, and (iii) c_1 if $g(c_1) \leq g(c_0)$, and c_0 if $g(c_0) \leq g(c_1)$, for $c_0 < \frac{1-n+\lambda}{-4k} < c_1$.

Case3. $\boxed{1-n+\lambda < 0}$ and $\boxed{k=0}$ Let $f \in B_\sigma$ be the approximated function, $b_0 > 0$ be the cube side as in Theorem2.1, and $d < \frac{b_0}{4\gamma_n(m+1)}$ be the fill distance. If $1-n+\lambda < 0$ and $k=0$, the optimal choice of c for $c \in [c_0, \infty)$ is the value minimizing $c^{\frac{1-n+\lambda}{4}} e^{\frac{c\sigma}{2}}$ on the interval $[c_1, \infty)$.

Reason: In this case $MN(c)$ is decreasing on $[c_0, c_1]$. Therefore the minimum value of $MN(c)$ happens on the interval $[c_1, \infty)$ where the essential part of $MN(c)$ is $c^{\frac{1-n+\lambda}{4}} e^{\frac{c\sigma}{2}}$.

Remark:(a) Although k rarely equals zero, we can make it zero by choosing σ in an appropriate way. It will make things easier for the optimal choice of c . (b) If $1-n+\lambda < 0$, it can be shown that $g(c) := c^{\frac{1-n+\lambda}{4}} e^{\frac{c\sigma}{2}}$ is decreasing on $(0, \frac{-1+n-\lambda}{2\sigma}]$ and increasing on $[\frac{-1+n-\lambda}{2\sigma}, \infty)$, with $g'(\frac{-1+n-\lambda}{2\sigma}) = 0$. Therefore the optimal choice of c for $c \in [c_0, \infty)$ in Case3 is in fact $\boxed{\max\{\frac{-1+n-\lambda}{2\sigma}, c_1\}}$.

Case4. $\boxed{1-n+\lambda < 0}$ and $\boxed{k > 0}$ Let $f \in B_\sigma$ be the approximated function, $b_0 > 0$ be the cube side as in Theorem2.1, and $d < \frac{b_0}{4\gamma_n(m+1)}$ be the fill distance. Let $g_1(c) := MN(c)|_{c \in [c_0, c_1]}$ and $g_2(c) := MN(c)|_{c \in [c_1, \infty)}$. If $1-n+\lambda < 0$ and $k > 0$, then

- (i) $g_1'(\frac{1-n+\lambda}{-4k}) = 0$, $g_1'(c) < 0$ for $c \in (0, \frac{1-n+\lambda}{-4k})$, and $g_1'(c) > 0$ for $c \in (\frac{1-n+\lambda}{-4k}, \infty)$, and
- (ii) $g_2'(\frac{-1+n-\lambda}{2\sigma}) = 0$, $g_2'(c) < 0$ for $c \in (0, \frac{-1+n-\lambda}{2\sigma})$, and $g_2'(c) > 0$ for $c \in (\frac{-1+n-\lambda}{2\sigma}, \infty)$.

In this case the minimum value of $g_1(c)$ on $[c_0, c_1)$ happens at (a) $c = c_0$ if $\frac{1-n+\lambda}{-4k} \leq c_0$, (b) $c = \frac{1-n+\lambda}{-4k}$ if $c_0 < \frac{1-n+\lambda}{-4k} < c_1$, and (c) $c = c_1$ if $c_1 \leq \frac{1-n+\lambda}{-4k}$. Also, the minimum value of $g_2(c)$ on $[c_1, \infty)$ happens at (a) $c = c_1$ if $\frac{-1+n-\lambda}{2\sigma} < c_1$, and (b) $c = \frac{-1+n-\lambda}{2\sigma}$ if $c_1 \leq \frac{-1+n-\lambda}{2\sigma}$.

Let $c^* \in [c_0, c_1]$ minimize $g_1(c)$ and $c^{**} \in [c_1, \infty)$ minimize $g_2(c)$. Then the optimal choice of $c \in [c_0, \infty)$ is (a) c^* if $g_1(c^*) \leq g_2(c^{**})$, and (b) c^{**} if $g_2(c^{**}) \leq g_1(c^*)$.

Reason: By direct differentiation, we get

$$g_1'(c) = c^{\frac{1-n+\lambda}{4}-1} e^{ck} \left(\frac{1-n+\lambda}{4} + ck \right)$$

and

$$g_2'(c) = c^{\frac{1-n+\lambda}{4}-1} e^{\frac{c\sigma}{2}} e^{\frac{b_0 \ln 2}{4\gamma_n d}} \left(\frac{1-n+\lambda}{4} + \frac{c\sigma}{2} \right).$$

The seemingly complicated criterion is then just a simple result of the two derivatives. $\#$

Case5. $\boxed{1-n+\lambda < 0}$ and $\boxed{k < 0}$ Let $f \in B_\sigma$ be the approximated function, $b_0 > 0$ be the cube side as in Theorem2.1, and $d < \frac{b_0}{4\gamma_n(m+1)}$ be the fill distance. If $1-n+\lambda < 0$ and $k < 0$, the optimal choice of c for $c \in [c_0, \infty)$ is (a) $c = c_1$ if $\frac{-1+n-\lambda}{2\sigma} < c_1$, and (b) $c = \frac{-1+n-\lambda}{2\sigma}$ if $c_1 \leq \frac{-1+n-\lambda}{2\sigma}$.

Reason: In this case $MN(c)$ is decreasing on $[c_0, c_1]$ and its minimum value happens at a number in $[c_1, \infty)$ which minimizes g_2 of Case4. Our criterion thus follows immediately from Case4. $\#$

3.2 b_0 not fixed

In Theorem2.1 one can easily find that if b_0 is not fixed and can be made arbitrarily large, C will become arbitrarily small by letting $c \rightarrow \infty$. This will make $\omega \rightarrow 0$ and $d_0 \rightarrow \infty$, a very beneficial situation. In fact some domain Ω does allow $b_0 \rightarrow \infty$. For example $\Omega = R^n$ or

$$\Omega = \{(x_1, \dots, x_n) \mid x_i > 0 \text{ for } i = 1, 2, \dots, n\}.$$

This kind of domain is called **dilation-invariant** by Madych in [17]. For this kind of domain we have a different set of criteria of choosing c .

By increasing the cube side b_0 , one can always keep $C = 2\rho'\sqrt{n}e^{2n\gamma_n}$ in Theorem2.1. Thus

$$MN(c) = c^{\frac{1-n+\lambda}{4}} e^{c\left(\frac{\sigma}{2} + \frac{\ln \frac{2}{3}}{12\rho\sqrt{n}e^{2n\gamma_n}d}\right)}$$

for all $c \in [c_0, \infty)$.

Note that we never decrease b_0 because it will make the error bound (7) worse.

As for the choice of d , there is no restriction. Once $d > 0$ is given, c_0 will be fixed. For any $c \in [c_0, \infty)$, the condition $d < d_0$ in Theorem2.1 will be satisfied. However the smaller d is, the larger $[c_0, \infty)$ is, making the criteria more meaningful.

Once c is chosen, we let $b_0 = \frac{c}{3\rho\sqrt{n}e^{2n\gamma_n}}$ such that $2\rho'\sqrt{n}e^{2n\gamma_n} = \frac{2}{3b_0}$ in the definition of C . Further increasing b_0 is not to be expected because more data points will be involved to keep the fill distance d fixed.

Note that by letting $k := \frac{\sigma}{2} + \frac{\ln \frac{2}{3}}{12\rho\sqrt{n}e^{2n\gamma_n}d}$, we have a more simple expression for $MN(c)$ as

$$MN(c) = c^{\frac{1-n+\lambda}{4}} e^{ck}.$$

The way of choosing c is then divided in the following cases.

Case1. $\boxed{1-n+\lambda > 0}$ and $\boxed{k \geq 0}$ Let $f \in B_\sigma$ be the approximated function, and the domain Ω in Theorem2.1 be dilation-invariant. For any fill distance $d > 0$, if $1-n+\lambda > 0$ and $k \geq 0$, the optimal choice of c for $c \in [c_0, \infty)$ is $c = c_0$.

Reason: In this case $MN(c)$ is increasing on $[c_0, \infty)$. ‡

Case2. $\boxed{1-n+\lambda > 0}$ and $\boxed{k < 0}$ Let $f \in B_\sigma$ be the approximated function, and the domain Ω in Theorem2.1 be dilation-invariant. For any fill distance $d > 0$, if $1-n+\lambda > 0$ and $k < 0$, the larger c is, the better it is.

Reason: In this situation,

$$\begin{aligned} MN'(c) &= c^{\frac{1-n+\lambda}{4}-1} e^{ck} \left(\frac{1-n+\lambda}{4} + kc \right) \\ &= 0 \end{aligned}$$

iff

$$c = \frac{1-n+\lambda}{-4k}.$$

The function $MN(c)$ is increasing and decreasing, respectively, depending on $c < \frac{1-n+\lambda}{-4k}$ or $c > \frac{1-n+\lambda}{-4k}$. Note that $MN(c_0)$ is a finite value and $MN(c) \rightarrow 0$ as $c \rightarrow \infty$. Our criterion thus follows.‡

Remark:(a)Usually $k \geq 0$. Hence Case2 rarely happens. (b)If $c_0 < \frac{1-n+\lambda}{-4k}$, $MN(c_0)$ may be small enough to be accepted. If it happens, $c = c_0$ is also a good choice.

Case3. $\boxed{1 - n + \lambda < 0}$ and $\boxed{k > 0}$ Let $f \in B_\sigma$ be the approximated function, and the domain Ω in Theorem2.1 be dilation-invariant. For any fill distance $d > 0$, if $1 - n + \lambda < 0$ and $k > 0$, the optimal choice of c for $c \in [c_0, \infty)$ is (a) $c = \frac{1-n+\lambda}{-4k}$ if $c_0 \leq \frac{1-n+\lambda}{-4k}$ and (b) $c = c_0$ if $c_0 > \frac{1-n+\lambda}{-4k}$.

Reason: By the structure of $MN'(c)$, one sees easily $MN(c)$ is decreasing on $(0, \frac{1-n+\lambda}{-4k}]$ and increasing on $[\frac{1-n+\lambda}{-4k}, \infty)$. Our criterion thus follows immediately. $\#$

Case4: $\boxed{1 - n + \lambda < 0}$ and $\boxed{k \leq 0}$ Let $f \in B_\sigma$ be the approximated function, and the domain Ω in Theorem2.1 be dilation-invariant. For any fill distance $d > 0$, if $1 - n + \lambda < 0$ and $k \leq 0$, the larger c is, the better it is.

Reason: In this case $MN(c)$ is decreasing on $(0, \infty)$. Moreover, $MN(c) \rightarrow 0$ as $c \rightarrow \infty$. $\#$

Remark:(a)Case1-4 are all based on Theorem2.1. Although c and b_0 appear before d in Theorem2.1, in the four cases d is given before c and b_0 are chosen. Once d is fixed, we chose c according to the four criteria. After c is chosen, we let $b_0 = \frac{c}{3\rho\sqrt{ne^{2n\gamma_n}}}$.(b)In order to keep the fill distance d fixed, more data points have to be added when the cube side b_0 increases. (c)Although in this paper we only explore c over the interval $[c_0, \infty)$, the curve of $MN(c)$ can be used as a conjecture for $c \in (0, c_0]$. It probably holds even if we cannot prove it.

References

- [1] M. Abramowitz and I. Stegun, *A Handbook of Mathematical Functions*, Dover Publications, New York, 1970.
- [2] M.D. Buhmann, *New Development in the Theory of Radial Basis Functions Interpolation*, Multivariate Approximation: From CAGD to Wavelets(K. Jetter, F.I. Utreras eds.), World Sciectific, Singapore, (1993), 35-75.
- [3] J. Duchon, *Sur l'erreur d'interpolation des fonctions de plusieurs variables par les D^m -splines*, RAIRO Analyse numerique 12(1978), 325-334.
- [4] N. Dyn, D. Levin, S. Rippa, *Numerical procedures for global surface fitting of scattered data by radial functions*, SIAM J. Sci. and Sta. Computing 7(1986), 639-659.
- [5] N. Dyn, *Interpolation and Approximation by Radial and Related Functions*, Approximation Theory VI,(C.K. Chui, L.L. Schumaker and J. Ward eds.), Academic press,(1989), 211-234.
- [6] I.M. Gelfand and G.E. Shilov, *Generalized Functions*, Vol.1, Academic Press, 1964.
- [7] L-T. Luh, *The Equivalence Theory of Native Spaces*, Approx. Theory Appl., Vol. 17, No.1,2001,76-96.
- [8] L-T. Luh, *The Embedding Theory of Native Spaces*, Approx. Theory Appl., Vol. 17, No.4,2001,90-104.

- [9] L-T. Luh, *The Mystery of the Shape Parameter I*, Math. ArXiv.
- [10] L-T. Luh, *The Mystery of the Shape Parameter II*, Math. ArXiv.
- [11] L-T. Luh, *The Mystery of the Shape Parameter III*, Math. ArXiv.
- [12] L-T. Luh, *The Mystery of the Shape Parameter IV*, Math. ArXiv.
- [13] L-T. Luh, *The Shape Parameter in the Gaussian Function*, Math. ArXiv.
- [14] L-T. Luh, *The High-level Error Bound for Shifted Surface Spline Interpolation*, Math. ArXiv.
- [15] W.R. Madych and S.A. Nelson, *Multivariate interpolation and conditionally positive definite function*, Approx. Theory Appl. 4, No. 4(1988), 77-89.
- [16] W.R. Madych and S.A. Nelson, *Multivariate interpolation and conditionally positive definite function, II*, Math. Comp. 54(1990), 211-230.
- [17] W.R. Madych, *Miscellaneous Error Bounds for Multiquadric and Related Interpolators*, Computers Math. Applic. Vol. 24, No. 12, pp. 121-138, 1992.
- [18] M.J.D. Powell, *The Theory of Radial Basis Functions Approximation in 1990*, Advances in Numerical Analysis Vol.II:Wavelets, Subdivision Algorithms and Radial Basis Functions(W.A. Light ed.), Oxford University Press,(1992), 105-210.
- [19] S. Rippa, *Interpolation and smoothing of scattered data by radial basis functions*, M.Sc. Thesis, Tel Aviv University, 1984.
- [20] H. Wendland, *Scattered Data Approximation*, Cambridge University Press, (2005).
- [21] J. Yoon, *Approximation in $L^p(\mathbb{R}^d)$ from a Space Spanned by the Scattered Shifts of Radial Basis Function*, Constr. Approx. 17(2001), no.2, 227-247.
- [22] J. Yoon, *Interpolation by Radial Basis Functions on Sobolev Space*, J. of Approx. Theory, 112(2001), pp. 1-15.
- [23] J. Yoon, *Spectral Approximation orders of Radial Basis Function Interpolation on the Sobolev Space*, SIAM J. Math. Anal. 2001, Vol.33, No.4, pp. 946-958.
- [24] J. Yoon, *L_p -Error Estimates for Shifted Surface Spline Interpolation on Sobolev Space*, Math. of Comp., 72(2003), 243, 1349-1367.