

# NEW SETS OF EXPERIMENTAL WAVENUMBER VALUES FOR TRIPLET-TRIPLET ROVIBRONIC TRANSITIONS OF $H_2$ AND $D_2$

B. P. Lavrov,<sup>1,\*</sup> A. S. Mikhailov,<sup>1</sup> and I. S. Umrikhin<sup>1</sup>

<sup>1</sup> *Faculty of Physics, St.-Petersburg State University,  
St.-Petersburg, 198904, Russia*

New sets of experimental wavenumber values for triplet-triplet rovibronic transitions of  $H_2$  and  $D_2$  for visible part of spectrum (400 ÷ 700 nm) have been obtained. The digital intensity registration providing a linear response of the detector gave us the opportunity of digital deconvolution of the recorded line profiles. For line centers that made it possible to reach an accuracy ( $< 0.0006$  nm) limited only by selfconsistency of various wavenumber standards. New sets of wavenumber values were obtained with accuracy  $0.006 \div 0.05$   $\text{cm}^{-1}$ .

At present almost all available experimental data on rovibronic line wavenumbers of  $H_2$  and  $D_2$  molecules were obtained from emission spectra by photographic recording. Our recent studies revealed that these data have significant differences with values predicted by Rydberg-Ritz combination principle as well as with fragmentary experimental data obtained by laser induced fluorescence and with our own emission spectroscopy data obtained with photoelectric recording by matrix detectors [1, 2]. The minority of the differences is caused by misprints and erroneous line assignments; they may amount to several wavenumbers. But vast majority of them amount  $0.01 \div 0.1$   $\text{cm}^{-1}$  and show random spread around "synthesized" values, calculated as differences of optimal energy level values. We suppose that they appear due to shifts of film blackening maxima for blended lines, finite precision of reading from photo plates, and round-up errors in calculating the wavenumbers from measured wavelengths in air.

To refine our knowledge of rovibronic term structure for isotopomers of the simplest neutral molecule we launched systematic measurements of the rovibronic line wavenumbers

---

\*Electronic address: lavrov@pobox.spbu.ru

in visible part of spectrum (400 ÷ 700 nm). The 2.65 m Ebert-Fastie spectrograph was equipped with additional camera lens and computer-controlled CMOS matrix (22.2x14.8 mm, 1728x1152 pixels) providing the reverse linear dispersion about  $10^{-3}$  nm/pixel. We used various light sources: glow discharge in  $Hg + Ar$  (for determination of instrumental broadening from the superfine structure of 404.6 nm and 546.1 nm mercury lines: Gaussian shape with FWHM = 0.183(9) and 0.143(9)  $\text{cm}^{-1}$ , much smaller than Doppler broadening of  $H_2$ ,  $D_2$  lines for our gas temperatures < 1500 K), a capillary-arc discharges filled with the mixture  $D_2 + H_2 + Ne$  (for studies of the precision of spectrometer calibration) and with pure  $H_2$  or  $D_2$  (for final wavenumber measurements).

We calibrated our spectrometer with available wavelengths in vacuum to avoid precise determination of refractive index of air in our conditions. The wavenumber values of finite number of strong unblended lines of  $H_2$  [3],  $D_2$  [4] and  $Ne$  [5] were used as standards. The dependence of line wavenumber on the coordinate (in pixels) is monotonic and close to linear. The calibration curve of the spectrometer was obtained by the least-squares fitting by a second-degree polynomial (a linear approximation was insufficient, while a third-degree polynomial superfluous).

The digital intensity registration providing a linear response of the detector gave us the opportunity of digital deconvolution of the recorded line profiles. For strong unblended lines profiles were close to the Gaussian shape except for insignificant far wings. Therefore, we approximated all parts of the spectrum by Gaussian multipeak fitting with fixed half-width and adjustable intensity and wavelength values for peak maxima. Thus we obtained wavenumbers of rovibronic radiative transitions for triplet-triplet band systems.

The contribution contains the details of experimental technique and results.

This work was financially supported in part by the Russian Foundation for Basic Research, project no. 10-03-00571a.

- 
- [1] Astashkevich S.A., Lavrov B.P., Modin A.V., Umrikhin I.S. The Radiative Characteristics of the Rovibronic States of the Hydrogen Molecule: IV. Relative Probabilities of the  $r^3\Pi_g, s^3\Delta_g - c^3\Pi_u$  Spontaneous Transitions of the  $H_2$  Molecule. *Russian Journal of Physical Chemistry B*, 2008, **2** (1) P.16–31

- [2] Lavrov B.P. and Umrikhin I.S. 2008 *J. Phys. B: At. Mol. Opt. Phys.* Optimal values of rovibronic energy levels for triplet electronic states of molecular deuterium. *J. Phys. B: At. Mol. Opt. Phys.*, 2008, **41** 105103 (25pp)
- [3] *The Hydrogen Molecule Wavelength Tables of G.H.Dieke / Ed. by H.M.Crosswhite*, 1972, (N.Y.: Wiley-Interscience) 616 pp.
- [4] Freund R.S., Schiavone J.A. and Crosswhite H.M. *J. Phys. Chem. Ref. Data.*, 1985, **14** P.235–383
- [5] <http://webbook.nist.gov/>