# Fermionic K-essence

Ratbay Myrzakulov\*

Dept. of Phys., California State University, Fresno, CA, 93740 USA

and

Eurasian International Center for Theoretical Physics, Dep. Gen. & Theor. Phys., Eurasian National University, Astana 010008, Kazakhstan

#### Abstract

In the present work, we study the cosmological model with fermionic field and with the non-canonical kinetic term (*fermionic k-essence* or *f-essence*). We also present some important reductions of the model as well as its some generalizations. We also found the exact solution of the model and examine the influence of such gravity-fermion interaction on the observed accelerated expansion of our universe.

## 1 Introduction

The observational evidence from different sources for the present stage of accelerated expansion of our universe has driven the quest for theoretical explanations of such feature. Assuming the validity of the theory of gravity, one attempt of explanation is the existence of an unregarded, but dominated at present time, ingredient of the energy content of the universe, known as dark energy, with unusual physical properties. The other possibility is modifying the general theory of relativity at large scales. In cosmology, the investigation for the constituents responsible for the accelerated periods in the evolution of the universe is of great interest. The mysterious dark energy has been proposed as a cause for the late time dynamics of the current accelerated phase of the universe.

During last years theories described by the action with non-standard kinetic terms, k-essence, attracted a considerable interest. Such theories were first studied in the context of k-inflation [1], and then the k-essence models were suggested as dynamical dark energy for solving the cosmic coincidence problem [2]-[4]. The action of the k-essence scalar field  $\phi$  minimally coupled to the gravitational field  $g_{\mu\nu}$  we write in the form (see e.g. [1]-[4])

$$S = \int d^4x \sqrt{-g} [R + K_1(X,\phi)], \qquad (1.1)$$

where

$$X = 0.5g^{\mu\nu}\nabla_{\mu}\phi\nabla_{\nu}\phi, \qquad (1.2)$$

is the canonical kinetic term,  $\nabla_{\mu}$  is the covariant derivative associated with metric  $g_{\mu\nu}$ . The important particular reductions of the scalar k-essence (1.1) are:

i)  $K_1 = A_1(X)$  (purely kinetic case),

ii) 
$$K_1 = A_1(X)B_1(\phi),$$

iii) 
$$K_1 = A_1(X) + B_1(\phi)$$
.

In the recent years several approaches were made to explain the accelerated expansion by choosing fermionic fields as the gravitational sources of energy (see e.g. refs. [5]-[18]). In particular,

<sup>\*</sup>Email: rmyrzakulov@csufresno.edu; cnlpmyra1954@yahoo.com

it was shown that the fermionic field plays very important role in: i) isotropization of initially anisotropic spacetime; ii) formation of singularity free cosmological solutions; iii) explaining late-time acceleration. In the present work, we study the cosmological model with fermionic field, the  $M_{33}$  - model, which has the non-canonical kinetic term (*f-essence*). We examine the influence of such gravity-fermionic interaction on the accelerated expansion of the Universe. The formulation of the gravity-fermionic theory has been discussed in detail elsewhere [19]-[22]., so we will only present the result here.

## 2 Einstein-Dirac equations

In order to have this work self-consistent, in this section we present briefly the techniques that are used to include fermionic sources in the Einstein theory of gravitation and for a more detailed analysis the reader is referred to [19]-[22]. The Einstein-Dirac action reads as

$$S = \int d^4x \sqrt{-g} [R + \epsilon Y - V], \qquad (2.1)$$

where  $\epsilon = \pm 1$  ( $\epsilon = 1$  is the usual case and  $\epsilon = -1$  is the phantom case) and

$$Y = 0.5i[\bar{\psi}\Gamma^{\mu}D_{\mu}\psi - (D_{\mu}\bar{\psi})\Gamma^{\mu}\psi], \quad V = V(\bar{\psi},\psi).$$
(2.2)

The closed system of the equations for this model looks like (see e.i. [8])

$$R_{\mu\nu} - 0.5Rg_{\mu\nu} + T_{\mu\nu} = 0, \qquad (2.3)$$

$$i\Gamma^{\mu}D_{\mu}\psi - \frac{dV}{d\bar{\psi}} = 0, \qquad (2.4)$$

$$iD_{\mu}\bar{\psi}\Gamma^{\mu} + \frac{dV}{d\psi} = 0, \qquad (2.5)$$

$$\dot{\rho}_f + 3H(\rho_f + p_f) = 0,$$
 (2.6)

where the density of energy and pressure are given by

$$\rho_f = -V, \quad p_f = -Y + V.$$
(2.7)

## **3** The $M_{33}$ - model

Let us now we consider the  $M_{33}$  - model, which has the action

$$S = \int d^4x \sqrt{-g} [R + K_2(Y, \psi, \bar{\psi})], \qquad (3.1)$$

where

$$Y = 0.5i[\bar{\psi}\Gamma^{\mu}D_{\mu}\psi - (D_{\mu}\bar{\psi})\Gamma^{\mu}\psi].$$
(3.2)

We work with a space-time metric of the form

$$ds^{2} = -dt^{2} + a^{2}(dx^{2} + dy^{2} + dz^{2}), \qquad (3.3)$$

that is the FRW metric. In this case, the equations of motion look like

$$3H^2 + 0.5[K_2 + 0.5(K_{2\psi}\psi + K_{2\bar{\psi}}\bar{\psi})] = 0, \qquad (3.4)$$

$$2\dot{H} + 3H^2 + 0.5K_2 = 0, (3.5)$$

$$K_{2Y}\psi + 0.5(3HK_{2Y} + K_{2Y})\psi - i\gamma^0 K_{2\bar{\psi}} = 0, \qquad (3.6)$$

$$K_{2Y}\bar{\psi} + 0.5(3HK_{2Y} + \dot{K}_{2Y})\bar{\psi} + iK_{2\psi}\gamma^0 = 0, \qquad (3.7)$$

$$\dot{\rho}_f + 3H(\rho_f + p_f) = 0,$$
 (3.8)

where  $Y = 0.5i(\bar{\psi}\gamma^0\dot{\psi} - \dot{\bar{\psi}}\gamma^0\psi)$  and

$$\rho_f = -[K_2 + 0.5(K_{2\psi}\psi + K_{2\bar{\psi}}\bar{\psi})], \quad p_f = K_2$$
(3.9)

are the energy density and pressure of the fermionic field. If  $K_2 = Y - V$ , then from the system (3.4)-(3.8) we get the equations corresponding to the Einstein-Dirac model that is the equations (2.3)-(2.6).

### 3.1 Submodels

### 3.1.1 The $M_{33A}$ - model

The Langrangian of the  $\mathrm{M}_{33A}$  - model has the form

$$K_2 = A_2(Y). (3.10)$$

So the  $M_{33A}$  - model is the purely kinetic fermionic k-essence.

#### 3.1.2 The $M_{33B}$ - model

The Langrangian of the  $M_{33B}$  - model reads as

$$K_2 = A_2(Y)B_2(\psi, \bar{\psi}). \tag{3.11}$$

### 3.1.3 The $M_{33C}$ - model

The Langrangian of the  $\mathrm{M}_{33A}$  - model has the form

$$K_2 = A_2(Y) + B_2(\psi, \bar{\psi}). \tag{3.12}$$

### 3.2 Solution

In this subsection we want to construct a solution of the M<sub>33</sub>-model. Let  $K_2 = K_2(Y, u)$ , where  $u = \bar{\psi}\psi$ . Then the system (3.4)-(3.8) becomes

$$3H^{2} + 0.5[K_{2} + K_{2}'u] = 0, (3.13)$$

$$2\dot{H} + 3H^2 + 0.5K_2 = 0, (3.14)$$

$$K_{2Y}\dot{\psi} + 0.5(3HK_{2Y} + \dot{K}_{2Y})\psi - i\gamma^{0}K_{2}^{'}\psi = 0, \qquad (3.15)$$

$$K_{2Y}\dot{\bar{\psi}} + 0.5(3HK_{2Y} + \dot{K}_{2Y})\bar{\psi} + iK_{2}'\bar{\psi}\gamma^{0} = 0, \qquad (3.16)$$

$$\dot{\rho}_f + 3H(\rho_f + p_f) = 0,$$
 (3.17)

where  $K_{2}^{'} = dK_{2}/du$  and

$$\rho_f = -[K_2 + K_2'u]. \quad p_f = K_2. \tag{3.18}$$

We now consider the submodel (3.12) that is the  $M_{33C}$ -model, where we assume that  $A_2 = \alpha Y^n$ ,  $B_2 = \beta u^m$  that is the case  $K_2 = \alpha Y^n + \beta u^m$ . Let  $a = a_0 t^{\lambda}$ . Then we have the following solution

$$Y = \left\{ \left[ -\frac{6m\lambda^2 - 4(m+1)\lambda}{\alpha m} \right] t^{-2} \right\}^{\frac{1}{n}}, \quad u = \left\{ \left[ \frac{-4\lambda}{\beta m} \right] t^{-2} \right\}^{\frac{1}{m}}, \quad (3.19)$$

where

$$m = \frac{1}{3\lambda - 1}, \quad n = \frac{1}{3\lambda}, \quad c = \alpha n a_0^3 \left[\frac{-4\lambda}{\beta m}\right]^{\frac{1}{m}} \left[-\frac{6m\lambda^2 - 4(m+1)\lambda}{\alpha m}\right]^{\frac{1-n}{n}}.$$
 (3.20)

Finally we present the following formulas

$$u = ca^{-3} K_{2Y}^{-1}, \quad \psi_j = c_j a^{-1.5} K_{2Y}^{-0.5} e^{i\gamma^0 \int K_2' K_{2Y}^{-1} dt}, \tag{3.21}$$

where  $c = |c|_1^2 + |c|_2^2 - |c|_3^2 - |c|_4^2$ ,  $c_j = cont$ .

## 4 The $M_{34}$ - model

We now would like to present the  $M_{34}$  - model which has the following action [23]

$$S = \int d^4x \sqrt{-g} [R + K(X, Y, \phi, \psi, \bar{\psi})].$$

$$(4.1)$$

It contents some important particular submodels. For example:

i) the scalar k-essence (1.1) as  $K = K_1(X, \phi)$ ;

ii) the M<sub>33</sub> - model (3.1) as  $K = K_2(Y, \psi, \psi)$ ;

iii) the M<sub>34A</sub> - model as  $K = K_1(X, \phi)K_2(Y, \psi, \psi);$ 

vi) the  $M_{34B}$  - model as  $K = K_1(X, \phi) + K_2(Y, \psi, \psi)$ .

Some properties of the  $M_{34}$  - model (4.1) were studied in [23]. In particular it is shown that it can describe the late-time acceleration of the universe.

## 5 Conclusion

We briefly summarize the present work. We first derived the equations of the M<sub>33</sub> - model for the FRW space-time. Then we found their exact solution for the *f*-essence  $K_2 = \alpha Y^n + \beta u^m$ . Finally, let us we present the expression for the equation of state parameter w. For the our particular solution (3.19) it takes the form

$$w_f = \frac{p_f}{\rho_f} = -1 - \frac{2}{3\lambda}.$$
 (5.1)

This formula tells us that the  $M_{33}$  - model can describes the observed accelerated expansion of our universe.

### References

- Armendariz-Picon C., Damour T., Mukhanov V.F. k-inflation, Phys. Lett. B458, 209-218 (1999) [hep-th/9904075].
- [2] Armendariz-Picon C., Mukhanov V.F., Steinhardt P.J. Essentials of k-essence, Phys. Rev. D63, 103510 (2001) [astro-ph/0006373].
- [3] Armendariz-Picon C., Mukhanov V.F., Steinhardt P.J. A dynamical solution to the problem of a small cosmological constant and late-time cosmic acceleration, Phys. Rev. Lett. 85, 4438-4441 (2000) [astro-ph/0004134].
- [4] Chiba T., Okabe T., Yamaguchi M. Kinetically driven quintessence, Phys. Rev. D62, 023511 (2000) [astro-ph/9912463].
- [5] Ribas M.O., Devecchi F.P., Kremer G.M. Fermions as sources of accelerated regimes in cosmology, [arXiv:gr-qc/0511099]
- [6] Cai Y.F., Wang J. Dark Energy Model with Spinor Matter and Its Quintom Scenario, Class. Quant. Grav., 25, 165014 (2008) [arXiv:0806.3890]
- [7] Wang J., Cui S.-W., Zhang C.-M. Thermodynamics of Spinor Quintom, Phys. Lett., B683, 101-107 (2010) [arXiv:0806.3890]
- [8] Ribas M.O., Devecchi F.P., Kremer G.M. Cosmological model with non-minimally coupled fermionic field, [arXiv:0710.5155]
- [9] Rakhi R., Vijayagovindan G.V., Indulekha K. A cosmological model with fermionic field, [arXiv:0912.1222]
- [10] Chimento L.P., Devecchi F.P., Forte M., Kremer G.M. Phantom cosmologies and fermions, [arXiv:0707.4455]

- [11] Anischenko S.V., Cherkas S.L., Kalashnikov V.L. Cosmological Production of Fermions in a Flat Friedmann Universe with Linearly Growing Scale Factor: Exactly Solvable Model, [arXiv:0911:0769]
- [12] Saha B., Shikin G.N. J. Math. Phys. 38, 5305 (1997)
- [13] Saha B. Phys. Rev. D, 64, 123501 (2001)
- [14] Saha B. Physics of Particles and Nuclei, 37, Suppl., S13 (2006)
- [15] Saha B. Phys. Rev. D, 74, 124030 (2006)
- [16] Vakili B., Sepangi H.R. Time reparameterization in Bianchi type I spinor cosmology, [arXiv:0709.2988]
- Balantekin A.B., Dereli T. An Exact Cosmological Solution of the Coupled Einstein-Majorana Fermion-Scalar Field Equations, [arXiv:gr-qc/0701025]
- [18] Armendariz-Picon C., Greene P. Gen.Relativ. Gravit., 35, 1637 (2003)
- [19] Weinberg S. Gravitation and Cosmology (John Wiley & Sons, New York, 1972), ibid. Cosmology (Cambridge, New York, 2007).
- [20] Wald R.M. General Relativity, (The University of Chicago Press, Chicago, 1984).
- [21] Ryder L.H. Quantum Field Theory (Cambridge University Press, Cambridge, 1996).
- [22] Birrell N.D., Davies P.C.W. Quantum Fields in Curved Space (Cambridge University Press, Cambridge, 1982).
- [23] Tsyba P., Myrzakulov R. et al. Generalized k-essence models, Vestnik MKEMU, V56, (2010).