

# Jordan derivations on $C^*$ -ternary algebras for a Cauchy-Jensen functional equation

Choonkil Park<sup>1</sup>, John Michael Rassias<sup>2</sup> and Won-Gil Park<sup>3</sup>

<sup>1</sup> Department of Mathematics, Research Institute for Natural Sciences  
Hanyang University, Seoul, 133–791, Republic of Korea

<sup>2</sup> Pedagogical Department, E.E., National and Capodistrian University of  
Athens 4, Agamemnonos Str., Aghia Paraskevi, Athens 15342, Greece

<sup>3</sup> Department of Mathematics Education, College of Education Mokwon  
University, Daejeon 302-729, Republic of Korea

<sup>1</sup>baak@@hanyang.ac.kr, <sup>2</sup>jrasias@@primedu.uoa.gr  
and <sup>3</sup>wgpark@@mokwon.ac.kr

## Abstract

In this paper, we proved the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -ternary algebras and of derivations on  $C^*$ -ternary algebras for the following Cauchy-Jensen functional equation

$$3f\left(\frac{x+y+z}{3}\right) = 2f\left(\frac{x+y}{2}\right) + f(z).$$

These were applied to investigate isomorphisms between  $C^*$ -ternary algebras.

## 1 Introduction and preliminaries

Ternary structures and their generalization, the so-called  $n$ -ary structures, raise certain hopes in view of their applications in physics. Some significant physical applications are as follows (see [13, 14]):

- (1) The algebra of ‘nonions’ generated by two matrices

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad \& \quad \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & \omega \\ \omega^2 & 0 & 0 \end{pmatrix} \quad (\omega = e^{\frac{2\pi i}{3}})$$

was introduced by Sylvester as a ternary analog of Hamilton’s quaternions (cf. [1]).

- (2) The quark model inspired a particular brand of ternary algebraic systems. The so-called ‘Nambu mechanics’ is based on such structures (see [5]).

There are also some applications, although still hypothetical, in the fractional quantum Hall effect, the non-standard statistics, supersymmetric theory, and Yang–Baxter equation (cf. [1, 14, 46]).

A  $C^*$ -ternary algebra is a complex Banach space  $A$ , equipped with a ternary product  $(x, y, z) \mapsto [x, y, z]$  of  $A^3$  into  $A$ , which is  $\mathbb{C}$ -linear in the outer variables, conjugate  $\mathbb{C}$ -linear in the middle variable, and associative in the sense that  $[x, y, [z, w, v]] = [x, [w, z, y], v] = [[x, y, z], w, v]$ , and satisfies  $\|[x, y, z]\| \leq \|x\| \cdot \|y\| \cdot \|z\|$  and  $\|[x, x, x]\| = \|x\|^3$  (see [2, 47]). Every left Hilbert  $C^*$ -module is a  $C^*$ -ternary algebra via the ternary product  $[x, y, z] := \langle x, y \rangle z$ .

If a  $C^*$ -ternary algebra  $(A, [\cdot, \cdot, \cdot])$  has an identity, i.e., an element  $e \in A$  such that  $x = [x, e, e] = [e, e, x]$  for all  $x \in A$ , then it is routine to verify that  $A$ , endowed with  $x \circ y := [x, e, y]$  and  $x^* := [e, x, e]$ , is a unital  $C^*$ -algebra. Conversely, if  $(A, \circ)$  is a unital  $C^*$ -algebra, then  $[x, y, z] := x \circ y^* \circ z$  makes  $A$  into a  $C^*$ -ternary algebra.

A  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  is called a  $C^*$ -ternary algebra homomorphism if

$$H([x, y, z]) = [H(x), H(y), H(z)]$$

for all  $x, y, z \in A$ . If, in addition, the mapping  $H$  is bijective, then the mapping  $H : A \rightarrow B$  is called a  $C^*$ -ternary algebra isomorphism. A  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a  $C^*$ -ternary derivation if

$$\delta([x, y, z]) = [\delta(x), y, z] + [x, \delta(y), z] + [x, y, \delta(z)]$$

for all  $x, y, z \in A$  (see [2], [15]–[18]).

In 1940, S. M. Ulam [45] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these was the following question concerning the stability of homomorphisms.

*We are given a group  $G$  and a metric group  $G'$  with metric  $\rho(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if  $f : G \rightarrow G'$  satisfies*

$$\rho(f(xy), f(x)f(y)) < \delta$$

*for all  $x, y \in G$ , then a homomorphism  $h : G \rightarrow G'$  exists with*

$$\rho(f(x), h(x)) < \epsilon$$

*for all  $x \in G$ ?*

In 1941, D. H. Hyers [8] considered the case of approximately additive mappings  $f : E \rightarrow E'$ , where  $E$  and  $E'$  are Banach spaces and  $f$  satisfies *Hyers inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \epsilon$$

for all  $x, y \in E$ . It was shown that the limit

$$L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n}$$

exists for all  $x \in E$  and that  $L : E \rightarrow E'$  is the unique additive mapping satisfying

$$\|f(x) - L(x)\| \leq \epsilon$$

for all  $x \in E$ .

In 1978, Th. M. Rassias [35] provided a generalization of the D. H. Hyers' theorem which allows the *Cauchy difference to be unbounded*.

**Theorem 1.1.** *(Th. M. Rassias) Let  $f : E \rightarrow E'$  be a mapping from a normed vector space  $E$  into a Banach space  $E'$  subject to the inequality*

$$(1) \quad \|f(x + y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p)$$

*for all  $x, y \in E$ , where  $\epsilon$  and  $p$  are constants with  $\epsilon > 0$  and  $p < 1$ . Then the limit*

$$L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n}$$

*exists for all  $x \in E$  and  $L : E \rightarrow E'$  is the unique additive mapping which satisfies*

$$(2) \quad \|f(x) - L(x)\| \leq \frac{2\epsilon}{2 - 2^p} \|x\|^p$$

for all  $x \in E$ . If  $p < 0$  then inequality (1) holds for  $x, y \neq 0$  and (2) for  $x \neq 0$ .

On the other hand, in 1982-1989, J. M. Rassias generalized the Hyers stability result by presenting a weaker condition controlled by a product of different powers of norms. The following is according to the J. M. Rassias' theorem.

**Theorem 1.2.** (J. M. Rassias) *If it is assumed that there exist constants  $\Theta \geq 0$  and  $p_1, p_2 \in \mathbb{R}$  such that  $p = p_1 + p_2 \neq 1$ , and  $f : E \rightarrow E'$  is a mapping from a normed space  $E$  into a Banach space  $E'$  such that the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon \|x\|^{p_1} \|y\|^{p_2}$$

for all  $x, y \in E$ , then there exists a unique additive mapping  $T : E \rightarrow E'$  such that

$$\|f(x) - T(x)\| \leq \frac{\Theta}{2 - 2^p} \|x\|^p$$

for all  $x \in E$ .

In 1990, Th. M. Rassias [36] during the 27<sup>th</sup> International Symposium on Functional Equations asked the question whether such a theorem can also be proved for  $p \geq 1$ . In 1991, Z. Gajda [6] following the same approach as in Th. M. Rassias [35], gave an affirmative solution to this question for  $p > 1$ . It was shown by Z. Gajda [6], as well as by Th. M. Rassias and P. Šemrl [41] that one cannot prove a Th. M. Rassias' type theorem when  $p = 1$ . The counterexamples of Z. Gajda [6], as well as of Th. M. Rassias and P. Šemrl [41] have stimulated several mathematicians to invent new definitions of *approximately additive* or *approximately linear* mappings, cf. P. Găvruta [7], S.-M. Jung [12], who among others studied the Hyers-Ulam stability of functional equations. The inequality (1) that was introduced for the first time by Th. M. Rassias [35] provided a lot of influence in the development of a generalization of the Hyers-Ulam stability concept. This new concept is known as *generalized Hyers-Ulam stability* of functional equations (cf. the books of P. Czerwik [4], D. H. Hyers et al. [9]).

P. Găvruta [7] provided a further generalization of Th. M. Rassias' Theorem. In 1996, G. Isac and Th. M. Rassias [11] applied the generalized Hyers-Ulam stability theory to prove fixed point theorems and study some new applications in Nonlinear Analysis. In [10], D. H. Hyers et al. studied the asymptoticity aspect of Hyers-Ulam stability of mappings. Several papers have been published on various generalizations and applications of Hyers-Ulam stability and generalized Hyers-Ulam stability to a number of functional equations and mappings, for example: quadratic functional equation, invariant means, multiplicative mappings - superstability, bounded  $n$ th

differences, convex functions, generalized orthogonality functional equation, Euler-Lagrange functional equation introduced by J. M. Rassias in 1992-1998, Navier-Stokes equations. Several mathematician have contributed works on these subjects (see [3], [19]-[44]).

In Section 2, we prove the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -ternary algebras for the Cauchy-Jensen additive mappings.

In Section 3, we investigate isomorphisms between unital  $C^*$ -ternary algebras associated with the Cauchy-Jensen additive mappings.

In Section 4, we prove the generalized Hyers-Ulam stability of derivations on  $C^*$ -ternary algebras for the Cauchy-Jensen additive mappings.

## 2 Stability of homomorphisms in $C^*$ -ternary algebras

Throughout this section, assume that  $A$  is a  $C^*$ -ternary algebra with norm  $\|\cdot\|_A$  and that  $B$  is a  $C^*$ -ternary algebra with norm  $\|\cdot\|_B$ .

For a given mapping  $f : A \rightarrow B$ , we define

$$D_\mu f(x, y, z) := 3f\left(\frac{\mu x + \mu y + \mu z}{3}\right) - 2\mu f\left(\frac{x+y}{2}\right) - \mu f(z)$$

for all  $\mu \in \mathbb{T}^1 := \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}$  and all  $x, y, z \in A$ .

**Lemma 2.1.** *Let  $f : A \rightarrow B$  be a mapping such that*

$$(3) \quad D_\mu f(x, y, z) = 0$$

*for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then  $f$  is  $\mathbb{C}$ -linear.*

*Proof.* Letting  $\mu = -1$  and  $x = y = z = 0$  in (3), we gain  $f(0) = 0$ . Putting  $\mu = 1$ ,  $y = -x$  and  $z = 2x$  in (3), we get  $3f(\frac{2}{3}x) = f(2x)$  for all  $x \in A$ . So we have  $3f(x) = f(3x)$  for all  $x \in A$ . Setting  $\mu = 1$ ,  $x = 0$  in (3), we gain

$$3f\left(\frac{y+z}{3}\right) = 2f\left(\frac{y}{2}\right) + f(z)$$

for all  $y, z \in A$ . So we get  $f(y+z) = 2f(\frac{y}{2}) + f(z)$  for all  $y, z \in A$ . Taking  $z = 0$  in the above equation, we have  $f(y) = 2f(\frac{y}{2})$  for all  $y \in A$ . Thus we obtain that  $f(y+z) = f(y) + f(z)$  for all  $y, z \in A$ . Hence  $f$  is additive.

Letting  $y = z = 0$  in (3), we gain  $3f(\frac{\mu}{3}x) = 2\mu f(\frac{x}{2})$  for all  $\mu \in \mathbb{T}^1$  and all  $x \in A$ . Since  $f$  is additive,  $f(\mu x) = f(3\frac{\mu}{3}x) = 3f(\frac{\mu}{3}x) = 2\mu f(\frac{x}{2}) = \mu f(2\frac{x}{2}) = \mu f(x)$  for all  $\mu \in \mathbb{T}^1$  and all  $x \in A$ . Now let  $\lambda \in \mathbb{C}$  and  $M$

an integer greater than  $2|\lambda|$ . Since  $|\frac{\lambda}{M}| < \frac{1}{2}$ , there is  $t \in (\frac{\pi}{3}, \frac{\pi}{2}]$  such that  $|\frac{\lambda}{M}| = \cos t = \frac{e^{it} + e^{-it}}{2}$ . Now  $\frac{\lambda}{M} = |\frac{\lambda}{M}|\mu$  for some  $\mu \in \mathbb{T}^1$ . Thus we have

$$\begin{aligned} f(\lambda x) &= f\left(M\frac{\lambda}{M}x\right) = Mf\left(\frac{\lambda}{M}x\right) = Mf\left(|\frac{\lambda}{M}|\mu x\right) \\ &= Mf\left(\frac{e^{it} + e^{-it}}{2}\mu x\right) = \frac{1}{2}Mf(e^{it}\mu x + e^{-it}\mu x) \\ &= \frac{1}{2}M[e^{it}\mu f(x) + e^{-it}\mu f(x)] = \lambda f(x) \end{aligned}$$

for all  $x \in A$ . So the mapping  $f : A \rightarrow B$  is  $\mathbb{C}$ -linear.  $\square$

We prove the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -ternary algebras for the functional equation  $D_\mu f(x, y, z) = 0$ .

**Theorem 2.2.** *Let  $r > 3$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying  $f(0) = 0$  such that*

$$\begin{aligned} (4) \quad & \|D_\mu f(x, y, z)\|_B \leq \theta(\|x\|_A^r + \|y\|_A^r + \|z\|_A^r), \\ (5) \quad & \|f([x, y, z]) - [f(x), f(y), f(z)]\|_B \leq \theta(\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) \end{aligned}$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique  $C^*$ -ternary algebra homomorphism  $H : A \rightarrow B$  such that

$$(6) \quad \|f(x) - H(x)\|_B \leq \theta \frac{3^r + 2}{3^r - 3} \|x\|_A^r$$

for all  $x \in A$ .

*Proof.* Letting  $\mu = 1$  and  $y = -x$  and  $z = 3x$  in (4), we obtain

$$(7) \quad \|3f(x) - f(3x)\|_B \leq \theta(2 + 3^r)\|x\|_A^r$$

for all  $x \in A$ . So we get

$$\left\|f(x) - 3f\left(\frac{x}{3}\right)\right\|_B \leq \theta\left(\frac{2}{3^r} + 1\right)\|x\|_A^r$$

for all  $x \in A$ . Thus we have

$$\begin{aligned} & \left\|3^l f\left(\frac{x}{3^l}\right) - 3^m f\left(\frac{x}{3^m}\right)\right\|_B \leq \sum_{j=l}^{m-1} \left\|3^j f\left(\frac{x}{3^j}\right) - 3^{j+1} f\left(\frac{x}{3^{j+1}}\right)\right\|_B \\ (8) \quad & \leq \theta\left(\frac{2}{3^r} + 1\right) \sum_{j=l}^{m-1} 3^{j(1-r)} \|x\|_A^r = \theta \frac{3^r + 2}{3^r - 3} [3^{l(1-r)} - 3^{m(1-r)}] \|x\|_A^r \end{aligned}$$

for all nonnegative integers  $m$  and  $l$  with  $m > l$  and all  $x \in A$ . It follows from (8) that the sequence  $\{3^n f(\frac{x}{3^n})\}$  is a Cauchy sequence for all  $x \in A$ .

Since  $B$  is complete, the sequence  $\{3^n f(\frac{x}{3^n})\}$  converges for all  $x \in A$ . Hence one can define a mapping  $H : A \rightarrow B$  by

$$H(x) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x}{3^n}\right)$$

for all  $x \in A$ . Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (8), we get (6).

It follows from (4) that

$$\begin{aligned} & \left\| 3H\left(\frac{x+y+z}{3}\right) - 2H\left(\frac{x+y}{2}\right) - H(z) \right\|_B \\ &= \lim_{n \rightarrow \infty} 3^n \left\| 3f\left(\frac{x+y+z}{3^{n+1}}\right) - 2f\left(\frac{x+y}{2 \cdot 3^n}\right) - f\left(\frac{z}{3^n}\right) \right\|_B \\ &\leq \lim_{n \rightarrow \infty} 3^{n(1-r)} \theta(\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) = 0 \end{aligned}$$

for all  $x, y, z \in A$ . So we get

$$3H\left(\frac{x+y+z}{3}\right) = 2H\left(\frac{x+y}{2}\right) + H(z)$$

for all  $x, y, z \in A$ . Since  $f(0) = 0$ , by the same methods as in proof of Lemma 2.1, the mapping  $H : A \rightarrow B$  is additive.

By the same reasoning as in the proof of Theorem 2.1 in [21], the mapping  $H : A \rightarrow B$  is  $\mathbb{C}$ -linear. It follows from (5) and (8) that

$$\begin{aligned} & \|H([x, y, z]) - [H(x), H(y), H(z)]\|_B \\ &= \lim_{n \rightarrow \infty} \left\| 3^n f\left(\frac{1}{3^n}[x, y, z]\right) - \left[3^n f\left(\frac{x}{3^n}\right), 3^n f\left(\frac{y}{3^n}\right), 3^n f\left(\frac{z}{3^n}\right)\right] \right\|_B \\ &= \lim_{n \rightarrow \infty} \left[ \left\| 3^n f\left(\frac{1}{3^n}[x, y, z]\right) - 3^{2n} f\left(\frac{1}{3^{2n}}[x, y, z]\right) \right\|_B \right. \\ &\quad + \left\| 3^{2n} f\left(\frac{1}{3^{2n}}[x, y, z]\right) - 3^{3n} f\left(\frac{1}{3^{3n}}[x, y, z]\right) \right\|_B \\ &\quad + \left\| 3^{3n} f\left(\left[\frac{x}{3^n}, \frac{y}{3^n}, \frac{z}{3^n}\right]\right) \right. \\ &\quad \left. \left. - \left[3^n f\left(\frac{x}{3^n}\right), 3^n f\left(\frac{y}{3^n}\right), 3^n f\left(\frac{z}{3^n}\right)\right] \right\|_B \right] \end{aligned}$$

$$\begin{aligned}
&\leq \lim_{n \rightarrow \infty} \left[ \theta \left( \frac{2}{3^r} + 1 \right) \sum_{j=n}^{2n-1} 3^{j(1-r)} \|[x, y, z]\|_A^r \right. \\
&\quad \left. + \theta \left( \frac{2}{3^r} + 1 \right) \sum_{j=2n}^{3n-1} 3^{j(1-r)} \|[x, y, z]\|_A^r \right. \\
&\quad \left. + 3^{n(3-r)} \theta (\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) \right] \\
&= \theta \frac{3^r + 2}{3^r - 3} \|[x, y, z]\|_A^r \lim_{n \rightarrow \infty} [3^{n(1-r)} - 3^{3n(1-r)}] \\
&\quad + \theta (\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) \lim_{n \rightarrow \infty} 3^{n(3-r)} \\
&= 0
\end{aligned}$$

for all  $x, y, z \in A$ . So

$$H([x, y, z]) = [H(x), H(y), H(z)]$$

for all  $x, y, z \in A$ .

Now, let  $T : A \rightarrow B$  be another additive mapping satisfying (6). Then we have

$$\begin{aligned}
\|H(x) - T(x)\|_B &= 3^n \left\| H\left(\frac{x}{3^n}\right) - T\left(\frac{x}{3^n}\right) \right\|_B \\
&\leq 3^n \left[ \left\| H\left(\frac{x}{3^n}\right) - f\left(\frac{x}{3^n}\right) \right\|_B + \left\| f\left(\frac{x}{3^n}\right) - T\left(\frac{x}{3^n}\right) \right\|_B \right] \\
&\leq \frac{2\theta}{3^{n(r-1)}} \frac{3^r + 2}{3^r - 3} \|x\|_A^r,
\end{aligned}$$

which tends to zero as  $n \rightarrow \infty$  for all  $x \in A$ . So we can conclude that  $H(x) = T(x)$  for all  $x \in A$ . This proves the uniqueness of  $H$ . Thus the mapping  $H : A \rightarrow B$  is a unique  $C^*$ -ternary algebra homomorphism satisfying (6).  $\square$

J. M. Rassias presents the following counterexample modified by the well-known counterexample of Z. Gajda [6] for the following Cauchy-Jensen functional equation:

$$3f\left(\frac{x+y+z}{3}\right) = 2f\left(\frac{x+y}{2}\right) + f(z).$$

Fix  $\theta > 0$  and put  $\mu := \frac{\theta}{6}$ . Define a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  given by

$$f(x) := \sum_{n=0}^{\infty} \frac{\phi(2^n x)}{2^n}$$

for all  $x \in \mathbb{R}$ , where

$$\phi(x) := \begin{cases} \mu & \text{if } x \geq 1 \\ \mu x & \text{if } -1 < x < 1 \\ -\mu & \text{if } x \leq -1 \end{cases}$$

for all  $x \in \mathbb{R}$ . It was proven in [6] that

$$|f(x+y) - f(x) - f(y)| \leq \theta(|x| + |y|)$$

for all  $x, y \in \mathbb{R}$ . From the above inequality, one can obtain that

$$\begin{aligned} (9) \quad & |f(x+y+z) - f(x) - f(y) - f(z)| \\ & \leq \frac{1}{3} [ |f(x+y+z) - f(x+y) - f(z)| \\ & \quad + |f(x+y+z) - f(x+z) - f(y)| \\ & \quad + |f(x+y+z) - f(y+z) - f(x)| \\ & \quad + |f(x+y) - f(x) - f(y)| + |f(x+z) - f(x) - f(z)| \\ & \quad + |f(y+z) - f(y) - f(z)| ] \\ & \leq \frac{5}{3} \theta(|x| + |y| + |z|) \end{aligned}$$

and

$$\begin{aligned} (10) \quad & \left| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right| \\ & \leq 2 \left| f\left(\frac{x}{2} + \frac{y}{2}\right) - f\left(\frac{x}{2}\right) - f\left(\frac{y}{2}\right) \right| \\ & \quad + \left| - \left[ f\left(\frac{x}{2} + \frac{x}{2}\right) - f\left(\frac{x}{2}\right) - f\left(\frac{x}{2}\right) \right] \right| \\ & \quad + \left| - \left[ f\left(\frac{y}{2} + \frac{y}{2}\right) - f\left(\frac{y}{2}\right) - f\left(\frac{y}{2}\right) \right] \right| \\ & \leq 2\theta(|x| + |y|) \end{aligned}$$

for all  $x, y, z \in \mathbb{R}$ . By the inequality (9), we see that

$$(11) \quad \left| 3f\left(\frac{x+y+z}{3}\right) - f(x) - f(y) - f(z) \right|$$

$$\begin{aligned}
&\leq 3 \left| f\left(\frac{x}{3} + \frac{y}{3} + \frac{z}{3}\right) - f\left(\frac{x}{3}\right) - f\left(\frac{y}{3}\right) - f\left(\frac{z}{3}\right) \right| \\
&\quad + \left| - \left[ f\left(\frac{x}{3} + \frac{x}{3} + \frac{x}{3}\right) - f\left(\frac{x}{3}\right) - f\left(\frac{x}{3}\right) - f\left(\frac{x}{3}\right) \right] \right| \\
&\quad + \left| - \left[ f\left(\frac{y}{3} + \frac{y}{3} + \frac{y}{3}\right) - f\left(\frac{y}{3}\right) - f\left(\frac{y}{3}\right) - f\left(\frac{y}{3}\right) \right] \right| \\
&\quad + \left| - \left[ f\left(\frac{z}{3} + \frac{z}{3} + \frac{z}{3}\right) - f\left(\frac{z}{3}\right) - f\left(\frac{z}{3}\right) - f\left(\frac{z}{3}\right) \right] \right| \\
&\leq \frac{10}{3} \theta (|x| + |y| + |z|)
\end{aligned}$$

for all  $x, y, z \in \mathbb{R}$ . From the inequalities (10) and (11), we obtain that

$$\begin{aligned}
&\left| 3f\left(\frac{x+y+z}{3}\right) - 2f\left(\frac{x+y}{2}\right) - f(z) \right| \\
&\leq \left| 3f\left(\frac{x+y+z}{3}\right) - f(x) - f(y) - f(z) \right| \\
&\quad + \left| - \left[ 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right] \right| \\
&\leq \frac{2}{3} \theta (8|x| + 8|y| + 5|z|) \leq \frac{16}{3} \theta (|x| + |y| + |z|)
\end{aligned}$$

for all  $x, y, z \in \mathbb{R}$ . But we observe from [6] that

$$\frac{f(x)}{x} \rightarrow \infty \quad \text{as } x \rightarrow \infty$$

and so

$$\frac{|f(x) - g(x)|}{|x|} \quad (x \neq 0) \text{ is unbounded,}$$

where  $g : \mathbb{R} \rightarrow \mathbb{R}$  is the function given by

$$g(x) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x}{3^n}\right)$$

for all  $x \in \mathbb{R}$ . Thus the function  $f$  provides an example to the effect that Theorem 2.2 fails to hold for  $r = 1$ .

**Theorem 2.3.** *Let  $r < 1$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying (4), (5) and  $f(0) = 0$ . Then there exists a unique  $C^*$ -ternary algebra homomorphism  $H : A \rightarrow B$  such that*

$$(12) \quad \|f(x) - H(x)\|_B \leq \theta \frac{2 + 3^r}{3 - 3^r} \|x\|_A^r$$

for all  $x \in A$ .

*Proof.* It follows from (7) that

$$\left\| f(x) - \frac{1}{3} f(3x) \right\|_B \leq \theta \frac{2 + 3^r}{3} \|x\|_A^r$$

for all  $x \in A$ . So

$$(13) \quad \begin{aligned} & \left\| \frac{1}{3^l} f(3^l x) - \frac{1}{3^m} f(3^m x) \right\|_B \leq \sum_{j=l}^{m-1} \left\| \frac{1}{3^j} f(3^j x) - \frac{1}{3^{j+1}} f(3^{j+1} x) \right\|_B \\ & \leq \theta \frac{2+3^r}{3} \sum_{j=l}^{m-1} 3^{j(r-1)} \|x\|_A^r = \theta \frac{2+3^r}{3-3^r} [3^{l(r-1)} - 3^{m(r-1)}] \|x\|_A^r \end{aligned}$$

for all nonnegative integers  $m$  and  $l$  with  $m > l$  and all  $x \in A$ . It follows from (13) that the sequence  $\{\frac{1}{3^n} f(3^n x)\}$  is a Cauchy sequence for all  $x \in A$ . Since  $B$  is complete, the sequence  $\{\frac{1}{3^n} f(3^n x)\}$  converges for all  $x \in A$ . So one can define the mapping  $H : A \rightarrow B$  by

$$H(x) := \lim_{n \rightarrow \infty} \frac{1}{3^n} f(3^n x)$$

for all  $x \in A$ . Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (13), we get (12).

By similar arguments to the proof of Theorem 2.2, the mapping  $H : A \rightarrow B$  is  $\mathbb{C}$ -linear. It follows from (5) and (13) that

$$\begin{aligned} & \|H([x, y, z]) - [H(x), H(y), H(z)]\|_B \\ & \leq \theta \frac{2+3^r}{3-3^r} \|[x, y, z]\|_A^r \lim_{n \rightarrow \infty} [3^{n(r-1)} - 3^{3n(r-1)}] \\ & \quad + \theta (\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) \lim_{n \rightarrow \infty} 3^{n(r-3)} \\ & = 0 \end{aligned}$$

for all  $x, y, z \in A$ . So

$$H([x, y, z]) = [H(x), H(y), H(z)]$$

for all  $x, y, z \in A$ . Now, let  $T : A \rightarrow B$  be another additive mapping satisfying (12). Then we have

$$\|H(x) - T(x)\|_B \leq 2\theta 3^{n(r-1)} \frac{2+3^r}{3-3^r} \|x\|_A^r,$$

which tends to zero as  $n \rightarrow \infty$  for all  $x \in A$ . So we can conclude that  $H(x) = T(x)$  for all  $x \in A$ . This proves the uniqueness of  $H$ . Thus the mapping  $H : A \rightarrow B$  is a unique  $C^*$ -ternary algebra homomorphism satisfying (12).  $\square$

**Theorem 2.4.** *Let  $r > \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying  $f(0) = 0$  such that*

$$(14) \quad \|D_\mu f(x, y, z)\|_B \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \cdot \|z\|_A^r,$$

$$(15) \quad \|f([x, y, z]) - [f(x), f(y), f(z)]\|_B \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \cdot \|z\|_A^r$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique  $C^*$ -ternary algebra homomorphism  $H : A \rightarrow B$  such that

$$(16) \quad \|f(x) - H(x)\|_B \leq \frac{3^r \theta}{27^r - 3} \|x\|_A^{3r}$$

for all  $x \in A$ .

*Proof.* Letting  $\mu = 1$  and  $y = -x$  and  $z = 3x$  in (14), we get

$$(17) \quad \|f(3x) - 3f(x)\|_B \leq 3^r \theta \|x\|_A^{3r}$$

for all  $x \in A$ . So

$$\left\| f(x) - 3f\left(\frac{x}{3}\right) \right\|_B \leq \frac{\theta}{9^r} \|x\|_A^{3r}$$

for all  $x \in A$ . Hence

$$(18) \quad \begin{aligned} & \left\| 3^l f\left(\frac{x}{3^l}\right) - 3^m f\left(\frac{x}{3^m}\right) \right\|_B \leq \sum_{j=l}^{m-1} \left\| 3^j f\left(\frac{x}{3^j}\right) - 3^{j+1} f\left(\frac{x}{3^{j+1}}\right) \right\|_B \\ & \leq \frac{\theta}{9^r} \sum_{j=l}^{m-1} 3^{j(1-3r)} \|x\|_A^{3r} = \frac{\theta}{9^r - 3^{1-r}} [3^{l(1-3r)} - 3^{m(1-3r)}] \|x\|_A^{3r} \end{aligned}$$

for all nonnegative integers  $m$  and  $l$  with  $m > l$  and all  $x \in A$ . It follows from (18) that the sequence  $\{3^n f(\frac{x}{3^n})\}$  is a Cauchy sequence for all  $x \in A$ . Since  $B$  is complete, the sequence  $\{3^n f(\frac{x}{3^n})\}$  converges. So one can define the mapping  $H : A \rightarrow B$  by

$$H(x) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x}{3^n}\right)$$

for all  $x \in A$ . Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (18), we get (16).

The rest of the proof is similar to the proof of Theorem 2.2.  $\square$

**Theorem 2.5.** Let  $r < \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying (14), (15) and  $f(0) = 0$ . Then there exists a unique  $C^*$ -ternary algebra homomorphism  $H : A \rightarrow B$  such that

$$(19) \quad \|f(x) - H(x)\|_B \leq \frac{3^r \theta}{3 - 27^r} \|x\|_A^{3r}$$

for all  $x \in A$ .

*Proof.* It follows from (17) that

$$\left\| f(x) - \frac{1}{3} f(3x) \right\|_B \leq 3^{r-1} \theta \|x\|_A^{3r}$$

for all  $x \in A$ . So

$$(20) \quad \left\| \frac{1}{3^l} f(3^l x) - \frac{1}{3^m} f(3^m x) \right\|_B \leq \sum_{j=l}^{m-1} \left\| \frac{1}{3^j} f(3^j x) - \frac{1}{3^{j+1}} f(3^{j+1} x) \right\|_B$$

$$\leq 3^{r-1} \theta \sum_{j=l}^{m-1} 3^{j(3r-1)} \|x\|_A^{3r} = \frac{3^{r-1} \theta}{1 - 3^{3r-1}} [3^{l(3r-1)} - 3^{m(3r-1)}] \|x\|_A^{3r}$$

for all nonnegative integers  $m$  and  $l$  with  $m > l$  and all  $x \in A$ . It follows from (20) that the sequence  $\{\frac{1}{3^n} f(3^n x)\}$  is a Cauchy sequence for all  $x \in A$ . Since  $B$  is complete, the sequence  $\{\frac{1}{3^n} f(3^n x)\}$  converges for all  $x \in A$ . So one can define the mapping  $H : A \rightarrow B$  by

$$H(x) := \lim_{n \rightarrow \infty} \frac{1}{3^n} f(3^n x)$$

for all  $x \in A$ . Moreover, letting  $l = 0$  and passing the limit  $m \rightarrow \infty$  in (20), we get (19).

The rest of the proof is similar to the proof of Theorem 2.2.  $\square$

### 3 Isomorphisms between $C^*$ -ternary algebras

Throughout this section, assume that  $A$  is a unital  $C^*$ -ternary algebra with norm  $\|\cdot\|_A$  and unit  $e$ , and that  $B$  is a unital  $C^*$ -ternary algebra with norm  $\|\cdot\|_B$  and unit  $e'$ .

We investigate isomorphisms between  $C^*$ -ternary algebras associated with the functional equation  $D_\mu f(x, y, z) = 0$ .

**Theorem 3.1.** *Let  $r > 1$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a bijective mapping satisfying (4) and  $f(0) = 0$  such that*

$$(21) \quad f([x, y, z]) = [f(x), f(y), f(z)]$$

for all  $x, y, z \in A$ . If  $\lim_{n \rightarrow \infty} 3^n f(\frac{e}{3^n}) = e'$ , then the mapping  $f : A \rightarrow B$  is a  $C^*$ -ternary algebra isomorphism.

*Proof.* By the same argument as in the proof of Theorem 2.2, one can obtain a  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  satisfying (6). The mapping  $H$  is given by

$$H(x) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x}{3^n}\right)$$

for all  $x \in A$ .

Since  $f([x, y, z]) = [f(x), f(y), f(z)]$  for all  $x, y, z \in A$ ,

$$\begin{aligned} H([x, y, z]) &= \lim_{n \rightarrow \infty} 3^{3n} f\left(\frac{1}{3^{3n}}[x, y, z]\right) = \lim_{n \rightarrow \infty} 3^{3n} f\left(\left[\frac{x}{3^n}, \frac{y}{3^n}, \frac{z}{3^n}\right]\right) \\ &= \lim_{n \rightarrow \infty} \left[3^n f\left(\frac{x}{3^n}\right), 3^n f\left(\frac{y}{3^n}\right), 3^n f\left(\frac{z}{3^n}\right)\right] \\ &= [H(x), H(y), H(z)] \end{aligned}$$

for all  $x, y, z \in A$ . So the mapping  $H : A \rightarrow B$  is a  $C^*$ -ternary algebra homomorphism.

It follows from (21) that

$$\begin{aligned} H(x) &= H([e, e, x]) = \lim_{n \rightarrow \infty} 3^{2n} f\left(\frac{1}{3^{2n}}[e, e, x]\right) = \lim_{n \rightarrow \infty} 3^{2n} f\left(\left[\frac{e}{3^n}, \frac{e}{3^n}, x\right]\right) \\ &= \lim_{n \rightarrow \infty} \left[3^n f\left(\frac{e}{3^n}\right), 3^n f\left(\frac{e}{3^n}\right), f(x)\right] = [e', e', f(x)] = f(x) \end{aligned}$$

for all  $x \in A$ . Hence the bijective mapping  $f : A \rightarrow B$  is a  $C^*$ -ternary algebra isomorphism.  $\square$

**Theorem 3.2.** *Let  $r < 1$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a bijective mapping satisfying (4), (21) and  $f(0) = 0$ . If  $\lim_{n \rightarrow \infty} \frac{1}{3^n} f(3^n e) = e'$ , then the mapping  $f : A \rightarrow B$  is a  $C^*$ -ternary algebra isomorphism.*

*Proof.* By the same argument as in the proof of Theorem 2.3, one can obtain a  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  satisfying (12).

The rest of the proof is similar to the proof of Theorem 3.1.  $\square$

**Theorem 3.3.** *Let  $r > \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a bijective mapping satisfying (14), (21) and  $f(0) = 0$ . If  $\lim_{n \rightarrow \infty} 3^n f\left(\frac{e}{3^n}\right) = e'$ , then the mapping  $f : A \rightarrow B$  is a  $C^*$ -ternary algebra isomorphism.*

*Proof.* By the same argument as in the proof of Theorem 2.4, one can obtain a  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  satisfying (16).

The rest of the proof is similar to the proof of Theorem 3.1.  $\square$

**Theorem 3.4.** *Let  $r < \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow B$  be a bijective mapping satisfying (14), (21) and  $f(0) = 0$ . If  $\lim_{n \rightarrow \infty} \frac{1}{3^n} f(3^n e) = e'$ , then the mapping  $f : A \rightarrow B$  is a  $C^*$ -ternary algebra isomorphism.*

*Proof.* By the same argument as in the proof of Theorem 2.5, one can obtain a  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  satisfying (19).

The rest of the proof is similar to the proof of Theorem 3.1.  $\square$

#### 4 Stability of $C^*$ -ternary derivations on $C^*$ -ternary algebras

Throughout this section, assume that  $A$  is a  $C^*$ -ternary algebra with norm  $\|\cdot\|_A$ .

We prove the generalized Hyers-Ulam stability of  $C^*$ -ternary derivations on  $C^*$ -ternary algebras for the functional equation  $D_\mu f(x, y, z) = 0$ .

**Theorem 4.1.** *Let  $r > 3$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying  $f(0) = 0$  such that*

$$(22) \quad \begin{aligned} \|D_\mu f(x, y, z)\|_A &\leq \theta(\|x\|_A^r + \|y\|_A^r + \|z\|_A^r), \\ \|f([x, y, z]) - [f(x), y, z] - [x, f(y), z] - [x, y, f(z)]\|_A \\ (23) \quad &\leq \theta(\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) \end{aligned}$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique  $C^*$ -ternary derivation  $\delta : A \rightarrow A$  such that

$$(24) \quad \|f(x) - \delta(x)\|_A \leq \theta \frac{3^r + 2}{3^r - 3} \|x\|_A^r$$

for all  $x \in A$ .

*Proof.* By the same argument as in the proof of Theorem 2.2, one can obtain a  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow B$  satisfying (24). The mapping  $\delta$  is given by

$$\delta(x) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x}{3^n}\right)$$

for all  $x \in A$ .

By the same reasoning as in the proof of Theorem 2.1 of [21], the mapping  $\delta : A \rightarrow A$  is  $\mathbb{C}$ -linear.

It follows from (23) that

$$\begin{aligned} &\|\delta([x, y, z]) - [\delta(x), y, z] - [x, \delta(y), z] - [x, y, \delta(z)]\|_A \\ &= \lim_{n \rightarrow \infty} \left\| 3^{3n} f\left(\frac{[x, y, z]}{3^{3n}}\right) - 3^{2n} \left[ 3^n f\left(\frac{x}{3^n}\right), \frac{y}{3^n}, \frac{z}{3^n} \right] \right. \\ &\quad \left. - 3^{2n} \left[ \frac{x}{3^n}, 3^n f\left(\frac{y}{3^n}\right), \frac{z}{3^n} \right] - 3^{2n} \left[ \frac{x}{3^n}, \frac{y}{3^n}, 3^n f\left(\frac{z}{3^n}\right) \right] \right\|_A \\ &\leq \lim_{n \rightarrow \infty} 3^{n(3-r)} \theta (\|x\|_A^r + \|y\|_A^r + \|z\|_A^r) = 0 \end{aligned}$$

for all  $x, y, z \in A$ . So

$$\delta([x, y, z]) = [\delta(x), y, z] + [x, \delta(y), z] + [x, y, \delta(z)]$$

for all  $x, y, z \in A$ .

By the same argument as in the proof of Theorem 2.2, the uniqueness of  $\delta$  is proved. Thus the mapping  $\delta$  is a unique  $C^*$ -ternary derivation satisfying (24).  $\square$

**Theorem 4.2.** *Let  $r < 1$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying (22), (23) and  $f(0) = 0$ . Then there exists a unique  $C^*$ -ternary derivation  $\delta : A \rightarrow A$  such that*

$$(25) \quad \|f(x) - \delta(x)\|_A \leq \theta \frac{2 + 3^r}{3 - 3^r} \|x\|_A^r$$

for all  $x \in A$ .

*Proof.* By the same argument as in the proof of Theorem 2.3, one can obtain a  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow B$  satisfying (25).

The rest of the proof is similar to the proof of Theorem 4.1.  $\square$

**Theorem 4.3.** *Let  $r > \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying  $f(0) = 0$  such that*

$$(26) \quad \|D_\mu f(x, y, z)\|_A \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \cdot \|z\|_A^r,$$

$$\|f([x, y, z]) - [f(x), y, z] - [x, f(y), z] - [x, y, f(z)]\|_A$$

$$(27) \quad \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \cdot \|z\|_A^r$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique  $C^*$ -ternary derivation  $\delta : A \rightarrow A$  such that

$$(28) \quad \|f(x) - \delta(x)\|_B \leq \frac{3^r \theta}{27^r - 3} \|x\|_A^{3r}$$

for all  $x \in A$ .

*Proof.* By the same argument as in the proof of Theorem 2.4, one can obtain a  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow B$  satisfying (28).

The rest of the proof is similar to the proof of Theorem 4.1.  $\square$

**Theorem 4.4.** *Let  $r < \frac{1}{3}$  and  $\theta$  be positive real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying (26), (27) and  $f(0) = 0$ . Then there exists a unique  $C^*$ -ternary derivation  $\delta : A \rightarrow A$  such that*

$$(29) \quad \|f(x) - \delta(x)\|_A \leq \frac{3^r \theta}{3 - 27^r} \|x\|_A^{3r}$$

for all  $x \in A$ .

*Proof.* By the same argument as in the proof of Theorem 2.5, one can obtain a  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow B$  satisfying (29).

The rest of the proof is similar to the proofs of Theorems 4.1.  $\square$

## References

- [1] V. Abramov, R. Kerner and B. Le Roy, *Hypersymmetry: a  $\mathbb{Z}_3$ -graded generalization of supersymmetry*, J. Math. Phys. **38** (1997), 1650–1669.
- [2] M. Amyari and M. S. Moslehian, *Approximately ternary semigroup homomorphisms*, Lett. Math. Phys. **77** (2006), 1–9.
- [3] C. Baak and M. S. Moslehian, *On the stability of  $J^*$ -homomorphisms*, Nonlinear Anal.-TMA **63** (2005), 42–48.
- [4] P. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific Publishing Company, New Jersey, Hong Kong, Singapore and London, 2002.
- [5] Y. L. Daletskii and L. Takhtajan, *Leibniz and Lie algebra structures for Nambu algebras*, Lett. Math. Phys. **39** (1997), 127–141.
- [6] Z. Gajda, *On stability of additive mappings*, Internat. J. Math. Math. Sci. **14** (1991), 431–434.
- [7] P. Găvruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184** (1994), 431–436.
- [8] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [9] D. H. Hyers, G. Isac and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [10] D. H. Hyers, G. Isac and Th. M. Rassias, *On the asymptoticity aspect of Hyers-Ulam stability of mappings*, Proc. Amer. Math. Soc. **126** (1998), 425–430.
- [11] G. Isac and Th. M. Rassias, *Stability of  $\psi$ -additive mappings : Applications to non-linear analysis*, Internat. J. Math. Math. Sci. **19** (1996), 219–228.
- [12] S.-M. Jung, *On the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **204** (1996), 221–226.
- [13] R. Kerner, *The cubic chessboard: Geometry and physics*, Classical Quantum Gravity **14** (1997), A203–A225.
- [14] R. Kerner, *Ternary algebraic structures and their applications in physics*, arXiv:math-ph/0011023v1.
- [15] M. S. Moslehian, *Almost derivations on  $C^*$ -ternary rings*, Bull. Belgian Math. Soc.–Simon Stevin **14** (2007), 135–142.
- [16] M. S. Moslehian, *Approximate  $C^*$ -ternary ring homomorphisms*, Bull. Brazilian Math. Soc. **38** (2007), 611–622.
- [17] M. S. Moslehian, *Ternary derivations, stability and physical aspects*, Acta Appl. Math. **100** (2008), 187–199.
- [18] M. S. Moslehian and L. Székelyhidi, *Stability of ternary homomorphisms via generalized Jensen equation*, Result. Math. **49** (2006), 289–300.

- [19] A. Najati, C. Park and J. R. Lee, *Homomorphisms and derivations in  $C^*$ -ternary algebras*, Abstract Appl. Anal. **2009** (2009), Article ID 612392.
- [20] A. Najati and A. Ranjbari, *On homomorphisms between  $C^*$ -ternary algebras*, J. Math. Inequal. **1** (2007), 387–407.
- [21] C. Park, *Homomorphisms between Poisson  $JC^*$ -algebras*, Bull. Braz. Math. Soc. **36** (2005), 79–97.
- [22] C. Park, *Isomorphisms between  $C^*$ -ternary algebras*, J. Math. Anal. Appl. **327** (2007), 101–115.
- [23] C. Park, *Generalized Hyers-Ulam stability of  $C^*$ -ternary algebra homomorphisms*, Dynam. Contin. Discrete Impuls. Systems Ser. A **16** (2009), 67–79.
- [24] C. Park and J. Cui, *Generalized stability of  $C^*$ -ternary quadratic mappings*, Abstract Appl. Anal. **2007** (2007), Article ID 23282.
- [25] C. Park and Th. M. Rassias, *Homomorphisms in  $C^*$ -ternary algebras and  $JB^*$ -triples*, J. Math. Anal. Appl. **337** (2008), 13–20.
- [26] K.-H. Park and Y.-S. Jung, *Perturbations of higher ternary derivations in Banach ternary algebras*, Commun. Korean Math. Soc. **23** (2008), 387–399.
- [27] J. M. Rassias, *On approximation of approximately linear mappings by linear mappings*, J. Funct. Anal. **46** (1982), 126–130.
- [28] J. M. Rassias, *On approximation of approximately linear mappings by linear mappings*, Bull. Sci. Math. **108** (1984), 445–446.
- [29] J. M. Rassias, *Solution of a problem of Ulam*, J. Approx. Theory **57** (1989), 268–273.
- [30] J. M. Rassias, *On the stability of the Euler-Lagrange functional equation*, Chinese J. Math. **20** (1992), 185–190.
- [31] J. M. Rassias, *On the stability of the non-linear Euler-Lagrange functional equation in real normed linear spaces*, J. Math. Phys. Sci. **28** (1994), 231–235.
- [32] J. M. Rassias, *On the stability of the general Euler-Lagrange functional equation*, Demonstratio Math. **29** (1996), 755–766.
- [33] J. M. Rassias, *Solution of the Ulam stability problem for Euler-Lagrange quadratic mappings*, J. Math. Anal. Appl. **220** (1998), 613–639.
- [34] J. M. Rassias and H.-M. Kim, *Approximate homomorphisms and derivations between  $C^*$ -ternary algebras*, J. Math. Phys. **49** 063507 (2008).
- [35] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [36] Th. M. Rassias, *Problem 16; 2*, Report of the 27<sup>th</sup> International Symp. on Functional Equations, Aequationes Math. **39** (1990), 292–293; 309.
- [37] Th. M. Rassias, *The problem of S.M. Ulam for approximately multiplicative mappings*, J. Math. Anal. Appl. **246** (2000), 352–378.
- [38] Th. M. Rassias, *On the stability of functional equations in Banach spaces*, J. Math. Anal. Appl. **251** (2000), 264–284.
- [39] Th. M. Rassias, *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62** (2000), 23–130.
- [40] Th. M. Rassias, *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, Boston and London, 2003.

- [41] Th. M. Rassias and P. Šemrl, *On the behaviour of mappings which do not satisfy Hyers–Ulam stability*, Proc. Amer. Math. Soc. **114** (1992), 989–993.
- [42] M. B. Savadkouhi, M. E. Gordji, J.M. Rassias and N. Ghobadipour, *Approximate ternary Jordan derivations on Banach ternary algebras*, J. Math. Phys. **50** (2009), Article ID 042303.
- [43] M. B. Savadkouhi, M. E. Gordji and N. Ghobadipour, *Stability of homomorphisms and derivations in  $C^*$ -ternary algebras* (preprint).
- [44] F. Skof, *Proprietà locali e approssimazione di operatori*, Rend. Sem. Mat. Fis. Milano **53** (1983), 113–129.
- [45] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ. New York, 1960.
- [46] L. Vainerman and R. Kerner, *On special classes of  $n$ -algebras*, J. Math. Phys. **37** (1996), 2553–2565.
- [47] H. Zettl, *A characterization of ternary rings of operators*, Adv. Math. **48** (1983), 117–143.