

Turán's problem and generalized Ramsey numbers

Zhi-Hong Sun

School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, P.R. China
E-mail: zhihongsun@yahoo.com
Homepage: <http://www.hytc.edu.cn/xsjl/szh>

Abstract

Let n, r, k, s be positive integers with $n, k \geq 2$. The generalized Ramsey number $R(n, r; k, s)$ is the smallest positive integer p such that for every graph G of order p , either G contains a subgraph induced by n vertices with at most $r - 1$ edges, or the complement \overline{G} of G contains a subgraph induced by k vertices with at most $s - 1$ edges. In this paper we completely determine $R(n, n(n - 1)/2 - r; k, 1)$ for $n \geq 4$ and $r \leq n - 2$, and pose several conjectures on Ramsey numbers.

MSC(2010): Primary 05C55, Secondary 05D10, 05C35

Keywords: Independence number; Ramsey number; Turán's theorem

1. Introduction

Let $n, k \geq 2$ be positive integers. The classical Ramsey number $R(n, k)$ is the minimum positive integer such that every graph on $R(n, k)$ vertices has a complete subgraph K_n or an independent set with k vertices. Up to now we only know the following exact values of Ramsey numbers (see [1, p.187] and [8, p.4]):

$$(1.1) \quad \begin{aligned} R(3, 3) = 6, \quad R(3, 4) = 9, \quad R(3, 5) = 14, \quad R(3, 6) = 18, \quad R(3, 7) = 23, \\ R(3, 8) = 28, \quad R(3, 9) = 36, \quad R(4, 4) = 18, \quad R(4, 5) = 25. \end{aligned}$$

In this paper we introduce the following generalized Ramsey numbers.

Definition 1.1. Let n, r, k, s be positive integers with $n, k \geq 2$. We define the generalized Ramsey number $R(n, r; k, s)$ to be the smallest positive integer p such that for every graph G of order p , either G contains a subgraph induced by n vertices with at most $r - 1$ edges, or the complement \overline{G} of G contains a subgraph induced by k vertices with at most $s - 1$ edges.

Clearly $R(n, 1; k, 1) = R(n, k)$. In 1981, Bolze and Harborth [2] introduced the generalized Ramsey number $r_{m,n}(s, t) = R(m, \binom{m}{2} - s + 1; n, \binom{n}{2} - t + 1)$ ($1 \leq s \leq \binom{m}{2}, 1 \leq t \leq \binom{n}{2}$). For some values of $r_{m,n}(s, t)$, see [2, 5, 7].

¹The author is supported by the National Natural Science Foundation of China (grant No. 11371163).

Let L be a family of forbidden graphs. As usual we use $ex(p; L)$ to denote the maximal number of edges in a graph of order p excluding any graphs in L . The general Turán's problem is to evaluate $ex(p; L)$.

For two positive integers p and k , let p_0 be the least nonnegative residue of p modulo k , and let

$$(1.2) \quad t_k(p) = \left(1 - \frac{1}{k}\right) \frac{p^2 - p_0^2}{2} + \frac{p_0^2 - p_0}{2}.$$

In 1941 Turán [11] showed that $ex(p; K_k) = t_{k-1}(p)$ for $k > 1$, where K_k is the complete graph with k vertices. This is now called Turán's theorem. In 1963, Dirac [4] proved a vast generalization of Turán's theorem, see (2.6). Let p, n and t be positive integers satisfying $2 \leq t \leq \frac{n}{2} + 2$ and $p \geq n \geq 4$. With the help of Dirac's generalization of Turán's theorem, in Section 3 we show that if G is a graph of order p and every induced subgraph of G by n vertices has at most $n - t$ edges, then

$$(1.3) \quad \alpha(G) \geq \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1,$$

where $\alpha(G)$ is the independence number of G and $[x]$ is the greatest integer not exceeding x . In Section 4 we show that

$$(1.4) \quad R(n, n(n-1)/2 - r; k, 1) = \begin{cases} \max\{n, k+r\} & \text{if } r \leq \frac{n}{2} - 1, \\ \max\{n, 2k-2 + \left\lfloor \frac{2r+4-n}{3} \right\rfloor\} & \text{if } r > \frac{n}{2} - 1, \end{cases}$$

where k, n, r are positive integers with $k \geq 2$, $n \geq 4$ and $r \leq n - 2$. In the special case $k = n$, (1.4) is known, see [5, Theorem 4]. In Section 5 we obtain an upper bound for $R(4, 3; k, 1)$, and in Section 6 we pose several conjectures on $R(n, k)$ and $R(n, r; k, s)$.

In addition to the above notation, all graphs in the paper are simple graphs, throughout the paper we use the following notation:

\mathbb{N} —the set of positive integers, $\{x\}$ —the fractional part of x , $V(G)$ —the set of vertices in the graph G , $e(G)$ —the number of edges in the graph G , $d(v)$ —the degree of the vertex v in a graph, $\Gamma(v)$ —the set of those vertices adjacent to the vertex v , $\delta(G)$ —the minimal degree of G , $\Delta(G)$ —the maximal degree of G , $g(G)$ —the girth of G , C_n —the cycle with n vertices, $G[V]$ —the subgraph of G induced by the vertices in V , $G - V$ —the subgraph of G obtained by deleting the vertices in V and all edges incident with them.

2. Some applications of the generalization of Turán's theorem

Let L be a family of forbidden graphs, and let $p \in \mathbb{N}$. The famous Erdős-Stone Theorem states that (see [1, pp.122-123])

$$(2.1) \quad ex(p; L) = \frac{1}{2} \left(1 - \frac{1}{\chi(L) - 1}\right) p^2 + o(p^2),$$

where $\chi(L) = \min\{\chi(G) : G \in L\}$ and $\chi(G)$ is the chromatic number of G .

Lemma 2.1. *Let L be a family of forbidden graphs. Let G be a graph of order $p \geq 2$ without any graphs in L and $e(G) = ex(p; L)$. Then*

$$\delta(G) \geq ex(p; L) - ex(p-1; L).$$

Proof. Suppose that v is a vertex in G such that $d(v) = \delta(G)$. Then clearly

$$ex(p; L) - \delta(G) = e(G) - d(v) = e(G - v) \leq ex(p-1; L),$$

which yields the result.

Theorem 2.1. *Let L be a family of forbidden graphs. Let $p > 2$ be a positive integer. Then*

$$ex(p; L) \leq \left\lceil \frac{p \cdot ex(p-1; L)}{p-2} \right\rceil.$$

Proof. Suppose that G is a graph of order p without any graphs in L and $e(G) = ex(p; L)$. Using Lemma 2.1 and Euler's theorem we see that

$$p(ex(p; L) - ex(p-1; L)) \leq p \cdot \delta(G) \leq \sum_{v \in V(G)} d(v) = 2e(G) = 2ex(p; L).$$

Thus,

$$ex(p; L) \leq \frac{p \cdot ex(p-1; L)}{p-2}.$$

As $ex(p; L)$ is an integer, the result follows.

Definition 2.1. *Let n and m be two integers such that $n \geq 1$ and $m \geq 0$. A graph G is said to be an (n, m) graph if every induced subgraph by n vertices has at most m edges. For $p \geq n$ we define $e(n, m; p)$ to be the maximal number of edges in (n, m) graphs of order p .*

By Theorem 2.1, we have

$$(2.2) \quad e(n, m; p) \leq \left\lceil \frac{p \cdot e(n, m; p-1)}{p-2} \right\rceil.$$

This has been given by the author in [9].

Let $p, m, n, r \in \mathbb{N}$ with $p \geq m \geq n \geq 3$ and $r \leq \binom{n}{2}$. Let G be an (n, r) graph of order p with $e(G) = e(n, r; p)$. Then clearly G is an $(m, e(n, r; m))$ graph of order p . As $e(G) = e(n, r; p)$, we have

$$(2.3) \quad e(n, r; p) \leq e(m, e(n, r; m); p).$$

In 1991, the author[9] proved that for $p \geq n \geq 3$,

$$(2.4) \quad e(n, n-2; p) = \left\lceil \frac{(n-2)p}{n-1} \right\rceil \quad \text{and} \quad e(n, n-1; p) = ex(p; \{C_3, \dots, C_n\}).$$

The first formula in (2.4) was first proved by Gol'berg and Gurvich in [6]. Let $t_k(p)$ be given by (1.2), and let $k \geq 2$ be the unique integer such that $t_{k-1}(n) \leq m < t_k(n)$. Using the Erdős-Stone Theorem, in 1991 the author proved that (see [9])

$$(2.5) \quad e(n, m; p) \sim \frac{k-2}{2(k-1)} p^2 \quad \text{as } p \rightarrow +\infty.$$

In [4] Dirac extended Turán's theorem by proving the following result (see also [9]):

$$(2.6) \quad e(n, t_{k-1}(n); p) = t_{k-1}(p) \quad \text{for } p \geq n \geq k \geq 2.$$

For $n \geq 2m \geq 2$ it is clear that $t_{n-m}(n) = \binom{n}{2} - m$. Thus,

$$(2.7) \quad e(n, \frac{n(n-1)}{2} - m; p) = t_{n-m}(p) \quad \text{for } p \geq n \geq 2m \geq 2.$$

Theorem 2.2. *Let $p, n, k \in \mathbb{N}$ with $p \geq n > k$, and let G be a graph of order p . If*

$$e(G) < \left\lfloor \frac{p}{k} \right\rfloor p - \frac{\left\lfloor \frac{p}{k} \right\rfloor (\left\lfloor \frac{p}{k} \right\rfloor + 1)}{2} k,$$

then G contains a subgraph induced by n vertices with at most $\left\lfloor \frac{n}{k} \right\rfloor (n - \frac{\left\lfloor \frac{n}{k} \right\rfloor + 1}{2} k) - 1$ edges.

Proof. Set $s = \left\lfloor \frac{p}{k} \right\rfloor$. Clearly $p = ks + r$ for some $r \in \{0, 1, \dots, k-1\}$. Thus

$$\begin{aligned} \binom{p}{2} - t_k(p) &= \binom{p}{2} - \frac{k-1}{k} \cdot \frac{p^2 - r^2}{2} - \frac{r^2 - r}{2} \\ &= \frac{p-r}{2} \left(\frac{p+r}{k} - 1 \right) = \frac{ks}{2} \left(\frac{p+p-ks}{k} - 1 \right) \\ &= sp - \frac{s(s+1)}{2} k. \end{aligned}$$

Hence, if $e(G) < sp - \frac{s(s+1)}{2} k$, then $e(G) < \binom{p}{2} - t_k(p)$ and so $e(\overline{G}) = \binom{p}{2} - e(G) > t_k(p) = e(n, t_k(n); p)$ by (2.6). Therefore, \overline{G} contains an induced subgraph by n vertices with at least $t_k(n) + 1$ edges. Hence G contains an induced subgraph by n vertices with at most $\binom{n}{2} - t_k(n) - 1$ edges. As $\binom{n}{2} - t_k(n) = \left\lfloor \frac{n}{k} \right\rfloor (n - \frac{\left\lfloor \frac{n}{k} \right\rfloor + 1}{2} k)$, we deduce the result.

Corollary 2.1. *Let $p, n, k \in \mathbb{N}$ with $2 \leq k+1 \leq n \leq 2k$ and $p \geq n$. Let G be a graph of order p satisfying*

$$e(G) < \left\lfloor \frac{p}{k} \right\rfloor p - \frac{\left\lfloor \frac{p}{k} \right\rfloor (\left\lfloor \frac{p}{k} \right\rfloor + 1)}{2} k.$$

Then G contains a subgraph induced by n vertices with at most $n - k - 1$ edges. In particular, for $p \geq n = k+1$ we have $\alpha(G) \geq k+1$.

Proof. Observe that $\left\lfloor \frac{n}{k} \right\rfloor = 1$ or 2 according as $k+1 \leq n < 2k$ or $n = 2k$. We then have $\left\lfloor \frac{n}{k} \right\rfloor (n - \frac{\left\lfloor \frac{n}{k} \right\rfloor + 1}{2} k) - 1 = n - k - 1$. Now applying Theorem 2.2 we deduce the result.

Corollary 2.2. *Let G be a graph of order $p \geq 4$. Let t be a nonnegative integer with $t < \frac{p}{4} - 3\{\frac{p-t}{3}\}$. If $e(G) < p + 2t + 6\{\frac{p-t}{3}\}$, then $\alpha(G) \geq \left\lfloor \frac{p-t}{3} \right\rfloor + 1$.*

Proof. Let $r = p - 3\left\lfloor \frac{p-t}{3} \right\rfloor$. Then $r = p - 3(\frac{p-t}{3} - \{\frac{p-t}{3}\}) = t + 3\{\frac{p-t}{3}\} \geq 0$. Since $t < \frac{p}{4} - 3\{\frac{p-t}{3}\}$ we have $4\{\frac{p-t}{3}\} < \frac{p-t}{3}$ and so

$$r = t + 3\left\lfloor \frac{p-t}{3} \right\rfloor < \frac{p-t}{3} - \left\lfloor \frac{p-t}{3} \right\rfloor = \left\lceil \frac{p-t}{3} \right\rceil.$$

Thus $\left\lfloor \frac{p}{\left\lceil \frac{p-t}{3} \right\rceil} \right\rfloor = 3$. Hence,

$$\left\lfloor \frac{p}{\left\lceil \frac{p-t}{3} \right\rceil} \right\rfloor \left(p - \frac{\left\lfloor \frac{p}{\left\lceil \frac{p-t}{3} \right\rceil} \right\rfloor + 1}{2} \left\lceil \frac{p-t}{3} \right\rceil \right)$$

$$= 3(p - 2\lceil \frac{p-t}{3} \rceil) = 3p - 6(\frac{p-t}{3} - \{\frac{p-t}{3}\}) = p + 2t + 6\{\frac{p-t}{3}\}.$$

Since $t \geq 0 > 3 - 2p$ we see that $p > \frac{p-t}{3} + 1 \geq \lceil \frac{p-t}{3} \rceil + 1$. We also have $\lceil \frac{p-t}{3} \rceil \geq \lceil \frac{p-p/4}{3} \rceil = \lceil \frac{p}{4} \rceil \geq 1$. Now applying the above and taking $k = \lceil \frac{p-t}{3} \rceil$ in Corollary 2.1 we obtain the result.

Theorem 2.3. *Let G be a graph of order $p \geq 4$. Let t be a nonnegative integer so that $3\{\frac{p+t}{3}\} - \frac{p}{4} < t \leq \frac{p}{2} + 3\{\frac{p+t}{3}\}$. If*

$$e(G) < p - 2t + 3\left\{\frac{p+t}{3}\right\} + \max\left\{t, 3\left\{\frac{p+t}{3}\right\}\right\},$$

then $\alpha(G) \geq \lceil \frac{p+t}{3} \rceil + 1$.

Proof. It is clear that

$$\begin{aligned} p - 2\lceil \frac{p+t}{3} \rceil &= p - 2(\frac{p+t}{3} - \{\frac{p+t}{3}\}) = \frac{1}{3}(p - 2t + 6\{\frac{p+t}{3}\}) \geq 0, \\ p - 3\lceil \frac{p+t}{3} \rceil &= p - 3(\frac{p+t}{3} - \{\frac{p+t}{3}\}) = 3\{\frac{p+t}{3}\} - t \end{aligned}$$

and

$$p - 4\lceil \frac{p+t}{3} \rceil = p - 4(\frac{p+t}{3} - \{\frac{p+t}{3}\}) = -\frac{p+4t-12\{\frac{p+t}{3}\}}{3} < 0.$$

Thus,

$$\left\lceil \frac{p}{\lceil \frac{p+t}{3} \rceil} \right\rceil = \begin{cases} 2 & \text{if } t > 3\{\frac{p+t}{3}\}, \\ 3 & \text{if } t \leq 3\{\frac{p+t}{3}\}. \end{cases}$$

Set $k = \lceil \frac{p+t}{3} \rceil$. We see that

$$\begin{aligned} &\left\lceil \frac{p}{k} \right\rceil \left(p - \frac{\lceil \frac{p}{k} \rceil + 1}{2} k \right) \\ &= \begin{cases} 2p - 3k = 2p - 3(\frac{p+t}{3} - \{\frac{p+t}{3}\}) = p - t + 3\{\frac{p+t}{3}\} & \text{if } t > 3\{\frac{p+t}{3}\}, \\ 3p - 6k = 3p - 6(\frac{p+t}{3} - \{\frac{p+t}{3}\}) = p - 2t + 6\{\frac{p+t}{3}\} & \text{if } t \leq 3\{\frac{p+t}{3}\}, \end{cases} \\ &= p - 2t + 3\left\{\frac{p+t}{3}\right\} + \max\left\{t, 3\left\{\frac{p+t}{3}\right\}\right\}. \end{aligned}$$

Since $p \geq 4$ and $0 \leq t < \frac{p}{2} + 3$ we see that $\lceil \frac{p+t}{3} \rceil \geq 1$ and $p \geq \frac{p+\frac{p}{2}+3}{3} + 1 \geq \lceil \frac{p+t}{3} \rceil + 1$. Now applying the above and Corollary 2.1 we obtain the result.

Putting $t = 0, 1, 2$ in Theorem 2.3 we deduce the following corollaries.

Corollary 2.3. *Let G be a graph of order $p \geq 9$. If $e(G) < p + 6\{\frac{p}{3}\}$, then $\alpha(G) \geq \lceil \frac{p}{3} \rceil + 1$.*

Corollary 2.4. *Let G be a graph of order $p \geq 5$. If*

$$e(G) < \begin{cases} p & \text{if } 3 \mid p, \\ p+2 & \text{if } 3 \mid p-1, \\ p-1 & \text{if } 3 \mid p-2, \end{cases}$$

then $\alpha(G) \geq \lceil \frac{p+1}{3} \rceil + 1$.

Corollary 2.5. Let G be a graph of order $p \geq 4$. If

$$e(G) < \begin{cases} p & \text{if } 3 \mid p, \\ p-2 & \text{if } 3 \mid p-1, \\ p-1 & \text{if } 3 \mid p-2, \end{cases}$$

then $\alpha(G) \geq \lceil \frac{p+2}{3} \rceil + 1$.

Theorem 2.4. Let $p, n, t \in \mathbb{N}$ with $t \leq \frac{n}{2} + 2$ and $p > n + 7 - 2t$. If G is a graph of order p and $e(G) < \frac{p+n}{2} + 1 - t$, then $\alpha(G) \geq \lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \rceil + 1$.

Proof. Set $r = p - 2 \lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \rceil$. Then

$$r \geq p - 2 \cdot \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} = \lceil \frac{n+4-2t}{3} \rceil \geq 0$$

and

$$r = p - 2 \left(\frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} - \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} \right) = 2 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} + \lceil \frac{n+4-2t}{3} \rceil.$$

Hence

$$\begin{aligned} & \left\lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\rceil - r \\ &= \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} - \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} - 2 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} - \lceil \frac{n+4-2t}{3} \rceil \\ &= \frac{p - 3 \lceil \frac{n+4-2t}{3} \rceil}{2} - 3 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} \\ &= \frac{p - 3 \left(\frac{n+4-2t}{3} - \left\{ \frac{n+4-2t}{3} \right\} \right)}{2} - 3 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} \\ &= \frac{p - (n+4-2t)}{2} + \frac{3}{2} \left\{ \frac{n+4-2t}{3} \right\} - 3 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} > \frac{3}{2} + 0 - 3 \cdot \frac{1}{2} = 0. \end{aligned}$$

So we have

$$\left\lceil \frac{p}{\lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \rceil} \right\rceil = 2.$$

As

$$\begin{aligned} 2p - \frac{2 \times 3}{2} \left\lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\rceil &= 2p - 3 \left(\frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} - \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} \right) \\ &= \frac{p}{2} + \frac{3}{2} \left(\frac{n+4-2t}{3} - \left\{ \frac{n+4-2t}{3} \right\} \right) + 3 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} \\ &= \frac{p}{2} + \frac{n+4-2t}{2} + 3 \left\{ \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\} - \frac{3}{2} \left\{ \frac{n+4-2t}{3} \right\} \\ &\geq \frac{p+n}{2} + 2 - t + 0 - 1, \end{aligned}$$

we see that

$$e(G) < \frac{p+n}{2} + 1 - t \leq 2p - \frac{2 \times 3}{2} \left\lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \right\rceil.$$

Since $\lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \rceil \geq \lceil \frac{p - \frac{n+4-2t}{3}}{2} \rceil = \lceil \frac{3p - (n+4-2t)}{6} \rceil \geq 1$ and $p \geq 2 \lceil \frac{p - \lceil \frac{n+4-2t}{3} \rceil}{2} \rceil$, by the above and Corollary 2.1 we deduce the result.

3. The best lower bound for independence numbers of (n, m) graphs ($m \leq n - 2$)

Theorem 3.1. *Let $p, m, n \in \mathbb{N}$ with $p \geq n \geq 4$ and $m \leq \frac{n}{2} - 1$. If G is an (n, m) graph of order p , then $\alpha(G) \geq p - m$.*

Proof. We prove the theorem by induction on p . Set $k = n - 1 - m$. Then $3 \leq k + 1 \leq n \leq 2k$ and so $\lfloor \frac{n}{k} \rfloor = 1$ or 2 . Thus, if G is a graph of order n with $e(G) \leq m$, then

$$e(G) \leq m < \left\lfloor \frac{n}{k} \right\rfloor n - \frac{\left\lfloor \frac{n}{k} \right\rfloor (\left\lfloor \frac{n}{k} \right\rfloor + 1)}{2} k.$$

Since $n \geq k + 1$, applying Corollary 2.1 we see that $\alpha(G) \geq k + 1 = n - m$. So the result holds for $p = n$.

Now assume $p > n$. Let G be an (n, m) graph of order p . We assert that G has an isolated vertex. Otherwise, we have

$$e(G) = \frac{1}{2} \sum_{v \in V(G)} d(v) \geq \frac{p}{2} > \frac{n}{2} \geq \left\lfloor \frac{n}{2} \right\rfloor.$$

We choose $\lfloor \frac{n}{2} \rfloor$ edges in G and consider a subgraph H induced by the $\lfloor \frac{n}{2} \rfloor$ edges. Clearly H has at most n vertices and $e(H) = \lfloor \frac{n}{2} \rfloor > \frac{n}{2} - 1 \geq m$. This contradicts the fact that G is an (n, m) graph. So G has an isolated vertex. Assume that v is an isolated vertex of G and the result holds for all (n, m) graphs of order $p - 1$. By the above we have

$$\alpha(G) = 1 + \alpha(G - v) \geq 1 + p - 1 - m = p - m.$$

Hence the theorem is proved by induction.

Lemma 3.1. *Let $n, m \in \mathbb{N}$ with $m \leq n - 2$. If G is an (n, m) graph of order $n + 1$, then $e(G) \leq m + 1$.*

Proof. By (2.2) we have

$$e(n, m; n + 1) \leq \left\lfloor \frac{(n + 1)e(n, m; n)}{n - 1} \right\rfloor = \left\lfloor \frac{(n + 1)m}{n - 1} \right\rfloor = m + \left\lfloor \frac{2m}{n - 1} \right\rfloor$$

As $m < n - 1$ we see that $\lfloor \frac{2m}{n - 1} \rfloor < 2$. Thus

$$e(G) \leq e(n, m; n + 1) \leq m + \left\lfloor \frac{2m}{n - 1} \right\rfloor \leq m + 1.$$

This is the result.

Lemma 3.2. *Let $p, m, n \in \mathbb{N}$ with $p \geq n \geq m + 2$. Let G be an (n, m) graph of order p . Then one of the components of G is a tree.*

Proof. Suppose that G_1, G_2, \dots, G_r are all components of G and $|V(G_i)| = p_i$ for $i = 1, 2, \dots, r$. Then $p_1 + \dots + p_r = p \geq n$. Since $m < n - 1$ and G is an (n, m) graph, G cannot contain a tree on n vertices as a subgraph. As every connected graph has a spanning tree, we must have $p_i < n$ for all $i \in \{1, 2, \dots, r\}$. Let us choose $j \in \{1, 2, \dots, r - 1\}$ so that $p_1 + \dots + p_j < n$ and $p_1 + \dots + p_j + p_{j+1} \geq n$. Then clearly G_{j+1} has a subtree T on $n - (p_1 + \dots + p_j)$ vertices. All the n vertices in $G_1 \cup \dots \cup G_j \cup T$ induce a subgraph

of G with at least $e(G_1) + \dots + e(G_j) + e(T)$ edges. If G_1, \dots, G_r are not trees, then $e(G_i) \geq p_i$ for $i = 1, 2, \dots, r$ and so

$$e(G_1) + \dots + e(G_j) + e(T) \geq p_1 + \dots + p_j + n - (p_1 + \dots + p_j) - 1 = n - 1.$$

Thus, the n vertices in $G_1 \cup \dots \cup G_j \cup T$ induce a subgraph of G with at least $n - 1$ edges. This contradicts the assumption. Hence one of G_1, \dots, G_r is a tree. This proves the lemma.

Lemma 3.3. *Let T be a tree with at least 3 vertices. Then there are two vertices u and v in T such that $\alpha(T) > \alpha(T - \{u, v\})$.*

Proof. It is well known that T has a vertex u such that $d(u) = 1$. Let v be the unique vertex of G adjacent to u . Let S be an independent set of $T - \{u, v\}$. Then clearly $S \cup \{u\}$ is an independent set of T . Thus $\alpha(T) > \alpha(T - \{u, v\})$.

Theorem 3.2. *Let $p, n, t \in \mathbb{N}, 2 \leq t \leq \frac{n}{2} + 2$ and $p \geq n \geq 4$. If G is an $(n, n - t)$ graph of order p , then*

$$\alpha(G) \geq \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1.$$

Proof. We prove the theorem by induction on p . Now suppose $p = n$ or $n + 1$. Let G be an $(n, n - t)$ graph of order p . For $t \geq 4$ and $p = n$ we have $e(G) \leq n - t < \frac{p+n}{2} + 1 - t$. For $t \geq 4$ and $p = n + 1$, by Lemma 3.1 we also have $e(G) \leq n - t + 1 < \frac{p+n}{2} + 1 - t$. Thus applying Theorem 2.4 we see that $\alpha(G) \geq \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1$. For $t = 3$ and $p \in \{n, n + 1\}$ it is easily seen that $\left\lceil \frac{p - \left\lfloor \frac{n-2}{3} \right\rfloor}{2} \right\rceil = \left\lceil \frac{p+2}{3} \right\rceil$. By Lemma 3.2 we have $e(G) \leq n - t + 1 = p - 3$ for $p = n + 1$ and $e(G) \leq n - t = p - 3$ for $p = n$. Thus, by Corollary 2.5 we have

$$\alpha(G) \geq \left\lceil \frac{p+2}{3} \right\rceil + 1 = \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1.$$

For $t = 2$ and $p \in \{n, n + 1\}$ it is easily seen that $\left\lceil \frac{p - \left\lfloor \frac{n}{3} \right\rfloor}{2} \right\rceil = \left\lceil \frac{p+1}{3} \right\rceil$. By Lemma 3.1 we have $e(G) \leq n - t + 1 = p - 2$ for $p = n + 1$ and $e(G) \leq n - t = p - 2$ for $p = n$. Thus, by Corollary 2.4 we have

$$\alpha(G) \geq \left\lceil \frac{p+1}{3} \right\rceil + 1 = \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1 \quad \text{for } p \geq 5.$$

When $p = n = 4$ and $t = 2$, we also have $\alpha(G) \geq 2 = \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1$. Summarizing the above we see that the result is true for $p \in \{n, n + 1\}$.

Now we assume $p \geq n + 2$ and the result holds for all $(n, n - t)$ graphs with at most $p - 1$ vertices. Let G be an $(n, n - t)$ graph of order p . By Lemma 3.2, one of the components of G is a tree. Let T be such a tree. If $T \cong K_1$, for any vertex u in $G - T$ we have $\alpha(G) = 1 + \alpha(G - T) \geq 1 + \alpha(G - \{T, u\})$. If $T \cong K_2$, it is clear that $\alpha(G) > \alpha(G - T)$. If T is a tree with at least three vertices, by Lemma 3.3 there are two vertices u and v in T such that $\alpha(G) > \alpha(G - \{u, v\})$. Hence, in any cases, there are two vertices u and v in G so that $\alpha(G) > \alpha(G - \{u, v\})$. Clearly $G - \{u, v\}$ is an $(n, n - t)$ graph of order $p - 2$. Thus, by the inductive hypothesis we have

$$\alpha(G) \geq 1 + \alpha(G - \{u, v\}) \geq 1 + \left\lceil \frac{p - 2 - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1 = \left\lceil \frac{p - \left\lfloor \frac{n+4-2t}{3} \right\rfloor}{2} \right\rceil + 1.$$

So the theorem is proved by induction.

4. Evaluation of $R(n, n(n-1)/2 - r; k, 1)$ ($r \leq n-2$)

Let n, r, k be positive integers with $r \leq \binom{n}{2}$. By Definition 1.1, $R(n, n(n-1)/2 - r; k, 1)$ is the smallest positive integer p such that for any (n, r) graph G of order p , we have $\alpha(G) \geq k$.

Now we are in a position to prove the following main result.

Theorem 4.1. *Let $k, n, r \in \mathbb{N}$ with $k \geq 2$, $n \geq 4$ and $r \leq n-2$. Then*

$$R(n, n(n-1)/2 - r; k, 1) = \begin{cases} \max\{n, k+r\} & \text{if } r \leq \frac{n}{2} - 1, \\ \max\{n, 2k-2 + \lceil \frac{2r+4-n}{3} \rceil\} & \text{if } r > \frac{n}{2} - 1. \end{cases}$$

Proof. We first assume $r \leq \frac{n}{2} - 1$. If $k+r \leq n$, putting $p = n$ and $m = r$ in Theorem 3.1 we see that $\alpha(G) \geq n-r \geq k$ for any graph G of order n with $e(G) \leq r$. Thus, $R(n, \binom{n}{2} - r; k, 1) \leq n$ and so $R(n, \binom{n}{2} - r; k, 1) = n = \max\{n, k+r\}$. Now suppose $k+r > n$. Then $k > n-r \geq n - (\frac{n}{2} - 1) = \frac{n}{2} + 1 > r$. Set $G = rK_2 \cup (k-r-1)K_1$. It is easily seen that G is a graph of order $k+r-1$ with $\alpha(G) = k-1$. For any n vertices in G , the corresponding induced subgraph by the n vertices must be given by $aK_2 \cup bK_1$, where a and b are nonnegative integers such that $2a+b = n$ and $a \leq r$. Thus, G is an (n, r) graph of order $k+r-1$. On the other hand, as $\alpha(G) = k-1$, G has no independent sets with k vertices. Hence, $R(n, n(n-1)/2 - r; k, 1) > k+r-1$. Since $k+r > n$, it follows from Theorem 3.1 that for any (n, r) graph G' of order $k+r$, $\alpha(G') \geq k+r-r = k$. Hence $R(n, n(n-1)/2 - r; k, 1) \leq k+r$ and so $R(n, n(n-1)/2 - r; k, 1) = k+r = \max\{n, k+r\}$.

Now assume $r \geq \frac{n-1}{2}$. We first suppose $r \leq 2n-3k+2$. Let G be a graph of order n with $e(G) \leq r$. Taking $p = n$ and $t = n-r$ in Theorem 3.2 we see that

$$\alpha(G) \geq \left\lceil \frac{n - \lceil \frac{n+4-2(n-r)}{3} \rceil}{2} \right\rceil + 1 = \left\lceil \frac{n - \lceil \frac{2r+4-n}{3} \rceil}{2} \right\rceil + 1 \geq \left\lceil \frac{n - \lceil \frac{2(2n-3k+2)+4-n}{3} \rceil}{2} \right\rceil + 1 = k.$$

Thus, $R(n, \frac{n(n-1)}{2} - r; k, 1) \leq n$. On the other hand, clearly $R(n, \frac{n(n-1)}{2} - r; k, 1) > n-1$. So $R(n, \frac{n(n-1)}{2} - r; k, 1) = n$. To see the result, we note that

$$2k-2 + \lceil \frac{2r+4-n}{3} \rceil \leq 2k-2 + \lceil \frac{2(2n-3k+2)+4-n}{3} \rceil = n.$$

Now we suppose $r > 2n-3k+2$. In this case, we have

$$2k-2 + \lceil \frac{2r+4-n}{3} \rceil \geq 2k-2 + \lceil \frac{2(2n-3k+2)+4-n}{3} \rceil = n$$

and

$$\begin{aligned} 0 &\leq \frac{2r+1-n}{3} = \frac{n+1-2(n-r)}{3} < \frac{3k-2-(n-r)+1-2(n-r)}{3} \\ &= k - \frac{1}{3} - (n-r) \leq k - \frac{1}{3} - 2 < k-1 \end{aligned}$$

and so

$$0 \leq \lceil \frac{2r+1-n}{3} \rceil \leq \frac{2r+1-n}{3} < k-1.$$

Set

$$G = \left\lceil \frac{2r+1-n}{3} \right\rceil K_3 \cup \left(k-1 - \left\lceil \frac{2r+1-n}{3} \right\rceil \right) K_2.$$

We claim that G is an (n, r) graph. Clearly any induced subgraph of G by n vertices can be written as $xK_1 \cup yK_2 \cup zK_3$, where x, y, z are nonnegative integers satisfying $x + 2y + 3z = n$ and $z \leq \left\lceil \frac{2r+1-n}{3} \right\rceil$. If $x + y \leq n - r - 1$, then

$$\begin{aligned} \frac{n+1-2(n-r)}{3} &\geq \left\lceil \frac{2r+1-n}{3} \right\rceil \geq z = \frac{n-2(x+y)+x}{3} \\ &\geq \frac{n-2(n-r-1)}{3} = \frac{n+2-2(n-r)}{3}. \end{aligned}$$

This is a contradiction. Hence $x + y \geq n - r$ and so

$$e(xK_1 \cup yK_2 \cup zK_3) = y + 3z = n - (x + y) \leq r.$$

This shows that G is an (n, r) graph. It is evident that

$$\alpha(G) = \left\lceil \frac{2r+1-n}{3} \right\rceil + k-1 - \left\lceil \frac{2r+1-n}{3} \right\rceil = k-1$$

and

$$|V(G)| = 3 \left\lceil \frac{2r+1-n}{3} \right\rceil + 2 \left(k-1 - \left\lceil \frac{2r+1-n}{3} \right\rceil \right) = 2k-2 + \left\lceil \frac{2r+1-n}{3} \right\rceil.$$

As G has no independent sets with k vertices and G does not contain any subgraphs with n vertices and at least $r+1$ edges, we must have

$$R\left(n, \frac{n(n-1)}{2} - r; k, 1\right) > |V(G)| = 2k-2 + \left\lceil \frac{2r+1-n}{3} \right\rceil.$$

On the other hand, if G is an (n, r) graph of order $2k-2 + \left\lceil \frac{2r+4-n}{3} \right\rceil$, by Theorem 3.2 we have

$$\alpha(G) \geq \left\lceil \frac{2k-2 + \left\lceil \frac{2r+4-n}{3} \right\rceil - \left\lceil \frac{n+4-2(n-r)}{3} \right\rceil}{2} \right\rceil + 1 = k.$$

Hence

$$R\left(n, \frac{n(n-1)}{2} - r; k, 1\right) \leq 2k-2 + \left\lceil \frac{2r+4-n}{3} \right\rceil$$

and so

$$R\left(n, \frac{n(n-1)}{2} - r; k, 1\right) = 2k-2 + \left\lceil \frac{2r+4-n}{3} \right\rceil = \max\left\{n, 2k-2 + \left\lceil \frac{2r+4-n}{3} \right\rceil\right\}.$$

Summarizing the above we prove the theorem.

Corollary 4.1. *Let k and n be positive integers with $k \geq 2$ and $n \geq 4$. Then*

$$R(n, n(n-1)/2 - n + 2; k, 1) = \begin{cases} n & \text{if } n \geq 3k-4, \\ 2k-2 + \left\lceil \frac{n}{3} \right\rceil & \text{if } n < 3k-4. \end{cases}$$

Proof. Putting $r = n - 2$ in Theorem 4.1 we obtain the result.

Definition 4.1. Let $k, s \in \mathbb{N}$ with $s \leq \binom{k}{2}$. Let G be a graph with at least k vertices. If every subgraph of G induced by k vertices has at least s edges, we say that G satisfies the (k, s) condition.

Lemma 4.1. Let $p, n, k, r, s \in \mathbb{N}$ with $n, k \geq 2$, $r < \binom{n}{2}$, $s < \binom{k}{2}$ and $p \geq \max\{n, k\}$. Let G be a $(n, \binom{n}{2} - r)$ graph of order p satisfying the (k, s) condition. Then for any $v \in V(G)$,

$$p - R(n, r; k - 1, s) \leq d(v) \leq R(n - 1, r; k, s) - 1.$$

Proof. Let x be a vertex in G such that $d(x) = \Delta(G)$. If $\Delta(G) \geq R(n - 1, r; k, s)$, then $\Delta(G) \geq n - 1$ and there are $n - 1$ vertices $x_1, \dots, x_{n-1} \in \Gamma(x)$ such that $e(G[\{x_1, \dots, x_{n-1}\}]) > \binom{n-1}{2} - r$. As $\binom{n-1}{2} - r + n - 1 = \binom{n}{2} - r$, we see that $e(G[\{x, x_1, \dots, x_{n-1}\}]) > \binom{n}{2} - r$. This contradicts the assumption. Hence $\Delta(G) < R(n - 1, r; k, s)$ and so $d(v) \leq \Delta(G) \leq R(n - 1, r; k, s) - 1$ for any vertex v in G .

Let y be a vertex in G such that $d(y) = \delta(G)$, and let $V_y = V(G) - \{y\} \cup \Gamma(y)$. Then clearly $|V_y| = p - 1 - d(y) = p - 1 - \delta(G)$. If $\delta(G) \geq p - R(n, r; k - 1, s)$, then $d(v) \geq \delta(G) \geq p - R(n, r; k - 1, s)$ for any $v \in V(G)$. Now assume $\delta(G) \leq p - R(n, r; k - 1, s) - 1$. As $R(n, r; k - 1, s) \geq k - 1$ we see that $\delta(G) \leq p - k$ and so $|V_y| = p - 1 - \delta(G) \geq k - 1$. For any $k - 1$ vertices v_1, \dots, v_{k-1} in V_y , by the assumption we have $e(G[\{v_1, \dots, v_{k-1}\}]) = e(G[\{y, v_1, \dots, v_{k-1}\}]) \geq s$. As G is a $(n, \binom{n}{2} - r)$ graph, we see that $G[V_y]$ is also a $(n, \binom{n}{2} - r)$ graph. Hence, by Definition 1.1 we have $p - 1 - \delta(G) < R(n, r; k - 1, s)$. Therefore, for any vertex v in G , $d(v) \geq \delta(G) \geq p - R(n, r; k - 1, s)$, which completes the proof.

Theorem 4.2. Let n, r, k, s be positive integers with $n, k \geq 3$. Then

$$R(n, r; k, s) \leq R(n - 1, r; k, s) + R(n, r; k - 1, s).$$

Moreover, the strict inequality holds when both $R(n - 1, r; k, s)$ and $R(n, r; k - 1, s)$ are even.

Proof. Clearly $R(n - 1, r; k, s) + R(n, r; k - 1, s) \geq \max\{n - 1 + n, k + k - 1\} > \max\{n, k\}$. Thus, if $R(n, r; k, s) = \max\{n, k\}$, the result is true. Now assume $R(n, r; k, s) > \max\{n, k\}$. By the definition of $R(n, r; k, s)$, there is a $(n, \binom{n}{2} - r)$ graph G of order $R(n, r; k, s) - 1$ satisfying the (k, s) condition. For any vertex v in G , by Lemma 4.1 we have

$$R(n, r; k, s) - 1 - R(n, r; k - 1, s) \leq d(v) \leq R(n - 1, r; k, s) - 1.$$

It then follows that

$$R(n, r; k, s) \leq R(n - 1, r; k, s) + R(n, r; k - 1, s).$$

Moreover, the equality holds if and only if G is a regular graph with degree $R(n - 1, r; k, s) - 1$. If the equality holds, we must have

$$\begin{aligned} 2e(G) &= (R(n, r; k, s) - 1)(R(n - 1, r; k, s) - 1) \\ &= (R(n - 1, r; k, s) + R(n, r; k - 1, s) - 1)(R(n - 1, r; k, s) - 1). \end{aligned}$$

This shows that either $R(n - 1, r; k, s)$ or $R(n, r; k - 1, s)$ is odd. Hence, if $R(n - 1, r; k, s)$ and $R(n, r; k - 1, s)$ are even, we have $R(n, r; k, s) < R(n - 1, r; k, s) + R(n, r; k - 1, s)$, which completes the proof.

Theorem 4.2 can be viewed as a generalization of the classical inequality ([1]) $R(n, k) \leq R(n-1, k) + R(n, k-1)$ for $n, k \geq 3$.

Theorem 4.3. *Let $k, n, r \in \mathbb{N}$ with $r \leq n-1$. Then*

(i) $R(n, r; k, 1) \geq R(n, r; k-1, 1) + n-1$.

(ii) $R(n, r; k, 1) \geq (n-1)(k-1) + 1$.

Proof. We first consider (i). By the definition of $R(n, r; k-1, 1)$, there is a $(k-1, \binom{k-1}{2} - 1)$ graph G of order $R(n, r; k-1, 1) - 1$ satisfying the (n, r) condition. Now we construct a new graph G' by adding $n-1$ new vertices to G and joining every new vertex and each vertex in G . Since G does not contain a copy of K_{k-1} , we see that G' does not contain a copy of K_k . For fixed $s \in \{1, 2, \dots, n-1\}$, if we choose n vertices in G' containing exactly s vertices in G , then the subgraph of G' induced by the n vertices has at least $s(n-s)$ edges. As $s(n-s) \geq n-1 \geq r$ and G satisfies the (n, r) condition, we see that G' also satisfies the (n, r) condition. Note that the order of G' is $R(n, r; k-1, 1) - 1 + n - 1$. We then have $R(n, r; k, 1) > R(n, r; k-1, 1) - 1 + n - 1$. This proves (i).

Now let us consider (ii). It is clear that $R(n, r; 2, 1) = n$. Thus, using (i) we see that

$$\begin{aligned} R(n, r; k, 1) &= R(n, r; 2, 1) + \sum_{s=3}^k (R(n, r; s, 1) - R(n, r; s-1, 1)) \\ &\geq n + (k-2)(n-1) = (n-1)(k-1) + 1. \end{aligned}$$

This proves (ii) and hence the theorem is proved.

5. The upper bound for $R(4, 3; k, 1)$

Let $k \in \{2, 3, \dots\}$. By Definition 1.1, $R(4, 3; k, 1)$ is the smallest positive integer p such that for any graph G of order p , either G has a subgraph induced by 4 vertices with at least 4 edges, or G contains an independent set with k vertices. Every subgraph of $(k-1)K_3$ induced by 4 vertices has at most three edges and the independence number of $(k-1)K_3$ is $k-1$. Thus $R(4, 3; k, 1) > 3(k-1)$.

Theorem 5.1. *For $k = 3, 4, 5, 6$ we have $R(4, 3; k, 1) = 3k - 2$.*

Proof. By the previous argument or Theorem 4.3(ii) we have $R(4, 3; k, 1) \geq 3k - 2$. Clearly $R(4, 3; 2, 1) = 4$ and $R(3, 3; 3, 1) = 3$. Thus, using Theorem 4.2 we see that

$$R(4, 3; 3, 1) \leq R(4, 3; 2, 1) + R(3, 3; 3, 1) = 4 + 3 = 7.$$

Hence $R(4, 3; 3, 1) = 7$.

For $k \in \{4, 5, 6\}$ let G be a $(4, 3)$ graph of order $3k - 2$. If $\alpha(G) < k$, for any vertex v in G , by Lemma 4.1 we have

$$(5.1) \quad 3k - 2 - R(4, 3; k-1, 1) \leq d(v) \leq R(3, 3; k, 1) - 1 = k - 1.$$

For a $(4, 3)$ graph G of order 10 with $\alpha(G) < 4$, it follows from (5.1) that G is a 3-regular graph with $g(G) \geq 5$. As $R(3, 4) = 9$, we must have $\alpha(G) \geq 4$. This is a contradiction. Hence for any $(4, 3)$ graph G of order 10 we have $\alpha(G) \geq 4$ and so $R(4, 3; 4, 1) = 10$.

Now let G be a $(4, 3)$ graph of order 13. If $\alpha(G) < 5$, from (5.1) we see that $3 \leq d(v) \leq 4$ for any $v \in V(G)$. If $d(v) = \Delta(G) = 4$, $\Gamma(v) = \{v_1, v_2, v_3, v_4\}$ and $x_1, x_2 \in \Gamma(v_1) - \{v\}$, then clearly $\{x_1, x_2, v_2, v_3, v_4\}$ is an independent set of G and so $\alpha(G) \geq 5$. If G is a

3-regular $(4, 3)$ graph of order 13, then $g(G) \geq 5$ and for given $v \in V(G)$ there are three vertices $u_i (i = 1, 2, 3)$ in G such that $d(u_i, v) \geq 3$ ($i = 1, 2, 3$). As $g(G) \geq 5$, we may assume that u_i is not adjacent to u_j , then clearly $\{u_i, u_j\} \cup \Gamma(v)$ is an independent set of G and so $\alpha(G) \geq 5$. By the above, for any $(4, 3)$ graph G of order 13 we have $\alpha(G) \geq 5$ and so $R(4, 3; 5, 1) = 13$.

Suppose that G is a $(4, 3)$ graph of order 16. If $\alpha(G) < 6$, from (5.1) we see that $3 \leq d(v) \leq 5$ for any $v \in V(G)$. If $d(v) = \Delta(G) = 5$, $\Gamma(v) = \{v_1, \dots, v_5\}$ and $x_1, x_2 \in \Gamma(v_1) - \{v\}$, then clearly $\{x_1, x_2, v_2, v_3, v_4, v_5\}$ is an independent set of G and so $\alpha(G) \geq 6$. If $d(v) = \Delta(G) = 4$, $\Gamma(v) = \{v_1, \dots, v_4\}$ and $\Gamma(v_1) = \{v, x_1, x_2, x_3\}$, then clearly $\{x_1, x_2, x_3, v_2, v_3, v_4\}$ is an independent set of G and so $\alpha(G) \geq 6$. If $d(v) = \Delta(G) = 4$, $\Gamma(v) = \{v_1, \dots, v_4\}$ and $d(v_1) = \dots = d(v_4) = 3$, there are three vertices $u_i (i = 1, 2, 3)$ in G such that $d(u_i, v) \geq 3$ ($i = 1, 2, 3$). As $g(G) \geq 5$, we may assume that u_i is not adjacent to u_j . Then clearly $\{u_i, u_j, v_1, \dots, v_4\}$ is an independent set of G and so $\alpha(G) \geq 6$. If $d(v) = \Delta(G) = 3$ and $\Gamma(v) = \{v_1, v_2, v_3\}$, then G is 3-regular and there are six vertices $u_i (i = 1, 2, \dots, 6)$ in G such that $d(u_i, v) \geq 3$ ($i = 1, 2, \dots, 6$). As $g(G) \geq 5$ and $R(3, 3) = 6$, there are three vertices $u_i, u_j, u_k \in \{u_1, \dots, u_6\}$ such that $\{u_i, u_j, u_k\}$ is an independent set and so $\{u_i, u_j, u_k, v_1, v_2, v_3\}$ is also an independent set of G . This shows that $\alpha(G) \geq 6$. Hence, by the above, for any $(4, 3)$ graph G of order 16 we have $\alpha(G) \geq 6$ and so $R(4, 3; 6, 1) = 16$. This completes the proof.

Theorem 5.2. *For $k = 3, 4, 5, \dots$ we have*

$$R(4, 3; k, 1) \leq R(4, 3; k-1, 1) + \frac{3}{2} + \sqrt{R(4, 3; k-1, 1) - \frac{3}{4}}.$$

Proof. By the definition of $R(4, 3; k, 1)$, there is a $(4, 3)$ graph G of order $R(4, 3; k, 1) - 1$ with $\alpha(G) < k$. For any vertex v in G , by Lemma 4.1 we have

$$(5.2) \quad R(4, 3; k, 1) - 1 - R(4, 3; k-1, 1) \leq d(v) \leq R(3, 3; k, 1) - 1 = k - 1.$$

If $\delta(G) \leq 2$, as $R(4, 3; k, 1) \geq R(4, 3; 3, 1) = 7$ we have $\delta(G) \leq \sqrt{R(4, 3; k, 1) - 2}$. If $\delta(G) \geq 3$, then clearly $g(G) \geq 5$ and so $\delta(G) \leq \sqrt{R(4, 3; k, 1) - 2}$ by [1, p.105] or [10, Proposition 1]. Now, from the above we deduce that

$$(5.3) \quad R(4, 3; k, 1) - 1 - R(4, 3; k-1, 1) \leq \delta(G) \leq \sqrt{R(4, 3; k, 1) - 2}.$$

Set $x_n = R(4, 3; n, 1)$. Then $(x_k - x_{k-1} - 1)^2 \leq x_k - 2$ and so $x_k^2 - (2x_{k-1} + 3)x_k + (x_{k-1}^2 + 2x_{k-1} + 3) \leq 0$. This yields

$$\begin{aligned} x_k &\leq \frac{2x_{k-1} + 3 + \sqrt{(2x_{k-1} + 3)^2 - 4(x_{k-1}^2 + 2x_{k-1} + 3)}}{2} \\ &= x_{k-1} + \frac{3}{2} + \sqrt{x_{k-1} - \frac{3}{4}}. \end{aligned}$$

This completes the proof.

Theorem 5.3. *Let $0 < \varepsilon \leq 1$ and $k \in \mathbb{N}$ with $k \geq 6$. Then*

$$R(4, 3; k, 1) \leq \frac{(k-6)(k+6+2a)}{4-\varepsilon} + 16 < \frac{(k+a)^2}{4-\varepsilon}$$

and

$$R(4, 3; k, 1) - R(4, 3; k - 1, 1) < 1 + \frac{k + a}{\sqrt{4 - \varepsilon}},$$

where

$$a = \frac{5 - 1.5\varepsilon}{2 - \sqrt{4 - \varepsilon}} - 6.$$

Proof. As

$$a + 6 = \frac{5 - 1.5\varepsilon}{2 - \sqrt{4 - \varepsilon}} \geq \frac{5 - 1.5}{2 - \sqrt{3}} > 8 > 4\sqrt{4 - \varepsilon},$$

we have $\frac{(a+6)^2}{4-\varepsilon} > 16$. Now we prove the first part by induction on k . Since $R(4, 3; 6, 1) = 16$, the result is true for $k = 6$. Suppose $k \geq 7$ and

$$R(4, 3; k - 1, 1) \leq \frac{(k - 1 - 6)(k - 1 + 6 + 2a)}{4 - \varepsilon} + 16 = \frac{(k - 1 + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon} + 16.$$

Then

$$\begin{aligned} & R(4, 3; k - 1, 1) + \frac{3}{2} + \sqrt{R(4, 3; k - 1, 1) - \frac{3}{4}} \\ & < \frac{(k - 1 + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon} + 16 + \frac{3}{2} + \frac{k - 1 + a}{\sqrt{4 - \varepsilon}} \\ & = \frac{(k + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon} + 16 + \frac{3}{2} - \left(\frac{2}{4 - \varepsilon} - \frac{1}{\sqrt{4 - \varepsilon}} \right) (k + a - 1) - \frac{1}{4 - \varepsilon}. \end{aligned}$$

As $k + a - 1 \geq a + 6 = \frac{5 - 1.5\varepsilon}{2 - \sqrt{4 - \varepsilon}}$, we see that $(k + a - 1)(2 - \sqrt{4 - \varepsilon}) \geq 5 - 1.5\varepsilon = \frac{3}{2}(4 - \varepsilon) - 1$ and so

$$\left(\frac{2}{4 - \varepsilon} - \frac{1}{\sqrt{4 - \varepsilon}} \right) (k + a - 1) + \frac{1}{4 - \varepsilon} > \frac{3}{2}.$$

Hence, by the above and Theorem 5.2 we obtain

$$\begin{aligned} R(4, 3; k, 1) & \leq R(4, 3; k - 1, 1) + \frac{3}{2} + \sqrt{R(4, 3; k - 1, 1) - \frac{3}{4}} \\ & < \frac{(k + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon} + 16 = \frac{(k - 6)(k + 6 + 2a)}{4 - \varepsilon} + 16 < \frac{(k + a)^2}{4 - \varepsilon}. \end{aligned}$$

From the above and (5.3) we also deduce that

$$\begin{aligned} & R(4, 3; k, 1) - R(4, 3; k - 1, 1) - 1 \\ & \leq \sqrt{R(4, 3; k, 1) - 2} < \sqrt{\frac{(k + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon}} + 14 < \frac{k + a}{\sqrt{4 - \varepsilon}}. \end{aligned}$$

This completes the proof.

Theorem 5.4. Let $0 < \varepsilon \leq 1$ and $a = \frac{5 - 1.5\varepsilon}{2 - \sqrt{4 - \varepsilon}} - 6$. Let G be a graph of order $p \geq 16$ with $g(G) \geq 5$. Then

$$\alpha(G) \geq \lceil \sqrt{(4 - \varepsilon)(p - 16) + (a + 6)^2} - a \rceil.$$

Proof. Set $k = \lceil \sqrt{(4 - \varepsilon)(p - 16) + (a + 6)^2} - a \rceil$. Then clearly $k \geq 6$. Using Theorem 5.3 we see that

$$p \geq \frac{(k + a)^2}{4 - \varepsilon} - \frac{(6 + a)^2}{4 - \varepsilon} + 16 \geq R(4, 3; k, 1).$$

Since $g(G) \geq 5$, G must be a $(4, 3)$ graph. Hence $\alpha(G) \geq k$. This proves the theorem.

6. Some open conjectures on Ramsey numbers

In this section we risk to pose the following conjectures on Ramsey numbers.

Conjecture 6.1. For any positive integer $n \geq 2$ we have

$$\frac{n-1}{R(3, n)-1} > \frac{n}{R(3, n+1)-1} \quad \text{and so} \quad R(3, n+1) > \frac{nR(3, n)-1}{n-1}.$$

As $\frac{1}{2} > \frac{2}{5} > \frac{3}{8} > \frac{4}{13} > \frac{5}{17} > \frac{6}{22} > \frac{7}{27} > \frac{8}{35}$, from (1.1) we know that Conjecture 6.1 is true for $n \in \{2, 3, \dots, 8\}$. If the conjecture is true, we have $R(3, 10) > \frac{9R(3, 9)-1}{8} > 40$. It is now known ([8]) that $40 \leq R(3, 10) \leq 43$.

Conjecture 6.2. Let $\{L_n\}$ be the Lucas sequence defined by $L_0 = 2$, $L_1 = 1$ and $L_{n+1} = L_n + L_{n-1}$ ($n \geq 1$). For $k = 3, 4, 5, \dots$ we have $R(k, k) = 4L_{2k-5} + 2$.

Conjecture 6.2 is true for $k = 3, 4$. By Conjecture 6.2, we have $R(5, 5) = 46$, $R(6, 6) = 118$ and $R(7, 7) = 306$. Since $L_{2(n+1)} = 3L_{2n} - L_{2(n-1)}$, Conjecture 6.2 is equivalent to

$$(6.1) \quad R(k, k) = 3R(k-1, k-1) - R(k-2, k-2) - 2 \quad \text{for} \quad k \geq 3.$$

It is well known that

$$L_n = \left(\frac{1+\sqrt{5}}{2}\right)^n + \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

Thus, by Conjecture 6.2,

$$\begin{aligned} R(k, k) &= 4\left\{\left(\frac{1+\sqrt{5}}{2}\right)^{2k-5} - \left(\frac{\sqrt{5}-1}{2}\right)^{2k-5}\right\} + 2 \\ &= 128\left\{\left(\frac{3+\sqrt{5}}{2}\right)^k - \left(\frac{3-\sqrt{5}}{2}\right)^k\right\} + 2. \end{aligned}$$

Hence,

$$(6.2) \quad R(k, k) \sim 128\left(\frac{3+\sqrt{5}}{2}\right)^k \quad \text{as} \quad k \rightarrow +\infty.$$

We note that $\frac{3+\sqrt{5}}{2} \approx 2.618$. It is known that $(\sqrt{2})^k < R(k, k) \leq 4^k$. P. Erdős offered \$350 to ask the value of $\lim_{k \rightarrow \infty} R(k, k)^{\frac{1}{k}}$ (see [3, p.10]). If the limit exists, it should be $\frac{3+\sqrt{5}}{2}$ by Conjecture 6.2.

Conjecture 6.3. For $n = 2, 3, 4, \dots$ we have

$$\sum_{r=1}^{n(n-1)/2} R(n, r; 3, 1) = R\left(3, \frac{n(n+1)}{2} - 1\right).$$

Conjecture 6.3 is true for $n = 2, 3, 4$. Since

$$\sum_{r=1}^{10} R(5, r; 3, 1) = 14 + 11 + 9 + 9 + 7 + 7 + 5 + 5 + 5 + 5 = 77,$$

by Conjecture 6.3 we have $R(3, 14) = 77$. It is known ([8]) that $66 \leq R(3, 14) \leq 78$.

Conjecture 6.4. For $k = 1, 2, 3, \dots$ we have $R(4, 3; k, 1) = 3k - 2$.

From Theorem 5.1 we know that Conjecture 6.4 is true for $k \leq 6$.

References

- [1] B. Bollobás, Modern Graph Theory, Springer-Verlag, New York, 1998.
- [2] R. Bolze, H. Harborth, The Ramsey number $r(K_4 - x, K_5)$, in The Theory and Applications of Graphs, (Kalamazoo, MI, 1980), Wiley, New York, 1981, 109-116.
- [3] F. Chung, R. Graham, Erdős on Graphs: His Legacy of Unsolved Problems, A K Peters, Massachusetts, 1999.
- [4] G. Dirac, Extensions of Turán's theorem on graphs, Acta Math. Hungar. **14**(1963), 417-422.
- [5] R. Faudree, H. Harborth, I. Mengersen, Ramsey numbers for graph sets versus complete graphs, Utilitas Math. **50**(1996), 85-95.
- [6] A.I. Gol'berg, V.A. Gurvich, On the maximum number of edges for a graph with n vertices in which every subgraph with k vertices has at most l edges, Soviet Math. Doklady **35**(1987), 255-260.
- [7] H. Harborth, Ramsey numbers for sets of five vertex graphs with fixed numbers of edges, Finite and Infinite Combinatorics in Sets and Logic, N.W. Sauer et. al. (eds), Kluwer Acad. Publ., Amsterdam, 1993, pp.169-174.
- [8] S.P. Radziszowski, Small Ramsey numbers, Dynamic Surveys of Electronic J. Combinatorics (2011), DS1.13, 84pp.
- [9] Z.H.Sun, A class of problems of Turán type (*Chinese*), J. Nanjing Univ. Math. Biquarterly **8**(1991), 87-98.
- [10] Z.H.Sun, Maximum size of graphs with girth not less than a given number (*Chinese*), J. Nanjing Univ. (Special Issue) **27** (1991), 43-50,146.
- [11] P. Turán, On an extremal problem in graph theory, Mat. Fiz. Lapok **48**(1941), 436-452.