Self-Duality of Green-Schwarz Sigma-Models

Amit Dekel and Yaron Oz

Raymond and Beverly Sackler School of Physics and Astronomy Tel-Aviv University, Ramat-Aviv 69978, Israel amitde@post.tau.ac.il, yaronoz@post.tau.ac.il

ABSTRACT: We study fermionic T-duality symmetries of integrable Green-Schwarz sigma-models on Anti-de-Sitter backgrounds with Ramond-Ramond fluxes, constructed as \mathbb{Z}_4 supercosets of superconformal algebras. We find three algebraic conditions that guarantee self-duality of the backgrounds under fermionic T-duality, we classify those that satisfy them and construct the map of the monodromy matrix. We introduce new T-duality directions, where some of them contain no bosonic directions, along which the backgrounds are self-dual. We find that the only self-dual backgrounds are $\mathrm{AdS}_n \times \mathrm{S}^n$ for n = 2, 3, 5. In addition we find that the backgrounds $\mathrm{AdS}_n \times \mathrm{S}^1$ for n = 2, 3, 5, $\mathrm{AdS}_4 \times \mathrm{S}^2$ and $\mathrm{AdS}_2 \times \mathrm{S}^4$ are self-dual at the level of the classical action, but have a non-trivial transformation of the dilaton.

KEYWORDS: Duality in Gauge Field Theories, String Duality.

Contents

1.	Introduction 2		
2.	2. Properties of Superconformal Al	gebras	3
	2.1 The conformal basis and \mathbb{Z} -grad	ation	3
	2.2 \mathbb{Z}_4 automorphism		5
	2.3 More on \mathbb{Z} -gradations		5
	2.3.1 \mathbb{Z} -gradation of type I SC	A's	6
	2.3.2 \mathbb{Z} -gradation of type II SC	CA's	8
3.	3. Green-Schwarz Sigma-models on	Semi-Symmetric backgrounds	10
	3.1 The action		10
	3.2 Integrability		10
4.	4. T-duality of the GS Sigma-Mode	els	11
	4.1 Assumptions		11
	4.2 T-duality of the Green-Schwarz	sigma-model	12
	4.3 Flat-connection transformation	under T-self-duality	15
	4.4 Quantum consistency of the T-S	elf-Duality transformation	16
5.	5. Classification of the backgrounds	3	17
	5.1 Type I		17
	5.2 Type II		18
	5.3 $\operatorname{AdS}_3 \times \mathcal{M}$ semi-symmetric space	es	20
	5.4 AdS_2 semi-symmetric space		20
	5.5 When does an AdS semi-symme	tric space satisfies $\Omega(U) = -U?$	21
6.	6. Discussion		21
А.	A. Notations		24
в.	B. Superalgebras		25
	B.1 $OSP(2N 2)$		25
	B.2 $OSP(2N 4)$		26
	B.3 $F(4;2)$		26
	B.4 $F(4;0)$		27
	B.5 SU(1,1 N), $N \neq 2$		27
	B.6 SU(2,2 N), $N \neq 4$		28
	B.7 PSU $(N, N 2N), N = 1, 2$		28

B.8 $D(2,1;\zeta)$	29
B.9 $OSP(4^* 2N)$	29
B.10 OSP $(8^* 2N)$	30
C. Rank of kappa symmetry	31
C.1 Type-U2	32
C.2 Type-U4	33
C.3 Type-O1	33
C.4 Type-O2	34
C.5 Type-Tu	35
C.6 Type-To	36

1. Introduction

Self-duality of the Green-Schwarz sigma model (GSSM) on $AdS_5 \times S^5$ background is used to explain the existence of the dual-superconformal symmetry of scattering amplitudes in $\mathcal{N} = 4$ SYM, and their connection to Wilson-loops [1][2]. The superconformal symmetry together with the dual one generate a Yangian symmetry algebra, which is related to the integrability properties of the theory.

It is well known that GSSM's on semi-symmetric spaces (\mathbb{Z}_4 supercoset spaces) exhibit an infinite set of conserved charges [3] which satisfy the Yangian algebra [4][5]. It is thus natural to ask whether GSSM's on other (than AdS₅ × S⁵) semisymmetric backgrounds are self-dual under T-duality. In previous papers [6][7][8], some backgrounds were checked to be self-dual, while other were found not to be self-dual. In those papers, the background's self-duality was checked on a case by case basis. A general argument for self-duality is still lacking. In the present paper we will take a rather general approach and formulate criteria for semi-symmetric backgrounds to be self-dual. We present three sufficient algebraic conditions for self-duality, and explain the lack of self-duality of backgrounds that do not satisfy them.

We denote the superconformal algebras (SCA's) by \mathfrak{g} , with the \mathbb{Z}_2 decomposition $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ to its even and parts respectively. We further decompose the SCA's according to a \mathbb{Z} -gradation with gradings $\pm 1, 0$ only, where the charges are assigned by a generator U. The T-duality is performed along all the directions associated with the grading 1 generators, which form an abelian subalgebra. We will prove that a background is self-dual if :

1. $\Omega(U) = -U$, where Ω is the \mathbb{Z}_4 automorphism map.

- 2. Rank(κ -symmetry) $\geq \dim(\mathfrak{g}_{\bar{1}})/4$.
- 3. The SCA's Killing-form vanishes.

The first condition ensures a non-singular coupling of the fermionic coordinates. The second condition allows a particular representation of the supergroup that is used in the T-duality procedure. The third condition guarantees the quantum consistency of the transformation, that is a non-trivial dilaton is not generated.

We find that the only self-dual GSSM's are the $\operatorname{AdS}_n \times \operatorname{S}^n$ for n = 2, 3, 5. All of them were found previously to be self-dual [1][2][6]. We find there are also backgrounds that are self-dual at the classical level, but at the quantum level their dilaton shifts, these are the $\operatorname{AdS}_n \times \operatorname{S}^1$ for n = 2, 3, 5 (the case of n = 5 was discussed in [8]), $\operatorname{AdS}_2 \times \operatorname{S}^4$, and $\operatorname{AdS}_4 \times \operatorname{S}^2$. In addition to the usual self-duality along the flat AdS directions followed by some odd directions, namely the directions associated with $\operatorname{span}\{P,Q\}$, we find other abelian subalgebras along which the GSSM is self-dual (one of them was discussed in [1]). Some of these directions involve only fermionic directions. We give the general transformation of the action and the flat-connection for any such abelian subalgebra. The transformation, as in the $\operatorname{AdS}_5 \times \operatorname{S}^5$ case, is a spectral parameter dependent automorphism, which is a composition of the \mathbb{Z}_4 automorphism map and an automorphism induced by the \mathbb{Z} -gradation.

The paper is organized as follows. In section 2 we briefly review some properties of the SCA's, including a discussion of their \mathbb{Z} -gradation structure. In section 3 we briefly discuss the GSSM and their basic integrability properties. In section 4 we prove T-self-duality of the GSSM's using the three algebraic conditions stated above. In section 5 we classify the SCA's according to the conditions for T-selfduality. In section 6 we discuss the results and various open questions. In appendix A we summarize our notations. In appendix B we provide technical computations concerning the SCA's and their classification according to the first condition. In appendix C we compute the kappa-symmetry needed for the second condition.

2. Properties of Superconformal Algebras

2.1 The conformal basis and \mathbb{Z} -gradation

The generators of the SCA in d-dimensions are $\mathfrak{g}_{\mathbb{C}} = \operatorname{span}\{P, K, D, L\}$ - the so(2, d-1) conformal subalgebra generators, span $\{R\}$ - the R-symmetry subalgebra generators, and span $\{Q\}$ and span $\{S\}$ - the (odd) supercharges and superconformal charges respectively. Altogether we have $\mathfrak{g}_{SC} = \operatorname{span}\{P, K, L, D; R; Q, S\}$. The SCA's supercommutation relations are given by the commutation relations of $\mathfrak{g}_{\mathbb{C}}$ and span $\{R\}$ together with

$$[P,Q] = 0, \quad [K,S] = 0, \quad [P,S] \sim Q, \quad [K,Q] \sim S, \tag{2.1}$$

$$[R,Q] \sim Q, \quad [R,S] \sim S,$$

$$\{Q,Q\} \sim P, \quad \{S,S\} \sim K,$$

$$\{Q,S\} \sim D + L + R.$$

These commutation relations can summarized using the charge of the generators under the dilatation generator D, see figure 1. This charge assignment is an example

Figure 1: The charge of the SCA's generators under D.

of \mathbb{Z} -gradation of the SCA, which is a decomposition of the algebra such that $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_i$ and $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$. In this case i = -2, -1, 0, 1, 2. Besides this \mathbb{Z} -gradation, the SCA may have others.

The superalgebras are classified according to their type, I or II [9]. The terminology type I and type II refers to the representation of the even part of the superalgebra on the odd part. If the representation is irreducible the superalgebra is called type II and if it is a direct sum of two irreducible representations the superalgebra is called type I. In the case of type I, the odd part decomposes according to another \mathbb{Z} -gradation, which is called the *distinguished gradation* [9]. This gradation is associated with the generator B (which we call the hypercharge) which is in the algebra for $A(m, n \neq m)$ and C(n + 1) and not for A(m, m). The generators are decomposed as

$$\mathfrak{g}_I = \mathfrak{g}_1 \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{-1} = \{Q, \hat{S}\} \oplus \{P, K, D, L; R\} \oplus \{\hat{Q}, S\}.$$
(2.2)

For type II SCA's we do not have such a gradation, but when the number of spacetime supersymmetries is even, $\mathcal{N} \in 2\mathbb{N}_1$, we do have another \mathbb{Z} -gradation associated with a generator of the R-symmetry subalgebra which we call λ . This decomposition, which further decomposes the R-symmetry generators to R, λ, \hat{R} , is given by

$$\mathfrak{g}_{II} = \mathfrak{g}_2 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} = \{R\} \oplus \{Q, \hat{S}\} \oplus \{P, K, D, L; \lambda\} \oplus \{\hat{Q}, S\} \oplus \{\hat{R}\}.$$
(2.3)

In order to present the SCA's, one has to work with real-forms of the SCA, since we have to take complex combinations of the odd generators and the R-symmetry generators. We summarize some relevant properties of the SCA's in table 1. Further decomposition of the commutation relations (2.1) should be obvious from the gradations introduced above.

Another characteristic of the SCA's is whether the Killing-form is degenerate or not, see table 1. The Killing-form is defined as the supertrace of every two generators

			1	1		
d	SCA	R-symmetry	$\dim(\mathfrak{g}_{\bar{1}})$	\mathcal{N}	type	Killing-form
1	osp(N 2)	$\operatorname{so}(N)$	2N	N	I for $N = 2$, else II	ND, except for $N = 4$
1	$\mathrm{su}(1,1 N\neq 2)$	$\mathrm{u}(N)$	4N	2N	Ι	ND
1	psu(1,1 2)	su(2)	8	4	Ι	Zero
1	$osp(4^* 2N)$	$\operatorname{su}(2) \times \operatorname{usp}(2N)$	8N	4N	II	ND
1	G(3)	g_2	14	7	II	ND
1	F(4;0)	so(7)	16	8	II	ND
1	$D(2,1;\alpha)$	so(4)	8	4	II	Zero
3	osp(N 4)	so(N)	4N	N	I for $N = 2$, else II	ND, except for $N = 6$
4	$\operatorname{su}(2,2 N\neq 4)$	$\mathrm{u}(N)$	8N	N	Ι	ND
4	psu(2,2 4)	su(4)	32	4	Ι	Zero
5	F(4;2)	su(2)	16	2	II	ND
6	$osp(8^* N)$	usp(N)	8N	N	II	ND, except for $N = 6$

Table 1: Some properties of SCA's

The table gives the SCA's as classified in [10]. The spinor representations for d = 3, 4, 5, 6 are $su(2), su(2) \times su(2), sp(4), su(4)$ respectively. ND- stands for non-degenerate. N is the number of space-time supersymmetries.

in the adjoint representation [11][9], $K_{ab} = \text{Str}(L_a^{\text{adj}}L_b^{\text{adj}})$. The SCA's with degenerate Killing-form are known to have special properties in the context of the Green-Schwarz sigma-models, e.g [12][13][14], and as we shall see they are also special with respect to the self-duality properties of the sigma-models.

2.2 \mathbb{Z}_4 automorphism

Every SCA has at least one \mathbb{Z}_4 automorphism [15]. A SCA is decomposed under this automorphism into four sets

$$\mathfrak{g} = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \mathcal{H}_3, \tag{2.4}$$

such that $[\mathcal{H}_i, \mathcal{H}_j] \subset \mathcal{H}_{i+j \mod 4}$, $B(\mathcal{H}_i, \mathcal{H}_j) \neq 0$ only if $i + j = 0 \mod 4$, and $\Omega(\mathcal{H}_k) = i^k \mathcal{H}_k$, where B represents the Cartan-Killing bilinear-form and $\Omega(\cdot)$ is the automorphism map.

Using the \mathbb{Z}_4 automorphism property we can define a semi-symmetric space by taking the quotient with respect to the invariant locus \mathcal{H}_0 (so the bosonic part is a symmetric-space). A SCA may have several different \mathbb{Z}_4 automorphisms and so one can identify a semi-symmetric space with respect to each automorphism. Some of the semi-symmetric spaces will have a bosonic AdS sub-space in which we are mainly interested in the present paper, although we will also consider non-AdS spaces.

2.3 More on \mathbb{Z} -gradations

As we have shown above, any SCA has a \mathbb{Z} -gradation, $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_i$ such that $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$. When *i* takes a finite number of values, say $i_{\min} \leq i \leq i_{\max}$, the set $\bigoplus_{i=i_{\max}/2}^{i_{\max}} \mathfrak{g}_i$ defines an abelian subalgebra if $i_{\max} > 0$, and similarly for the set $\bigoplus_{i=i_{\min}}^{i_{\min}/2} \mathfrak{g}_i$ if $i_{\min} < 0$.

In the present paper we will be interested in Z-gradations with $|i| \leq 1$, that is $\mathfrak{g} = A_1 \oplus B_0 \oplus A_{-1}$, with $A_{\pm 1}$ abelian subalgebras¹ and B_0 a subalgebra, so these are the Z-gradations we will consider from now on. Any such decomposition can be induced by introducing a U(1) generator U, with respect to \mathfrak{g}_0 (which may or may not be part of the SCA), satisfying $\mathrm{ad}_U(L_a) = [U, L_a] = aL_a, \forall L_a \in \mathfrak{g}$, where $a = \pm 1, 0$ is the charge of the generator. The decomposition under ad_U induces a one-parameter dependent automorphism

$$\sigma_{\lambda}(L_a) = \lambda^U L_a \lambda^{-U} = \lambda^a L_a, \quad \lambda \in \mathbb{C}.$$
 (2.5)

The various \mathbb{Z}_4 automorphisms [15] of the SCA's may have different relations with the U(1) generator inducing the \mathbb{Z} -gradation. We are interested in those satisfying

$$\Omega(U) = -U. \tag{2.6}$$

 Ω is defined to act on a commutator as $\Omega([L_a, L_b]) = [\Omega(L_a), \Omega(L_b)]$, thus

$$\sigma_{\lambda}(\Omega(L_a)) = \lambda^{-a}(\Omega(L_a)), \qquad (2.7)$$

and the non-trivial bilinear-form is of the form $B(L_a\Omega(L_b))$ with a = b. For backgrounds satisfying (2.6) and some other conditions (to be discussed later) we will be able to prove T-self-duality.

Next we consider four classes of \mathbb{Z} -gradations that will be of interest in the study of T-self-duality of GS-sigma-models. We study the type I and type II SCA's separately.

Comment: when talking about the grading, one usually use integer labeling, while the charges with respect to the U(1) generators we will use $(D, B, \check{R}, \check{\lambda})$ are integer for the even generators and half integer for the odd generator (e.g there should really be a factor of 2 in figure 1, and later in figure 2). In the next sections when we will write the charges with respect to the generators we will use integer numbers although they should be understood to be divided by 2, so at the end of the day when we will discuss the gradations with charges $\pm 1, 0$ only, these will really be the charges under the combination of the U(1) generators.

2.3.1 Z-gradation of type I SCA's

The type I SCA's include the su(1, $1|N \neq 2$), su(2, $2|N \neq 4$), psu(1, 1|2), psu(2, 2|4), osp(2|2) \simeq su(1, 1|1), osp(2|4). Generally, the bosonic part of these SCA's is $\mathfrak{g}_{\bar{0}} =$ so(2, d) \oplus su(N) \oplus u(1). We will refer to the last u(1) as the hypercharge, which in the case of the 'psu' SCA's decouples from the SCA. The two 'osp' SCA's are missing the su(N) subalgebra.

¹The \mathbb{Z} -gradation considered should not necessarily be consistent, namely $\mathfrak{g}_{\pm 1}$ and \mathfrak{g}_0 may contain even and odd generators respectively [11][9].

We will consider three u(1)'s, generating consistent \mathbb{Z} -gradations of the SCA's, and their combinations which generates \mathbb{Z} -gradations with $|i| \leq 1$. We already considered D, which induces the following consistent \mathbb{Z} -gradation

$$\mathfrak{g}_{2} = \operatorname{span}\{P\}, \quad \mathfrak{g}_{1} = \operatorname{span}\{Q, \hat{Q}\}, \qquad (2.8)$$
$$\mathfrak{g}_{0} = \operatorname{span}\{D, L\} \oplus \mathrm{u}(N),$$
$$\mathfrak{g}_{-1} = \operatorname{span}\{S, \hat{S}\}, \quad \mathfrak{g}_{-2} = \operatorname{span}\{K\},$$

and B which induces the distinguished \mathbb{Z} -gradation

$$\mathfrak{g}_1 = \operatorname{span}\{Q, \hat{S}\}, \quad \mathfrak{g}_0 = \operatorname{su}(M, M) \oplus \operatorname{u}(N), \quad \mathfrak{g}_{-1} = \operatorname{span}\{\hat{Q}, S\}.$$
 (2.9)

Lastly, we decompose the $\operatorname{su}(N)$ generators, R_l^k with k, l = 1, ..., N and $\sum_k R_k^k = 0$, under $\operatorname{su}(N) \to \operatorname{su}(P) \times \operatorname{u}(N-P)$. This divides the R-symmetry indices to k, l, ... = 1, ..., P and k', l', ... = P + 1, ..., N, so the $\operatorname{u}(P)$ generators are R_l^k and the $\operatorname{u}(N-P)$ generators are $R_{l'}^{k'}$, with the relation $\sum_k R_k^k + \sum_{k'} R_{k'}^{k'} = 0$. We define the generator, $\check{R} = \sum_k R_k^k - \sum_{k'} R_{k'}^{k'}$, inducing the \mathbb{Z} -gradation

$$\mathfrak{g}_{2} = \operatorname{span}\{R_{l}^{k'}\}, \quad \mathfrak{g}_{1} = \operatorname{span}\{Q^{k}, \hat{Q}_{l'}, S_{l'}, \hat{S}^{k}\},$$
(2.10)
$$\mathfrak{g}_{0} = \operatorname{s}(\operatorname{u}(P) \times \operatorname{u}(N-P)) \oplus \operatorname{su}(M, M),$$

$$\mathfrak{g}_{-1} = \operatorname{span}\{Q^{k'}, \hat{Q}_{l}, S_{l}, \hat{S}^{k'}\}, \quad \mathfrak{g}_{-2} = \operatorname{span}\{R_{l'}^{k}\}.$$

Next we consider the combinations of D, B and \hat{R} that give the \mathbb{Z} -gradations with $|i| \leq 1$ (up no normalization). First we have the well known U = D + B [2][1] which gives the non-consistent \mathbb{Z} -grading decomposition

$$\mathfrak{g} = (P, Q)_1 \oplus (L, D, \hat{Q}, \hat{S}, R)_0 \oplus (K, S)_{-1}.$$
(2.11)

This decomposition holds for all the type I SCA's. In this case the invariant subalgebras are (s)u(M|N) for (p)su(2M|N), and u(1|2) for osp(2|4). Similarly for U = D - B we get the same gradation with the hatted and unhatted generators interchanged.

Next, we consider $U = D + \check{R}$ which generates the decomposition

$$\mathfrak{g} = (P, Q^k, \hat{Q}_{k'}, R_{k'}{}^l)_1 \oplus (L, D, Q^{k'}, \hat{Q}_k, S_{k'}, \hat{S}^k, R_k{}^l, R_{k'}{}^l')_0 \oplus (K, S_k, \hat{S}^{k'}, R_k{}^{l'})_{-1}$$
(2.12)

which was mentioned in [1] for psu(2, 2|4). This decomposition cannot be applied for the 'osp' SCA's. For this decomposition the invariant subalgebras are $(p)s(u(M|P) \times u(M|N-P))$ for (p)su(2M|N).

Next, we consider U = 2B which generates the consistent distinguished-gradation

$$\mathfrak{g} = (Q, \hat{S})_1 \oplus (P, K, D, L; R)_0 \oplus (\hat{Q}, S)_{-1}.$$
(2.13)

This decomposition was not considered before in the context of T-duality, and implies the model may be self-dual under T-duality along fermionic directions only.

Lastly, we consider $U = B + \check{R}$ which generates the inconsistent-gradation

$$\mathfrak{g} = (Q^k, \hat{S}^k, R_{k'}{}^l)_1 \oplus (P, K, L, D, Q^{k'}, \hat{S}^{k'}, S_{k'}, \hat{Q}_{k'}, R_k{}^l, R_{k'}{}^l)_0 \oplus (S_k, \hat{Q}_k, R_k{}^{l'})_{-1}.$$
(2.14)

This decomposition also was not considered before in the context of T-duality. It is similar to the decomposition (2.12), interchanging the roles of the AdS and the sphere.

We summarize the above decompositions in figure 2 (a-c). The results for $AdS_5 \times S^5$ background are also given in table 3. Note that in cases where the SCA is 'psu', the hypercharge is not a part of the SCA and so the Z-gradation automorphisms are outer. The rest of the automorphisms are inner. Decompositions with respect to combination, different from the ones presented by changing the relative signs of the generators are obvious and addressed in the figure. One can find more decompositions of the SCA's, which we find less interesting with respect to the AdS backgrounds².

2.3.2 Z-gradation of type II SCA's

The type II SCA's include the $osp(N \neq 2|2)$, $osp(N \neq 2|4)$, $osp(4^*|2)$, $osp(8^*|N)$, $D(2, 1; \alpha)$, F(4) and G(3), but we'll consider only $osp(2N \neq 2|2)$, $osp(2N \neq 2|4)$, $osp(4^*|2)$, $osp(8^*|2N)$, $D(2, 1; \alpha)$ and F(4) which can be decomposed according to (2.3) (the ones with even number of space-time supersymmetries).

As mentioned above the type II SCA's decompose under the charge assignment of D,

$$\mathfrak{g}_{2} = \operatorname{span}\{P\}, \quad \mathfrak{g}_{1} = \operatorname{span}\{Q, \hat{Q}\}, \quad (2.15)$$

$$\mathfrak{g}_{0} = \operatorname{span}\{D, L\} \oplus \operatorname{span}\{R, \lambda, \hat{R}\}, \quad (2.15)$$

$$\mathfrak{g}_{-1} = \operatorname{span}\{S, \hat{S}\}, \quad \mathfrak{g}_{-2} = \operatorname{span}\{K\}.$$

and according to the gradation (2.3), where the R-symmetry decomposes to $R_2 \oplus \lambda_0 \oplus \hat{R}_{-2}$ (the subscript indicates the gradation). The R-symmetry decomposition for all type II SCA's with such decomposition is given in table 2. The Z-gradation of (2.3) is induced by the generator $\check{\lambda}$ given in the table.

Combining the two, $U = D + \check{\lambda}$, we find that all type II SCA's have the inconsistent Z-gradation

$$\mathfrak{g} = (P, Q, R)_1 \oplus (L, D, \hat{Q}, \hat{S}, \lambda)_0 \oplus (K, S, \hat{R})_{-1}.$$
(2.16)

²For example, in the notation of $AdS_5 \times S^5$ we have

 $[\]begin{array}{c} (P^{\alpha\dot{\alpha}}, K^{\alpha\dot{\alpha}}, L^{\alpha\alpha}, L^{\dot{\alpha}\dot{\alpha}}, Q^{i\alpha}, \hat{Q}^{\dot{\alpha}}_{i'}, S^{i\dot{\alpha}}_{i'}, \hat{S}^{i\dot{\alpha}}, R_{i'}{}^{i}) \oplus (P^{\bar{\alpha}\dot{\alpha}}, P^{\alpha\bar{\alpha}}, K^{\alpha\bar{\alpha}}, K^{\alpha\bar{\alpha}}, L^{\alpha\bar{\alpha}}, L^{\dot{\alpha}\dot{\alpha}}, Q^{i\bar{\alpha}}, Q^{i\bar{\alpha}}, \hat{Q}^{\dot{\alpha}}_{i}, \hat{Q}^{\dot{\alpha}}_{i'}, \hat{Q}^{\dot{\alpha}}_{i'}, \hat{S}^{\dot{\alpha}}_{i'}, \hat{S}^{i\prime\dot{\alpha}}, \hat{S}^{i\dot{\alpha}}, R_{i}{}^{i}, R_{i'}{}^{i'}) \oplus (K^{\bar{\alpha}\bar{\alpha}}, P^{\bar{\alpha}\bar{\alpha}}, L^{\bar{\alpha}\bar{\alpha}}, L^{\bar{\alpha}\bar{\alpha}}, S^{\bar{\alpha}}_{i}, \hat{S}^{i'\bar{\alpha}}, Q^{i'\bar{\alpha}}, \hat{Q}^{\bar{\alpha}}_{i}, R_{i}{}^{i'}) \end{array}$

where $(\alpha, \dot{\alpha}) = (1, 1), (1, 2), (2, 1), \text{ or } (2, 2), (\bar{\alpha}, \bar{\dot{\alpha}})$ take different value then $(\alpha, \dot{\alpha})$, and $i = 1, ..., n \leq 4$, i' = n + 1, ..., 4. The abelian subalgebra involves unphysical directions, L. The invariant subalgebra is $ps(u(2|n) \times u(2|4 - n))$.



Figure 2: Z-gradation of type-I SCA's with 4N-odd generators and R-symmetry $SU(2M) \times U(1)$. The abelian subalgebras are circled. (a) Decomposition under B and D. In this case the relevant U(1)'s are $(\pm)(D \pm B)$ (circled with solid blue contours) where the abelian subalgebra contains d-bosonic and N-fermionic generators, and $\pm 2B$ (circled with dashed red contours) where the abelian subalgebra contains 2N-fermionic generators. (b) Decomposition under B and \check{R} . In this case the relevant U(1)'s are $(\pm)(\check{R} \pm B)$ (circled with solid blue contours) where the abelian subalgebra contains M^2 -bosonic and 2M-fermionic generators, and $\pm 2B$ (circled with dashed red contours) where the abelian subalgebra contains M^2 -bosonic and 2M-fermionic generators, and $\pm 2B$ (circled with dashed red contours) where the abelian subalgebra contains M^2 -bosonic and 2M-fermionic generators. (c) Decomposition under D and \check{R} . In this case the relevant U(1)'s are $(\pm)(\check{R} \pm D)$ where the abelian subalgebra contains $d + M^2$ -bosonic and N-fermionic generators. (d) Z-gradation of type-II SCA's with 4N-odd generators and R-symmetry $R_1 \oplus \lambda_0 \oplus R_{-1}$. The abelian subalgebras are circled. We have the decomposition under $\check{\lambda}$ and D. In this case the relevant U(1)'s are $(\pm)(D \pm \check{\lambda})$ where the abelian subalgebra contains $d + \dim(R_1)$ -bosonic and N-fermionic generators.

The invariant subalgebras of the decomposition are u(N|1), u(2|1), $u(1) \oplus osp(2|4)$, u(2|N), u(N|2) and u(4|N) for osp(2N|2), $D(2,1;\alpha)$, F(4), $osp(4^*|2N)$, osp(2N|4)and $osp(8^*|2N)$ respectively

Note that all type II SCA's have no analog of the hypercharge, which assigns non-trivial charge to all odd generators and only to them.

SCA	R-symmetry	$[\lambda_0]$	$[R_1]$	λ
osp(2N 2)	so(2N)	$\lambda_l^k \in \mathbf{u}(N)$	$R_{kl} = -R_{lk}$	$\sum_k \lambda_k^k$
osp(2N 4)	so(2N)	$\lambda_l^k \in \mathrm{u}(N)$	$R_{kl} = -R_{lk}$	$\sum_k \lambda_k^k$
$osp(4^* 2N)$	$\operatorname{su}(2) \times \operatorname{usp}(2N)$	$(\lambda_l^k \in \mathrm{u}(N)) \times \mathrm{su}(2)$	$R_{kl} = R_{lk}$	$\sum_k \lambda_k^k$
		$(J_3 \in \mathbf{u}(1)) \times \mathbf{usp}(2N)$	J_+	J_3
$osp(8^* 2N)$	usp(2N)	$\lambda_l^k \in \mathrm{u}(N)$	$R_{kl} = R_{lk}$	$\sum_k \lambda_k^k$
$D(2,1;\alpha)$	$su(2) \times su(2)$	$J_3 \in \mathrm{u}(1)$	J_+	J_3
F(4;0)	so(7)	$N_{ab} \times \lambda \in \mathrm{so}(5) \times \mathrm{u}(1)$	R_a	λ
F(4; 2)	su(2)	$J_3 \in \mathrm{u}(1)$	J_+	J_3

 Table 2:
 Type II SCA's R-symmetry decomposition.

 $[\lambda_0]$ and $[R_1]$ indicates the set of generators of the R-symmetry decomposed under λ with charges 0 and 1 respectively. The indices take the values: k, l = 1, ..., N and a = 1, ..., 5. We gave only $[R_1]$ where $[\hat{R}_{-1}]$ should be understood.

We summarize the decomposition in figure 2 (d).

3. Green-Schwarz Sigma-models on Semi-Symmetric backgrounds

3.1 The action

It is well known how to construct the Green-Schwarz sigma-model (GSSM) action on semi-symmetric spaces backgrounds G/H with RR-flux, as was first done for the $AdS_5 \times S^5$ background in [16] (see e.g [17] for other backgrounds). We introduce the left-invariant-one-form $\mathbf{j} = g^{-1}dg$ where $g \in G$, which take values in the SCA, and so decomposes under the \mathbb{Z}_4 automorphism to $\mathbf{j} = \mathbf{j}_0 + \mathbf{j}_1 + \mathbf{j}_2 + \mathbf{j}_3$. We shall work in the 2d-conformal basis where $\mathbf{j} = g^{-1}\partial g$ and $\mathbf{\bar{j}} = g^{-1}\partial g$, so the GS sigma-model action is given by

$$S_{\rm GS} = \int d^2 \sigma \text{Str} \left(j_2 \bar{j}_2 + \frac{1}{2} \left(j_1 \bar{j}_3 - j_3 \bar{j}_1 \right) \right).$$
(3.1)

The action is invariant under left G-global multiplications of g and right H-local multiplications. It is also invariant under the local fermionic κ -symmetry transformation, where the rank of the transformation depends on the coset background [16][14].

3.2 Integrability

The sigma-model is known to be integrable [3] by introducing a flat-connection depending on a spectral parameter. In the present paper we shall work with the flat connection

$$A(z) = \mathbf{j}_{(0)} + z\mathbf{j}_{(1)} + \frac{1}{2}(z^2 + z^{-2})\mathbf{j}_{(2)} + z^{-1}\mathbf{j}_{(3)} - \frac{1}{2}(z^2 - z^{-2}) * \mathbf{j}_{(2)}$$
(3.2)

where $z \in \mathbb{C}$ is the spectral parameter. Noting that

$$j_{(0)} = \frac{1}{2}(1+\Omega)j_B, \quad j_{(2)} = \frac{1}{2}(1-\Omega)j_B,$$

$$j_{(1)} = \frac{1}{2}(1-i\Omega)j_F, \quad j_{(3)} = \frac{1}{2}(1+i\Omega)j_F,$$
(3.3)

where j_B and j_F are the even and odd parts of the current j, we can rewrite the flat-connection

$$A(z) = \frac{1}{4}(z+z^{-1})^2 \mathbf{j}_B - \frac{1}{4}(z-z^{-1})^2 \Omega(\mathbf{j}_B) - \frac{1}{4}(z^2-z^{-2}) * (\mathbf{j}_B - \Omega(\mathbf{j}_B))$$
(3.4)
+ $\frac{1}{2}(z+z^{-1})\mathbf{j}_F - \frac{i}{2}(z-z^{-1})\Omega(\mathbf{j}_F).$

Once we have the flat-connection we can construct an infinite tower of conserved charges using the monodromy matrix, which is given by

$$M(z) = \overrightarrow{\mathcal{P}} \exp \int_{\gamma} a(z) \tag{3.5}$$

where \mathcal{P} stands for path ordering and a(z) is the gauge invariant flat-connection, related to A(z) by $d + a(z) = g(d + A(z))g^{-1}$. Expanding M(z) around $z = \pm 1$, one can extract the conserved n-local charges.

4. T-duality of the GS Sigma-Models

4.1 Assumptions

In this section we summarize our assumptions and motivate them. First of all, we T-dualize along all directions associated with the generators of A_1 (see section 2.3). This means we T-dualize along the coordinates that couple to the generators in A_1 in the expression for $j = g^{-1}dg$ when expanded to first order in all coordinates. Thus, these coordinates should appear in the action only through their derivatives, in order to have translation isometry. For this reason we parameterize the group element representative as

$$g(x,y) = e^{a(x)}e^{b(y)},$$
(4.1)

with $a(x) = x^I L_I \in A_1$ and assume $b(y) = y^{\alpha} L_{\alpha} \notin A_1$.

As will be shown in the next subsection the WZW piece of the Lagrangian is given by $L_{WZW} = -\frac{i}{2} \text{Str}(j_F \Omega(\bar{j}_F))$. To zeroth order in the fermions, keeping only their derivatives, this is the term that should produce the quadratic term for the fermionic coordinates. In order to perform the T-duality transformation this term must be non-singular, namely the matrix that couples the fermions which we want to T-dualize along, should be invertible. For type I SCA's, when we want to Tdualize, for example, along span{P, Q} directions, it means that in order to have a non-singular coupling for the fermionic coordinates, $Str(Q\Omega(Q))$ must be non-trivial, namely $\Omega(Q) \sim S$, which is the case if (2.6) is satisfied.

This condition also implies the existence of a non-singular fermionic quadratic term for type II SCA's. But for type II, in principle, one can also have a non-singular quadratic term also if (2.6) is not satisfied, by conjugating Q with $g_{\hat{R}} = \exp(z \cdot \hat{R})$, so $g_{\hat{R}}^{-1}Qg_{\hat{R}} \sim Q + z\hat{Q}$. This means that if $\Omega(\hat{Q}) \sim S$, $\operatorname{Str}(Q\Omega(\hat{Q}))$ is not trivial. As found in [6], the coupling can still be singular if, for example, the dimension of the representation of \hat{R} is odd. We note that this automorphism relation implies $\Omega(R) \sim R$ and $\Omega(\hat{R}) \sim \hat{R}$, while the non-trivial bilinear form for these generators is of the form $B(R, \hat{R})$, so if we want also to T-dualize along the R direction (as we do according to the first assumption) we will have to parameterize $g \sim e^{yR}e^{z\hat{R}}$. This will give the kinetic term $L_{\text{kin}} \sim \partial y(z^2 \bar{\partial} y + \bar{\partial} z) + \text{c.c}$ which after T-duality along ygives $\tilde{L}_{\text{kin}} \sim \frac{\partial \tilde{y} \bar{\partial} \tilde{y}}{z^2} + dy \wedge dz$. Thus, we do not get back the same background, since we did not have this B-field in the original action. For these reasons we impose that (2.6) is satisfied.

We note that if the parametrization includes terms in A_{-1} , it is unlikely to get self-duality, where again T-duality will produce a B-field instead of part of the metric. For example, if we take $g \sim e^{xP}e^{yK}$, then $j = \partial x + \partial y + ...$ (where the ellipsis stands for higher powers of y which will not affect the point we are making), and the kinetic term is $L_{kin} \sim \partial x \bar{\partial} y + \bar{\partial} x \partial y - \partial x \bar{\partial} x - \partial y \bar{\partial} y$. Under T-duality along x, $\tilde{L}_{kin} \sim dx \wedge dy - \partial x \bar{\partial} x - \partial y \bar{\partial} y$. So the background is not invariant under this transformation. This motivates us to consider only parameterizations of the form

$$g(x,y) = e^{a(x)}e^{b(y)},$$
(4.2)

with $a(x) = x^{I}L_{I} \in A_{1}$ and $b(y) = y^{\alpha}L_{\alpha} \in B_{0}$. The indices I, J, ... will denote generators in A_{1} . The indices $\alpha, \beta, ...$ will denote generators in B_{0} which are eigenstates of $\Omega(\cdot)$. The current decomposes to

$$j = g^{-1}dg = e^{-b(y)}dx^{I}L_{I}e^{b(y)} + e^{-b(y)}de^{b(y)} \equiv J(x,y) + j(y)$$
(4.3)

where $J \in A_1$ and $j \in B_0$. In order to parameterize g as in (4.2), we will have to use the local H-gauge-symmetry and κ -symmetry. As we will see later the problem will reduce to computing the rank of the κ -symmetry. For the \mathbb{Z} -gradations introduced in section 2.3, we have to gauge away at least quarter of the odd degrees of freedom and in one case at least half. The decomposition of the current (4.3) implies that the action will not have mixed terms of J and j.

To summarize, we assume the backgrounds to satisfy (2.6) and that we can parameterize the action as in (4.2).

4.2 T-duality of the Green-Schwarz sigma-model

We rewrite the action (3.1) in terms of j_B and j_F defined in (3.3). First we consider

the WZW term (assuming A_1 containing odd generators)

$$L_{\text{WZW}} = \frac{1}{2} \text{Str}(\mathbf{j}_1 \bar{\mathbf{j}}_3 - \mathbf{j}_3 \bar{\mathbf{j}}_1) = \frac{1}{8} \text{Str}((1+i\Omega)\mathbf{j}_F(1-i\Omega)\bar{\mathbf{j}}_F - (1-i\Omega)\mathbf{j}_F(1+i\Omega)\bar{\mathbf{j}}_F) \quad (4.4)$$
$$= -\frac{i}{2} \text{Str}(J_F \Omega(\bar{J}_F) + j_F \Omega(\bar{J}_F)).$$

It is natural to define a bilinear-form³ $\eta_{AB} = \text{Str}(L_A\Omega(L_B)), L_A, L_B \in \mathfrak{g}$ and its inverse satisfying $\eta^{AB}\eta_{BC} = \delta_C^A$. We will raise and lower indices using the bilinearform. Note that it satisfies $\eta_{AB} = \eta_{BA}$ for both bosonic and fermionic parts⁴. We further introduce the notation $\mathbf{j} = \mathbf{j}^A L_A = \eta^{AB} \text{Str}(\mathbf{j}\Omega(L_B))L_A$. Returning to the WZW term we find

$$L_{\rm WZW} = \frac{i}{2} J_F^I \bar{J}_F^J \eta_{IJ}^F + \frac{i}{2} j_F^{\alpha} \bar{j}_F^{\beta} \eta_{\alpha\beta}^{(1,3)}, \qquad (4.5)$$

where $\eta_{\alpha\beta}^{(1,3)}$ means that $\alpha \in \mathcal{H}_1$ and $\beta \in \mathcal{H}_3$ or $\alpha \in \mathcal{H}_3$ and $\beta \in \mathcal{H}_1$.

Next, we consider the kinetic part of the GS action

$$L_{\rm kin} = \operatorname{Str}(j_2 \bar{j}_2) = \frac{1}{4} \operatorname{Str}((1 - \Omega) j_B (1 - \Omega) \bar{j}_B)$$

$$= \frac{1}{2} \operatorname{Str}(J_B \bar{J}_B - J_B \Omega(\bar{J}_B) + j_B \bar{j}_B - j_B \Omega(\bar{j}_B)).$$

$$(4.6)$$

Since we work with $\Omega(A_1) \in A_{-1}$, the first term vanishes and we get

$$L_{\rm kin} = \frac{1}{2} \operatorname{Str} \left(-J_B^I \bar{J}_B^J L_I \Omega(L_J) + j_B^{\alpha} \bar{j}_B^{\beta} L_{\alpha} (1-\Omega)(L_{\beta}) \right)$$

$$= -\frac{1}{2} J_B^I \bar{J}_B^J \eta_{IJ}^B - j_B^{\alpha} \bar{j}_B^{\beta} \eta_{\alpha\beta}^{(2)},$$

$$(4.7)$$

where the superscript in $\eta^{(2)}$ reminds us that we take only generators in \mathcal{H}_2 .

All in all we find the GS sigma-model Lagrangian is given by

$$L_{\rm GS} = -\frac{1}{2} J_B^I \bar{J}_B^J \eta_{IJ}^B - j_B^{\alpha} \bar{j}_B^{\beta} \eta_{\alpha\beta}^{(2)} + \frac{i}{2} J_F^I \bar{J}_F^J \eta_{IJ}^F + \frac{i}{2} j_F^{\alpha} \bar{j}_F^{\beta} \eta_{\alpha\beta}^{(1,3)}.$$
(4.8)

We want to T-dualize the sigma-model along the directions x^{I} which appear only through their derivatives in J, so we introduce the gauge fields $A = \partial x^{I} L_{I}$ and $\bar{A} = \bar{\partial} x^{I} L_{I}$, and add the Lagrange multiplier term

$$L_{\rm LM} = -\frac{1}{2} \partial \tilde{x}^I \operatorname{Str}(e^b \bar{A}' e^{-b} \Omega(L_I)) + \frac{1}{2} \bar{\partial} \tilde{x}^I \operatorname{Str}(e^b A' e^{-b} \Omega(L_I))$$
(4.9)

³This Bilinear form is consistent $(B(X,Y) = 0 \forall X \in \mathfrak{g}_{\bar{0}} \text{ and } Y \in \mathfrak{g}_{\bar{1}})$, but neither supersymmetric $(B(X,Y) = (-)^{|X|+|Y|}B(Y,X))$ nor invariant (B([X,Y],Z) = B(X,[Y,Z])), see [11][9].

⁴This is a bit unusual property due to our special definition of the bilinear-form, which further implies $\psi^A \psi^B \eta_{AB} = 0$ if ψ is odd. Usually the fermionic part of the bilinear-form is anti-symmetric.

where $A' = e^{-b}Ae^b = A'_B + A'_F \in A_1$ (note that the index *I* runs over both bosonic and fermionic generators of A_1). We define

$$W_J = \operatorname{Str}(e^{-b}\partial \tilde{x}^I \Omega(L_I) e^b L_J) = \operatorname{Str}(e^{-\Omega(b)}\partial \tilde{x}^I \Omega^2(L_I) e^{\Omega(b)} \Omega(L_J)), \qquad (4.10)$$

and rewrite

$$L_{\rm LM} = \frac{1}{2} A^{\prime I} \bar{W}_I - \frac{1}{2} \bar{A}^{\prime I} W_I.$$
(4.11)

The EOM for J give

$$\bar{A}_B^{\prime J} \eta_{IJ}^B = \bar{W}_I, \quad A_B^{\prime I} \eta_{IJ}^B = -W_J, \qquad (4.12)$$
$$\bar{A}_F^{\prime J} \eta_{IJ}^F = i \bar{W}_I, \quad A_F^{\prime I} \eta_{IJ}^F = i W_J,$$

or by using the metric definitions

$$\bar{A}_{B}^{\prime I} = \bar{W}^{I}, \quad A_{B}^{\prime I} = -W^{I},$$

$$\bar{A}_{F}^{\prime I} = i\bar{W}^{I}, \quad A_{F}^{\prime I} = iW^{I}.$$
(4.13)

Plugging these into the action we get

$$\tilde{L}_{GS} = -\frac{1}{2} W_B^I \bar{W}_B^J \eta_{IJ}^B - j_B^{\alpha} \bar{j}_B^{\beta} \eta_{\alpha\beta}^{(2)} + \frac{i}{2} W_F^I \bar{W}_F^J \eta_{IJ}^F + \frac{i}{2} j_F^{\alpha} \bar{j}_F^{\beta} \eta_{\alpha\beta}^{(1,3)}.$$
(4.14)

We can also construct \tilde{L}_{GS} , if instead of the original parametrization we take $\tilde{g} = e^{\Omega(a)}e^b$ and replace x with \tilde{x} , so $\tilde{j} = j$ is unaffected as desired, and

$$\tilde{J} \equiv \tilde{J}^K \Omega(L_K) = \operatorname{Str}(e^{-b} \partial \tilde{x}^J \Omega(L_J) e^b \Omega^2(L^K)) \Omega(L_K)$$
(4.15)

 \mathbf{SO}

$$\tilde{J}_K = \operatorname{Str}(e^{-b}\partial \tilde{x}^J \Omega(L_J) e^b \Omega^2(L_K)) = (-)^K \operatorname{Str}(e^{-b}\partial \tilde{x}^J \Omega(L_J) e^b L_K) = (-)^K W_K.$$
(4.16)

Note that the minus factor $(-)^{K}$ doesn't affect the action since the \tilde{J} 's appear quadratically. We also used $\operatorname{Str}(\Omega(L_{I})\Omega^{2}(L_{J})) = \operatorname{Str}(L_{I}\Omega(L_{J}))$ in the $W\bar{W}$ part of the WZW term. These two parametrization are related by the automorphism Ω if we redefine the coordinates in e^{b} . If $b = y^{\alpha}L_{\alpha}$, then the dual coordinates should be $\tilde{y}^{\alpha} = i^{\alpha}y^{\alpha}$ (where the α in i^{α} indicates the \mathbb{Z}_{4} grading of L_{α}), so

$$\Omega(g(x,y)) = e^{\Omega(a(x))} e^{\Omega(b(y))} = e^{\Omega(a(\tilde{x}))} e^{b(\tilde{y})} = \tilde{g}(\tilde{x},\tilde{y}).$$
(4.17)

We also note that $\Omega(\tilde{b}) = \Omega(\tilde{y}^{\alpha}L_{\alpha}) = y^{\alpha}L_{\alpha} = b$, so (from now on a tilde over a current means we take the original current and plug the dual coordinates)

$$\tilde{J}_K \equiv J_K(\tilde{x}, \tilde{y}) = \operatorname{Str}(e^{-\tilde{b}}\partial \tilde{x}^I L_I e^{\tilde{b}} \Omega(L_K))$$

$$= \operatorname{Str}(e^{-b}\partial \tilde{x}^I \Omega(L_I) e^b \Omega^2(L_K)) = (-)^K W_K.$$
(4.18)

Similarly

$$\tilde{j}_{L_{\alpha}} \equiv \tilde{j}^{\alpha} L_{\alpha} \equiv j^{\alpha}(\tilde{y}) L_{\alpha} = \operatorname{Str}(e^{-\tilde{b}} \partial e^{\tilde{b}} \Omega(L^{\alpha})) L_{\alpha}$$

$$= i^{-\alpha} \operatorname{Str}(e^{-b} \partial e^{b} \Omega^{2}(L^{\alpha})) \Omega(L_{\alpha}) = \Omega(j_{L_{\alpha}}).$$

$$(4.19)$$

To summarize, using (4.13) we find that the left-invariant-one-form transforms as

$$\widetilde{J}_B = *J_B,
\widetilde{J}_F = iJ_F,
\widetilde{j}_{L_{\alpha}} = \Omega(j_{L_{\alpha}}).$$
(4.20)

4.3 Flat-connection transformation under T-self-duality

Now that we know how the left-invariant-one-form transforms under T-duality, see (4.20), we can find the general transformation of the flat-connection (3.2). For simplicity of notation we define $b(z) = (z + z^{-1})/2$, so the flat connection takes the form

$$A(z) = b(z)^{2}j_{B} + b(iz)^{2}\Omega(j_{B}) + ib(z)b(iz) * (j_{B} - \Omega(j_{B})) + b(z)j_{F} - b(iz)\Omega(j_{F}).$$
(4.21)
$$= b(z)^{2}(J+j)_{B} + b(iz)^{2}\Omega((J+j)_{B}) + ib(z)b(iz) * ((J+j)_{B} - \Omega((J+j)_{B})) + b(z)(J+j)_{F} - b(iz)\Omega((J+j)_{F}).$$

Let now write down the dual flat-connection, plugging (4.20) into (4.21)

$$\tilde{A}(z) = b(z)^{2}(*J_{B} + \Omega(j_{B})) + b(iz)^{2}(\Omega(*J_{B}) + j_{B}) + ib(z)b(iz)(J_{B} + *\Omega(j_{B}) - \Omega(J_{B}) - *j_{B})$$

$$(4.22)$$

$$+b(z)(iJ_{F} + \Omega(j_{F})) - b(iz)(\Omega(iJ_{F}) - j_{F}).$$

As in [2], one can relate the two flat-connection by using a z-dependent automorphism $U_z(\cdot)$ which acts as follows

$$U_{z}(J) = f(z)\Omega(J), \quad U_{z}(\Omega(J)) = (-)^{J} f^{-1}(z)J,$$

$$U_{z}(j) = \Omega(J), \quad U_{z}(\Omega(j)) = (-)^{j} j,$$
(4.23)

with f(z) = -ib(iz)/b(z), so

$$U_z(A(z)) = \tilde{A}(z). \tag{4.24}$$

This automorphism is a composition of the \mathbb{Z}_4 automorphism and the one-parameter automorphism induced by the \mathbb{Z} -gradation (2.5), with $\lambda = f(z)$. That is

$$U_z(\cdot) = \sigma_{f(z)}(\Omega(\cdot)). \tag{4.25}$$

The U(1) used in [2] is U = D + B where B is the hypercharge. In table 3 we give more examples of possible U(1) charges for the GS sigma-model on $AdS_5 \times S^5$, as explained in section 2.3.

Repeating the arguments of [2][18][1], the A_1 Noether charges becomes trivial, the A_{-1} charges gets lifted and become non-local, and the B_0 generators remains local and transforms into themselves up to commutators and boundary terms.

U(1)	P	$Q^{i\alpha}$	$Q^{i'\alpha}$	$\hat{Q}^{\dot{lpha}}_i$	$\hat{Q}^{\dot{lpha}}_{i'}$	$R^i_{i'}$	R^i_i	D
D	1	1/2	1/2	1/2	1/2	0	0	0
В	0	1/2	1/2	-1/2	-1/2	0	0	0
Ř	0	1/2	-1/2	-1/2	1/2	1	0	0
D + B	1	1	1	0	0	0	0	0
D-B	1	0	0	1	1	0	0	0
$D + \check{R}$	1	1	0	0	1	1	0	0
$D - \tilde{R}$	1	0	1	1	0	-1	0	0
$B + \check{R}$	0	1	0	-1	0	1	0	0
$B - \check{R}$	0	0	1	0	-1	-1	0	0
2B	0	1	1	-1	-1	0	0	0

Table 3: U(1) charges.

We give U(1) generators, D, B and \check{R} , with respect to gradations of the SCA psu(2, 2|4). The charge of the SCA's generators under the U(1)'s and under their combinations that decomposes the SCA to $A_1 \oplus B_0 \oplus A_{-1}$ are given. i and i' are the su(4) indices when broken to su(2) × su(2). The charges of the generators related to those in the table by Ω , have the opposite charge, while the rest of the generators not given in the table have zero charge.

4.4 Quantum consistency of the T-Self-Duality transformation

In previous subsections we have shown that the GSSM is self-dual under T-duality where we made some assumption regarding the Z-gradation and Z₄ automorphisms of the SCA, and the possibility to kappa-gauge fix the action in a certain way. Though it is enough at the level of the classical action, at the quantum level we should also worry about the dilaton transformation under the T-duality transformation [19][20]. In order to have self-duality at the quantum level we need the dilaton to be left invariant under the transformation [1]. This means that the super-Jacobian of the transformation should equal one. This depends on how the generators in A_1 transform under conjugation with e^b .

For the Z-gradation induced by B+D and $D+\tilde{\lambda}$ for the type I and II respectively, a necessary condition for invariance of the dilaton is that the number of Q's will be twice the number of P's, since their charge under D is half of the charge of the P's, where $D \in \mathcal{H}_2$ and so should appears in the parametrization as in [1]. For type II, the number of Q's should also be twice the number of R's for the same reason (but with respect to $\tilde{\lambda} \in \mathcal{H}_2$), namely the number of P's should also equal the number of R's. We find that all SCA's satisfying this condition have vanishing Killing-form. It is known that GSSM on backgrounds based on supergroups with vanishing Killing-form are special, e.g these models are conformal invariant at one-loop [12][13][14]. This means that self-duality might be related to conformal invariance of the sigma-model. This condition is also required by the other Z-gradations considered in section 2.3. For example when U = 2B, if the supergroup does not have vanishing Killing-form, the hypercharge - $B \in \mathfrak{g}_{\bar{0}}$ and also $B \in \mathcal{H}_2$ so we can't gauge it away while all generators in A_1 (span $\{Q, \hat{S}\}$) have the same charge under it. As we will see later, there are cases where the action is self-dual classically, but the dilaton transforms non-trivially.

5. Classification of the backgrounds

In this section we would like to find all the GSSM on semi-symmetric backgrounds which are self-dual under T-duality along the directions of the abelian-subalgebras found in section 2.3. In order to do so, we have to find the backgrounds based on SCA's and Ω 's satisfying:

- $\Omega(U) = -U$.
- Rank(κ -symmetry) $\geq \dim(\mathfrak{g}_{\bar{1}})/4$.
- The Killing-form is degenerate.

We discuss separately the type I and type II SCA's. We consider backgrounds of dimension ≤ 10 with an AdS bosonic subspace. For $AdS_{n>2}$ we T-dualize along even number of bosonic directions or else we will switch type IIA with type IIB and vice versa.

For later use we note that the \mathbb{Z}_4 's satisfying (2.6), act on the SCA in the same way, namely

$$\Omega(P) \sim K, \quad \Omega(D) \sim D, \quad \Omega(L) \sim L,$$

$$\Omega(Q) \sim S, \quad \Omega(\hat{Q}) \sim \hat{S},$$
(5.1)

and for the type I R-symmetry

 $\Omega(R) \sim R,$

and for type II

 $\Omega(R) \sim \hat{R}, \quad \Omega(\lambda) \sim \lambda.$

5.1 Type I

In table 4 we give the \mathbb{Z}_4 automorphisms which satisfy the condition (2.6) for AdSsemi-symmetric spaces. As one can see, such a \mathbb{Z}_4 automorphism does not always exists. For each semi-symmetric space induced by Ω , we give the rank of the kappasymmetry, which should be \geq then quarter of the number of odd generators of the SCA (or half of then in case where $A_1 = \operatorname{span}\{Q, \hat{S}\})^5$. We also write if the super-Jacobian is unity or not based on the degeneracy of the Killing-form. By these

⁵Note that whenever the rank of kappa-symmetry is large enough to eliminate quarter of the odd degrees of freedom, it actually large enough to eliminate half of them. This can expected since one can interpret the charges transformation under the T-duality as a sort of rotation in the Yangian space, and in some sense the (Q, \hat{S}) duality can be thought of as a composition of (P, Q) and (K, \hat{S}) , so these two exist then we expect the other to also exist.

criteria we determine whether the sigma-model is self-dual or not at the classical and quantum levels.

In table 5 we give the \mathbb{Z}_4 automorphisms which satisfy (2.6) but induce semisymmetric space which in not a product of AdS background with some other compact space. In all of these cases the rank of kappa-symmetry is zero.

To summarize, we find there are two models, the $AdS_5 \times S^5$ and $AdS_2 \times S^2$ which are self-dual at the quantum level, these results were proven in [1][2] and in [6] respectively. There are two more backgrounds which are self-dual only at the classical level, these are the $AdS_5 \times S^1$ and $AdS_2 \times S^1$, the first one was given in [8] (although the dilaton shift was not discussed) and the second is new. These models should not be conformal invariant by themselves and one have to add D-branes in order to make them conformal invariant [21]. It might be that after adding open strings degrees of freedom one would find quantum consistent self-duality.

	PSU(1, 1 2)	$\begin{array}{c} \mathrm{SU}(1,1 N)\\ N>2 \end{array}$	PSU(2,2 4)	$ SU(2,2 2N) N \neq 2 $	OSP(2 2)	OSP(2 4)
B_0	SU(1 2)	$\mathrm{U}(1 N)$	SU(2 4)	U(2 2N)	U(1 1)	U(1 2)
Invariant $\operatorname{sub} - \operatorname{alg}.$	$U(1)^{2}$	$\mathrm{SO}(1,1) \times \mathrm{SO}(N)$	$USp(2,2) \times USp(4)$	$\mathrm{USp}(2,2) imes \mathrm{USp}(2N)$	U(1)	ø
Bosonic— subspace	$AdS_2 \times S^2$	$\mathrm{AdS}_2 \times \mathrm{S}^1 \times \mathrm{AI}(N)$	$AdS_5 \times S^5$	$\mathrm{AdS}_5 \times \mathrm{S}^1 \times \mathrm{AII}(N)$	$AdS_2 \times S^1$	ø
Rank of $\kappa - \text{symm.}$	4	0	16	8 for N=1, else 0	2	Ø
sJacobian	1	Ø	1	$\neq 1$	$\neq 1$	Ø
Self-dual	Yes-Q	No	Yes-Q	$\begin{array}{c} \text{Yes} - \text{C for N=1,} \\ \text{else No} \end{array}$	Yes-C	No

Table 4: AdS Semi-symmetric spaces based on type-I SCA's.

When the bosonic symmetric space is not AdS or a sphere, we symbolize it according to the Cartan classification, see [22]. Note that $\dim(AI(N)) = \frac{(N-1)(N+2)}{2}$ and $\dim(AII(N)) = (N-1)(2N+1)$. We symbolize with \emptyset the cases where we don't have relevant \mathbb{Z}_4 automorphism generating AdS space, and when the calculation of the super-Jacobian is not relevant. Yes-Q means the model is self-dual at the quantum level, and Yes-C means the model is self-dual only at the classical level. Note that $SU(1, 1|1) \simeq OSP(2|2)$.

Finally, we note that the background $\operatorname{AdS}_2 \times \mathbb{CP}^n$ has been claimed to be selfdual in [8]. This background does not appear in our classification as self-dual. The calculation in [8] has a flaw: the authors redefine the odd generators below (eq. 27) in that paper, but the redefinition is not one-to-one. Thus, the generators are no longer independent, and the algebra does not close and does not represent $\operatorname{SU}(1,1|N)$. The coset space is therefore not a quotient of a super-Lie-algebra and of-course does not give the $\operatorname{AdS}_2 \times \mathbb{CP}^n$ background.

5.2 Type II

In table 6 we give the \mathbb{Z}_4 automorphisms which satisfy the condition (2.6) and gives

	PSU(1, 1 2)	$\begin{array}{c} \mathrm{SU}(1,1 N)\\ N\neq 2 \end{array}$	PSU(2,2 4)	$ \begin{aligned} & \operatorname{SU}(2,2 N) \\ & N \neq 4 \end{aligned} $	OSP(2 2)	OSP(2 4)
B_0	SU(1 2)	$\mathrm{U}(1 N)$	SU(2 4)	$\mathrm{U}(2 N)$	U(1 1)	U(1 2)
Invariant $\operatorname{sub} - \operatorname{alg}.$	ø	Ø	$SO(2,2) \times SO(4)$	$SO(2,2) \times SO(N)$	Ø	U(2)
Bosonic— subspace	ø	Ø	$AI(2,2) \times AI(4)$	$AI(2,2) \times AI(N) \times S^1$	Ø	$\mathrm{CI}(4) \times \mathrm{S}^1$
Rank of $\kappa - \text{symm.}$	ø	Ø	0	0	Ø	0
Self-dual	No	No	No	No	No	No

Table 5: Non-AdS Semi-symmetric spaces based on type-I SCA's.

The notations are the same as in table 5. Note that $\dim(AI(N)) = \frac{(N-1)(N+2)}{2}$ and $\dim(C(4)) = 6$. We symbolize with \emptyset the cases where we don't have a \mathbb{Z}_4 automorphism generating non-AdS space.

 $AdS_2 \times \mathcal{M}$ backgrounds, and in table 7 backgrounds with $AdS_{n>3} \times \mathcal{M}$. The notations are the same as for the type I SCA's.

In table 8 we give the \mathbb{Z}_4 automorphisms which satisfy (2.6) but induce semisymmetric space which in not a product of AdS background with some other compact space.

For the F(4; 0) SCA, the semi-symmetric space satisfying (2.6) is $AdS_2 \times BDI(3; 4)$ with the invariant subalgebra $SO(3) \times SO(4) \times U(1)$ with 14-dimensional space-time so we don't treat it in table 6. Similarly we omit F(4; 2) from table 8 were the relevant semi-symmetric space is $BDI(2, 1; 3, 1) \times S^2$ with the invariant subalgebra $SO(2, 1) \times SO(3, 1) \times U(1)$ with 14-dimensional space-time.

To summarize, we find there are two models which are self-dual only at the classical level, these are the $AdS_2 \times S^4$ and $AdS_4 \times S^2$, these two model were not considered before in the context of T-self-duality and are not self-dual at the quantum level.

Table	Table 0. Ad52 benn-symmetric spaces based on type-11 SOA S.				
	OSP(2N 2), N > 1	$D(2,1;\alpha)$	$OSP(4^* 2N)$	$OSP(4^* 4N)$	
B_0	U(N 1)	U(2 1)	$\mathrm{U}(2 N)$	U(2 2N)	
Invariant sub – alg.	$SO(N)^2 \times U(1)$	$U(1)^3$	$\mathrm{U}(1)^2 \times \mathrm{U}(N)$	$U(2) \times USp^2(2N)$	
Bosonic— subspace	$AdS_2 \times BDI(N; N)$	$AdS_2{\times}S^2{\times}S^2$	$AdS_2 \times S^2 \times CI(N)$	$\mathrm{AdS}_2 \times \mathrm{CII}(N;N)$	
Rank of κ – symm.	0	0	0	8 for $N = 1$	
sJacobian	Ø	Ø	Ø	$\neq 1$ for $N = 1$	
Self-dual	No	No	No	$\begin{array}{c} \text{Yes} - \text{C for N=1,} \\ \text{else No} \end{array}$	

Table 6: AdS₂ Semi-symmetric spaces based on type-II SCA's.

Note that $\dim(\text{BDI}(p,q)) = pq$, $\dim(\text{CI}(N)) = N(N+1)$ and $\dim(\text{CII}(N;N)) = 4N^2$, for N = 1 we have $\text{CII}(1;1) \simeq S^4$.

	OSP(4N 4)	F(4;2)	$OSP(8^* 2N)$
B_0	U(2N 2)	$U(1) \times OSP(2 4)$	U(4 N)
Invariant $\operatorname{sub} - \operatorname{alg}.$	$U(2N) \times SP(2)^2$	Ø	Ø
Bosonic— subspace	$AdS_4 \times DIII(2N)$	Ø	Ø
Rank of κ – symm.	8 for $N = 1$	Ø	Ø
sJacobian	$\neq 1$ for $N = 1$	Ø	Ø
Self-dual	Yes - C for N=1,else No	No	No

Table 7: $AdS_{n>3}$ Semi-symmetric spaces based on type-II SCA's.

Note that $\dim(\text{DIII}(N)) = N(N-1)$, for N = 2 we have $\text{DIII}(2) = S^2$.

 Table 8: Non-AdS Semi-symmetric spaces based on type-I SCA's.

	OSP(2N 4), N > 1	$OSP(8^* 2N)$	$OSP(8^* 4N)$
B_0	U(N 2)	$\mathrm{U}(4 N)$	U(4 2N)
Invariant $\operatorname{sub} - \operatorname{alg}.$	$U(2) \times SO(N)^2$	$SO(3,1) \times SO(3,1) \times U(N)$	$U(2,2) \times USp(2N)^2$
Bosonic— subspace	$\operatorname{CI}(4) \times \operatorname{BDI}(N; N)$	$BDI(3,1;3,1) \times CI(N)$	$\mathrm{DIII}(2,2)\times\mathrm{CII}(N;N)$
Rank of κ – symm.	0	0	0
Self-dual	No	No	No

5.3 $AdS_3 \times M$ semi-symmetric spaces

Semi-symmetric spaces with AdS₃ subspace are generated by supergroups which are not simple, by taking a product of two supergroups with a bosonic subgroup of SU(1,1), so we have SU(1,1)× SU(1,1) \simeq SO(2,2) as a subgroup. Generally we always have a Z₄ automorphism for these products by taking the coset $\frac{G \times G}{G_{\text{bosonic}}}$ [23, 14]. The cases of $\frac{\text{PSU}(1,1|2)^2}{\text{SU}(1,1) \times \text{SU}(2)} \simeq \text{AdS}_3 \times \text{S}^3$ and $\frac{\text{D}(2,1;\alpha)^2}{\text{SU}(1,1) \times \text{SU}(2)^2} \simeq \text{AdS}_3 \times \text{S}^3 \times \text{S}^3$ are discussed in [12] and [23] respectively. Let us discuss the self-duality of these models for semi-symmetric spaces with irreducible sub-symmetric spaces. The candidates are the type I supergroups SU(1,1|N) and PSU(1,1|2), and the type II supergroups OSP(N|2), D(2,1;\alpha), F(4;0) and G(3;p).

The quotient satisfying (2.6) with dimension ≤ 10 are given in table 9. We find one self-dual model at the quantum level, $AdS_3 \times S^3$, which was proven to be self-dual in [6], and another new model self-dual only at the classical level, $AdS_3 \times S^1$.

5.4 AdS_2 semi-symmetric space

The case where the full bosonic space AdS_2 is somewhat degenerate. Usually one can gauge away all fermionic degrees of freedom using kappa-symmetry, and so it comes down to T-dualizing only along one bosonic coordinate. So classically the sigmamodel is self-dual, but as explained above, the dilaton will shift. For example one

Background	$AdS_3 \times S^1$	$\mathrm{AdS}_3 \times \mathrm{S}^3$	$AdS_3 \times S^3 \times S^3$
Quotient	$\frac{\mathrm{SU}(1,1 1)^2}{\mathrm{SU}(1,1)\times\mathrm{U}(1)}$	$\frac{\mathrm{PSU}(1,1 1)^2}{\mathrm{SU}(1,1)\times\mathrm{SU}(2)}$	$\frac{\mathrm{D}(2,1,\alpha)^2}{\mathrm{SU}(1,1)\times\mathrm{SO}(4)}$
kappa-rank	4	8	0
sJacobian	$\neq 1$	1	Ø
Type	Ι	Ι	II
Self-dual	Yes-C	Yes-Q	No

Table 9: AdS_3 Semi-symmetric spaces.

The osp(4|2) is a special case of the $D(2, 1; \alpha)$ with $\alpha = 1$.

can construct these models using $OSP(2|2)/(SO(2) \times U(1))$ [24] or $OSP(4|2)/(SO(4) \times U(1))$ [17].

5.5 When does an AdS semi-symmetric space satisfies $\Omega(U) = -U$?

As one can see from the tables above, we cannot find an automorphism generating AdS subspace which also satisfies (2.6), for all SCA's. We can see how the problem comes about when we look at the anti-commutation relations of the odd generators, and how they close on the Lorentz subalgebra, M_{ab} , plus noting the relations (5.1) for the \mathbb{Z}_4 automorphism. In both cases (type-I and type-II) we have $\{Q_{\alpha}^l, S_{\beta}^k\} \sim$ $\delta^{kl}M_{\alpha\beta}+\ldots$ where α,β are the spinor indices in d-dimensions and k, l are R-symmetry indices. When d < 6 there are two cases where $M_{\alpha\beta} = M_{\beta\alpha}$ or $M_{\alpha\beta} = -M_{\beta\alpha}$. When $M_{\alpha\beta}$ is antisymmetric we can just take $\Omega(Q^l_{\alpha}) = iS^l_{\alpha}$ and $\Omega(S^l_{\alpha}) = iQ^l_{\alpha}$ which gives the desired invariant bosonic subalgebra to induce AdS_{d+1} space times some internal space. This is the case for d = 2. When $M_{\alpha\beta}$ is symmetric we have to take $\Omega(Q^l_{\alpha}) = i C^l_{\ k} S^k_{\alpha}$ and $\Omega(S^l_{\alpha}) = -i C^l_{\ k} Q^k_{\alpha}$ where C is antisymmetric full rank matrix in order to get an AdS space. This is the case for $d = 3, 4, 5^6$. Such a matrix exists only for even dimension of the R-symmetry representation of the odd generators. For d = 6 the Lorentz generators are combinations of symmetric and anti-symmetric parts, so it is not possible to find automorphism satisfying (2.6) which will give AdS subspace. The details for all SCA's are found in appendix B.

6. Discussion

In the paper we analyzed properties of GSSM's on semi-symmetric spaces under Tduality. For SCA's with Z-gradation under U, with gradings $\pm 1, 0$ only, we found three algebraic conditions that guarantee T-self-duality of the sigma-models. These are:

⁶For d = 3 we have $M_{\alpha\beta} = M_{\beta\alpha}$, $\alpha, \beta = 1, 2$, spin(3) \simeq SU(2), for d = 4 we have self and antiself-dual generators, $M_{\alpha\beta} = M_{\beta\alpha}$ and $M_{\dot{\alpha}\dot{\beta}} = M_{\dot{\beta}\dot{\alpha}}$, $\alpha, \beta, \dot{\alpha}, \dot{\beta} = 1, 2$, spin(4) \simeq SU(2)×SU(2) and for d = 5 we have $M_{ab} = M_{ba}$, a, b = 1, 2, 3, 4, spin(5) \simeq SP(4). For d = 6 we have M_{ab} traceless, a, b = 1, 2, 3, 4, spin(6) \simeq SU(4).

- 1. $\Omega(U) = -U$, where Ω is the \mathbb{Z}_4 automorphism map.
- 2. Rank(κ -symmetry) $\geq \dim(\mathfrak{g}_{\bar{1}})/4$.
- 3. The SCA's Killing-form vanishes.

We found that only three backgrounds are consistent with all three conditions. These are the $AdS_n \times S^n$ for n = 2, 3, 5, which were found previously to be T-self-dual [1][2][6]. All of these backgrounds are constructed from the type I SCA's PSU(N, N|2N) with N = 1, 2.

The last condition is necessary for quantum consistency of the transformation; it implies the unity of the super-Jacobian of the transformation. We found that there are backgrounds that satisfy the first two conditions, namely they are self-dual at the level of the classical action, but their dilaton transforms non-trivially under the transformation. These are the $AdS_n \times S^1$ for n = 2, 3, 5 which are constructed from the type I SCA's SU(N, N|N) with N = 1, 2 (the N = 2 case was studied in [8]), and the $AdS_4 \times S^2$ and $AdS_2 \times S^4$ which are constructed from the type II SCA OSP(4|4)(and its real-forms). All other backgrounds satisfying the first condition have rank zero kappa-symmetry.

The last condition, for quantum consistency of the transformation, also implies the vanishing of the beta-function at one-loop [13]. This might mean that having self-duality only at classical level is related to lack of worldsheet conformal invariance. If true, adding D-branes degrees of freedom [21] may be a way to fix it.

The classification of backgrounds satisfying the first condition follows from a relation between the spinor representation of the SCA and the R-symmetry representation of the odd generators. Besides the PSU SCA's, among the backgrounds constructed by SCA's with zero Killing-form, generating AdS spaces, (namely, D(2, 1; α), OSP(6|4) and their direct products⁷) only the D(2, 1; α) and D(2, 1; α)² SCA's admit a semi-symmetric space satisfying the first condition (AdS₂×S²×S² and AdS₃×S³×S³ respectively), but these background's kappa-symmetry rank vanishes [23][14]. The semi-symmetric space constructed from OSP(6|4) that satisfies the first condition is not an AdS space, where the AdS₄ × CP³ backgrounds doesn't satisfy it. There is also one semi-symmetric space constructed from OSP(4|2) which doesn't satisfy the condition which is AdS₂ × S³.

We found there are several new families of coordinate directions (or equivalently abelian subalgebras), along which the self-dual sigma-models also admit self-duality under T-duality summarized in figure 2, one of them includes only fermionic directions. These families are different for type I and type II SCA's. The new families should generate a dual-SCA similar to the well known one [25] which is associated with T-dualizing along the P and Q directions, but the heuristic transformation in

⁷We include OSP(4|2) as a special case of $D(2, 1; \alpha)$.

the Yangian charges space given in [18] should be with respect to the \mathbb{Z} -gradation charges of the generators.

For the scattering amplitudes on the gauge theory side, one associates dual variables, such that the scattering amplitudes in terms of theses variables shows manifest invariance under the dual-superconformal symmetry [25]. The self-duality along different directions introduces other dual-superconformal symmetries (which ofcourse related to the same Yangian), so in principle we can construct dual variables in analogy to the one constructed to for T-duality along span{P, Q} [25] (see the first line in table 10). We give these dual-variables in table 10. The problem is that in terms of the on-shell variables, the dual variables include derivatives, so in order to construct them we fourier transform them. After fourier transforming the scattering amplitude, the dual variables do not appear in delta-functions as in case of T-duality along span{P, Q}, and the dual-symmetry is not manifest.

	Table 10: Dual variables.				
U	Dual variables				
D+B	$(x_i - x_{i+1})_{\alpha \dot{\alpha}} = \lambda_{i\alpha} \tilde{\lambda}_{i\dot{\alpha}}, (\theta_i - \theta_{i+1})^A_{\alpha} = \lambda_{i\alpha} \eta^A_i$				
2B	$(\theta_i - \theta_{i+1})^A_{\alpha} = \lambda_{i\alpha} \eta^A_i, (\xi_i - \xi_{i+1})^A_{\dot{\alpha}} = \mu_{i\dot{\alpha}} \eta^A_i$				
$D + \check{R}$	$(x_i - x_{i+1})_{\alpha\dot{\alpha}} = \lambda_{i\alpha}\tilde{\lambda}_{i\dot{\alpha}}, (\theta_i - \theta_{i+1})^A_{\alpha} = \lambda_{i\alpha}\eta^A_i, (\hat{\theta}_i - \hat{\theta}_{i+1})_{\dot{\alpha}A'} = \tilde{\lambda}_{i\dot{\alpha}}\psi_{iA'}, (r_i - r_{i+1})^A_{A'} = \eta^A_i\psi_{iA'}$				
$B + \check{R}$	$(\theta_i - \theta_{i+1})^A_\alpha = \lambda_{i\alpha}\eta^A_i, (\xi_i - \xi_{i+1})^A_\alpha = \mu_{i\dot\alpha}\eta^A_i, (r_i - r_{i+1})^A_{A'} = \eta^A_i\psi_{iA'}$				

The on-shell variables of the $\mathcal{N} = 4$ SYM scattering amplitudes are $\lambda, \tilde{\lambda}$ and η . μ and ψ are the fourier transforms of $\tilde{\lambda}$ and η respectively.

It is interesting to find if there are objects on the gauge theory side that can be related based on the other dual-superconformal algebras (similar to the scattering amplitudes/Wilson loops duality).

We showed that the flat-connection transformation under T-duality is a parameter dependent automorphism, which is the \mathbb{Z}_4 automorphism Ω followed by conjugation with $f(z)^U$ where U is the generator inducing the \mathbb{Z} -gradation and f is always the same function depending on the spectral parameter z.

For the case of the $\operatorname{AdS}_4 \times \mathbb{CP}^3$ background, where the first condition is not satisfied, namely $\Omega(U) \neq -U$, there is evidence on the gauge theory side that the background may have similar properties to the $\operatorname{AdS}_5 \times \operatorname{S}^5$ background which were interpreted as a consequence of T-self-duality of the background, [26][27][28][29]. On the other hand, there is also other evidence that this background is not self-dual [6][7][30][31].

The condition $\Omega(U) = -U$ seems also to be related to the Pohlmeyer-reduction of the $\operatorname{AdS}_n \times \operatorname{S}^n$ sigma-models introduced in [32]. A key property of the SCA used in the procedure of [32] was to further decompose the SCA (on top of the \mathbb{Z}_4 decomposition), such that a generator $T \in \mathcal{H}_2$ forms the projection $\mathcal{P}(\cdot) = [T, [T, \cdot]]$ of a \mathbb{Z}_2 -decomposition. Since our U(1) gives charges ± 1 and 0 to all generators, taking T = U the projection is $\mathcal{P}(L_a) = |a|L_a$ with $a = \pm 1, 0$, so the sets $A_1 \oplus A_{-1}$ have grading 1 and B_0 grading 0. The condition $T \in \mathcal{H}_2$ was essential in the reduction procedure where elements in \mathcal{H}_2 were gauge fixed to T. So actually $\Omega(U) = -U$ is not enough, but we also need the \mathbb{Z} -gradation automorphism to be inner, which is possible for all SCA's which were found to be self-dual⁸. In cases where $T \in \mathcal{H}_0$ one might expect Pohlmeyer-reduction procedure to fail, e.g for the $\mathrm{AdS}_4 \times \mathbb{CP}^3$ background. The Pohlmeyer-reduction also relies heavily on the possibility to use kappa-symmetry as does the T-duality procedure.

Acknowledgements

We would like to thank Ido Adam for valuable discussions and comments on the manuscript. The work is supported in part by the Israeli Science Foundation center of excellence, by the Deutsch-Israelische Projektkooperation (DIP), by the US-Israel Binational Science Foundation (BSF), and by the German-Israeli Foundation (GIF).

A. Notations

In this section we summarize our notations used throughout the main text.

We denote the superconformal algebras (SCA's) by \mathfrak{g} , with the \mathbb{Z}_2 decomposition $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ to its even and odd parts respectively. \mathfrak{g}_I , \mathfrak{g}_{II} will denote type-I and Type-II SCA's respectively.

 Ω is the \mathbb{Z}_4 automorphism map, which decomposes the SCA as $\mathfrak{g} = \bigoplus_{i=0}^3 \mathcal{H}_i$, where *i* denotes the grading.

The \mathbb{Z} -gradation decomposition with gradings $\pm 1, 0$ only is induced by U, with the charge given as the eigenvalue of ad_U . We denote the decomposition as follows, $\mathfrak{g} = A_{-1} \oplus B_0 \oplus A_1$, where the subscript indicating the grading. The \mathbb{Z} -gradation also induces the map $\sigma_{\lambda}(L_a) = \lambda^U L_a \lambda^{-U} = \lambda^a L_a$, where $[U, L_a] = aL_a, L_a \in \mathfrak{g}$, and $a = \pm 1, 0$.

We use the left-invariant-one-form, $j = g^{-1}dg$, which in the (2d-worldsheet) conformal basis is given by $j = g^{-1}\partial g$, $\bar{j} = g^{-1}\bar{\partial}g$. We use two decompositions of j, one according to the \mathbb{Z} -gradation, $j \equiv J + j$ where $J \in A_1$ and $j \in B_0$, and the other according to the \mathbb{Z}_2 -grading, $j = j_B + j_F$ where $j_B \in \mathfrak{g}_{\bar{0}}$ and $j_F \in \mathfrak{g}_{\bar{1}}$.

Our indices conventions are

- I, J, K, \dots Indices of generators in A_1 .
- $\alpha, \beta, \gamma, \dots$ Indices of generators in B_0 diagonal with respect to Ω .

⁸That is, for the PSU SCA's we can take $U = D + \check{R}$.

• A, B, C, \dots - Indices of any generator in the SCA.

We use the bilinear-form $\eta_{AB} = \text{Str}(L_A \Omega(L_B))$ which is symmetric for both even and odd generators.

B. Superalgebras

In this section we find the \mathbb{Z}_4 automorphisms satisfying (2.6) for all SCA's with the gradations discussed in section 2. For simplicity we will not quote the entire commutation relations of the SCA's. Instead we will give only the anti-commutation relations that will suffice to constrain the automorphism. For type I and type II SCA's we assume the \mathbb{Z}_4 automorphism transformations (5.1). Since theses automorphisms interchanges P with K, we must require $\Omega(D) = -D$. We also note that in order to get an AdS subspace we need $\Omega(M_{ab}) = M_{ab}$ for the Lorentz rotations. Throughout the subsections we use the matrices C and F, defined such that $C_{ij}C^{jk} = \delta_i^k$ and $F_a{}^bF_b{}^c = \delta_a^c$ where l, k are R-symmetry indices and a, b are spinor indices. We assume $C_{ij} = (-)^{s_c}C_{ij}$ and $F_{ab} = (-)^{s_f}F_{ba}$ where $s_c, s_f = 0$ or 1. We raise and lower the spinor indices using the charge conjugation matrix $\epsilon, \psi^a = \epsilon^{ab}\psi_b$ and $\psi_a = \epsilon_{ab}\psi^b$ and $\epsilon_{ab}\epsilon^{bc} = -\delta_a^c$. We use the standard semi-symmetric spaces notations [22] whenever the space is not a sphere or AdS, with a superscript/subscript indicating the dimensionality.

B.1 OSP(2N|2)

The relevant commutation relations are

$$\{Q_l, S^k\} = \delta_l^k D + \lambda_l^k, \quad \{Q_l, \hat{S}_k\} = R_{lk}, \quad \{\hat{Q}^l, S^k\} = \hat{R}^{lk}.$$
(B.1)

where l = 1, ..., N is the R-symmetry index, λ_l^k form U(N) subalgebra of SO(2N), and $R_{lk} = -R_{kl}$, $\hat{R}^{lk} = -\hat{R}^{kl}$. The automorphism transformation is

$$\Omega(Q_l) = iC_{lk}S^k, \quad \Omega(S^l) = iC^{lk}Q_k.$$
(B.2)

Because $\Omega(D) = -D$, we must have $C_{lk} = C_{kl}$. This implies the transformation

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \mp(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}),$$
(B.3)
$$\Omega(C^{[pl}\lambda_{l}{}^{k]}) = C^{[pl}\lambda_{l}{}^{k]}, \quad \Omega(C^{(pl}\lambda_{l}{}^{k)}) = -C^{(pl}\lambda_{l}{}^{k)}.$$

So the semi-symmetric space induced by the automorphism is

$$\operatorname{AdS}_{2} \times \operatorname{BDI}(N; N)^{N^{2}} \simeq \frac{\operatorname{OSP}(2N|2)}{\operatorname{SO}(N)^{2} \times \operatorname{U}(1)},$$
(B.4)

which for N = 1 is

$$\operatorname{AdS}_2 \times \operatorname{S}^1 \simeq \frac{\operatorname{OSP}(2|2)}{\operatorname{U}(1)}.$$
 (B.5)

B.2 OSP(2N|4)

The relevant commutation relations are

$$\{Q_{l\alpha}, S^k_{\beta}\} = \delta^k_l(\epsilon_{\alpha\beta}D + M_{\alpha\beta}) + \epsilon_{\alpha\beta}\lambda^k_l, \quad \{Q_{l\alpha}, \hat{S}_{k\beta}\} = \epsilon_{\alpha\beta}R_{lk}, \quad \{\hat{Q}^l_{\alpha}, S^k_{\beta}\} = \epsilon_{\alpha\beta}\hat{R}^{lk}.$$
(B.6)

where l = 1, ..., N is the R-symmetry index, λ_l^k form U(N) subalgebra of SO(2N), and $R_{lk} = -R_{kl}$, $\hat{R}^{lk} = -\hat{R}^{kl}$. $\alpha = 1, 2$ is the spinor index in the representation spin(3) \simeq SU(2) and $M_{\alpha\beta} = M_{\beta\alpha}$. The automorphism transformation is

$$\Omega(Q_{l\alpha}) = iC_{lk}F_{\alpha}{}^{\beta}S_{\beta}^{k}, \quad \Omega(S_{\alpha}^{l}) = iC^{lk}F_{\alpha}{}^{\beta}Q_{k\beta}.$$
(B.7)

We get $\Omega(D) = (-)^{1+s_c+s_f}D$, so we must have $C_{lk} = (-)^s C_{kl}$ and $F_{\alpha\beta} = (-)^s F_{\beta\alpha}$. Only for s = 1 we get AdS subspace (i.e, $F = \epsilon$ and $\Omega(M_{\alpha\beta}) = M_{\alpha\beta}$). If we take s = 1 we must work with even N. The transformations of the R-symmetry are,

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \pm (C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}), \qquad (B.8)$$
$$\Omega(C^{[pl}\lambda_l{}^{k]}) = -C^{[pl}\lambda_l{}^{k]}, \quad \Omega(C^{(pl}\lambda_l{}^{k)}) = C^{(pl}\lambda_l{}^{k)}.$$

So the semi-symmetric space induced by the automorphism is

$$AdS_4 \times DIII(2N)^{2N(2N-1)} \simeq \frac{OSP(4N|4)}{U(2N) \times SP(2)^2}.$$
 (B.9)

If we take s = 0 we have

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \mp (C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}), \qquad (B.10)$$
$$\Omega(C^{[pl}\lambda_l^{k]}) = C^{[pl}\lambda_l^{k]}, \quad \Omega(C^{(pl}\lambda_l^{k)}) = -C^{(pl}\lambda_l^{k)}, \qquad \Omega(M_{\alpha\beta}) = -F_{\alpha}{}^{\gamma}F_{\beta}{}^{\delta}M_{\delta\gamma}.$$

and we get the non-AdS background semi-symmetric space induced by the automorphism

$$\operatorname{CI}(4)_6 \times \operatorname{BDI}(N; N)^{N^2} \simeq \frac{\operatorname{OSP}(2N|4)}{\operatorname{SO}(N) \times \operatorname{SO}(N) \times \operatorname{U}(2)}.$$
 (B.11)

B.3 F(4;2)

The relevant commutation relations are

$$\{Q_{\alpha}, S_{\beta}\} = \epsilon_{\alpha\beta}D + M_{\alpha\beta} + \epsilon_{\alpha\beta}\lambda, \quad \{Q_{\alpha}, \hat{S}_{\beta}\} = \epsilon_{\alpha\beta}R, \quad \{\hat{Q}_{\alpha}, S_{\beta}\} = \epsilon_{\alpha\beta}\hat{R}.$$
(B.12)

 $\alpha = 1, 2, 3, 4$ is the spinor index in the representation spin(5) \simeq SP(4), $\epsilon^T = -\epsilon$ and $M_{\alpha\beta} = M_{\beta\alpha}$. The automorphism transformation is

$$\Omega(Q_{\alpha}) = iF_{\alpha}{}^{\beta}S_{\beta}, \quad \Omega(S_{\alpha}) = iF_{\alpha}{}^{\beta}Q_{k\beta}.$$
(B.13)

We get $\Omega(D) = (-)^{1+s_f} D$, so we must have $F_{\alpha\beta} = F_{\beta\alpha}$. But then we have $\Omega(F_{[\gamma}{}^{\alpha}M_{\alpha\beta]}) = F_{[\gamma}{}^{\alpha}M_{\alpha\beta]}$, so the invariant subalgebra contains just part of the Lorentz subalgebra so(1, 4), namely the subalgebra $F_{[\gamma}{}^{\alpha}M_{\alpha\beta]} \simeq u(1, 1) \in \mathcal{H}_0$. So we get the non-AdS semi-symmetric space

BDI
$$(3, 1; 2, 1)_{12} \times S^2 \simeq \frac{F(4; 2)}{SO(3, 1) \times SO(2, 1) \times U(1)}$$
. (B.14)

B.4 F(4;0)

The relevant commutation relations are

$$\{Q_{\alpha}, S_{\beta}\} = \epsilon_{\alpha\beta}D + \lambda_{\alpha\beta} + \epsilon_{\alpha\beta}\lambda, \quad \{Q_{\alpha}, \hat{S}_{\beta}\} = R_{\alpha\beta}, \quad \{\hat{Q}_{\alpha}, S_{\beta}\} = \hat{R}_{\alpha\beta}.$$
(B.15)

 $\alpha, \beta = 1, 2, 3, 4$ is the spinor index in the representation of the R-symmetry spin(5) \simeq SP(4), $\lambda_{\alpha\beta} = (\epsilon\Gamma^a\Gamma^b)_{\alpha\beta}N_{ab} = \lambda_{\beta\alpha}$ is ten-dimensional and $R_{\alpha\beta} = (\epsilon\Gamma^a)_{\alpha\beta}A_a = -R_{\beta\alpha}$ and $\hat{R}_{\alpha\beta} = (\epsilon\Gamma^a)_{\alpha\beta}B_a = -\hat{R}_{\beta\alpha}$ are five-dimensional (the Γ 's are 4×4 gamma-matrices in five dimensions, a = 1, ..., 5, we have the constraint $\text{Tr}(\epsilon R) = \text{Tr}(\epsilon \hat{R}) = 0$). The automorphism transformation is

$$\Omega(Q_{\alpha}) = iF_{\alpha}{}^{\beta}S_{\beta}, \quad \Omega(S_{\alpha}) = iF_{\alpha}{}^{\beta}Q_{\beta}.$$
(B.16)

We get $\Omega(D) = (-)^{1+s_f} D$, so we must have $F_{\alpha\beta} = F_{\beta\alpha}$. Then we have $\Omega(F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}) = F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}$, so the invariant subalgebra contains just part of the subalgebra $\lambda_{\alpha\beta} \simeq$ so(5), namely the subalgebra $F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]} \simeq u(2) \in \mathcal{H}_0$. We also have $\Omega(\lambda) = -\lambda$ and $\Omega(F_{\gamma}{}^{\alpha}R_{\alpha\beta}) = -F_{\beta}{}^{\alpha}\hat{R}_{\alpha\gamma}$. So the semi-symmetric space is

$$AdS_2 \times BDI(3;4)^{12} \simeq \frac{F(4;0)}{SO(3) \times SO(4) \times U(1)}.$$
 (B.17)

B.5 SU(1, 1|N), $N \neq 2$

The relevant commutation relations are

$$\{Q_l, S^k\} = \delta_l^k (D+A) + \lambda_l^k, \quad \{Q_l, S_k\} = 0, \quad \{\hat{Q}^l, \hat{S}^k\} = 0, \quad (B.18)$$

where l = 1, ..., N is the R-symmetry index, $(\lambda_l^k)^{\dagger} = -\lambda_k^l$ are the SU(N) generators, and A is the U(1). The automorphism transformation is

$$\Omega(Q_l) = iC_{lk}S^k, \quad \Omega(S^l) = iC^{lk}Q_k.$$
(B.19)

In order to have $\Omega(D) = -D$, we must take $C_{lk} = C_{kl}$. This implies the transformation

$$\Omega(C^{[pl}\lambda_l^{k]}) = C^{[pl}\lambda_l^{k]}, \quad \Omega(C^{(pl}\lambda_l^{k)}) = -C^{(pl}\lambda_l^{k)}, \quad \Omega(A) = -A.$$
(B.20)

So SO(N) $\simeq C^{[pl}\lambda_l^{k]} \in H_0$. This implies the semi-symmetric space induced by the DIA is

$$\operatorname{AdS}_{2} \times \operatorname{AI}(N)^{\frac{(N-1)(N+2)}{2}} \times \operatorname{S}^{1} \simeq \frac{\operatorname{SU}(1,1|N)}{\operatorname{U}(1) \times \operatorname{SO}(N)}.$$
 (B.21)

B.6 SU(2, 2|N), $N \neq 4$

The relevant commutation relations are

$$\{Q_{l\alpha}, S^k_{\beta}\} = \delta^k_l(\epsilon_{\alpha\beta}(D+A) + M_{\alpha\beta}) + \epsilon_{\alpha\beta}\lambda_l^k, \quad \{Q_{l\alpha}, S_{k\beta}\} = 0, \quad \{\hat{Q}^l_{\dot{\alpha}}, \hat{S}^k_{\dot{\beta}}\} = 0, \quad (B.22)$$

where l = 1, ..., N is the R-symmetry index, $(\lambda_l{}^k)^{\dagger} = -\lambda_k{}^l$ are the SU(N) generators, and A is the U(1). $\alpha = 1, 2$ and $\dot{\alpha} = 1, 2$ are the spinor indices in the representation spin(4) \simeq SU(2) \times SU(2) and $M_{\alpha\beta} = M_{\beta\alpha}$ and we also have $M_{\dot{\alpha}\dot{\beta}} = M_{\alpha\beta}^{\dagger}$. The automorphism transformation is

$$\Omega(Q_{l\alpha}) = iC_{lk}F_{\alpha}{}^{\beta}S_{\beta}^{k}, \quad \Omega(S_{\alpha}^{l}) = iC^{lk}F_{\alpha}{}^{\beta}Q_{k\beta}.$$
(B.23)

We get $\Omega(D) = (-)^{1+s_c+s_f}D$, so we must have $C_{lk} = (-)^s C_{kl}$ and $F_{\alpha\beta} = (-)^s F_{\beta\alpha}$. Only for s = 1 we get AdS subspace $(F = \epsilon)$. If we take s = 1 we must work with even N. The transformations of the R-symmetry are,

$$\Omega(C^{(ml}\lambda_l^{k)}) = C^{(ml}\lambda_l^{k)}, \quad \Omega(C^{[ml}\lambda_l^{k]}) = -C^{[ml}\lambda_l^{k]}, \quad \Omega(A) = -A.$$
(B.24)

So $\mathrm{USp}(N) \simeq C^{(pl}\lambda_l^{(k)} \in H_0$, and the semi-symmetric space induced by the transformation is

$$AdS_5 \times AII(N)^{(N-1)(2N+1)} \times S^1 \simeq \frac{SU(2,2|2N)}{USp(2,2) \times USp(2N)}.$$
 (B.25)

If we take s = 0 we have

$$\Omega(C^{(ml}\lambda_l{}^{k)}) = -C^{(ml}\lambda_l{}^{k)}, \quad \Omega(C^{[ml}\lambda_l{}^{k]}) = C^{[ml}\lambda_l{}^{k]}, \quad \Omega(A) = -A, \quad (B.26)$$
$$\Omega(M_{\alpha\beta}) = -F_{\alpha}{}^{\gamma}F_{\beta}{}^{\delta}M_{\delta\gamma}, \quad \Omega(M_{\dot{\alpha}\dot{\beta}}) = -F_{\dot{\alpha}}{}^{\dot{\gamma}}F_{\dot{\beta}}{}^{\dot{\delta}}M_{\dot{\delta}\dot{\gamma}},$$

so $SO(N) \simeq C^{[pl}\lambda_l^{k]} \in H_0$, and we get the non-AdS background semi-symmetric space induced by the automorphism

$$\operatorname{AI(2,2)}_{6} \times \operatorname{AI(N)}^{\frac{(N-1)(N+2)}{2}} \times \operatorname{S}^{1} \simeq \frac{\operatorname{SU}(2,2|N)}{\operatorname{SO}(2,2) \times \operatorname{SO}(N)}$$
 (B.27)

B.7 PSU(N, N|2N), N = 1, 2

The PSU superalgebras have the same structure as the SU superalgebras with the (important) modification of eliminating the S¹. For PSU(1, 1|2) we get

$$AdS_2 \times S^2 \simeq \frac{PSU(1,1|2)}{U(1) \times SO(2)}.$$
(B.28)

For PSU(2, 2|4) we get

$$AdS_5 \times S^5 \simeq \frac{PSU(2,2|4)}{USp(2,2) \times USp(4)},$$
(B.29)

or

$$\operatorname{AI}(2,2)_6 \times \operatorname{AI}(4)^9 \simeq \frac{\operatorname{PSU}(2,2|N)}{\operatorname{SO}(2,2) \times \operatorname{SO}(4)}.$$
 (B.30)

In the case of $AdS_2 \times S^2$ the rank of the κ -symmetry is 4 [12]. In the case of $AdS_5 \times S^5$ the rank of the κ -symmetry is 16 [16]. In the case of $AI(2,2)_6 \times AI(4)^9$ the rank of the κ -symmetry is 0.

B.8 $D(2,1;\zeta)$

The relevant commutation relations are

$$\{Q_{\alpha}, S_{\beta}\} = \epsilon_{\alpha\beta}D + \epsilon_{\alpha\beta}\lambda + \lambda_{\alpha\beta}, \qquad (B.31)$$

$$\{Q_{\alpha}, \hat{S}_{\beta}\} = \epsilon_{\alpha\beta} R, \quad \{\hat{Q}_{\alpha}, S_{\beta}\} = \epsilon_{\alpha\beta} \hat{R}, \tag{B.32}$$

where $\zeta = \cos^2(\phi)$, $\lambda = \cos^2 \phi L_3$ and $\lambda_{\alpha\beta} = \lambda_{\beta\alpha} = -\sin^2 \phi (\epsilon \gamma^a R_a)_{\alpha\beta} \simeq \text{su}(2)$, $R = -\cos^2 \phi L_+$ and $\hat{R} = \cos^2 \phi L_-$. $\alpha = 1, 2$ is a spinor index of the R-symmetry spin(2). The automorphism transformation is

$$\Omega(Q_{\alpha}) = i F_{\alpha}{}^{\beta} S_{\beta}, \quad \Omega(S_{\alpha}) = i F_{\alpha}{}^{\beta} Q_{k\beta}.$$
(B.33)

We get $\Omega(D) = (-)^{1+s_f} D$, so we must have $F_{\alpha\beta} = F_{\beta\alpha}$. Then we have $\Omega(F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}) = F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}$, so $U(1) \simeq F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]} \in \mathcal{H}_0$. We also have $\Omega(\lambda) = -\lambda$ and $\Omega(R) = -\hat{R}$. The semi-symmetric space is

$$AdS_2 \times S^2 \times S^2 \simeq \frac{D(2,1;\alpha)}{U(1) \times U(1) \times U(1)}.$$
(B.34)

The κ -symmetry rank is zero [14].

B.9 OSP $(4^*|2N)$

The relevant commutation relations are

$$\{Q_{l\alpha}, S^k_\beta\} = \epsilon_{\alpha\beta} \delta^k_l D + \epsilon_{\alpha\beta} \lambda_l^{\ k} + \delta^k_l \lambda_{\alpha\beta}, \tag{B.35}$$

$$\{Q_{l\alpha}, \hat{S}_{k\beta}\} = \epsilon_{\alpha\beta} R_{lk}, \quad \{\hat{Q}^l_{\alpha}, S^k_{\beta}\} = \epsilon_{\alpha\beta} \hat{R}^{lk}, \tag{B.36}$$

 $\lambda_{\alpha\beta} = \lambda_{\beta\alpha}$ form SU(2) subalgebra of SO^{*}(4), λ_l^k forms U(N) subalgebra of USp(2N), $R_{lk} = R_{kl}$ and $\hat{R}^{lk} = \hat{R}^{kl}$ (l, k = 1, ..., N). $\alpha = 1, 2$ is a spinor index of the R-symmetry spin(2). The automorphism transformation is

$$\Omega(Q_{l\alpha}) = iC_{lk}F_{\alpha}{}^{\beta}S_{\beta}^{k}, \quad \Omega(S_{\alpha}^{l}) = iC^{lk}F_{\alpha}{}^{\beta}Q_{k\beta}.$$
(B.37)

We get $\Omega(D) = (-)^{1+s_c+s_f}D$, so we must have $C_{lk} = (-)^s C_{kl}$ and $F_{\alpha\beta} = (-)^s F_{\beta\alpha}$. Only for s = 1 we get AdS subspace $(F = \epsilon)$. If we take s = 1 we must work with even N. The transformations of the R-symmetry are,

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \pm(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}),$$
(B.38)

$$\Omega(C^{(ml}\lambda_l{}^{k)}) = C^{(kl}\lambda_l{}^{m)}, \quad \Omega(C^{[ml}\lambda_l{}^{k]}) = -C^{[kl}\lambda_l{}^{m]},$$

$$\Omega(F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}) = -F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}, \quad \Omega(F_{(\gamma}{}^{\alpha}\lambda_{\alpha\beta)}) = F_{(\gamma}{}^{\alpha}\lambda_{\alpha\beta)}.$$

So the invariant subalgebra under the automorphism includes $SP(N) \simeq C^{(ml}\lambda_l^{k)}$ and $SU(2) \simeq F_{(\gamma}{}^{\alpha}\lambda_{\alpha\beta)}$. Thus, the semi-symmetric space induced by the automorphism is

$$\operatorname{AdS}_{2} \times \operatorname{CII}(N; N)^{4N^{2}} \simeq \frac{\operatorname{OSP}(4^{*}|4N)}{\operatorname{U}(2) \times \operatorname{SP}(2N)^{2}},$$
(B.39)

note that $\operatorname{CII}(1; 1) \simeq \mathrm{S}^4$. If we take s = 0 we get

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \mp (C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}), \qquad (B.40)$$
$$\Omega(C^{(ml}\lambda_l{}^{k)}) = -C^{(kl}\lambda_l{}^{m)}, \quad \Omega(C^{[ml}\lambda_l{}^{k]}) = C^{[kl}\lambda_l{}^{m]}, \\\Omega(F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}) = F_{[\gamma}{}^{\alpha}\lambda_{\alpha\beta]}, \quad \Omega(F_{(\gamma}{}^{\alpha}\lambda_{\alpha\beta)}) = -F_{(\gamma}{}^{\alpha}\lambda_{\alpha\beta)}.$$

So now the invariant subalgebra under the automorphism includes $SO(N) \simeq C^{[ml} \lambda_l^{k]}$ and $U(1) \simeq F_{[\gamma}{}^{\alpha} \lambda_{\alpha\beta]}$. the semi-symmetric space

$$\operatorname{AdS}_{2} \times \operatorname{S}^{2} \times \operatorname{CI}(N)^{N(N+1)} \simeq \frac{\operatorname{OSP}(4^{*}|2N)}{\operatorname{U}(1) \times \operatorname{U}(1) \times \operatorname{U}(N)},$$
(B.41)

note that $CI(1) \simeq S^2$ which is similar to the $D(2, 1; \zeta)$ result.

B.10 OSP $(8^*|2N)$

The relevant commutation relations are

$$\{Q_{l\alpha}, S^k_\beta\} = \delta^k_l (\epsilon_{\alpha\beta} D + M_{\alpha\beta}) + \epsilon_{\alpha\beta} \lambda^k_l, \qquad (B.42)$$

$$\{Q_{l\alpha}, \hat{S}_{k\beta}\} = \epsilon_{\alpha\beta} R_{lk}, \quad \{\hat{Q}^l_{\alpha}, S^k_{\beta}\} = \epsilon_{\alpha\beta} \hat{R}^{lk}, \tag{B.43}$$

 λ_l^k forms U(N) subalgebra of USp(2N), $R_{lk} = R_{kl}$ and $\hat{R}^{lk} = \hat{R}^{kl}$ (l, k = 1, ..., N). $\alpha = 1, ..., 4$ is a spinor index of the R-symmetry spin(6) \simeq SU(4). $M_{\alpha\beta}$ is the 15-dimensional SO(1,5) \simeq SU^{*}(4) Lorentz subalgebra, so $\Sigma M_{\alpha\alpha} = 0$. Since $M_{\alpha\beta}$ is neither symmetric nor antisymmetric we'll not be able to get an AdS space satisfying (2.6). The automorphism transformation is

$$\Omega(Q_{l\alpha}) = iC_{lk}F_{\alpha}{}^{\beta}\hat{S}^{k}_{\beta}, \quad \Omega(\hat{S}^{l}_{\alpha}) = iC^{lk}F_{\alpha}{}^{\beta}Q_{k\beta}.$$
(B.44)

We get $\Omega(D) = (-)^{1+s_c+s_f}D$, so we must have $C_{lk} = (-)^s C_{kl}$ and $F_{\alpha\beta} = (-)^s F_{\beta\alpha}$. If we take s = 1 we must work with even N. The transformations of the R-symmetry are,

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \pm (C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}),$$
(B.45)
$$\Omega(C^{(ml}\lambda_{l}^{k)}) = C^{(ml}\lambda_{l}^{k)}, \quad \Omega(C^{[ml}\lambda_{l}^{k]}) = -C^{[ml}\lambda_{l}^{k]},$$

$$\Omega(F_{(\zeta}{}^{\alpha}M_{\alpha\beta})) = F_{(\zeta}{}^{\alpha}M_{\alpha\beta}), \quad \Omega(F_{[\zeta}{}^{\alpha}M_{\alpha\beta]}) = -F_{[\zeta}{}^{\alpha}M_{\alpha\beta]}.$$

So the invariant subalgebra contains $F_{(\zeta}^{\alpha}M_{\alpha\beta)} \simeq \text{SP}(4)$ and $C^{(ml}\lambda_l^{k)} \simeq \text{USp}(N)$. Thus, the semi-symmetric space induced by the automorphism is

DIII(2,2)₁₂ × CII(N; N)^{4N²} ≃
$$\frac{\text{OSP}(8^*|4N)}{\text{U}(2,2) \times \text{USp}(2N)^2}$$
, (B.46)

note that $CII(1; 1) \simeq S^4$. If we take s = 0 we get

$$\Omega(C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}) = \mp (C^{pl}R_{lk} \pm C_{kl}\hat{R}^{lp}), \qquad (B.47)$$

$$\Omega(C^{(ml}\lambda_l^{\ k)}) = -C^{(ml}\lambda_l^{\ k)}, \quad \Omega(C^{[ml}\lambda_l^{\ k]}) = C^{[ml}\lambda_l^{\ k]}, \qquad (B.47)$$

$$\Omega(F_{(\zeta}^{\ \alpha}M_{\alpha\beta)}) = -F_{(\zeta}^{\ \alpha}M_{\alpha\beta)}, \quad \Omega(F_{[\zeta}^{\ \alpha}M_{\alpha\beta]}) = F_{[\zeta}^{\ \alpha}M_{\alpha\beta]}.$$

So the invariant subalgebra contains $F_{[\zeta}^{\alpha}M_{\alpha\beta]} \simeq \mathrm{SO}(2,1) \times \mathrm{SO}(3)$ and $C^{[ml}\lambda_l^{k]} \simeq \mathrm{SO}(N)$. Thus, the semi-symmetric space

$$BDI(3,1;3,1)_{16} \times CI(N)^{N(N+1)} \simeq \frac{OSP(8^*|2N)}{SO(3,1) \times SO(3,1) \times U(N)},$$
(B.48)

note that $CI(1) \simeq S^2$.

C. Rank of kappa symmetry

In this section we calculate the rank of the kappa symmetry following [14]. We shall also use the same notations for the different type of \mathbb{Z}_4 's. Most of the calculations are the same as in [14] where we just have to modify the dimension of the matrices.

The rank of the kappa-symmetry is $N_{\kappa} + N_{\tilde{\kappa}}$ where

$$N_{\kappa} = \dim \ker \operatorname{ad} K|_{\mathcal{H}_3}, \quad N_{\tilde{\kappa}} = \dim \ker \operatorname{ad} \bar{K}|_{\mathcal{H}_1},$$
 (C.1)

with K and \overline{K} general elements in \mathcal{H}_2 , which will be denoted as

$$K(\text{or }\bar{K}) = \begin{pmatrix} A & 0\\ 0 & B \end{pmatrix}.$$
 (C.2)

We'll compute

$$\left[\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}, \begin{pmatrix} 0 & \Theta \\ \Psi & 0 \end{pmatrix} \right] = \begin{pmatrix} 0 & A\Theta - \Theta B \\ B\Psi - \Psi A & 0 \end{pmatrix}$$
(C.3)

so the commutator vanish if $A\Theta = \Theta B$ and $B\Psi = \Psi A$, and the number of solution to this equation is the rank of the kappa-symmetry. In addition we should impose the Virasoro constraint

$$trA^2 = trB^2. (C.4)$$

We use the notation of [14] for the different types of semi-symmetric spaces. The notation type-U2 and type-U4 refers to the \mathbb{Z}_4 gradings of $\mathrm{SU}(M|N)$ or $\mathrm{PSU}(M|N)$ superalgebras, where the invariant locos is $\mathrm{SO}(M) \times \mathrm{SO}(N)$ and $\mathrm{SP}(M) \times \mathrm{SP}(N)$ respectively, type-O1 and type-O2 refers to $\mathrm{OSP}(M|N)$ with the invariant locus $\mathrm{SO}(2M-P) \times \mathrm{SO}(P) \times \mathrm{U}(N)$ and $\mathrm{U}(M) \times \mathrm{SP}(2N-2P) \times \mathrm{SP}(2P)$ respectively, and type-Tu and type-To to $(\mathrm{P})\mathrm{SU}(M|N)^2/\mathrm{SU}(M) \times \mathrm{SU}(N)$ and $\mathrm{OSP}(M|2N)^2/\mathrm{SO}(M) \times \mathrm{SP}(2N)$ respectively.

C.1 Type-U2

The coset space is

$$\operatorname{AI}(N)_{\frac{(N-1)(N+2)}{2}} \times \operatorname{S}^{1} \times \operatorname{AI}(M)^{\frac{(M-1)(M+2)}{2}} \simeq \frac{\operatorname{SU}(M|N)}{\operatorname{SO}(M) \times \operatorname{SO}(N)}.$$
 (C.5)

The relevant models in our classification are

$$\operatorname{AdS}_{2} \times \operatorname{S}^{1} \times \operatorname{AI}(N)^{\frac{(N-1)(N+2)}{2}} \simeq \frac{\operatorname{SU}(1,1|N)}{\operatorname{SO}(1,1) \times \operatorname{SO}(N)}, \quad N \neq 2,$$

$$\operatorname{AdS}_{2} \times \operatorname{S}^{2} \simeq \frac{\operatorname{PSU}(1,1|2)}{\operatorname{SO}(1,1) \times \operatorname{SO}(2)},$$

$$\operatorname{AI}(2,2)_{9} \times \operatorname{S}^{1} \times \operatorname{AI}(N)^{\frac{(N-1)(N+2)}{2}} \simeq \frac{\operatorname{SU}(2,2|N)}{\operatorname{SO}(2,2) \times \operatorname{SO}(N)}, \quad N \neq 4,$$

$$\operatorname{AI}(2,2)_{9} \times \operatorname{AI}(4)^{9} \simeq \frac{\operatorname{PSU}(2,2|4)}{\operatorname{SO}(2,2) \times \operatorname{SO}(4)}.$$

(C.6)

The calculation of the rank goes the same as in [14], where one have to solve

$$A\Theta = \Theta B, \quad A = A^t, \quad B = B^t, \quad \operatorname{Tr}(A) = \operatorname{Tr}(B) (= 0 \text{ for } \operatorname{PSU's}),$$
(C.7)

$$\operatorname{Tr}(A^2) = \operatorname{Tr}(B^2).$$

There are no solutions in general to this equation and the rank is zero, with the two exceptions:

- PSU(1,1|2) in which there are two solutions, so the rank is four, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 2$ [14]. The coset space is AdS₂ × S².
- SU(1, 1|1) in which there is one solution, so the rank is two, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 1$. The eigenvalues of A and B are $\{0, \alpha\}$ and $\{\alpha\}$ respectively. The coset space is $\operatorname{AdS}_2 \times \operatorname{S}^1$.

In both cases the kappa symmetry rank is half the number of fermionic d.o.f.

C.2 Type-U4

The coset space is

$$\operatorname{AII}(M)_{(M-1)(2M+1)} \times \operatorname{S}^{1} \times \operatorname{AII}(N)^{(N-1)(2N+1)} \simeq \frac{\operatorname{SU}(M|N)}{\operatorname{SP}(M) \times \operatorname{SP}(N)}.$$
 (C.8)

The relevant models in our classification,

$$\operatorname{AdS}_{5} \times \operatorname{S}^{1} \times \operatorname{AII}(N)^{(N-1)(2N+1)} \simeq \frac{\operatorname{SU}(2,2|2N)}{\operatorname{USp}(2,2) \times \operatorname{USp}(2N)}, \quad N \neq 2,$$

$$\operatorname{AdS}_{5} \times \operatorname{S}^{5} \simeq \frac{\operatorname{PSU}(2,2|4)}{\operatorname{USp}(2,2) \times \operatorname{USp}(4)}.$$
(C.9)

The calculation of the rank goes the same as in [14], where one have to solve

$$A\Theta = \Theta B, \quad A = -JA^t J, \quad B = -\tilde{J}B^t \tilde{J}, \quad \operatorname{Tr}(A) = \operatorname{Tr}(B) (= 0 \text{ for PSU's}),$$

$$(C.10)$$

$$\operatorname{Tr}(A^2) = \operatorname{Tr}(B^2).$$

There are no solutions in general to this equation and the rank is zero, with the two exceptions:

- PSU(2,2|4) in which there are eight solutions, so the rank is sixteen, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 8$ [14].
- SU(2, 2|2) in which there are four solution, so the rank is eight, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 4$. The eigenvalues of A and B are $\{0, 0, \alpha, \alpha\}$ and $\{\alpha, \alpha\}$ respectively.

In both cases the kappa symmetry rank is half the number of fermionic d.o.f.

C.3 Type-O1

The relevant coset is

$$BDI(2M-P;P)_{P(2M-P)} \times CI(N)^{N(N+1)} \simeq \frac{OSP(2M|2N)}{SO(2M-P) \times SO(P) \times U(N)}.$$
 (C.11)

We have several models relevant for our classification (both type I and type II),

$$CI(4)_{6} \times S^{1} \simeq \frac{OSP(2|4)}{U(2)},$$

$$AdS_{2} \times BDI(N; N)^{N^{2}} \simeq \frac{OSP(2N|2)}{SO(N)^{2} \times U(1)},$$

$$AdS_{2} \times S^{2} \times CI(N)^{N(N+1)} \simeq \frac{OSP(4^{*}|2N)}{SO(2)^{2} \times U(N)},$$

$$CI(4)_{6} \times BDI(N; N)^{N^{2}} \simeq \frac{OSP(2N|4)}{SO(N)^{2} \times U(2)},$$

$$BDI(3, 1; 3, 1)_{16} \times CI(N)^{N(N+1)} \simeq \frac{OSP(8^{*}|2N)}{SO(3, 1)^{2} \times U(N)}.$$

$$(C.12)$$

The calculation of the rank is very similar to the one in [14]. In all of our cases we have P = M in (C.11), so we'll analyze the rank of these model only. One has the \mathbb{Z}_4 decomposition

$$\mathcal{H}_{2} : A = \begin{pmatrix} 0 & A_{1(M \times M)} \\ -A_{1(M \times M)}^{t} & 0 \end{pmatrix}, \quad B = \begin{pmatrix} B_{1(N \times N)} & B_{2(N \times N)} \\ B_{2(N \times N)} & -B_{1(N \times N)} \end{pmatrix}, \quad B_{i} = B_{i}^{t},$$

$$\mathcal{H}_{1} : \qquad \begin{pmatrix} \Theta_{1(M \times N)} & -i\Theta_{1(M \times N)} \\ \Theta_{2(M \times N)} & i\Theta_{1(M \times N)} \end{pmatrix}, \\
\mathcal{H}_{3} : \qquad \begin{pmatrix} \Theta_{1(M \times N)} & i\Theta_{1(M \times N)} \\ \Theta_{2(M \times N)} & -i\Theta_{1(M \times N)} \end{pmatrix}.$$
(C.13)

The equation $A\Theta = \Theta B$ and the Virasoro constraint comes down to solving

$$-A_1 A_1^t \Theta_1 = \Theta_1 B_+ B_-, \quad B_{\pm} = B_1 \pm i B_2, \quad -2 \operatorname{Tr}(A_1 A_1^t) = 2 \operatorname{Tr}(B_+ B_-). \quad (C.14)$$

In general there are no solutions to this equation and hence the rank of kappasymmetry is zero. There is one exception

• OSP(2|2) - in this case A_1 and B_{\pm} are numbers and there is one solution so the rank of kappa-symmetry is two, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 1$ (this is actually the same result as for the type-U2 SU(1,1|1)). In this case the kappa symmetry rank is half the number of fermionic d.o.f.

C.4 Type-O2

The relevant coset is

$$\mathrm{DIII}(M)_{M(M-1)} \times \mathrm{CII}(N-P;P)^{4P(N-P)} \simeq \frac{\mathrm{OSP}(2M|2N)}{\mathrm{U}(M) \times \mathrm{SP}(2N-2P) \times \mathrm{SP}(2P)}.$$
(C.15)

We have several models relevant for our classification,

$$AdS_{2} \times CII(N; N)^{4N^{2}} \simeq \frac{OSP(4^{*}|4N)}{U(2) \times SP(2N)^{2}},$$

$$AdS_{4} \times DIII(2N)^{2N(2N-1)} \simeq \frac{OSP(4N|4)}{U(2N) \times SP(2)^{2}},$$

$$DIII(2; 2)_{12} \times CII(N; N)^{4N^{2}} \simeq \frac{OSP(8^{*}|4N)}{U(4) \times SP(2N)^{2}}.$$
(C.16)

The calculation of the rank is very similar to the one in [14]. In all of our cases we have N and M-even and P = N/2 in (C.15), so we'll analyze the rank of these model only. One has the \mathbb{Z}_4 decomposition

$$\mathcal{H}_{2} : A = \begin{pmatrix} A_{1(M \times M)} & A_{2(M \times M)} \\ A_{2(M \times M)} & -A_{1(M \times M)} \end{pmatrix}, \quad A_{i} = -A_{i}^{t}, \quad B = \begin{pmatrix} 0_{(N \times N)} & B_{1(N \times N)} \\ JB_{1(N \times N)}^{t} J & 0_{(N \times N)} \end{pmatrix},$$
$$\mathcal{H}_{1} : \qquad \begin{pmatrix} \Theta_{1(M \times N)} & \Theta_{2(M \times N)} \\ -i\Theta_{1(M \times N)} & i\Theta_{2(M \times N)} \end{pmatrix},$$
$$\begin{pmatrix} \Theta_{1(M \times N)} & \Theta_{2(M \times N)} \\ -i\Theta_{1(M \times N)} & i\Theta_{2(M \times N)} \end{pmatrix},$$

$$\mathcal{H}_{3}: \qquad \begin{pmatrix} \Theta_{1(M\times N)} & \Theta_{2(M\times N)} \\ i\Theta_{1(M\times N)} & -i\Theta_{2(M\times N)} \end{pmatrix}.$$
(C.17)

The equation $A\Theta = \Theta B$ and the Virasoro constraint comes down to solving

$$-A_{\pm}A_{\mp}^{t}\Theta_{1} = \Theta_{1}B_{1}JB_{1}^{t}J, \quad \text{Tr}(A_{1}^{2} + A_{2}^{2}) = \text{Tr}(B_{1}JB_{1}^{t}J).$$
(C.18)

In general there are no solutions to this equation and hence the rank of kappasymmetry is zero. There is one exception

• OSP(4|4) - in this case the eigenvalues of $-A_{\pm}A_{\mp}^{t}$ and $B_{1}JB_{1}^{t}J$ are $\{\alpha, \alpha\}$ and $\{\alpha, \alpha\}$, so there are four solutions, so the rank of the kappa-symmetry is eight, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 4$. The backgrounds in this case are $AdS_{4} \times S^{2}$ or $AdS_{2} \times S^{4}$ if we take the real form of the superalgebra. In this case the kappa-symmetry rank is half the number of fermionic d.o.f.

C.5 Type-Tu

The coset space is

$$\frac{\mathrm{SU}(M|N)^2}{\mathrm{SU}(M) \times \mathrm{SU}(N) \times \mathrm{U}(1)}.$$
(C.19)

In our classification we are interested only in AdS_3 models, so we take M = 2 (or more precisely SU(1,1)). The relevant models in our classification are

$$\operatorname{AdS}_{3} \times \operatorname{S}^{1} \times \frac{\operatorname{SU}(N)^{2}}{\operatorname{SU}(N)} \simeq \frac{\operatorname{SU}(1,1|N)^{2}}{\operatorname{SU}(1,1) \times \operatorname{U}(N)}, \quad N \neq 2,$$

$$\operatorname{AdS}_{3} \times \operatorname{S}^{3} \simeq \frac{\operatorname{PSU}(1,1|2)}{\operatorname{SU}(1,1) \times \operatorname{SU}(2)}.$$
(C.20)

According to [14] we have to find the number of solutions to the two equations $A\Theta = \Theta B$ and $B\Psi = \Psi A$, so we need to find out how many eigenvalues of A = U(2) and B = U(M) coincide upon the Virasoro constraint and Tr(A) = Tr(B). In general the eigenvalues are different and we don't have kappa-symmetry. There are two exceptions

• PSU(1,1|2) - in this case the eigenvalues of A and B are $\{\alpha, -\alpha\}$ and $\{\alpha, -\alpha\}$, so there are two solutions for Θ and two for Ψ , so the rank of the kappasymmetry is eight, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 4$ [14]. The background in this case is AdS₃ × S³. SU(1, 1|1) - in this case the eigenvalues of A and B are {α} and {α, 0}, so there is one solution for Θ and one for Ψ, so the rank of the kappa-symmetry is four, namely, N_κ = N_{κ̃} = 2. The background in this case is AdS₃ × S¹.

In both cases the kappa symmetry rank is half the number of fermionic d.o.f.

C.6 Type-To

The coset background is

$$\frac{\operatorname{OSP}(M|2N)^2}{\operatorname{SO}(M) \times \operatorname{SP}(2N)}.$$
(C.21)

In our classification we are interested only in AdS_3 models, so we take N = 1. The relevant models in our classification are

$$\operatorname{AdS}_3 \times \frac{\operatorname{SO}(M)^2}{\operatorname{SO}(M)} \simeq \frac{\operatorname{OSP}(M|2)^2}{\operatorname{SO}(M) \times \operatorname{SP}(2)},$$

According to [14] we have to find the number of solutions to the equation $A\Theta = \Theta B$, so we need to find out how many eigenvalues of A = SO(M) and B = SP(2) coincide upon the Virasoro constraint. In general the eigenvalues are different and we don't have kappa-symmetry. There are three exceptions

- OSP(1|2) in this case the eigenvalues of A and B are $\{0\}$ and $\{0,0\}$, so there are two solutions for Θ , so the rank of the kappa-symmetry is four, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 2$ [14]. The background in this case is AdS₃. In This case the kappa-symmetry rank is equal to the number of fermionic d.o.f.
- OSP(2|2) that is the same case as SU(1,1|1) of $AdS_3 \times S^1$ with kappasymmetry of rank four.
- OSP(3|2) in this case the eigenvalues of A and B are $\{0, \alpha, -\alpha\}$ and $\{\alpha, -\alpha\}$, so there are two solutions for Θ , so the rank of the kappa-symmetry is four, namely, $N_{\kappa} = N_{\tilde{\kappa}} = 2$ [14]. The background in this case is AdS₃ × S³. In This case the kappa-symmetry rank is one third of the number of fermionic d.o.f.

References

- N. Berkovits and J. Maldacena, Fermionic T-Duality, Dual Superconformal Symmetry, and the Amplitude/Wilson Loop Connection, JHEP 09 (2008) 062, [0807.3196].
- [2] N. Beisert, R. Ricci, A. A. Tseytlin, and M. Wolf, *Dual Superconformal Symmetry* from AdS5 x S5 Superstring Integrability, 0807.3228.
- [3] I. Bena, J. Polchinski, and R. Roiban, Hidden symmetries of the AdS(5) x S**5 superstring, Phys. Rev. D69 (2004) 046002, [hep-th/0305116].

- [4] L. Dolan, C. R. Nappi, and E. Witten, A relation between approaches to integrability in superconformal Yang-Mills theory, JHEP 10 (2003) 017, [hep-th/0308089].
- [5] L. Dolan, C. R. Nappi, and E. Witten, Yangian symmetry in D = 4 superconformal Yang-Mills theory, hep-th/0401243.
- [6] I. Adam, A. Dekel, and Y. Oz, On Integrable Backgrounds Self-dual under Fermionic T- duality, JHEP 04 (2009) 120, [0902.3805].
- [7] I. Adam, A. Dekel, and Y. Oz, On the fermionic T-duality of the $AdS_4 \times CP^3$ sigmamodel, 1008.0649.
- [8] C.-g. Hao, B. Chen, and X.-c. Song, On Fermionic T-duality of Sigma modes on AdS backgrounds, JHEP 12 (2009) 051, [0909.5485].
- [9] L. Frappat, P. Sorba, and A. Sciarrino, *Dictionary on Lie superalgebras*, hep-th/9607161.
- [10] W. Nahm, Supersymmetries and their representations, Nucl. Phys. B135 (1978) 149.
- [11] V. G. Kac, Lie Superalgebras, Adv. Math. 26 (1977) 8–96.
- [12] N. Berkovits, M. Bershadsky, T. Hauer, S. Zhukov, and B. Zwiebach, Superstring theory on AdS(2) x S(2) as a coset supermanifold, Nucl. Phys. B567 (2000) 61-86, [hep-th/9907200].
- [13] D. Kagan and C. A. S. Young, Conformal Sigma-Models on Supercoset Targets, Nucl. Phys. B745 (2006) 109–122, [hep-th/0512250].
- [14] K. Zarembo, Strings on Semisymmetric Superspaces, JHEP 05 (2010) 002, [1003.0465].
- [15] V. V. Serganova, Classification of real simple lie superalgebras and symmetric superspaces, Funct. Anal. Appl. 17 (1983) 200–207.
- [16] R. R. Metsaev and A. A. Tseytlin, Type IIB superstring action in AdS(5) x S(5) background, Nucl. Phys. B533 (1998) 109–126, [hep-th/9805028].
- [17] I. Adam, A. Dekel, L. Mazzucato, and Y. Oz, Integrability of type II superstrings on Ramond-Ramond backgrounds in various dimensions, JHEP 06 (2007) 085, [hep-th/0702083].
- [18] N. Beisert, T-Duality, Dual Conformal Symmetry and Integrability for Strings on AdS 5 x S 5, Fortschr. Phys. 57 (2009) 329–337, [0903.0609].
- [19] T. H. Buscher, Path Integral Derivation of Quantum Duality in Nonlinear Sigma Models, Phys. Lett. B201 (1988) 466.
- [20] T. H. Buscher, A Symmetry of the String Background Field Equations, Phys. Lett. B194 (1987) 59.

- [21] I. R. Klebanov and J. M. Maldacena, Superconformal gauge theories and non-critical superstrings, Int.J.Mod.Phys. A19 (2004) 5003-5016, [hep-th/0409133].
- [22] S. Helgason, Differential geometry, Lie groups, and symmetric spaces. American Mathematical Society, 2001.
- [23] A. Babichenko, B. Stefanski, Jr., and K. Zarembo, Integrability and the AdS(3)/CFT(2) correspondence, JHEP 03 (2010) 058, [0912.1723].
- [24] H. Verlinde, Superstrings on ads_2 and superconformal matrix quantum mechanics, 2004.
- [25] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, Dual superconformal symmetry of scattering amplitudes in N=4 super-Yang-Mills theory, Nucl. Phys. B828 (2010) 317–374, [0807.1095].
- [26] T. Bargheer, F. Loebbert, and C. Meneghelli, Symmetries of Tree-level Scattering Amplitudes in N=6 Superconformal Chern-Simons Theory, 1003.6120.
- [27] Y.-t. Huang and A. E. Lipstein, Dual Superconformal Symmetry of N=6 Chern-Simons Theory, 1008.0041.
- [28] S. Lee, Yangian Invariant Scattering Amplitudes in Super-Chern-Simons Theory, Phys. Rev. Lett. 105 (2010) 151603, [arXiv:1007.4772].
- [29] D. Gang, Y.-t. Huang, E. Koh, S. Lee, and A. E. Lipstein, Tree-level Recursion Relation and Dual Superconformal Symmetry of the ABJM Theory, 1012.5032.
- [30] P. A. Grassi, D. Sorokin, and L. Wulff, Simplifying superstring and D-brane actions in AdS(4) x CP(3) superbackground, JHEP 08 (2009) 060, [0903.5407].
- [31] I. Bakhmatov, On $AdS_4 \times CP^3$ T-duality, 1011.0985. * Temporary entry *.
- [32] M. Grigoriev and A. A. Tseytlin, Pohlmeyer reduction of ads_5 x s5 superstring sigma model, Nuclear Physics B 800 (2008) 450.