THE MAPLE PACKAGE FOR SL_2 -INVARIANTS AND KERNEL OF WEITZENBÖCK DERIVATIONS.

LEONID BEDRATYUK

ABSTRACT. We offer a Maple package SL_2_Inv_Ker for calculating of minimal generating sets for the algebras of joint invariants/semi-invariants of binary forms and for calculations of the kernels of Weitzenböck derivations.

1. Introduction

Let $V_d \cong \langle v_0, v_1, \ldots, v_d \rangle$ be d+1-dimensional SL_2 -module of binary forms of degree d and let $V_d = V_{d_1} \oplus V_{d_2} \oplus \cdots \oplus V_{d_n}$, $d = (d_1, d_2, \ldots, d_n)$. Denote by $\mathcal{O}(V_d)^{SL_2}$ the algebra of polynomial SL_2 -invariant functions on V. Denote by $\mathcal{O}(V_d)^{U_2}$ the algebra of polynomial U_2 -invariant functions on V, where U_2 is the maximal unipotent subgroup of SL_2 . We have the obvious inclusion $\mathcal{O}(V_d)^{SL_2} \subset \mathcal{O}(V_d)^{U_2}$. Moreover $\mathcal{O}(V_d)^{U_2} \cong \mathcal{O}(V_1 \oplus V_d)^{SL_2}$. The elements of the finitely generated algebras can be identified with the algebras of joint invariants and joint semi-invariants of the binary forms of degrees d.

A linear derivation \mathcal{D} of a polynomial algebra is called Weitzenböck derivation if its matrix (as linear map on the vector space generated by variables of the polynomial algebra) is nilpotent. Any Weitzenböck derivation \mathcal{D} is completely determined by the Jordan normal form of its matrix. We will denote by \mathcal{D}_d , $d := (d_1, d_2, \ldots, d_n)$ the Weitzenbök derivation if its matrix consisting of n Jordan blocks of size $d_1 + 1$, $d_2 + 1$, \ldots , $d_n + 1$, respectively.

As is well known, the kernel of the derivations \mathcal{D}_d , $d := (d_1, d_2, \dots, d_n)$ is isomorphic to the algebra of joint semi-invariants (and to the algebra of covariants) for n binary forms of degrees d_1, d_2, \dots, d_n . Thus, the calculations of a minimal generated sets of the algebras $\mathcal{O}(V_d)^{SL_2}$ or $\mathcal{O}(V_d)^{U_2}$ can be reduced to calculation of minimal generating set of the kernel of suitable Weitzenböck derivation. For instance, the algebra $\mathcal{O}(V_d)^{U_2}$ can be identified with ker \mathcal{D} , where the derivation \mathcal{D} is defined by $\mathcal{D}(v_i) = iv_{i-1}$, $i = 0, \dots, d$.

2. Algorithm

The algebra $\mathcal{O}(V_d)^{U_2}$ is a finitely generated multigraded algebra under the multidegree-order:

$$\mathcal{O}(V_d)^{U_2} = (\mathcal{O}(V_d)^{U_2})_{m,0} + (\mathcal{O}(V_d)^{U_2})_{m,1} + \dots + (\mathcal{O}(V_d)^{U_2})_{m,j} + \dots,$$

where each subspace $(\mathcal{O}(V_d)^{U_2})_{d,j}$ of joint semi-invariants of order j and multidegree $m := (m_1, m_2, \ldots, m_n)$ is finite-dimensional. The formal power series

$$\mathcal{P}(\mathcal{O}(V_d)^{U_2}, z_1, z_2, \dots, z_n, t) = \sum_{m, j=0}^{\infty} \dim((\mathcal{O}(V_d)^{U_2})_{m,j}) z_1^{m_1} z_2^{m_2} \cdots z_n^{m_n} t^j,$$

is called the multivariariate Poincaré series of the algebra of join semi-variants C_d . Note that each semi-invariant of order zero is an invariant.

Suppose that $\mathcal{P}(\mathcal{O}(V_d)^{SL_2}, t) = \frac{P(t)}{Q(t)}$ and P(t), Q(t) are coprime polynomials. Denote by β_d the degree of the denominator Q(t). Many experimental data leads to the following conjecture:

Conjecture. A degree upper bound for irreducible invariants of the algebra $\mathcal{O}(V_d)^{SL_2}$ does not exceed β_d .

In the first version of the package we used the following algorithm:

- 1. Calculate multivariate Poincaré series of the algebras $\mathcal{O}(V_d)^{SL_2}$ (or $\mathcal{O}(V_d)^{SL_2}$ or $\ker \mathcal{D}_m$.)
- 2. For every term $z_1^{m_1} z_2^{m_2} \cdots z_n^{m_n} t^j$ of the Poincaré series calculate (by linear algebra method) a basis of the vector space of semi-invariants (or invariants or elements of kernel) and the multidegree (m_1, m_2, \ldots, m_n) .
 - 3. Separate irreducible polynomials.
 - 4. Stop calculation if $m_1 + m_2 + \ldots + m_n > 18$.

For the package procedures Min_Gen_Set_Invariants_S we use also the following simplified algorithm:

- 1. Calculate Poincaré series of the algebras $\mathcal{O}(V_d)^{SL_2}$.
- 2. For every term z^m of the Poincaré series calculate a basis of the vector space of invariants of the degree m.
 - 3. Separate irreducible polynomials.
 - 4. Stop calculation if m > 18.

The second algorithm works some fast for small values d_i and n > 4.

The package calculate the set of irreducible invariants up to degree $\min(18, \beta_d)$, but in all known computable cases this set coincides with a minimal generating set, see, for example, Brouwer's webpage http://www.win.tue.nl/~aeb/math/invar/invarm.html

3. Installation.

The package can be downloaded from the web http://sites.google.com/site/bedratyuklp/.

- (1) download the file SL_2_Inv_Ker.mpl and save it into your Maple directory.
- (2) download the Xin's file (see link at the web page) Ell2.mpl and save it into your Maple directory.
- (3) run Maple
- (4) > restart: read "SL_2_Inv_Ker.mpl":read "Ell2.mpl":
- (5) If necessary use > HelP();

4. Package Procedures and Syntax

Procedure name: Min_Gen_Set_Semi_Invariants

Feature: Computes irreducible invariants of the algebra $\mathcal{O}(V_d)^{U_2}$ up to degree $\min(18, \beta_d)$.

Calling sequence: $Min_Gen_Set_Semi_Invariants([d_1, d_2, \dots, d_n]);$ Parameters:

 $[d_1, d_2, \dots, d_n]$ - a list of degrees of n binary forms. n - an integer, n < 11.

Procedure name: Min_Gen_Set_Invariants

Feature: Computes irreducible invariants of the algebra $\mathcal{O}(V_d)^{SL_2}$ up to degree

```
THE MAPLE PACKAGE FOR SL_2-INVARIANTS AND KERNEL OF WEITZENBÖCK DERIVATIONS
```

```
\min(18, \beta_d).
Calling sequence: Min\_Gen\_Set\_Invariants([d_1, d_2, ..., d_n]);
Parameters:
 [d_1, d_2, \ldots, d_n] - a list of degrees of n binary forms.
                - an integer, n < 11.
Procedure name: Kernel_LLN_Der
Feature: Computes irreducible elements of the kernel of Weitzenböck derivation \mathcal{D}_d up
to degree min(18, \beta_d).
Calling sequence: Kernel_LLN_Der([d_1, d_2, \dots, d_n]);
Parameters:
 [d_1, d_2, \ldots, d_n] - a list of degrees of n binary forms.
         - an integer, n < 11.
Procedure name: Min_Gen_Set_Invariants_S
Feature: By using the second algorithm the procedure computes a set of irreducible
invariants of the algebra \mathcal{O}(V_d)^{SL_2} up to degree min(18, \beta_d).
Calling sequence: Min_Gen_Set_Invariants_S([d_1, d_2, \dots, d_n]);
Parameters:
[d_1, d_2, \ldots, d_n] - a list of degrees of n binary forms.
   - an integer, n < 11.
                                   5. Examples
5.1. Compute \mathcal{O}(V_4)^{SL_2}. Use the command
  > dd:=[4]:INV:=Min_Gen_Set_Invariants(dd):
                     "calculating multivariate Poincare series...."
  "done!, upper bound", 13
  " irreducible invariant of multidegree ", [2], "found"
        " irreducible invariant of multidegree ", [3], "found"
         -----degree-
                    -degree-
                    -degree-
                    -degree-
                    -degree-
                    -degree-
                    -degree-
```

"Total number of irreducible invariants in minimal generating set ", 2

-degree—

-degree— -degree—

To extract the two invariants from the set INV one should to use the commands

-----, 12 ------, 13 >INV[2];

$$\{6x_2^2 + 2x_0x_4 - 8x_1x_3\}$$

>INV[3];

$$\left\{-6 x_2^3 - 6 x_0 x_3^2 - 6 x_1^2 x_4 + 6 x_0 x_2 x_4 + 12 x_1 x_2 x_3\right\}$$

5.2. Compute $\mathcal{O}(V_3 \oplus V_4)^{SL_2}$. Use the command

```
> dd:=[3,4]:INV:=Min_Gen_Set_Invariants(dd):
                 "calculating multivariate Poincare series...."
                          "done!, upper bound", 13
                       " irreducible invariant of multidegree ", [0, 2], "found"
                "------degree-----", 3
            " irreducible invariant of multidegree ", [0, 3], "found"
                "-----degree-----", 4
             " irreducible invariant of multidegree", [4, 0], "found"
                "_____degree______". 5
            " irreducible invariant of multidegree ", [4, 1], "found"
             " irreducible invariant of multidegree", [2, 3], "found"
                "------degree-----", 6
             " irreducible invariant of multidegree", [4, 2], "found"
             " irreducible invariant of multidegree", [4, 2], "found"
                " irreducible invariant of multidegree", [4, 3], "found"
             " irreducible invariant of multidegree", [4, 3], "found"
             " irreducible invariant of multidegree", [4, 3], "found"
                                -degree--
            " irreducible invariant of multidegree ", [6, 2], "found"
            " irreducible invariant of multidegree", [4, 4], "found" "irreducible invariant of multidegree", [4, 4], "found"
                    -----degree----". 9
             " irreducible invariant of multidegree", [6, 3], "found"
             " irreducible invariant of multidegree", [6, 3], "found"
             " irreducible invariant of multidegree", [6, 3], "found"
             " irreducible invariant of multidegree", [4, 5], "found"
                                -degree---
             "irreducible invariant of multidegree", [6, 4], "found"
             " irreducible invariant of multidegree", [6, 4], "found"
                               --degree----
             " irreducible invariant of multidegree", [6, 5], "found"
               degree ", 12
                               --degree-
                                                      -". 13
```

[&]quot;Total number of invariants in a minimal generating set ", 20

THE MAPLE PACKAGE FOR SL_2 -INVARIANTS AND KERNEL OF WEITZENBÖCK DERIVATIONS

To get the number of invariants of degree i use the command nops(INV[i]), for instance

```
> nops(INV[5]);
```

2

To extract and manipulate by these two invariants use the following commands

> Inv5_1:=INV[5][1]:Inv5_2:=INV[5][2];

$$Inv5_2 := 6x_1^2x_3^2y_0 + 18x_1^2x_2^2y_2 - 12x_1^3x_3y_2 - 12x_1^3x_2y_3 + 6x_0^2x_2^2y_4 + 6x_2^4y_0 + 6x_1^4y_4 + 12x_1^2x_2x_3y_1 - 12x_0x_1x_3^2y_1 + 12x_0x_2^2x_3y_1 + 12x_0x_1^2x_3y_3 - 12x_0x_1^2x_2y_4 - 12x_0^2x_2x_3y_3 - 12x_1x_2^2x_3y_0 + 12x_0x_1x_2^2y_3 - 12x_0x_2^3y_2 + 6x_0^2x_3^2y_2 - 12x_1x_2^3y_1$$

5.3. Compute $\mathcal{O}(V_1 \oplus V_1 \oplus V_2)^{U_2}$. Use the command

> dd:=[1,1,2]:COV:=Min_Gen_Set_Semi_Invariants(dd):

"calculating multivariate Poincare series...."

"done!, upper bound", 13

"_____degree_____", 2

"irreducible semi-invariant of multidegree", [0, 0, 2], "and order ", 0, "found" "irreducible semi-invariant of multidegree", [1, 1, 0], "and order ", 0, "found"

"irreducible semi-invariant of multidegree", [0, 1, 1], "and order ", 1, "found"

"irreducible semi-invariant of multidegree", [2, 0, 1], "and order ", 0, "found" "irreducible semi-invariant of multidegree", [1, 1, 1], "and order ", 0, "found"

"irreducible semi-invariant of multidegree", [0, 2, 1], "and order ", 0, "found"

"	degree	, 4
11	degree	", 5
11	degree	", 6
11	degree	
11	degree	, 8
!!	degree	
II	degree	", 10
II	degree	", 11
II	degree	", 12
!!	degree	" 13

"number of semi-invariant of minimal generating set", 10

Below is this minimal generating set listed by degree

> COV[1]; nops(%);

$$\{x_0, y_0, u_0\}$$

> COV[2];nops(%);

$$\{-x_0y_1 + x_1y_0, -y_1u_0 + y_0u_1, -x_1u_0 + x_0u_1, 2u_0u_2 - 2u_1^2\}$$

4

> COV[3]; nops(%);


```
\{-2y_0y_1u_1 + y_1^2u_0 + y_0^2u_2, -x_1y_1u_0 + x_1y_0u_1 + x_0y_1u_1 - x_0y_0u_2, 
                                                                          x_1^2u_0 + x_0^2u_2 - 2x_0x_1u_1
```

3

command

```
5.4. Compute the kernel of the Weitzenböck derivation \mathcal{D}_d, d = (1,3). Use the
  > dd:=[1,3]:Ker:=Kernel_LLN_Der(dd):
                   "calculating multivariate Poincare series...."
                           "done!, upper bound", 13
                       "irreducible element of multidegree", [1, 1], "and order ", 2, "found"
        "irreducible element of multidegree", [0, 2], "and order ", 2, "found"
                  "irreducible element of multidegree", [2, 1], "and order ", 1, "found"
        "irreducible element of multidegree", [0, 3], "and order ", 3, "found"
        "irreducible element of multidegree", [1, 2], "and order ", 1, "found"
                        -----degree----". 4
        "irreducible element of multidegree", [2, 2], "and order ", 0, "found"
        "irreducible element of multidegree", [3, 1], "and order ", 0, "found"
        "irreducible element of multidegree", [0, 4], "and order ", 0, "found"
        "irreducible element of multidegree", [1, 3], "and order ", 2, "found"
                        "irreducible element of multidegree", [2, 3], "and order ", 1, "found"
                 "irreducible element of multidegree", [3, 3], "and order ", 0, "found"
                      -----degree-
                   -----degree-
                                  -degree-
                                 -degree-
                                 -degree-
                                                   -----, 12
                                 -degree-
                                          ------, 13
                                 -degree-
             "number of semi-invariant in minimal generating set", 13
  > Ker[3];
 \{-6y_0^2y_3+6y_0y_1y_2-2y_1^3, 4x_1y_0y_2-6x_0y_0y_3+2x_0y_1y_2-2x_1y_1^2, -2x_0^2y_2-x_1^2y_0+2x_0x_1y_1\}
5.5. Compute \mathcal{O}(V_1 \oplus V_1 \oplus V_1 \oplus V_2 \oplus V_2)^{SL_2}. Use the command
  > dd:=[1,1,1,2,2]:Inv:=Min_Gen_Set_Invariants_S(dd):
                         "calculating Poincare series...."
                            "done!, upper bound", 6
```

THE MAPLE PACKAGE FOR SL_2 -INVARIANTS AND KERNEL OF WEITZENBÖCK DERIVATIONS7



> Ker[3];

$$\{3\,y_0^2v_2 - 6\,y_0y_1v_1 + 3\,y_1^2v_0, 3\,y_0^2w_2 + 3\,y_1^2w_0 - 6\,y_0y_1w_1, 6\,u_0u_1w_1 - 3\,u_0^2w_2 - 3\,u_1^2w_0, \\ -6\,y_0u_0v_2 - 6\,y_1u_1v_0 + 6\,y_1u_0v_1 + 6\,y_0u_1v_1, -6\,y_0u_0w_2 - 6\,y_1u_1w_0 + 6\,y_0u_1w_1 + 6\,y_1u_0w_1, \\ 6\,u_0u_1v_1 - 3\,u_1^2v_0 - 3\,u_0^2v_2, 6\,u_1x_0v_1 + 6\,u_0x_1v_1 - 6\,u_0x_0v_2 - 6\,u_1x_1v_0, \\ 6\,y_1x_0w_1 - 6\,y_1x_1w_0 - 6\,y_0x_0w_2 + 6\,y_0x_1w_1, -6\,u_1x_1w_0 - 6\,u_0x_0w_2 + 6\,u_0x_1w_1 + 6\,u_1x_0w_1, \\ 6\,y_0x_1v_1 - 6\,y_1x_1v_0 + 6\,y_1x_0v_1 - 6\,y_0x_0v_2, -3\,x_1^2v_0 + 6\,x_0x_1v_1 - 3\,x_0^2v_2, -3\,x_0^2w_2 + 6\,x_0x_1w_1 - 3\,x_1^2w_0 \}$$