

Fundamental groups of locally connected subsets of the plane

Greg Conner and Curt Kent

Abstract

We show that every homomorphism from a one-dimensional Peano continuum to a planar Peano continuum is induced by a continuous map up to conjugation. We then prove that the topological structure of the space of points at which a planar Peano continuum is not semi-locally simply connected is determined solely by the algebraic structure of its fundamental group. Furthermore, we demonstrate how to reconstruct the topological structure of the space of points at which a planar Peano continuum is not semi-locally simply connected using only the subgroup lattice of its fundamental group.

Introduction

The Hawaiian earring is the one-point compactification of countably many disjoint open arcs. Its fundamental group, which we will denote by \mathbb{H} , is called the Hawaiian earring group.

The fundamental groups of planar and one-dimensional Peano continua have been studied extensively. In [Hig], G. Higman studies the inverse limit of finite rank free groups which he calls the unrestricted free product of countably many copies of \mathbb{Z} . There he proves that this group is not free and that each of its free quotients has finite rank. He considers a subgroup P of the unrestricted product which turns out to be a Hawaiian earring group but he does not prove it there. In [CF], Curtis and Fort show that the Menger curve has fundamental group which embeds in the inverse limit of finite rank free groups. As well in [MM], Morgan and Morrison study the Hawaiian earring group. Cannon and Conner in [CC1] and [CC2] studied the fundamental groups of one-dimensional spaces. In [ADTW], Akiyama, Dorfer, Thuswaldner, and Winkler studied the fundamental group of the Sierpinski-Gasket.

Katsuya Eda [Eda2] was the first to prove results concerning homomorphisms between the fundamental groups of one-dimensional Peano continua. He proved that every homomorphism from \mathbb{H} to \mathbb{H} is conjugate to a homomorphism which is induced by a continuous map. Summers gave a combinatorial proof of the same result in [CS]. In [Eda], Eda was able to extend his original result to show that a homomorphism from \mathbb{H} to the fundamental group of a one-dimensional metric space is conjugate to a homomorphism which is induced by a continuous map.

In Section 2, we prove that homomorphisms from \mathbb{H} to the fundamental group of a planar Peano continuum are conjugate to homomorphisms induced by continuous maps. This is then used to prove the following result.

Theorem 3.5 *Let X be a one-dimensional Peano continuum and Y a planar Peano continuum. Then for every homomorphism $\varphi : \pi_1(X, x) \rightarrow \pi_1(Y, y_0)$ then there exists a path*

$\alpha : (I, 0, 1) \rightarrow (Y, y_0, y)$ and a continuous function $f : X \rightarrow Y$ such that $\hat{\alpha} \circ \varphi = f_*$.

Eda ^{eda3}[E3] proved the analogue of Theorem ^{cont1dim}3.5 when both X and Y are one-dimensional Peano continua. In Section ^{sec2}3, we will show the necessary modifications of his proof to be able to extend it to the case when Y is planar. The second author has also shown that the theorem holds when X is allowed to be a planar Peano continuum and Y is either a planar or one-dimensional Peano continuum ^{ck1}[K1].

In ^{eda}[E2], Eda showed that the fundamental group of a one-dimensional continuum which is not semilocally simply connected at any point determines the homeomorphism type of the space. Eda and Conner show that the fundamental group of a one-dimensional continuum which is not semilocally simply connected at any point can be used to reconstruct the original space with its topology ^{ce}[CE]. We prove the following analogue for planar Peano continua.

Theorem ^{bad}4.17 *If X is a planar Peano continuum, the homeomorphism type of $B(X)$ is completely determined by $\pi_1(X, x_0)$, where $B(X)$ is the subspace of points at which X is not semilocally simply connected.*

The proof that we present in Section ^{sec3}4 is for a planar Peano continuum; however, with a few insignificant changes our proof works for both planar and one-dimensional Peano continua. Our proof shows that what is really necessary for a results of this type isn't the one-dimensionality of the spaces involved but the fact that essential loops are cannot be homotoped to much (see Lemma ^{homotoped off}A6 for an example of what this means in the planar case). Thus types of results are possible for any class of spaces where essential loops have this property.

1 Definitions

A Peano continuum is a compact, connected, locally path connected, metric space. For a metric space (X, d) , let $B_\epsilon^X(x_0) = \{x \in X \mid d(x, x_0) < \epsilon\}$ be the open ϵ -ball about x_0 and $S_\epsilon^X(x_0) = \{x \in X \mid d(x, x_0) = \epsilon\}$ be the ϵ -sphere about x_0 . When the underling space is understood, we will simply write $B_\epsilon(x_0)$ or $S_\epsilon(x_0)$.

The one-point compactification of a sequence of disjoint arcs can be realized in the plane as the union of circles centered at $(0, \frac{1}{n})$ with radius $\frac{1}{n}$. We will use \mathbf{E} to denote this subspace of the plane and \mathbf{a}_n to denote the circle centered at $(0, \frac{1}{n})$ with radius $\frac{1}{n}$. Then $\pi_1(\mathbf{E}, (0, 0)) = \mathbb{H}$. At times it will be convenient to consider certain subspaces of \mathbf{E} and certain subgroups of \mathbb{H} . We will let $\mathbf{E}_n = \bigcup_{i \geq n} \mathbf{a}_i$, $\mathbf{E}_e = \bigcup_{i \text{ even}} \mathbf{a}_i$ and $\mathbf{E}_o = \bigcup_{i \text{ odd}} \mathbf{a}_i$. Then let \mathbb{H}_n , \mathbb{H}_e , and \mathbb{H}_o be their respective fundamental groups which are all isomorphic to \mathbb{H} .

Cannon and Conner have shown that \mathbb{H} is generated in the sense of infinite products by a countable set of generators corresponding to the circles \mathbf{a}_n , where an infinite product is legal if each \mathbf{a}_n is transversed only finitely many times. For more details see ^{cc1}[CC1], ^{cc3}[CC2]. We will refer to this infinite generating set for the fundamental group of \mathbf{E} as $\{a_n\}$, i.e. a_n represents the canonical path which transverses counterclockwise \mathbf{a}_n one time. We will frequently denote the base point $(0, 0)$ of \mathbf{E} by just 0.

Let $\{\mathbf{b}_n\}$ be a null sequence of loops based at x_0 in some metric space X . Then any function $f : (\mathbf{E}, 0) \rightarrow (X, x)$ which sends \mathbf{a}_n continuously to \mathbf{b}_n and sends 0 to x is continuous. Often when defining functions from \mathbf{E} to X , we will only be interested in the property that $f_*(a_n) = [\mathbf{b}_n]$. Therefore to simplify the construction of such functions; when defining a function f we will only state that f maps \mathbf{a}_n to \mathbf{b}_n (for every n) meaning that f maps \mathbf{a}_n continuously to \mathbf{b}_n and sends 0 to the base point of \mathbf{b}_n such that $f_*(a_n) = [\mathbf{b}_n]$.

We will use $[g]$ to represent the homotopy class relative to endpoints of the path g . Then a continuous function f induces an homomorphism on fundamental groups which we will denote by f_* . A loop f is a *reparametrization* of g if there exists continuous maps $\theta_1, \theta_2 : [0, 1] \rightarrow [0, 1]$ such that θ_i is nondecreasing, $\theta_i(0) = 0$ and $\theta_i(1) = 1$ for $i = 1, 2$, and $f \circ \theta_1 = g \circ \theta_2$. This is an equivalence relation and we will write $f =_p g$.

Suppose that $f_i : I \rightarrow X$ is a sequence of paths such that $f_i(1) = f_{i+1}(0)$. Then $f_1 * f_2$ will denote the standard concatenation of paths and $\prod_i f_i$ the infinite concatenation when it is defined. We use $\overline{f}_1(t)$ to denote the path $\overline{f}_1(t) = f_1(1-t)$. For a path $\alpha : (I, 0, 1) \rightarrow (X, x_0, x_1)$, let $\widehat{\alpha} : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ by the standard change of base point isomorphism, i.e. $\widehat{\alpha}([g]) = [\overline{\alpha} * g * \alpha]$. This isomorphism has inverse $\widehat{\alpha}$.

Definition 1.1. Let X be a one-dimensional space. We say that a path $f : I \rightarrow X$ is *reducible* if there is an open arc $(x, y) \subset I$ such that $f|_{[x, y]}$ is a non-degenerate null-homotopic loop. A loop f is *reduced* if it is not reducible. A constant loop is, by definition reduced and every reparametrization of a reduced path is reduced.

reducedloop-thm

Theorem 1.2 (Theorem 3.9 ^{CC3} [CC2]). *Let $f : I \rightarrow X$ be a continuous map into a one-dimensional space. There exist a maximal (under set inclusion) set of open intervals $\{(s_i, t_i)\}_{i \in J}$ with disjoint closures such that, for each $i \in J$, $f|_{[s_i, t_i]}$ is a null-homotopic subloop of f .*

Define $f_r : I \rightarrow X$ by $f_r(t) = f(t)$ for $t \notin \bigcup_{i \in J} (s_i, t_i)$ and $f_r(t) = f(s_i)$ for $t \in (s_i, t_i)$. Then f_r is the unique (up to reparametrization) reduced loop which is homotopic to f .

We will use f_r to denote the path where every null-homotopic subloop of f is replaced by a constant path, in which case we will call f_r a *reduced representative* of f or $[f]$. When considering one-dimensional spaces, we will use $[\cdot]_r$ to denote a reduced representative of the path class of $[\cdot]$.

Definition 1.3. Let X be a one-dimensional space. Let $g : I \rightarrow X$ be a reduced representative for the path class $[g]$. Then we say that $a : I \rightarrow X$ is a *head* for g if there exists $b : I \rightarrow X$ such that $g =_p a * b$. We will write $a \xrightarrow{h} g$. Additionally, we say that $b : I \rightarrow X$ is a *tail* for g if there exist $c : I \rightarrow X$ such that $g =_p c * \overline{b}$. We will write $b \xrightarrow{t} g$.

We say that $t : I \rightarrow X$ is a *head-tail* for a reduced path $g : I \rightarrow X$ if there exists $b : I \rightarrow X$ such that $g =_p t * g * \overline{t}$, i.e. t is a head and a tail for g . We will write $t \xrightarrow{h-t} g$.

Since g is a reduced path; the paths a , b , and c are necessarily reduced paths.

2 Homomorphisms from \mathbb{H}

sec1

We will use the following theorem for one-dimensional spaces.

hto1d

Theorem 2.1 (Eda [E2]). Let $\varphi : \mathbb{H} \rightarrow \pi_1(X, x_0)$ a homomorphism into the fundamental group of a one-dimensional Peano continuum X . Then there exists a continuous function $f : (\mathbf{E}, 0) \rightarrow (X, x_1)$ and a path $T : (I, 0, 1) \rightarrow (X, x_0, x_1)$, with the property that $f_* = \widehat{T} \circ \varphi$. Additionally; if the image of φ is uncountable, then T is unique up to homotopy rel endpoints.

2.1 Delineation

delin

To prove Theorem ^{contplanar}2.22, we will use an upper semicontinuous decomposition of the planar Peano continuum to get a continuous map into a one-dimensional Peano continuum which is injective on fundamental groups. This allows us to, in some degree, reduce the planar case to the one-dimensional case. If π_k is this decomposition map, we show how to lift the path T such that $\widehat{T} \circ \pi_{k*} \circ \varphi$ is induced by a continuous map. Then for α a lift of T , we show that $\widehat{\alpha} \circ \varphi$ is induced by a continuous function.

Definition 2.2. Let k be a line in the plane and X a planar set. Let $\pi_k : X \rightarrow X/G$ be a decomposition map where the nontrivial decomposition elements of G are the maximal line segments contained in X which are parallel to k . We will use X_k to denote the decomposition space corresponding to π_k and refer to π_k as a *delineation map*.

The following is Theorem 1.4 in [cc4].

Theorem 2.3 (Cannon & Conner). If X is a planar set and k is any line in the plane; then π_k is upper semicontinuous and X_k is a one-dimensional Peano continuum. In addition; if X is a planar Peano continuum, then the induced homomorphism of fundamental groups is injective.

Let $h : C \rightarrow D$ be a function. Then a subset C' of C is *h-saturated* if $C' = h^{-1}(D')$ for some $D' \subset D$.

continuous maps

Lemma 2.4. Let X be a planar set and $\pi_k : X \rightarrow X_k, \pi_l : X \rightarrow X_l$ be delineation maps for non-parallel lines k, l respectively.

If $f : Z \rightarrow X$ is a map from a first countable space Z such that $\pi_k \circ f$ and $\pi_l \circ f$ are continuous, then f is continuous.

Proof. Let k, l be non-parallel lines in the plane with slopes m_k, m_l respectively. Let U be an open parallelogram in the plane with sides parallel to k and l . Chose $c_{k,1} < c_{k,2}$ and $c_{l,1} < c_{l,2}$ such that $U = U_k \cap U_l$ where $U_k = \{(x, y) \mid m_k x + c_{k,1} \leq y \leq m_k x + c_{k,2}\}$ and $U_l = \{(x, y) \mid m_l x + c_{l,1} \leq y \leq m_l x + c_{l,2}\}$.

Since $U_k \cap X$ is π_k -saturated and open in X , $\pi_k(U_k \cap X)$ is open in X_k . Similarly, $\pi_l(U_l \cap X)$ is open in X_l . Then

$$\begin{aligned} f^{-1}(U \cap X) &= f^{-1}(U_k \cap X) \cap f^{-1}(U_l \cap X) \\ &= (\pi_k \circ f)^{-1}(\pi_k(U \cap X)) \cap (\pi_l \circ f)^{-1}(\pi_l(U \cap X)) \end{aligned}$$

which is open since $\pi_k \circ f$ and $\pi_l \circ f$ are continuous.

This completes the proof since such sets form a basis for the topology of X . \square

maps-corollary

Corollary 2.5. *Let X be a planar set and $\pi_k : X \rightarrow X_k$, $\pi_l : X \rightarrow X_l$ be delineation maps for non-parallel lines k, l respectively.*

A sequence x_n of points in X converges to x_0 if and only if $\pi_k(x_n)$ and $\pi_l(x_n)$ converge to $\pi_k(x_0)$ and $\pi_l(x_0)$, respectively.

Proof. The one direction is trivial since delineation maps are continuous. Suppose that $\pi_k(x_n)$ and $\pi_l(x_n)$ converge to $\pi_k(x_0)$ and $\pi_l(x_0)$, respectively. Let $Z = \{0\} \cup \{\frac{1}{n}\}$ and $f : Z \rightarrow X$ such that $f(\frac{1}{i}) = x_i$ and $f(0) = x_0$. Then $\pi_k \circ f$ and $\pi_l \circ f$ are continuous. Hence by f is continuous and x_n converges to x_0 . \square

kreduced

Lemma 2.6. *Let X be a planar Peano continuum. If $g : I \rightarrow X$ is a path and $\pi_k \circ g$ has reduced representative f then there exists $\tilde{g} : I \rightarrow X$ such that $\pi_k \circ \tilde{g} =_p f$ and g is homotopic rel-endpoints to \tilde{g} .*

Proof. If $\pi_k \circ g$ is reduced, we are done. Otherwise there exists an interval $[c, d]$ such that $\pi_k \circ g|_{[c, d]}$ is a non-constant null-homotopic subloop. Then $\pi_k \circ g(c) = \pi_k \circ g(d)$ which implies that the line segment $\overline{g(c)g(d)}$ is contained in X . The loop $g|_{[c, d]} * \overline{g(d)g(c)}$ maps to $\pi_k \circ g|_{[c, d]}$ and hence must be null-homotopic since π_{k*} is injective. Therefore g is homotopic to g' where the subpath $g|_{[c, d]}$ is replaced by the line segment $\overline{g(c)g(d)}$.

Suppose that $\{J_i = (c_i, d_i)\}$ is a maximal set of disjoint open subintervals of I such that $\pi_k \circ g|_{J_i}$ is a non-constant null-homotopic loop. Let l_i be a parametrization of the line segment from $g(c_i)$ to $g(d_i)$. Let \tilde{g} be the path g where each subpath $g|_{[c_i, d_i]}$ is replaced by l_i . Then \tilde{g} defines a continuous path such that $g(t) = \tilde{g}(t)$ for $t \notin \cup_i J_i$.

We need to show that g is homotopic to \tilde{g} . Since g is uniformly continuous, $\text{diam}(g|_{[c_i, d_i]})$ must converge to zero.

By Lemma ^{cutoff}A7, there exists homotopies $H_i : I \times [c_i, d_i] \rightarrow X$ with the property that $H_i|_{\{0\} \times [c_i, d_i]} = g|_{[c_i, d_i]}$, $H_i|_{\{1\} \times [c_i, d_i]} = l_i$, and $\text{diam} H_i(I \times [c_i, d_i]) \rightarrow 0$.

Lemma ^{lemma}A3 implies that g is homotopic to \tilde{g} . By our choice of $\{J_i\}$, $\pi_k \circ \tilde{g}$ is reduced which completes the proof. \square

Definition 2.7. If g maps to a reduced path under π_k , we will say that g is reduced with respect to k or g is k -reduced. For any path g ; if \tilde{g} is obtained from g as in Lemma ^{kreduced}2.6 then we will say \tilde{g} is obtained by reducing g with respect to k .

Lemma 2.8. *If $g : S^1 \rightarrow X$ is a loop and $\pi_k \circ g$ has cyclically reduced representative f then there exists $\tilde{g} : S^1 \rightarrow X$ such that $\pi_k \circ \tilde{g} =_p f$ and g is freely homotopic to \tilde{g} .*

Proof. The proof is the same as in Lemma ^k2.6, except we choose $\{J_i = (c_i, d_i)\}$ to be a maximal set of disjoint open subsets of S^1 which are homeomorphic to open intervals such that $\pi_k \circ g|_{J_i}$ is a non-constant null-homotopic loop. \square

2.2 Weight Function

Before we proceed, we need to introduce a weight function. This weight function is a discrete version of the oscillation function defined by Cannon, Conner, and Zastrow in [CCZ]. The discrete version was also used in [K], since it is preserved under nerve approximations.

2.2.1 Weight function

Let X be a normal topological space. For a path $f : I \rightarrow X$ and U, V disjoint open subsets of X , let $r_f : f^{-1}(U \cup V) \rightarrow \{-1, 1\}$ by $r_f(s) = 1$ if $f(s) \in U$ and $r_f(s) = -1$ if $f(s) \in V$. Let $\bar{w}_U^V(f) = \sup \left(- \sum_i r_f(s_i) \cdot r_f(s_{i+1}) \right)$ taken over all increasing countable subsets of $f^{-1}(U \cup V)$. For any collection consisting of 0 or 1 point, we will consider the sum to be 0.

If the image of two consecutive points in our countable subset of $f^{-1}(U \cup V)$ are contained in the same open set, then the sum would increase by deleting one. Thus the supremum is obtained by choosing an increasing sequence of points from $f^{-1}(U \cup V)$ whose image alternates between U and V . Therefore \bar{w} counts the number of times that the image of f alternates between U and V . If f is continuous and U, V have disjoint closures, then its image is compact and can only alternate between sets with disjoint closures finitely many times. So the supremum is actually realized for some finite set of points. If $U' \subset U$ and $V' \subset V$, then $\bar{w}_{U'}^{V'}(f) \leq \bar{w}_U^V(f)$.

Definition 2.9. The *weight* of f with respect to subsets A and B of X with disjoint closures is $w_B^A(f) = \inf \bar{w}_U^V(f)$ taken over all possible separations U and V of \bar{A} and \bar{B} . If $[f]$ is a homotopy equivalence class of functions, then $w_B^A([f]) = \inf_{f \sim f'} \{w_B^A(f')\}$.

Suppose that $\theta : I \rightarrow I$ is nondecreasing function such that $\theta(0) = 0$ and $\theta(1) = 1$. Then $\bar{w}_V^U(f) = \bar{w}_V^U(f \circ \theta)$. Thus \bar{w} is preserved under reparameterization.

If $f : I \rightarrow X$ is a map into a one-dimensional space and $f|_{J_i}$ is a null-homotopic subloop. Then $\bar{w}_V^U(f) \geq \bar{w}_V^U(f')$ where f' is the path obtained by replacing the subpath $f|_{J_i}$ of f by a constant subpath. Thus we obtain $\bar{w}_V^U([f]) = \bar{w}_V^U(f_r)$.

The set $\{\bar{w}_V^U(f) \mid U, V \text{ are a separation of } \bar{A}, \bar{B}\}$ is a subset of the natural numbers and hence has a minimum. Thus we may chose an open separation U, V such that $w_B^A(f) = \bar{w}_U^V(f)$. For continuous f , $f^{-1}(U \cup V)$ can be partitioned into a finite collection of disjoint open sets, I_i , in I with a natural ordering ($I_i \leq I_j$ if $x \leq y$ for all $x \in I_i$ and $y \in I_j$) such that $f(I_i) \subset U$ or $f(I_i) \subset V$. If for some i , $f(I_i) \subset U \setminus \bar{A}$ (or $f(I_i) \subset V \setminus \bar{B}$), then there would exist an open set contained in U (or V) containing \bar{A} (or \bar{B}) which did not intersect $f(I_i)$ and thus alternate fewer

times. Therefore, $f(I_i)$ must intersect \overline{A} (or \overline{B}). So points which realize the weight can be chosen in the closures of A and B . Thus there exists a finite increasing set of points $\{s_i\}$ which can be chosen to have image in the closures of A and B such that $w_B^A(f) = \overline{w}_U^V(f) = \sum_i -r_f(s_i) \cdot r_f(s_{i+1})$. We will sometimes write $w_B^A(f) = \sum_i -r_f(s_i) \cdot r_f(s_{i+1})$. This implicitly implies a choice of U and V to define r_f . However; if $\{s_i\}$ are chosen to have image in the closure of A and B , $r_f(b_i)$ is the same for every choice of U and V . Thus we will ignore this choice at times.

preserve0

Lemma 2.10. *Let X be a planar Peano continuum and k a line in the plane. Suppose that A and B are disjoint closed π_k -saturated sets and A_k, B_k their respective images under π_k . Then $w_B^A(g) = w_{B_k}^{A_k}(\pi_k \circ g)$.*

This follows directly from the fact that the weight can be realized by a finite set of points.

lem6

Lemma 2.11. *If g is an essential loop in a one-dimensional space X , there exist sets O', O'' with disjoint closures such that, for all r , $w_{O''}^{O'}([g]^r) \geq r$. In addition; if $X = \pi_k(Y)$ for some planar Peano continuum Y and some line k in the plane, then O' and O'' can be chosen to be the images of disjoint closed half planes of Y with boundaries parallel to k .*

Proof. We may assume g is a reduced path since the weight of the reduced path is less than or equal to the weight of all paths in its homotopy class.

Since the set of head-tails for g has a natural ordering which is bounded there exists a maximal head-tail, t for g where $g = t * f * \bar{t}$ such that $f * f$ is reduced. Hence $w_B^A([g]^r) \geq w_B^A([f]^r) = r(w_B^A([f]))$ for all A and B . Since g is essential, f is essential which implies f is a non-degenerate path. Then for any two distinct points O' and O'' in the image of f , we have $w_{O''}^{O'}([f]) \neq 0$. Thus $w_{O''}^{O'}([g]^r) \geq r(w_{O''}^{O'}([f])) \geq r$, for any r .

Suppose that $X = \pi_k(Y)$ for some planar Peano continuum Y and some line k in the plane. Then $\pi_k^{-1}(O'), \pi_k^{-1}(O'')$ are disjoint line segments in the plane. Hence there exists disjoint closed half planes A, B such that $\pi_k^{-1}(O') \subset A$ and $\pi_k^{-1}(O'') \subset B$. Thus $w_{\pi_k(B)}^{\pi_k(A)}([g]^r) \geq r(w_{\pi_k(B)}^{\pi_k(A)}([f])) \geq r$, for any r . \square

For f and g as in the proof of Lemma [2.11](#), we will say that f is the *core* of g .

preserve

Lemma 2.12. *Let X be a planar Peano continuum and k a line in the plane. The delineation map, π_k , preserve weights of homotopy classes on disjoint closed saturated sets, i.e. $w_B^A([g]) = w_{B_k}^{A_k}([\pi_k \circ g])$.*

Proof. Let \tilde{g} be the path homotopic to g such that $\pi_k \circ \tilde{g}$ is reduced. Then $w_{B_k}^{A_k}([\pi_k \circ g]) = w_{B_k}^{A_k}(\pi_k \circ \tilde{g}) = w_B^A(\tilde{g}) \geq w_B^A([g])$. For g' homotopic to g , $w_B^A(g') = w_{B_k}^{A_k}(\pi_k \circ g') \geq w_{B_k}^{A_k}(\pi_k \circ \tilde{g}) = w_{B_k}^{A_k}([\pi_k \circ g])$. Then $w_B^A([g]) \geq w_{B_k}^{A_k}([\pi_k \circ g])$.

Thus $w_B^A([g]) = w_{B_k}^{A_k}([\pi_k \circ g])$.

\square

Lemma ^{preserve} 2.12 implies that a necessary condition for g to be k -reduced is that it have minimal weight in its path class with respect to all disjoint half-planes A and B with boundaries parallel to k .

In fact this condition is also sufficient. Suppose g has minimal weight in its path class with respect to all subsets A and B with boundaries parallel to k . If $\pi_k \circ g$ is not reduced, then there exists $g(c)$ and $g(d)$ such that $\overline{g(c)g(d)}$ is contained in X and $\pi_k \circ g|_{[c,d]}$ is null-homotopic but not constant. Then $g|_{[c,d]}$ must not be contained in the line segment $\overline{g(c)g(d)}$. However, then g is homotopic to \tilde{g} where $g|_{[c,d]}$ is replaced by $\overline{g(c)g(d)}$ and the weight of \tilde{g} is strictly less than the weight of g for some disjoint half-planes with boundaries parallel to k .

This characterization of being reduced implies that if \tilde{g} is obtained by reducing g , a k -reduced path, with respect to l ; then \tilde{g} is (k, l) -reduced (reduced with respect to both k and l).

Lemma 2.13. *Suppose that $f : [0, 3] \rightarrow Y$ is a path into a one-dimensional space Y for $i \in 0, 1, 2$. Then for any A and B with disjoint closures, $w_B^A([f]) \leq w_B^A(f|_{[0,1]}) + w_B^A(f|_{[1,2]}) + w_B^A(f|_{[2,3]}) + 2$.*

Proof. Fix a set of maximal disjoint open intervals $\{(a_i, b_i)\}_{i \in J}$ such that $f|_{[a_i, b_i]}$ is a non-degenerate null homotopic loop. Then define f_r as in Theorem ^{reduced loop - thm} 1.2. Choose $\{s_j\}$ such that $w_B^A(f_r) = \sum_{j=1}^n -r_{f_r}(s_j) \cdot r_{f_r}(s_{j+1})$. Since f_r is constant on the intervals (a_i, b_i) , we may assume that $s_j \notin \bigcup_{i \in J} (a_i, b_i)$ for all j . Then $\sum_{j=1}^n -r_{f_r}(s_j) \cdot r_{f_r}(s_{j+1}) = \sum_{j=1}^n -r_f(s_j) \cdot r_f(s_{j+1})$ and

$$\begin{aligned} \sum_{j=1}^n -r_f(s_j) \cdot r_f(s_{j+1}) &\leq \sum_{s_{j+1} \leq 1} -r_f(s_j) \cdot r_f(s_{j+1}) + \sum_{\substack{s_j \geq 1 \\ s_{j+1} \leq 2}} -r_f(s_j) \cdot r_f(s_{j+1}) \\ &\quad + \sum_{s_j \geq 2} -r_f(s_j) \cdot r_f(s_{j+1}) + 2 \\ &\leq w_B^A(f|_{[0,1]}) + w_B^A(f|_{[1,2]}) + w_B^A(f|_{[2,3]}) + 2 \end{aligned}$$

□

nt delineations

Lemma 2.14. *Suppose that $\alpha, \beta : I \rightarrow X$ are (k, l) -reduced paths such that $\pi_k \circ \alpha =_p \pi_k \circ \beta$. Then $w_B^A(\alpha) \leq w_B^A(\beta) + 4$ for any A and B which are disjoint closed half-planes with boundary parallel to l .*

Proof. Since $\pi_k \circ \alpha =_p \pi_k \circ \beta$, the line segments $\overline{\alpha(0)\beta(0)}$ and $\overline{\beta(1)\alpha(1)}$ are contained in X .

Claim 1. $\overline{\alpha(0)\beta(0)}$ and $\overline{\beta(1)\alpha(1)}$ are (k, l) -reduced paths and $w_B^A(\overline{\alpha(0)\beta(0)}), w_B^A(\overline{\beta(1)\alpha(1)}) \leq 1$ for any A and B which are disjoint closed half-planes with boundary parallel to l .

Proof of claim. Notice that $\overline{\alpha(0)\beta(0)}$ and $\overline{\beta(1)\alpha(1)}$ are line segments in X parallel to k ; hence, their images in X_k are degenerate and reduced. Any two non-parallel line segments intersect

at most at a single point. Thus the line segments are also l -reduced and the weight condition follows. \square

However, $\overline{\alpha(0)\beta(0)} * \beta * \overline{\beta(1)\alpha(1)}$ is not necessarily l -reduced.

Fix A and B which are disjoint closed half-planes with boundary parallel to l . Define $f : [0, 3] \rightarrow X$ by $f|_{[0,1]} =_p \overline{\alpha(0)\beta(0)}$, $f|_{[1,2]} =_p \beta$, and $f|_{[2,3]} =_p \overline{\beta(1)\alpha(1)}$. The previous lemma together with the claim gives us the desired inequality. \square

2.3 Induced by a continuous map

For $\varphi : \mathbb{H} \rightarrow \pi_1(X, x_0)$ a fixed homomorphism into the fundamental group of a planar Peano continuum and each line k in the plane, we will use T_k to denote the path such that $\widehat{T}_k \circ (\pi_{k*} \circ \varphi)$ is induced by a continuous map.

The key to being able to reduced the planar case to the one-dimensional case is the following proposition.

phedtail **Proposition 2.15.** *For k a line in the plane, there exists α_k a path in X such that $\pi_k(\alpha_k) = T_k$.*

To prove this proposition we will construct a single word $a \in \mathbb{H}$ such that $T_k \xrightarrow{t} (\pi_k \circ \varphi(a))_r$.

toobig1 **Lemma 2.16.** *Let X be a one-dimensional space. Suppose $\alpha, \beta, \gamma : [0, 1] \rightarrow X$ are reduced paths such that $\alpha(1) = \beta(0)$, $\beta(1) = \gamma(0)$, and $\beta * \gamma$ is a reduced path. If there exists A, B such that $w_B^A(\alpha) < w_B^A(\beta)$ then $(\alpha * \beta)_r * \gamma$ is a reduced path.*

Proof. The only way $(\alpha * \beta)_r * \gamma$ might not be a reduced path is if $\alpha = \gamma' * \overline{\beta}$ for some non-constant path γ' . This would imply $w_B^A(\beta) \leq w_B^A(\alpha)$ for all A, B . \square

toobig2 **Lemma 2.17.** *Let X be a one-dimensional space. Suppose $\alpha, \beta, \gamma : [0, 1] \rightarrow X$ are reduced paths such that $\alpha(1) = \beta(0)$, $\beta(1) = \gamma(0)$, and $\alpha * \beta$ is a reduced path. If $\text{diam}(\gamma) < \text{diam}(\beta)$ then $\alpha * (\beta * \gamma)_r$ is a reduced path.*

The proof follows is similar to the proof of Lemma **toobig1** 2.16.

In Lemma **toobig1** 2.16, the condition that there exists A, B such that $w_B^A(\alpha) < w_B^A(\beta)$ can be replaced by $\text{diam}(\alpha) < \text{diam}(\beta)$. The analogous change for Lemma **toobig2** 2.17 is also valid.

key **Lemma 2.18.** *Let Y be a one-dimensional Peano continuum. Let $f : \mathbf{E} \rightarrow Y$ be a continuous function enjoying the property that $\text{im}(f_*)$ is uncountable and let T be any path from $f(0)$ to $y \in Y$. Then there exists $a \in \mathbb{H}$ such that no non-constant head of T is a tail for $f_*(a)_r$, i.e. $f_*(a)_r * T$ is reduced.*

This can be derived as a result of Lemma 6.1 in **leda** [E2]. However; the proof of Proposition **pmaximalheadtail** 2.20 depends heavily on the construction of a , so we will present a proof here for completeness.

Proof. Let b_n and t_n be subpaths of $f_*(a_n)_r$ such that $f_*(a_n)_r =_p t_n * b_n * \bar{t}_n$ where b_n is the core of $f_*(a_n)_r$. We may assume that b_n is an essential loop, by passing to a subsequence if necessary.

Since f is continuous, the paths t_n and $f_*(a_n)_r$ must both form null sequences. Hence, we may choose an increasing sequence i_n such that $\text{diam}(t_{i_n})$ is decreasing and $\text{diam}(f_*(a_n)_r) < \frac{\text{diam}(t_{i_{n-1}})}{2}$.

There exists an r_n such that for some A_n and B_n , $w_{B_n}^{A_n}(T) < w_{B_n}^{A_n}(b_{i_n} * \cdots * b_{i_n})$, the concatenation of r_n copies of b_{i_n} .

We will inductively define integers s_n and homotopy classes c_n in \mathbb{H} . Let $s_1 = 0$, c_1 be the identity in \mathbb{H} , and suppose s_i and c_i are defined for $1 \leq i < n$.

Again, there exists an s_n such that for some A'_n and B'_n , $w_{B'_n}^{A'_n}(f_*(\prod_{i=1}^{n-1} (c_i))_r * t_{i_n}) < w_{B'_n}^{A'_n}(b_{i_n} * \cdots * b_{i_n})$, the concatenation of s_n copies of b_{i_n} . Let $c_n = a_{i_n}^{r_n + s_n}$. Then $f_*(c_n)_r = t_n * b_n^{r_n + s_n} * \bar{t}_n$. To simplify notation, let t_{i_0}, b_{i_0} be constant paths and $r_{i_0}, s_{i_0} = 0$.

Let $a = \prod_{n=1}^{\infty} c_n$ and note that this is a well-define homotopy class in \mathbb{H} . Since f is a continuous maps; $\prod_{n=1}^{\infty} t_n * b_n^{r_n + s_n} * \bar{t}_n$ is a well defined path and $f_*(a) = \left[\prod_{n=1}^{\infty} t_n * b_n^{r_n + s_n} * \bar{t}_n \right]$. Our objective with the next three claims is to understand the backtracking that appears in the path $\prod_{n=1}^{\infty} t_n * b_n^{r_n + s_n} * \bar{t}_n$.

claim1 **Claim 2.** For $m \geq 2$,

$$t_{i_1} * b_{i_1}^{s_1} * \left(\prod_{j=2}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r \right) \quad (2.1) \quad \boxed{1}$$

is a reduced path.

Since t_{i_0}, b_{i_0} are constant paths and $r_{i_0} = 0$, Equation (2.1) is the same (up to reparameterization) as

$$\prod_{j=1}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r.$$

Proof of Claim 2. claim1 We will proceed by induction on m . The case $m = 1$ is trivial. By inducting on m , suppose that that $\prod_{j=1}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r$ is reduced.

Then

$$\left(\prod_{j=1}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r \right) * b_{i_m}^{r_{i_m}} * \bar{t}_{i_m}$$

is a reduced path by Lemma toobig1 2.16 with $\alpha = \left(\left(\prod_{j=1}^{m-1} b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r \right) * b_{i_{m-1}}^{r_{i_{m-1}}} * \bar{t}_{i_{m-1}} * t_{i_m} \right)_r$, $\beta = b_{i_m}^{s_{i_m}}$, and $\gamma = b_{i_m}^{r_{i_m}} * \bar{t}_{i_m}$. Here we are also using the inductive hypothesis.

We can complete the proof by applying Lemma toobig2 2.17, one more time, with $\alpha = \left(\prod_{j=1}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r \right) * b_{i_m}^{r_{i_m}}$, $\beta = \bar{t}_{i_m}$, and $\gamma = t_{i_2} * b_{i_2}^{s_{i_2}}$.

□

claim3 **Claim 3.** For $m \geq 2$,

$$b_{i_m}^{r_{i_m}} * \bar{t}_{i_m} * f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r \quad (2.2) \quad \boxed{4}$$

is a reduced path.

This follow by applying Lemma [too big 2](#) [2.17](#) with $\alpha = b_{i_m}^{r_{i_m}}$, $\beta = t_{i_m}$ and $\gamma = f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r$.

claim2 **Claim 4.** For all $m \geq 2$,

$$f_*(a)_r =_p t_{i_1} * b_{i_1}^{s_1} * \left(\prod_{j=2}^m b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r \right) * b_{i_m}^{r_{i_m}} * \bar{t}_{i_m} * f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r \quad (2.3) \quad \boxed{2}$$

where the equality implies that the right hand side is reduced.

Proof of Claim [claim2](#) [4](#). Since b_{i_m} is cyclically reduced and Equation [\(2.2\)](#) is reduced,

$$b_{i_m}^{s_{i_m}} * \left(b_{i_m}^{r_{i_m}} * \bar{t}_{i_m} * f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r \right)$$

is a reduced path.

This give us that

$$\left(t_{i_1} * b_{i_1}^{s_1} * \left[\prod_{j=2}^m (b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r) \right] \right)_r * \left(b_{i_m}^{r_{i_m}} * \bar{t}_{i_m} * f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r \right) \quad (2.4) \quad \boxed{3}$$

is a reduced path by applying Lemma [too big 1](#) [2.16](#) with

$$\alpha = \left(t_{i_1} * b_{i_1}^{s_1} * \left(\prod_{j=2}^{m-1} (b_{i_{j-1}}^{r_{j-1}} * (\bar{t}_{i_{j-1}} * t_{i_j} * b_{i_j}^{s_{i_j}})_r) \right) * b_{i_{m-1}}^{r_{m-1}} * \bar{t}_{i_{m-1}} * t_{i_m} \right)_r, \beta = b_{i_m}^{s_{i_m}}, \text{ and}$$

$$\gamma = \left(b_{i_m}^{r_{i_m}} * \bar{t}_{i_m} * f_* \left(\prod_{i=m+1}^{\infty} c_i \right)_r \right).$$

Since Equations [\(2.1\)](#), [\(2.2\)](#), and [\(2.4\)](#) are reduced; Equation [\(2.3\)](#) is reduced.

□

We can now complete the proof of Lemma [key](#) [2.18](#). Suppose that $T|_{[0,s]}$ is a non-constant tail of $f_*(a)_r$. Then we can choose m_0 sufficiently large such that $\text{diam} \left(b_{i_{m_0-1}}^{r_{i_{m_0-1}}} * \bar{t}_{i_{m_0-1}} * f_* \left(\prod_{i=m_0}^{\infty} c_i \right)_r \right) < \text{diam}(T|_{[0,s]})$. However, then Claim [claim2](#) [4](#) for $m = m_0$ would imply that $b_{m_0}^{r_{m_0}}$ is a subpath of T which contradicts our choice of r_{m_0} .

□

corkey **Corollary 2.19.** Let X be a planar Peano continuum and $\varphi : \mathbb{H} \rightarrow \pi_1(X, x_0)$ be a homomorphism with uncountable image. If k is a line in the plane and $\widehat{T}_k \circ \pi_{k*} \circ \varphi = f_*$ for some continuous map $f : \mathbf{E} \rightarrow X_k$, then T_k is a tail for $(\pi_{k*} \circ \varphi(a))_r$ where a is chosen as in Lemma [key](#) [2.18](#).

Proof. Note that $(\pi_{k_*} \circ \varphi(a))_r = (T_k * f_*(a))_r * \overline{T_k}_r$. Then Lemma [2.16](#) and Proposition [2.18](#) imply that $(T_k * f_*(a))_r * \overline{T_k}$ is a reduced path. □

Then Proposition [2.15](#) follows from Corollary [2.19](#) and Lemma [2.6](#).

Proposition 2.20. *Let X be a planar Peano continuum, $\varphi : \mathbb{H} \rightarrow \pi_1(X, x_0)$ be a homomorphism with uncountable image, and k_1, k_2 lines in the plane. Then there exists a path α in X such that $\pi_k(\alpha) = T_k$ where T_k is the path such that $\widehat{T_k} \circ \pi_{k_*} \circ \varphi$ is induced by a continuous map and $k \in \{k_1, k_2\}$.*

Proof. If k_1, k_2 are parallel this is an immediate consequence of Corollary [2.19](#). Hence we will assume that k_1, k_2 are non-parallel.

Let f_j be the continuous map such that $f_{j*} = \widehat{T_{k_j}} \circ (\pi_{k_{j*}} \circ \varphi)$.

We will now show how to modify the proof of Proposition [2.18](#) to find a single a such that T_{k_j} is tail for $(\pi_{k_{j*}} \circ \varphi(a))_r$ for either $j = 1, 2$.

Let $b_{n,j}$ and $t_{n,j}$ be subpaths of $f_{j*}(a_n)_r$ such that $f_{j*}(a_n)_r =_p t_{n,j} * b_{n,j} * \overline{t_{n,j}}$ where $b_{n,j}$ is the core of $f_{j*}(a_n)_r$.

We may assume that $b_{n,j}$ is an essential loop for both $j = 1, 2$; by passing to a subsequence if necessary. Since f_j is continuous, the paths $t_{n,j}$ and $f_{j*}(a_n)_r$ must both form null sequences. We may choose an increasing sequence i_n such that $\text{diam}(t_{i_n,j})$ is decreasing and $\text{diam}(f_{j*}(a_n)_r) < \frac{\text{diam}(t_{i_n-1,j})}{2}$ for all $n \geq i_n$. The subsequence i_n does not depend on j .

There exists an $r_{n,j}$ such that for some $A_{n,j}$ and $B_{n,j}$,

$$w_{B_{n,j}}^{A_{n,j}}(T_{k_j}) < r_{n,j} \leq w_{B_{n,j}}^{A_{n,j}}(b_{i_n,j} * \cdots * b_{i_n,j}),$$

the concatenation of $r_{n,j}$ copies of $b_{i_n,j}$.

Fix β_{k_j} a (k_1, k_2) -reduced path such that $\pi_{k_j} \circ \beta_{k_j} =_p T_{k_j}$. There exists an $r'_{n,j}$ such that for some $A'_{n,j}$ and $B'_{n,j}$ disjoint half planes with boundary parallel to k_j , we have

$$w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_j} \circ \beta_{k_j}) + 4 < r'_{n,j} \leq w_{B'_{n,j}}^{A'_{n,j}}(b_{i_n,j} * \cdots * b_{i_n,j})$$

the concatenation of $r'_{n,j}$ copies of $b_{i_n,j}$, where $j^c = \{1, 2\} \setminus \{j\}$

Let $r_n = \max\{r_{n,1}, r_{n,2}, r'_{n,1}, r'_{n,2}\}$.

We will inductively define integers $s_{n,j}$ and homotopy classes c_n in \mathbb{H} . Let $s_{1,j} = 0$ for $j = 1, 2$ and c_1 be the identity in \mathbb{H} . Suppose $s_{n,j}$ and c_i are defined for $1 \leq i < n$ and $j = 1, 2$.

Again, there exists an $s_{n,j}$ such that for some $A''_{n,j}$ and $B''_{n,j}$,

$$w_{B''_{n,j}}^{A''_{n,j}}(f_{j*}(\prod_{i=1}^{n-1} (c_i))_r * t_{i_n,j}) < w_{B''_{n,j}}^{A''_{n,j}}(b_{i_n,j} * \cdots * b_{i_n,j}),$$

the concatenation of $s_{n,j}$ copies of $b_{i_n,j}$ for $j = 1, 2$. Let $s_n = \max\{s_{n,1}, s_{n,2}\}$ and $c_n = a_{i_n}^{r_n + s_n}$. Let $a = \prod_{n=1}^{\infty} c_n$.

Fix $h : I \rightarrow X$, a (k_1, k_2) -reduced path representative of $\varphi(a)$. We can then apply the remainder of the proof of Lemma [2.18](#) without change to find terminal supaths α_1, α_2 of h mapping to $\overline{T}_{k_1}, \overline{T}_{k_2}$ respectively.

We may assume that $\alpha_1 =_p \gamma * \alpha_2$ for some subpath γ of h . Suppose that $\pi_{k_2} \circ \gamma$ is non-degenerate. By our construction of a , $b_{i_n,2}^{r_n}$ is a subpath of $\pi_{k_2} \circ \gamma$ for some n .

Since $\pi_{k_1} \circ \alpha_1 =_p \pi_{k_1} \circ \beta_{k_1}$, we can apply Lemma [2.14](#) and Lemma [2.10](#) to obtain $w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_2} \circ \alpha_1) \leq w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_2} \circ \beta_{k_1}) + 4$.

Hence $w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_2} \circ \gamma) \leq w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_2} \circ \alpha_1) < r_n$ and $r_n \leq w_{B'_{n,j}}^{A'_{n,j}}(b_{i_n,2}^{r_n}) \leq w_{B'_{n,j}}^{A'_{n,j}}(\pi_{k_2} \circ \gamma)$ which is the desired contradiction.

Thus $\pi_{k_2} \circ \alpha_1 =_p T_{k_2}$ and $\pi_{k_1} \circ \alpha_1 =_p T_{k_1}$ as desired. □

We will now show that α is the path such that $\widehat{\alpha} \circ \varphi$ is induced by a continuous map.

Lemma 2.21. *Let $\alpha_k : I \rightarrow X$ be a path with the property that $\pi_k \circ \alpha_k = T_k$, up to reparametrization. Let U be a π_k -saturated neighborhood of $\pi_k^{-1}(T_k(1))$. For sufficiently large n , every element of $\widehat{\alpha} \circ \varphi(\mathbb{H}_n)$ has a representative in U .*

Proof. If g is a loop based at a point $y \in \pi_k^{-1}(x)$ and $w_{U^c}^{\pi_k^{-1}(x)}([g]) = 0$, then g has a representative in U .

Let U be a π_k -saturated neighborhood of $\pi_k^{-1}(T_k(1))$. Let U' be an open π_k -saturated neighborhood of $\pi_k^{-1}(T_k(1))$ with closure contained in the interior of U . We must show that, for some sufficiently large n , $w_B^A(\widehat{\alpha} \circ \varphi(b)) = 0$ for all $b \in \mathbb{H}_n$ where $A = \pi_k^{-1}(T_k(1))$ and $B = \overline{U'}^c$.

Since $\widehat{T}_k \circ \pi_{k*} \circ \varphi$ is induced by a continuous map, there exists an N such that, for all $n > N$, $\widehat{T}_k \circ \pi_{k*} \circ \varphi(H_n) \subset \pi_1(\pi_k(U'), T_k(1))$. Hence $w_{B_k}^{A_k}(\widehat{T}_k \circ \pi_{k*} \circ \varphi(b)) = 0$ for all $b \in \mathbb{H}_n$ where $n > N$.

For $b \in \mathbb{H}_n$, where $n > N$, let f be a k -reduced representative of $\widehat{\alpha} \circ \varphi(b)$. Then the $w_B^A(\widehat{\alpha} \circ \varphi(b)) \leq w_B^A(f) = w_{B_k}^{A_k}(\pi_k \circ f) = w_{B_k}^{A_k}(\widehat{T}_k \circ \pi_{k*} \circ \varphi(b)) = 0$. □

contplanar

Theorem 2.22. *Let $\varphi : \mathbb{H} \rightarrow \pi_1(X, x_0)$ a homomorphism into the fundamental group of a planar Peano continuum X . Then there exists a continuous function $f : (\mathbf{E}, 0) \rightarrow (X, x)$ and a path $\alpha : (I, 0, 1) \rightarrow (X, x_0, x)$, such that $f_* = \widehat{\alpha} \circ \varphi$. Additionally, if the image of φ is uncountable then α is unique up to homotopy relative to its endpoints.*

Proof. For k and l nonparallel lines in the plane, there exists a path α in X such that $\pi_k(\alpha) = T_k$ and $\pi_l(\alpha) = T_l$, by Lemma [2.20](#).

It is sufficient to show that for any neighborhood U of $\alpha(1)$ there exists an N such that every element of $\widehat{\alpha} \circ \varphi(\mathbb{H}_n)$ has a representative in U .

This is done by finding U_l and U_k such that $U_k \cap U_l \subset U$ and U_l is a π_l -saturated neighborhood of $\pi_l^{-1} \circ \pi_l(\alpha(1))$ and U_k is a π_k -saturated neighborhood of $\pi_k^{-1} \circ \pi_k(\alpha(1))$. The uniqueness follows from the uniqueness of T_k and the injectivity of π_{k*} . \square

3 Homomorphisms induced by continuous functions

sec2

Throughout this section, we will assume all paths are from the unit interval $[0, 1]$. We will use j for the inclusion map of subsets of X into X . When there is no chance of confusion, we will allow the domain of j to change without comment.

Definition 3.1. Let $\varphi : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ be a homomorphism. We will say x is φ -bad if $\varphi \circ \widehat{\alpha} \circ j_*(\pi_1(B_\epsilon(x), x))$ is nontrivial for all $\epsilon > 0$ where α is a path from x_0 to x . We will say that X is φ -bad, if it is at each point. We will write $B_\varphi(X)$ for the set of φ -bad points of X . If $X = Y$ and $\varphi = \text{id}$, then this definition is the definition given in Section [sec3](#) for the *bad* set of X and we will simply write $B(X)$.

Note this is independent of the chosen α .

Cannon and Conner in [cc3](#) defined $B(X)$ for a certain class of subset of the plane which where homotopy equivalent to planar Peano continua. It is a trivial exercise to see that these definitions are compatible. Conner and Eda in [ce](#) defined a similar set for one-dimensional spaces which they referred to as the set of *wild* points. While both terms have appeared in the literature, here we will use the term bad.

The following theorem, while not explicitly stated, follows from the proof of Theorem 1.2 in [eda3](#) [E3](#).

retractEda-thm

Theorem 3.2 (Eda, [eda3](#) [E3](#)). *Let $\varphi : \pi_1(X, x_0) \rightarrow G$ be a homomorphism from the fundamental group of a one-dimensional Peano continuum X to a group G which is a subgroup of an inverse limit of free groups. Suppose that $x_0 \in B_\varphi(X)$. Then there exists a retraction $r : (X, x_0) \rightarrow (X, x_0)$ such that $B_\varphi(X) = B(r(X))$ and $\varphi([f]) = \varphi([r \circ f])$ for all loops f based at x_0 .*

corr

Proposition 3.3. *Let X and Y be one-dimensional or planar Peano continua. Suppose that $\varphi : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ is a homomorphism between their fundamental groups. Then for every path $\alpha : (I, 0, 1) \rightarrow (X, x_0, x)$ such that $x \in B_\varphi(X)$, there exists a unique (up to homotopy rel endpoints) path $\psi(\alpha) : (I, 0, 1) \rightarrow (Y, y_0, y)$ where $\psi(\alpha)$ depends only on the homotopy class (rel endpoints) of α and y depends only on x which enjoys the following property.*

$$\forall g : (\mathbf{E}, 0) \rightarrow (X, \alpha(1)), h : (\mathbf{E}, 0) \rightarrow (Y, \psi(\alpha)(1)) \text{ such that } \varphi \circ \widehat{\alpha} \circ g_* = \widehat{\psi(\alpha)} \circ h_*. \quad (3.1)$$

univ_proper

If X and Y one-dimensional, then this is Lemma 5.1 in ^{eda}[E2]. The proof we present here is essentially that same as Eda's proof. We give the proof for the planar case here to establish notation and to show the insignificant changes required in the planar situation.

Proof. Since $x \in B_\varphi(X)$, there exists a sequence of loops, $\{\mathbf{b}_i\}$, based at x which converge to x with the property that $[\alpha * \mathbf{b}_i * \bar{\alpha}]$ (or the conjugation by any other path from x_0 to x) map non-trivially under φ . Let g be a continuous map which sends \mathbf{a}_i to \mathbf{b}_i . Theorem ^{contplanar}2.22 implies that there exists a unique $\psi(\alpha) : (I, 0, 1) \rightarrow (Y, y_0, y)$ and an $h : (\mathbf{E}, 0) \rightarrow (Y, y)$ such that $\varphi \circ \widehat{\alpha} \circ g_* = \widehat{\psi(\alpha)} \circ h_*$.

Suppose $g' : (\mathbf{E}, 0) \rightarrow (X, x)$ is any continuous function. Let $g'' : (\mathbf{E}, 0) \rightarrow (X, x)$ be the function which sends \mathbf{E}_e to X by g and sends \mathbf{E}_o to X by g' . Notice g'' is continuous since $g(0) = g'(0) = x$. Then there exists \mathbf{c} and $h'' : (\mathbf{E}, 0) \rightarrow (Y, y)$ such that $\varphi \circ \widehat{\alpha} \circ g''_* = \widehat{\mathbf{c}} \circ h''_*$.

Then $\widehat{\psi(\alpha)} \circ h_* = \varphi \circ \widehat{\alpha} \circ g_* = \varphi \circ \widehat{\alpha} \circ (g''|_{\mathbf{E}_e})_* = \widehat{\mathbf{c}} \circ (h''|_{\mathbf{E}_e})_*$. Then the uniqueness in Theorem ^{contplanar}2.22 implies that $\psi(\alpha)$ is homotopic to \mathbf{c} . Hence $\varphi \circ \widehat{\alpha} \circ g'_* = \varphi \circ \widehat{\alpha} \circ (g''|_{\mathbf{E}_o})_* = \widehat{\psi(\alpha)} \circ (h''|_{\mathbf{E}_o})_*$.

Now we must only show that y depends only on x . Suppose $\alpha' : (I, 0, 1) \rightarrow (X, x_0, x)$ is a path which is not necessarily homotopic to α . Then for some path $\psi(\alpha')$ and h' , we have $\varphi \circ \widehat{\alpha'} \circ g_* = \widehat{\psi(\alpha')} \circ h'_*$.

$$\begin{aligned} \widehat{\psi(\alpha')} \circ h'_*(u) &= \varphi \circ \widehat{\alpha'} \circ g_*(u) \\ &= \varphi \circ \widehat{\alpha' * \bar{\alpha}} \circ \widehat{\alpha} \circ g_*(u) \\ &= \varphi(\alpha' * \bar{\alpha}) \cdot \varphi \circ \widehat{\alpha} \circ g_*(u) \cdot (\varphi(\alpha' * \bar{\alpha}))^{-1} \\ &= \varphi(\alpha' * \bar{\alpha}) \cdot \widehat{\psi(\alpha)} \circ h_*(u) \cdot \varphi^{-1}(\alpha' * \bar{\alpha}) \end{aligned}$$

Hence $h'(\mathbf{a}_n)$ is freely homotopic to $h(\mathbf{a}_n)$ for all i .

If $\psi(\alpha')(1) \neq \psi(\alpha)(1)$, then we can find n sufficiently large such that $h(\mathbf{a}_n)$ and $h'(\mathbf{a}_n)$ can be separated by disjoint balls in Y . This contradicts Lemma ^{homotoped off}A6. (We can choose n sufficiently large such that the disjoint balls satisfy the hypothesis of Lemma ^{homotoped off}A6.) \square

arc-reduced

Theorem 3.4 (Meistrup, ^{precm}[CM]). *Every one-dimensional Peano continuum is homotopy equivalent to a one-dimensional Peano continuum X such that X is a finite connected graph or $X \setminus B(X)$ is null sequence of disjoint open arcs with end points in $B(X)$.*

Such a continuum is called *arc-reduced*.

We are now ready to prove:

cont1dim

Theorem 3.5. *Let $\varphi : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ be a homomorphism where X is a one-dimensional Peano continuum and Y is a planar Peano continua. Then there exists a path $T : (I, 0, 1) \rightarrow (Y, y_0, y)$ and a continuous function $f : X \rightarrow Y$ such that $\widehat{T} \circ \varphi = f_*$.*

When both X and Y are one-dimensional, this is Theorem 1.2 in Eda's recent paper ^{eda3}[E3]. We will show how to adapt Eda's to the case where Y is a planar Peano continuum.

3.1 Proof of Theorem ^{cont1dim}3.5

Let $\varphi : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ be a homomorphism where X is a one-dimensional Peano continuum and Y is a planar Peano continua.

Observation: Let $r : X \rightarrow X$ be any retraction of X such that $\ker r_* \subset \ker \varphi$. Then there exists a unique homomorphism $\varphi' : \pi_1(r(X), r(x_0)) \rightarrow \pi_1(Y, y_0)$ such that $\varphi([\alpha]) = \varphi'([r \circ \alpha])$ for every loop α in X . In addition, if there exists a path $T : (I, 0, 1) \rightarrow (Y, y_0, y)$ and a continuous function $f' : r(X) \rightarrow Y$ such that $\hat{T} \circ \varphi' = f'_*$; then

$$\hat{T} \circ \varphi = \hat{T} \circ (\varphi' \circ r_*) = f'_* \circ r_* = (f' \circ r)_*.$$

Thus by Theorem ^{retractEda-thm}3.2 and Theorem ^{arc-reduced}3.4, we may assume that $B(X) = B_\varphi(X)$ and X is arc-reduced. Since we are only concerned about φ up to composition with a change of base point isomorphism, we may assume that $x_0 \in B(X)$.

Fix k, l non-parallel lines in the plane. Then there exists paths T_k, T_l in $\pi_k(Y), \pi_l(Y)$ respectively and functions $f_k : X \rightarrow \pi_k(Y), f_l : X \rightarrow \pi_l(Y)$ such that $\hat{T}_k \circ \pi_{k*} \circ \varphi = f_{k*}$ and $\hat{T}_l \circ \pi_{l*} \circ \varphi = f_{l*}$.

Since X is arc-reduced, $X \setminus B(X)$ is a null sequence of disjoint open arcs, A_i . For each i , fix a homeomorphism $m_i : (0, 1) \rightarrow A_i$. Then extend m_i to $[0, 1]$ by sending $0, 1$ to the corresponding endpoints of A_i in $B(X)$. Without loss of generality, we may assume that f_k, f_l take each m_i to a reduced path in $\pi_k(Y), \pi_l(Y)$ respectively.

To simplify notation; we will use $[f]_r$ to represent the reduced representative of $[f]$ when f is a path in a one-dimensional continuum and $[f]_{k,l}$ to represent a (k, l) -reduced representative of f when f is a path a planar Peano continuum.

Proposition ^{corr}3.3 gives us a function ψ from the set of reduced paths with initial point x_0 and terminal point in $B(X)$ to the set of paths in Y from y_0 to points in $B(Y)$. We want to extended ψ to reduced paths from x_0 to arbitrary points of X .

Let \mathbf{m}_i be a (k, l) -reduced representative of $[\overline{\psi(\beta_i) * \psi(\beta_i * m_i)}]$ where β_i is a path from x_0 to $m_i(0)$. The next two lemmas will show that \mathbf{m}_i is independent of our chosen β_i .

The proofs we present here for the next two lemmas is essentially that same as Eda's proofs of the corresponding lemmas.

eda3.3 **Lemma 3.6.** [^{eda3}Lemma 3.3, ^[E3]] Let α, β be paths from x_0 to $x \in B(X)$. Then $\varphi(\alpha * \overline{\beta}) = [\psi(\alpha) * \overline{\psi(\beta)}]$.

Proof. For all continuous $h : (\mathbf{E}, 0) \rightarrow (X, x)$, there exists $g, g' : (\mathbf{E}, 0) \rightarrow (Y, y)$ such that

$\varphi \circ \widehat{\beta} \circ h_* = \overline{\widehat{\psi}(\beta)} \circ g_*$ and $\varphi \circ \widehat{\alpha} \circ h_* = \overline{\widehat{\psi}(\alpha)} \circ g'_*$. Then for all $u \in \mathbb{H}$,

$$\begin{aligned} \overline{\widehat{\psi}(\alpha)} \circ g'_*(u) &= \varphi \circ \widehat{\alpha} \circ h_*(u) \\ &= \varphi \circ \overline{\widehat{\alpha} * \widehat{\beta}} \circ \widehat{\beta} \circ h_*(u) \\ &= \varphi([\alpha * \overline{\beta}]) \cdot \varphi \circ \widehat{\beta} \circ h_*(u) \cdot (\varphi([\alpha * \overline{\beta}]))^{-1} \\ &= \varphi(\alpha * \overline{\beta}) \cdot \overline{\widehat{\psi}(\beta)} \circ g_*(u) \cdot (\varphi(\alpha * \overline{\beta}))^{-1} \\ &= \overline{\varphi(\alpha * \overline{\beta})_{k,l} * \widehat{\psi}(\beta)} \circ g_*(u) \end{aligned}$$

Then Theorem [contplanar 2.22](#) implies that $\psi(\alpha)$ is homotopic to $\varphi(\alpha * \overline{\beta})_{k,l} * \psi(\beta)$ which completes the proof. \square

eda3.4 **Lemma 3.7.** [[Lemma 3.4](#), [eda3 \[E3\]](#)] The path m_i is independent (up to homotopy rel endpoints) of the path β_i from x_0 to the initial point of m_i .

Proof. Let β, γ be paths from x_0 to the initial point of m_i . Then by Lemma [eda3.3 3.6](#),

$$\begin{aligned} \left(\overline{\psi(\gamma)} * \psi(\gamma * m_i) * \overline{\psi(\beta * m_i)} * \psi(\beta) \right) &=_{0,1} \overline{\psi(\gamma)} * \left(\psi(\gamma * m_i) * \overline{\psi(\beta * m_i)} \right) * \psi(\beta) \\ &=_{0,1} \overline{\psi(\gamma)} * \varphi([\gamma * m_i * \overline{m_i} * \overline{\beta}])_{k,l} * \psi(\beta) \\ &=_{0,1} \overline{\psi(\gamma)} * \varphi([\gamma * \overline{\beta}])_{k,l} \cdot \psi(\beta) \\ &=_{0,1} \overline{\psi(\gamma)} * \psi(\gamma) * \psi(\overline{\beta}) * \psi(\beta) =_{0,1} x_0 \end{aligned}$$

where x_0 is the constant path. \square

We can now extend ψ to reduced paths from x_0 to arbitrary points of X . Let $\alpha : (I, 0, 1) \rightarrow (X, x_0, x)$ be a reduced path. If $x \in B(X)$, then $\psi(\alpha)$ is already defined. Otherwise, $x = m_i(t_0)$ for some i and t_0 . Then α , up to re-parametrization, is $\alpha' * m_i|_{[0,t_0]}$ or $\alpha' * \overline{m_i}|_{[0,1-t_0]}$ where α' is uniquely determined and is a path between points in $B(X)$. Then we will define $\psi(\alpha)$ to be $\psi(\alpha') * m_i|_{[0,t_0]}$ if $\alpha = \alpha' * m_i|_{[0,t_0]}$ or $\psi(\alpha') * \overline{m_i}|_{[0,1-t_0]}$ if $\alpha = \alpha' * \overline{m_i}|_{[0,1-t_0]}$.

We are now ready to define our function between X and Y which will induce φ up to conjugation. Let $f : X \rightarrow Y$ by $f(x) = \psi(\alpha)(1)$ for a path α from x_0 to x . If $x \in B(X)$, this is independent of α by Proposition [corr 3.3](#). If $x \in X \setminus B(X)$, this is independent of α by Lemma [eda3.4 3.7](#). By construction, $f|_{m_i}$ is continuous.

composition **Lemma 3.8.** Let $x \in B(X)$ and $\alpha : (I, 0, 1) \rightarrow (X, x_0, x)$. Then $\pi_k \circ \psi(\alpha) =_{0,1} T * f_k \circ \alpha$ and $\pi_k \circ f(x) = f_k(x)$.

Proof. By construction $\psi(\alpha)$ is the unique path satisfying [univ_property \(3.1\)](#) for the homomorphism φ . By construction $f(x) = \psi(\alpha)(1)$.

Fix $g : (\mathbf{E}, 0) \rightarrow (X, \alpha(1))$.

By Proposition [3.3](#)^{corr}, there exists $h : (\mathbf{E}, 0) \rightarrow (Y, \psi(\alpha)(1))$ such that $\varphi \circ \hat{\alpha} \circ g_* = \overline{\hat{\psi}(\alpha)} \circ h_*$. Hence

$$\begin{aligned} \pi_{k_*} \circ \varphi \circ \hat{\alpha} \circ g_* &= \pi_{k_*} \circ \overline{\hat{\psi}(\alpha)} \circ h_* \\ &= \overline{\pi_k \circ \hat{\psi}(\alpha)} \circ \pi_{k_*} \circ h_* \\ &= \overline{\pi_k \circ \hat{\psi}(\alpha)} \circ (\pi_k \circ h)_* \end{aligned}$$

and, as well,

$$\begin{aligned} \pi_{k_*} \circ \varphi \circ \hat{\alpha} \circ g_* &= \hat{T} \circ f_{k_*} \circ \hat{\alpha} \circ g_* \\ &= \hat{T} \circ \overline{f_k \circ \alpha} \circ (f_k \circ g)_* \\ &= \overline{T * f_k \circ \alpha} \circ (f_k \circ g)_*. \end{aligned}$$

The uniqueness conditions from Proposition [3.3](#)^{corr} imply that $f_k(x) = f_k \circ \alpha(1) = \pi_k \circ \psi(\alpha)(1) = \pi_k \circ f(x)$ and $\pi_k \circ \psi(\alpha) =_{0,1} T * f_k \circ \alpha$. \square

ost composition

Lemma 3.9. $\pi_k \circ f \circ m_i =_{0,1} f_k \circ m_i$

Proof. By Lemma [3.8](#)^{composition},

$$\begin{aligned} \pi_k \circ f \circ m_i &= \pi_k \circ \mathbf{m}_i =_{0,1} \overline{\pi_k \circ \psi(\beta_i) * \pi_k \circ \psi(\beta_i * m_i)} \\ &=_{0,1} \overline{(T * f_k \circ \beta_i)} * (T * (f_k \circ \beta_i) * (f_k \circ m_i)) \\ &=_{0,1} f_k \circ m_i \end{aligned}$$

\square

Note that both Lemma [3.8](#)^{composition} and Lemma [3.9](#)^{almost composition} hold when k is replaced by l .

Lemma 3.10. f is continuous.

Proof. By Lemma [2.4](#) and [3.8](#)^{composition}, we know that $\pi_k \circ f$ and $\pi_l \circ f$ are continuous of $B(X)$. Since $m_i(I)$ from a null sequence in X , $f_k(m_i(I))$ and $f_l(m_i(I))$ both form null sequence of sets in Y_k, Y_l respectively. Since \mathbf{m}_i is (k, l) -reduced and $f_k \circ m_i$ is reduced, $\pi_k(f(m_i(I))) = f_k(m_i(I))$ and $\pi_l(f(m_i(I))) = f_l(m_i(I))$. Then Lemma [2.5](#)^{continuous maps-corrollary} implies that $f(m_i(I))$ forms a null sequence of sets in Y . Thus f is continuous on $\bigcup_i m_i(I)$ which completes the proof. \square

Lemma 3.11. $\pi_{k_*} \circ f_* = f_{k_*}$

Proof. Suppose that $\alpha : (I, 0, 1) \rightarrow (X, x_0, x_0)$. Choose open subinterval I_j of I with the canonical ordering such that $\bigcup I_j = I \setminus \alpha^{-1}(B(X))$. Then $\pi_k \circ f \circ \alpha|_{I_j}$ is a reparametrization of $f_k \circ \alpha|_{I_j}$ and $\pi_k \circ f = f_k$ on $B(X)$ which completes the proof. \square

All that remains to show is that $\varphi = \hat{\alpha} \circ f_*$ for some path α from y_0 to $f(x_0)$. As in Proposition [2.20](#), we can find a path α such that $\pi_k(\alpha) =_{0,1} T_k$. Then

$$\begin{aligned} \pi_{k*} \circ \varphi &= \hat{T}_k \circ f_{k*} \\ &= \hat{T}_k \circ \pi_{k*} \circ f_* \\ &= \pi_{k*} \circ \hat{\alpha} \circ f_*. \end{aligned}$$

Since π_{k*} is injective $\varphi = \hat{\alpha} \circ f_*$ and the uniqueness of α follows from the uniqueness of T_k which completes the proof of Theorem [2.22](#).

4 Recovering the bad set

sec3

Recall that Eda and Conner in [\[CE\]](#) showed that the fundamental group of a one-dimensional continuum which is not semilocally simply connected at any point can be used to reconstruct the original space. The construction that was used there is similar to the one we will present here for the planar case.

Definition 4.1. We will denote the set of points at which X is not semilocally simply connected by $B(X)$, the *bad set* of X .

fixed

Lemma 4.2. *Let be X a planar Peano continuum. Then $B(X)$ is fixed by every self-homotopy of X .*

Proof. Let $f : X \rightarrow X$ be homotopic to the identity on X . Then any loop is homotopic to its image under f . Suppose that $x \in B(X)$ such that $f(x) \neq x$. Since X is not semilocally simple connected at x , there exists a closed disk neighborhood D of x in the plane with the property that ∂D is not contained in X and $D \cap f(D) = \emptyset$. Then any loop contained in D can be homotoped into $f(D)$. However; this is a contradiction of Lemma [A6](#), since there exist essential loops contained in the interior of $D \cap X$.

□

Definition 4.3. We will say that H is a *pseudo-Hawaiian earring subgroup* of G , if H is a subgroup of G and the uncountable homomorphic image of \mathbb{H} .

It will sometimes be convenient to talk about a generating set, in the sense of infinite products, for a (pseudo-)Hawaiian earring subgroup of $\pi_1(X, x_0)$. For each (pseudo-)Hawaiian earring subgroup $H \in \pi_1(X, x_0)$, we may choose an (homomorphism) isomorphism $\varphi : \mathbb{H} \rightarrow H$. We will say that $\{\varphi(a_i)\}$ *generates (in the sense of infinite products) H* and will write $H = \langle\langle \varphi(a_n) \rangle\rangle$.

In this section, we will show that there exists a natural equivalence relation on pseudo-Hawaiian earring subgroups. Natural in the sense that it is determined by the topology and determines the topology of $B(X)$.

A natural question is what are the possible isomorphism types of pseudo-Hawaiian earring subgroups. Conner has conjectured that every uncountable homomorphic image of an Hawaiian earring group which embeds in the fundamental group of a planar or one-dimensional Peano continuum is a Hawaiian earring group.

4.1 Equivalence classes of pseudo-Hawaiian earring subgroups

We are now ready to prove:

Theorem 4.17 ^{bad} *The homeomorphism type of $B(X)$ is completely determined by $\pi_1(X, x_0)$, where $B(X)$ is the subspace of points at which X is not semilocally simply connected.*

Definition 4.4. We will say that x is a *wedge point* for $H \subset \pi_1(X, x_0)$ a pseudo-Hawaiian earring subgroup if, there exists a surjective homomorphism $\psi : \mathbb{H} \rightarrow H$ with the property that $\psi = \hat{\alpha} \circ f$ for a continuous function $f : \mathbb{E} \rightarrow X$ such that $f(0) = x$.

The following two lemmas are immediate from the definitions.

sur **Lemma 4.5.** *If X is a planar Peano continuum and $x \in B(X)$, then there exists H a pseudo-Hawaiian earring subgroup of $\pi_1(X, x_0)$ with x as its wedge point.*

wedgepoint **Lemma 4.6.** *If x is a wedge point for H a pseudo-Hawaiian earring subgroup, then x is a wedge point for $\hat{\alpha}(H)$.*

We should show the existence and uniqueness of wedge points.

Lemma 4.7. *There exists a unique wedge point for each pseudo-Hawaiian earring subgroup.*

Proof. The existence follows Theorem ^{contplanar} 2.22.

Suppose the a pseudo-Hawaiian earring subgroup H has two distinct wedge points x and x' which correspond to two homomorphism ψ and ψ' with image H . Then there exists $f : (\mathbf{E}, 0) \rightarrow (X, x)$, $f' : (\mathbf{E}, 0) \rightarrow (X, x')$, $\alpha : (I, 1) \rightarrow (X, x)$, and $\alpha' : (I, 1) \rightarrow (X, x')$ such that $\hat{\alpha} \circ f_* = \psi$ and $\psi' = \hat{\alpha}' \circ f'_*$.

Let U and U' be disjoint disks in the plane whose boundaries are not contained in X such that $U \cap X$ and $U' \cap X$ are neighborhoods of x and x' respectively. Since f and f' are continuous, the image of $f(a_i)$ is eventually contained in $U \cap X$ and the image of $f'(a_i)$ is eventually contained in $U' \cap X$. Hence there exists N such that $f_*(\mathbb{H}_N)$ is contained in $U \cap X$ and $f'_*(\mathbb{H}_N)$ is contained in $U' \cap X$.

A countability argument shows that $\psi(\mathbb{H}_n) \cap \psi'(\mathbb{H}_n)$ is nontrivial for all n . However, by the previous paragraph any element in the intersection for $n > N$ can then be freely homotoped from U to U' which contradicts Lemma ^{homotoped off} A6. □

This lemma also implies that the definition of a wedge point is independent of our choice of homomorphism.

We will let \mathcal{H} be the set of pseudo-Hawaiian earring subgroups of $\pi_1(X, x_0)$. Then we can now use the previous lemmas to define $\Upsilon : \mathcal{H} \rightarrow B(X)$, by taking a pseudo-Hawaiian earring subgroup to its wedge point. This function is surjective by Lemma [4.5](#)^{sur}. However, Lemma [4.6](#)^{conjugatewedgepoint} implies that it is highly non-injective. We will now define an equivalence relation on \mathcal{H} to make Υ injective on equivalency classes.

Definition 4.8. Let H_1, H_2 be pseudo-Hawaiian earring subgroups of $\pi_1(X, x_0)$. Then we will say that H_1 is *similar* to H_2 if there exists a $g \in \pi_1(X, x_0)$ and an $H \in \pi_1(X, x)$ a pseudo-Hawaiian earring subgroup such that $H_1, gH_2g^{-1} \subset H$. We will write $H_1 \sim H_2$.

If X is one-dimensional these equivalence classes are the same as maximal compatible nonempty subfamilies of the set of subgroups of $\pi_1(X, x_0)$ in [\[CE\]](#)^{ce}.

The following is well know.

art **Theorem 4.9.** For $\psi : \mathbb{H} \rightarrow \mathbb{F}$ a homomorphism where \mathbb{F} is a free group, there exists an i such that ψ factors through P_{i*} .

If we consider homomorphisms from the natural inverse limit containing \mathbb{H} to free groups, then this is a theorem of Higman [\[Hig\]](#)^{hig}. Morgan and Morrison also gave a topological proof in [\[MM\]](#)^{mm}. When we consider homomorphism from \mathbb{H} , this is a result of Theorem 4.4 in [\[CC2\]](#)^{cc3} and a proof can be found in [\[S\]](#)^{cs}.

ifwedge **Lemma 4.10.** If H_1 and H_2 have the same wedge point, then $H_1 \sim H_2$.

Proof. Let x be a wedge point for H_1 and H_2 . Let $\psi_i : \mathbb{H} \rightarrow \pi_1(X, x_0)$ be a homomorphism with image H_i for $i \in \{1, 2\}$. Then there exists continuous functions $f_i : (\mathbf{E}, 0) \rightarrow (X, x)$ and $\alpha_i : (I, 1) \rightarrow (X, x)$ such that $\widehat{\alpha}_i \circ f_{i*} = \psi_i$.

Let $g : (\mathbf{E}, 0) \rightarrow (X, x)$ be sending \mathbf{E}_o to $f_1(\mathbf{E})$ and \mathbf{E}_e to $f_2(\mathbf{E})$, i.e. a_i goes to $f_1(a_{\frac{i+1}{2}})$ for odd i and $f_2(a_{\frac{i}{2}})$ for even i .

Hence $\widehat{\alpha}_i(H_i) \subset f_*(\mathbb{H})$ and $f_*(\mathbb{H})$ is a pseudo-Hawaiian earring subgroup. Then after conjugating H_2 by $\alpha_1 * \overline{\alpha_2}$ and $f_*(\mathbb{H})$ by α_1 , we have necessary inclusion to show that $H_1 \sim H_2$. \square

onlyifwedge **Lemma 4.11.** If $H_1 \sim H_2$, then H_1 and H_2 have the same wedge point.

Proof. It is sufficient to show that if H_1 and H_2 are pseudo-Hawaiian earring subgroups with the property that $H_1 \subset H_2$, then H_1 and H_2 have the same wedge point. There exists continuous functions $f_i : (\mathbf{E}, 0) \rightarrow (X, x_i)$ and paths $\alpha_i : (I, 0, 1) \rightarrow (X, x_0, x_i)$ such that $\widehat{\alpha}_i \circ f_i(\mathbb{H}) = H_i$.

Suppose that $f_1(0) = x_1 \neq x_2 = f_2(0)$. By Lemma [A5](#)^{alem2}, we may find r_i such that $B_{2r_1}^X(x_1) \cap B_{2r_2}^X(x_2) = \emptyset$. Then we may choose N such that $f_i(\mathbf{E}_N) \subset B_{r_i}^X(x_i)$ for $i = 1, 2$.

Let $X' = X \cup B_{r_2}^{\mathbb{R}}(x_2)$. Let $f'_i : (\mathbf{E}, 0) \rightarrow (X', x_i)$ denote the map obtained by extending the range of f_i and $H'_i = \widehat{\alpha}_i \circ f'_i(\mathbb{H})$. Then H'_2 is countable which implies that H'_1 is countable. Hence, $f_{2*}(a_n)$ is eventually trivial.

Fix $n > N$ such that $f'_2(\mathbf{a}_n)$ is null-homotopic. Choose a continuous map $h : \mathbb{D} \rightarrow X'$ such that $h|_{\partial\mathbb{D}} = f'_2(\mathbf{a}_n)$. Hence by Lemma [A7](#), h can be modified to a map with image contained in $B_{2r_2}^{X'}(x_2)$. Since $B_{2r_1}^X(x_1)$ and $B_{2r_2}^X(x_2)$ are disjoint, h maps into X and $f_2(\mathbf{a}_n)$ is null-homotopic. This holds for any $n > N$; thus $f_{2*}(\mathbb{H})$ is countable which contradicts our choice that H_2 was a pseudo-Hawaiian earring group.

Therefore $x_1 = x_2$. □

Clearly, \sim is reflexive and symmetric. Then Lemmas [4.I0](#) and [4.I1](#) show that \sim is transitive. Hence, \sim is actually an equivalency relation. We will use $[H]$ to denote the equivalency class of pseudo-Hawaiian earring subgroups generated by the equivalency relation \sim .

We can now give an equivalent definition for the wedge point of a pseudo-Hawaiian earring subgroup. Let $H = \langle\langle g_n \rangle\rangle$ be a pseudo-Hawaiian earring subgroup. Then the wedge point of H is the unique point, x , which enjoys the following property: *For every $\epsilon > 0$, there exists an N such that the set of homotopy classes $\{g_n\}_{n>N}$ have representatives contained in $B_\epsilon(x)$.*

bijjective

Proposition 4.12. *$B(X)$ corresponds bijectively to equivalence classes of pseudo-Hawaiian earring subgroups of $\pi_1(X, x_0)$, i.e. $\Upsilon^{-1}(x) = [H]$*

Then it makes sense to talk about the *wedge point of a pseudo-Hawaiian earring subgroup equivalence class* being the wedge point of any of its members.

Definition 4.13. Let $\{[H_n]\}$ be a sequence of equivalency classes of pseudo-Hawaiian earring subgroups of the fundamental group of a planar Peano continuum. We will say that $[H_n]$ *converges* to $[H]$, a pseudo-Hawaiian earring subgroup equivalence class, if there exists a choice of representatives $\langle\langle g_{n,i} \rangle\rangle_i \in [H_n]$ and a sequence of natural numbers $\{M_n\}$ so that for any sequence $\{k_n\}$, with $k_n > M_n$ the pseudo-Hawaiian earring subgroup $\langle\langle g_{n,k_n} \rangle\rangle_n \in [H]$. We will use all the standard terminology when referring to sequence of pseudo-Hawaiian earring subgroup equivalence classes.

Actually since we have to be able to choose the representative for the equivalence class, we may as well choose the representative such that $M_n = 1$. This can be done by removing the first finitely many big loops which will not change the equivalence class.

Initially there is no reason to suppose that a convergent sequence has a unique limit point. However, this will be a consequence of Proposition [4.14](#) and Lemma [4.I0](#).

limitpoint

Proposition 4.14. *If $[H_n] \rightarrow [H]$, then $x_n \rightarrow x$ where x_n is the wedge point of $[H_n]$ and x is the wedge point of $[H]$.*

Proof. Suppose that $[H_n]$ *converges* to $[H]$ a pseudo-Hawaiian earring subgroup equivalence class. Then there exists a choice of representatives $\langle\langle g_{n,i} \rangle\rangle_i \in [H_n]$ and $\{M_n\}$ such that if $k_n \geq M_n$, then $\langle\langle g_{n,k_n} \rangle\rangle_n \in [H]$.

We will proceed by way of contradiction. Let x_n be the wedge point for $[H_n]$. Suppose that $x_n \not\rightarrow x$. Then there is a subsequence x_{n_i} with the property that $x_{n_i} \rightarrow x' \neq x$. Fix A, A' disjoint open disk neighborhoods of x and x' respectively, whose boundaries are not contained in X . Then we may choose $\{M'_{n_i}\}_i$ and N' such that g_{n_i, k_i} can be freely homotoped into A' for $n_i > N'$ and $k_i > M'_{n_i}$. Let $M'_k = 0$ if $k \neq n_i$ for any i .

Fix $g_{n, k(n)}$ such that $k(n) > \max\{M'_n, M_n\}$. Then $\langle\langle g_{n, k(n)} \rangle\rangle_n \in [H]$ and there exists an N such that, for $n > N$, $g_{n, k(n)}$ can be freely homotoped into A . Then for $n_i > \max\{N', N\}$ $g_{n_i, k(n_i)}$ can be freely homotoped into A and A' , a contradiction. \square

The uniqueness of the limit of a convergent sequence then follows by considering the corresponding sequence of wedge points.

Let $\overline{\mathcal{H}}$ be the set of equivalency classes of pseudo-Hawaiian earring subgroups. We will now construct a topology on $\overline{\mathcal{H}}$.

Definition 4.15. We will say that $A \subset \overline{\mathcal{H}}$ is *closed* if it contains all of its limit points. Then a set is *open* in $\overline{\mathcal{H}}$ if its complement is closed.

It is an easy exercise to show that this defines a topology on $\overline{\mathcal{H}}$.

Let $\overline{\Upsilon} : \overline{\mathcal{H}} \rightarrow B(X)$ by $\overline{\Upsilon}([H]) = \Upsilon(H)$. This is well defined by Lemma 4.11 and a ^{only if wedge} bijection by Proposition 4.12. ^{bijective}

Proposition 4.14 ^{limitpoint} is exactly what is required for $\overline{\Upsilon}$ to be continuous. Suppose $C \subset B(X)$ which is closed. Consider $\{[H_n]\} \subset \overline{\Upsilon}^{-1}(C)$ which converges to $[H]$. If x_n is the wedge point of $[H_n]$, then x_n is in C and x_n must converge to x the wedge point of $[H]$. Thus $x \in C$ and $[H] \in \overline{\Upsilon}^{-1}(C)$.

The next lemma will give the continuity of $\overline{\Upsilon}^{-1}$ immediately since X is metric.

Lemma 4.16. *If $x_n \in B(X)$ and $x_n \rightarrow x$, then the corresponding pseudo-Hawaiian Earring subgroup equivalency classes converge.*

Proof. Let H_n be a pseudo-Hawaiian Earring subgroup of $\pi_1(X, x_0)$ with wedge point x_n . Then there exists a continuous function $f_n : (\mathbf{E}, 0) \rightarrow (X, x_n)$ which induces an injective homomorphism. There exists a path $T : I \rightarrow X$ with the property that $T(0) = x$, $T(1) = x_1$, and $T(\frac{1}{n}) = x_n$. Let $\alpha : I \rightarrow X$ be any path from x to x_0 . Let β_n be a parametrization of the path $\alpha * T|_{[0, \frac{1}{n}]}$. Then $\langle\langle \overline{\beta_n} * f_n(a_i) * \beta_n \rangle\rangle_i$ has wedge point x_n . Let $g_{n, i} = \overline{\beta_n} * f_n(a_i) * \beta_n$. There exists M_n such that $f_n(a_i) \subset B_{\frac{1}{n}}(x_n)$ for $i > M_n$. If $k \geq M_n$ for each n , then $\langle\langle g_{n, k} \rangle\rangle_n \in [H]$. \square

Therefore we have now proved the following result.

bad **Theorem 4.17.** *The function $\overline{\Upsilon} : \overline{\mathcal{H}} \rightarrow B(X)$ is a homeomorphism, i.e. the homeomorphism type of $B(X)$ is completely determined by $\pi_1(X, x_0)$.*

APPENDIX

Lemma A1. *Every non-degenerate null-homotopic loop in a one-dimensional space contains a proper non-degenerate subloop which is also null-homotopic.*

Proof. Suppose that $f : S^1 \rightarrow X$ is a non-degenerate null-homotopic loop in a one-dimensional space X . Then f factors through a map $f' : S^1 \rightarrow D$ where D is a one-dimensional planar dendrite (see [CC2, Theorem 3.7]). Since $f(S^1)$ is non-degenerate, so is D and there exists $x \in D$ such that $D \setminus \{x\}$ has at least two components. Hence $S^{-1} \setminus \{f'^{-1}(x)\}$ has at least two components. Let A be the closure of a component of $S^{-1} \setminus \{f'^{-1}(x)\}$. Then $f|_A$ is a proper null-homotopic subpath of f . Since $f'|_A$ maps onto the closure of a component of $D \setminus \{x\}$ and the components of $D \setminus \{x\}$ are non-degenerate, $f|_A$ is non-degenerate. \square

Lemma A2. *Suppose that $f : [0, 3] \rightarrow X$ is a path into a one-dimensional space X such that $f|_{[i, i+2]}$ is a reduced path for $i \in 0, 1$. If $f|_{[1, 2]}$ is a non-degenerate then f is reduced.*

Proof. Suppose $f|_{[s, t]}$ is a non-degenerate null-homotopic subloop of f . Since $f|_{[i, i+2]}$ is a reduced path for $i \in 0, 1$; $s \leq 1 < 2 \leq t$.

If $s_1 \leq s_2 \leq 1 < 2 \leq t_1 \leq t_2$ and $f|_{[s_i, t_i]}$ is a non-degenerate null-homotopic subloop of f , then $f|_{[s_2, t_1]}$ is a non-degenerate null-homotopic subloop of f .

Suppose that $s_1 \leq s_2 \leq \dots \leq 1 < 2 \leq \dots \leq t_2 \leq t_1$. Let $s = \lim s_i$ and $t = \lim t_i$. It is not hard to see that $f|_{[s, t]}$ is also a non-degenerate null-homotopic subloop of f (for details see the proof of Theorem 3.9 in [CC2]).

Then there exists a unique minimal interval $[s_0, t_0]$ such that $f|_{[s_0, t_0]}$ is a non-degenerate null-homotopic subloop of f . But then f contains no a proper non-degenerate null-homotopic subloops which contradicts the previous lemma. \square

Greg Conner and Mark Meilstrup proved the following lemma in [CM].

aalem1 **Lemma A3.** *Let H be a function from the first-countable space $X \times Y$ into Z . Let $\{C_i\}$ be a null sequence of closed sets whose union is X . Suppose that $\{D_i = H(C_i \times Y)\}$ is a null sequence of sets in Z and H is continuous on each $C_i \times Y$. If for every subsequence $C_{i_k} \rightarrow x_0$ there exists a $z_0 \in Z$ such that $D_{i_k} \rightarrow z_0$ then H is continuous on $X \times Y$.*

Proof. Consider a sequence $(x_n, y_n) \rightarrow (x_0, y_0)$. For each n , choose an i_n such that $x_n \in C_{i_n}$. If $\{C_{i_n}\}$ is finite then by restricting H to $\cup_n C_{i_n} \times Y$ we have $H(x_n, y_n) \rightarrow H(x_0, y_0)$ be a finite application of the pasting lemma. If $\{C_{i_n}\}$ is infinite, then since $\{C_i\}$ is a null sequence and $x_n \in C_{i_n}$, we have $C_{i_n} \rightarrow x_0$ and thus $H(x_n, y_n) \in D_{i_n} \rightarrow z_0 = H(x_0, y_0)$. Thus H is continuous on all of $X \times Y$. \square

alem1 **Lemma A4.** *If A is an annulus and C is a closed subset of $\text{int } A$ that separates the boundary components J_1 and J_2 of A , then some component of C separates J_1 from J_2 in A .*

This is a consequence of Phragmén-Brouwer properties. (see ^{hw}[HW].)

Let $B_r^X(x) = \{y \in X \mid d(x, y) < r\}$ and $S_r^X(x) = \{y \in X \mid d(x, y) = r\}$. If X is a planar set then $B_r^X(x) = B_r^{\mathbb{R}^2}(x) \cap X$ and $S_r^X(x) = S_r^{\mathbb{R}^2}(x) \cap X$.

alem2 **Lemma A5.** *If X is a planar set which is not semilocally simply connected at x , then there exists arbitrarily small r such that $S_r^{\mathbb{R}^2}(x) \not\subset X$.*

Proof. Suppose that there exists an $\epsilon > 0$ such that for all $r < \epsilon$, $S_r^{\mathbb{R}^2}(x) \subset X$. Then $B_\epsilon^{\mathbb{R}^2}(x) \subset X$ which implies that $B_\epsilon^{\mathbb{R}^2}(x) = B_\epsilon^X(x)$ and thus X is simply connected at x . \square

homotoped off **Lemma A6.** *Let X be a subset of \mathbb{R}^2 and U a closed disk in the plane whose boundary is not contained in X . If α is an essential loop contained in $X \cap \text{int}(U)$, then α cannot be freely homotoped out of $U \cap X$.*

Proof. Let X be a subset of \mathbb{R}^2 and U a closed disk in the plane whose boundary is not contained in X . Suppose α is an essential loop contained in $X \cap \text{int}(U)$ such that α can be homotoped out of $U \cap X$.

Let A be an annulus with boundary components J_1 and J_2 . Then there exists a map $h : A \rightarrow X$ which takes J_1 to α and J_2 to a loop in X not intersecting $U \cap X$. Then $h^{-1}(\partial U \cap X)$ is a closed subset contained in the int A which separates J_1 from J_2 in A . Hence by Lemma ^{alem1}A4, some component C of $h^{-1}(\partial U \cap X)$ separates J_1 from J_2 in A . Since $\partial U \not\subset X$, $h|_C$ maps to an arc in X . Hence by Tietze Extension Theorem, we may alter h to a map of a disk into X . Therefore α is null-homotopic, a clear contradiction. \square

cutoff **Lemma A7.** *Suppose the $f : \partial \mathbb{D} \rightarrow X$ is a continuous map into a planar set. If f is null-homotopic then there exists a map $h : \mathbb{D} \rightarrow X$ such that $\text{diam } h(\mathbb{D}) \leq 2 \text{diam } f(\partial \mathbb{D})$.*

Proof. Let $x_0 \in f(\partial \mathbb{D})$ and $r_0 = \text{diam } f(\partial \mathbb{D})$. Let $r_1 = \inf\{r \geq r_0 \mid S_r^{\mathbb{R}^2}(x_0) \not\subset X\}$. Then $S_{r_1}^X(x)$ is the disjoint union of closed intervals (some of which are possible degenerate). Let $h : \mathbb{D} \rightarrow X$ be a null homotopy of f . Consider the components of $h^{-1}(S_{r_1}^X(x))$. Then as before we can cut h off at $S_{r_1}^X(x)$. Then the modified h can be contracted into $B_{r_0}^X(x_0)$ which completes the proof. \square

alem3 **Lemma A8.** *If $x_n \rightarrow x$ in a Peano continuum, then there exists a path $T : I \rightarrow X$ with the property that $T(0) = x_1$, $T(1) = x$, and $T(1 - \frac{1}{n}) = x_n$.*

Proof. Let U_n be a path connected neighborhood of x contained in $B_{\frac{1}{n}}(x) \cap U_{n-1}$. Let $T_n : [1 - \frac{1}{n}, 1 - \frac{1}{n+1}] \rightarrow X$ be any path from x_n to x_{n+1} with the property that if $x_n, x_{n+1} \in U_k$ then $T_n([1 - \frac{1}{n}, 1 - \frac{1}{n+1}]) \subset U_k$. T_n will exist since there is a minimal U_k containing both x_n and x_{n+1} . Then $T : I \rightarrow X$ by $T(x) = T_n(x)$ for $x \in [1 - \frac{1}{n}, 1 - \frac{1}{n+1}]$ and $T(1) = x$ is a continuous path. \square

References

- [ADTW] S. Akiyama and G. Dorfer and J.M. Thuswaldner R. Winkler, *On the fundamental group of the Sierpiński-gasket*, *Topology Appl.* 156 (2009), 1655–1672.
- [CC1] J.W. Cannon and G.R. Conner, *The combinatorial structure of the Hawaiian earring group*, *Topology Appl.* 106 (2000), 225–271.
- [CC2] J.W. Cannon and G.R. Conner, *On the fundamental groups of one-dimensional spaces*, *Topology Appl.* 153 (2006), 2648–2672.
- [CC3] J.W. Cannon and G.R. Conner, *The homotopy dimension of codiscrete subsets of the 2-sphere S^2* , *Fund. Math.* 197 (2007), 35–66.
- [CCZ] J.W. Cannon and G.R. Conner and A. Zastrow, *One-dimensional sets and planar sets are aspherical*, *Topology Appl.* 120 (2002), 23–45.
- [CE] G. Conner and K. Eda, *Fundamental groups having the whole information of spaces*, *Topology Appl.* 146/147 (2005), 317–328.
- [CM] G. Conner and M. Meilstrup, *Deforestation of Peano continua and minimal deformation retracts*, *Topology Appl.* 159 (2012), 3253–3262.
- [CF] M.L. Curtis and M.K. Fort, *Singular homology of one-dimensional spaces*, *Ann. of Math.* (2) 69 (1959), 309–313.
- [E1] Katsuya Eda, *Free σ -products and fundamental groups of subspaces of the plane*, *Topology Appl.* 84 (1998), 283–306.
- [E2] Katsuya Eda, *The fundamental groups of one-dimensional spaces and spatial homomorphisms*, *Topology Appl.* 123 (2002), 479–505.
- [E3] Katsuya Eda, *Homotopy types of one-dimensional Peano continua*, *Fund. Math.* 209 (2010), 27–42.
- [H] G. Higman, *Unrestricted free products, and varieties of topological groups*, *J. London Math. Soc.* 27 (1952), 73–81.
- [HW] W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton Mathematical Series, v. 4, Princeton University Press, Princeton, N. J., 1941.
- [K] C. Kent, *Homomorphisms into the fundamental group of one-dimensional and planar Peano continua*, Masters Thesis (2008).

- [K1] C. Kent, *A note on homomorphisms from the fundamental group of planar sets*, preprint.
- [vM] J. van Mill, *Infinite-dimensional topology*, North-Holland Mathematical Library, North-Holland Publishing Co., Amsterdam, 1989.
- [dS] B. de Smit, *The fundamental group of the Hawaiian earring is not free*, *Internat. J. Algebra Comput.* 2 (1992), 33–37.
- [MM] J. Morgan and I. Morrison, *A van Kampen theorem for weak joins*, *Proc. London Math. Soc.* (3) 53 (1986), 562–576.
- [S] E. Summers, *There is no surjective map from the hawaiian earring group to the double hawaiian earring group*, Masters Thesis (2002).