

# Origin of craters on Phoebe: comparison with Cassini's data.

R. P. Di Sisto <sup>\*</sup> and A. Brunini

Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata and  
Instituto de Astrofísica de La Plata, CCT La Plata-CONICET-UNLP  
Paseo del Bosque S/N (1900), La Plata, Argentina.

Received / Accepted

## ABSTRACT

*Context.* Phoebe is one of the irregular satellites of Saturn; the images taken by Cassini-Huygens spacecraft allowed us to analyze its surface and the craters on it.

*Aims.* We study the craters on Phoebe produced by Centaur objects from the Scattered Disk (SD) and plutinos escaped from the 3:2 mean motion resonance with Neptune and compare our results with the observations by Cassini.

*Methods.* We use previous simulations on trans-Neptunian Objects and a method that allows us to obtain the number of craters and the cratering rate on Phoebe.

*Results.* We obtain the number of craters and the greatest crater on Phoebe produced by Centaurs in the present configuration of the Solar System. Moreover, we obtain a present normalized rate of encounters of Centaurs with Saturn of  $\dot{F} = 7.1 \times 10^{-11}$  per year, from which we can infer the current cratering rate on Phoebe for each crater diameter.

*Conclusions.* Our study and the comparison with the observations suggest that the main crater features on Phoebe are unlikely to have been produced in the present configuration of the Solar System and that they must have been acquired when the SD were depleted in the early Solar System. If this is what happened and the craters were produced when Phoebe was a satellite of Saturn, then it had to be captured, very early in the evolution of the Solar System.

**Key words.** methods: numerical – Kuiper Belt: general

## 1. Introduction

Phoebe is one of the irregular satellites of Saturn. It has a retrograde orbit, which suggests that it was captured by Saturn instead of being formed “in situ” (e.g Pollack et al. 1979). Moreover, Phoebe’s composition is close to that derived from bodies such as Triton and Pluto, and it is different from that of the regular satellites of Saturn supporting Phoebe’s origin as a captured body of the outer Solar System (Johnson & Lunine 2005).

On 11 June 2004, Cassini-Huygens spacecraft encountered Phoebe in a fly-by within 2000 km of Phoebe’s surface. This encounter allowed Cassini to analyze Phoebe’s surface in detail

---

Send offprint requests to: R. P. Di Sisto

\* romina@fcaglp.unlp.edu.ar

which improved the previous Voyager data. Buratti et al. (2004) analyzed and characterized the physical properties of the surface from photometric data from Cassini VIMS (Visual and Infrared Spectrometer), concluding that it is rough and dusty, perhaps from a history of out-gassing or a violent collisional history suggested by Nesvorny et al. (2003). Hendrix & Hansen (2008) analyzed the first UV spectra of Phoebe with the Cassini Ultraviolet Imaging Spectrograph (UVIS) during the Cassini spacecraft fly-by and detected water ice. Using VIMS data, Cruikshank et al. (2004) concluded that Phoebe's surface is rich in organics, which is compatible with the low albedo of Phoebe. Porco et al (2005) calculated a mean density of Phoebe of  $1.63 \text{ gr/cm}^3$  from calculation of the volume and the determination of the mass from tracking the spacecraft. If Phoebe's surface was a mixture of rock and ice, the previous density is compatible with a porosity lower than  $\sim 40\%$  (Porco et al. 2005). Johnson and Lunine (2005) analyzed the relation between composition and probable porosity of Phoebe and they found that if Phoebe was derived from the same compositional reservoir as Pluto and Triton, Phoebe's measure density is consistent with a porosity of  $\sim 15\%$ .

Giese et al. (2006) presented the results of a photogrametric analysis of the high-resolution stereo images of Phoebe. In particular they obtained a mean figure radius of 107.2 km and a digital terrain model of the surface reveals significant morphological detail. The images revealed that Phoebe basically exhibits simple crater shapes with the only exception of the greatest impact crater Jason with a diameter of  $\sim 100$  km. Several of the smaller craters present pronounced conical shapes which could indicate the presence of porous, low compacting material on the surface of Phoebe.

Kirchoff & Schenk (2010) reexamined the impact of crater distribution of the mid-sized saturnian satellites. For Phoebe they found that the crater size frequency distribution has relatively constant values for crater diameters  $D \leq 1$  km, but then it has a sudden and confined dip around  $D \sim 1.5$  km. Beyond this dip, the crater size frequency distribution has a slow increase. This behaviour is unique in the saturnian satellite system and it is probably connected with Phoebe's origin.

Zhanle et al. (2003) write a previous paper that calculates cratering rates in the satellites of the outer planets. They used impact rates on the giant planets obtained by Levison & Duncan (1997) and independent constraints on ecliptic comets. Their results will be compared with ours.

As we have seen, the origin of craters on Phoebe is unclear. But the main population of objects that can produce craters on Phoebe are Centaurs, since they are the small body objects that cross the orbit of the giant planets, in particular the orbit of Saturn, and then its satellites.

Centaurs are transient bodies between their source in the trans-Neptunian population and the Jupiter Family Comets. They come mainly from a sub population in the trans-Neptunian zone, the Scattered Disk Objects (SDOs). The SDOs are bodies with perihelion distances  $q$  greater than 30 AU and smaller than  $\sim 39$  AU that can cross the orbit of Neptune and eventually evolve into the giant planetary zone, crossing the orbits of those planets, and then the orbits of their satellites (Di Sisto & Brunini 2007, Levison & Duncan 1997). The secondary source of Centaurs are plutinos and the low eccentricity trans-Neptunian objects (Di sisto et al. 2010, Levison & Duncan 1997). Plutinos are those trans-Neptunian objects located in the 3:2 mean motion resonance with Neptune at  $a \sim 39.5$  AU. They are "protected" by the 3:2 mean motion resonance with Neptune but some of them are long term escapers that are presently escaping from the resonance (Morbidelli 1997). In

this paper we will study the production of craters on Phoebe from Centaur objects from SDOs and plutinos escaped from the 3:2 mean motion resonance with Neptune, as the two main populations of impactors. We use here previous simulations on trans-Neptunian Objects (TNOs) and a method that allows us to obtain directly the cratering rate on Phoebe. This study, and the comparison with the observations of Cassini images may help us to determine the origin of craters on Phoebe, in order to determine the history of Phoebe's surface and also constrain its origin.


## 2. The number of SDOs

Cratering rates depend on the number and sizes of the impactor population. Thus we must know the real initial number of SDOs to calculate the total number of collisions on Phoebe. Then, we are going to estimate the total number of present SDOs.

Parker & Kavelaars (2010a) re-characterized the orbital sensitivity of several published pencil-beam surveys. They found that these surveys were sensitive to distant populations like SDOs and Sedna-like objects. Using this result Parker & Kavelaars (2010b) derived new upper limits on those distant populations and used this new limits to obtain the number of SDOs. To do this they performed a model that considered two laws for the radial distance distribution of SDOs. On the one hand they took a radial distance distribution of SDOs  $\propto r^{-1.5}$  and obtained a maximum population of  $N(d > 100 \text{ km}) = 3.5 \times 10^5$ . On the other hand they took a uniform radial distance distribution obtaining in this case a maximum population of  $N(d > 100 \text{ km}) = 25 \times 10^5$ . In this paper we take the number of SDOs greater than  $d = 100 \text{ km}$  equal to  $N(d > 100 \text{ km}) = 3.5 \times 10^5$  since the considered radial distance distribution is consistent with the one obtained by Di Sisto & Brunini (2007). Then the total population of SDOs with diameter greater than  $d_0$  will be given by  $N(d > d_0) = 3.5 \times 10^5 (d_0/100)^{-s+1}$ , where  $d_0$  must be in km and  $s$  is the index of the differential size distribution. There are some authors who have found a single power law size distribution for TNOs (Petit et al. 2000, Fraser et al. 2008). However, other papers suggest that the size distribution function (SDF) of TNOs could have a break at a diameter of  $\sim 60 \text{ km}$  (Bernstein et al. 2004, Gil Hutton et al. 2009, Fraser & Kavelaars 2009, Fuentes & Holman 2008, Fuentes et al. 2009). The differential power law indexes for smaller TNOs (this is  $d < 60 \text{ km}$ ) found by those surveys are  $s_2 = 2.8, 2.4, 1.9, 2.5$  and  $2$ , respectively. It seems to be enough evidence for a break in the size distribution of the TNO population. In particular we are going to assume that this break is also valid for all the dynamical classes of TNOs.

Elliot et al. (2005) accounts for the SDF for each dynamical class in the TN region. Specifically for SDOs they found that the differential size distribution index is  $s_1 = 4.7$  for the brightest objects. Then taking into account the assumed break in the SDF of SDOs we consider that the power law SDF of SDOs breaks at  $d \sim 60 \text{ km}$  to an index of between 3.5 and 2.5. We analyze those indexes as limit cases that give higher and lower ranges for the population of SDOs and then the production of craters on Phoebe. The higher value of  $s_2 = 3.5$  corresponds to a population in steady-state (Dohnanyi 1969) which could be the case for the smallest SDOs (Gil Hutton et al. 2009). Considering all this, the number of SDOs greater than a diameter  $d_0$  will be given by

$$\begin{aligned} N(d > d_0) &= C_0 \left( \frac{1\text{km}}{d} \right)^{s_2-1} & \text{for} & \quad d \leq 60\text{km}, \\ N(d > d_0) &= 3.5 \times 10^5 \left( \frac{100\text{km}}{d} \right)^{s_1-1} & \text{for} & \quad d > 60\text{km}, \end{aligned} \quad (1)$$



**Fig. 1.** Cumulative number of SDOs and Plutinos according to the size distribution laws described in the text.

where  $C_0 = 3.5 \times 10^5 100^{s_1-1} (60)^{s_2-s_1}$  by continuity for  $d = 60$  km,  $s_1 = 4.7$  and  $s_2 = 2.5$  and  $3.5$ . This law is plotted in Fig. 1 with the two breaks considered.

### 3. SDO collisions on Phoebe

To study the collisions of SDOs on Phoebe and the contribution of that population to the cratering history of the satellite we use some of the outputs of the numerical simulation performed in a previous paper by Di Sisto & Brunini (2007). In that work we integrated numerically 1000 objects from the SD (95 real + 905 fictitious) and studied their evolution in the Centaur zone under the gravitational action of the Sun and the four giant planets. The computations were followed for 4.5 Gyr, or until the test body collided with a planet, was ejected, or entered the region inside Jupiter's orbit ( $r < 5.2$  AU). In that paper we also stored in a file the encounters of the fictitious SDOs with the planets and registered the time of the encounter, the minimum distance to the planet ( $q$ ) and the relative velocity at this distance ( $v(q)$ ). From these data we can calculate, the number of encounters with Saturn within the Hill's sphere of the planet. Using the particle in a box approximation and assuming that the geometry of the encounters is isotropic it is possible to calculate the number of collisions on Phoebe ( $N_c$ ) through the relation:

$$\frac{N_c}{N_e} = \frac{v_i R_p^2}{v(R) R^2} \quad (2)$$

Where  $N_e$  is the number of encounters with Saturn inside its Hill's sphere of radius  $R$ ,  $R_p$  is the radius of Phoebe,  $v(R)$  is the mean relative encounter velocity of SDOs when entering the Hill's sphere of the planet and  $v_i$  is the collision velocity of SDOs on Phoebe.  $v(R)$  can be calculated from  $v(q)$  registered in our outputs from the relation:

$$v(R) = v^2(q) + 2Gm\left(\frac{1}{R} - \frac{1}{q}\right) \quad (3)$$

where  $G$  is the constant of gravitation and  $m$  is the mass of Saturn.

The collision velocity on Phoebe is computed assuming that the geometry of collisions is isotropic and then:

$$v_i = \sqrt{v_p^2 + v_0^2} \quad (4)$$

**Table 1.** Radius  $R_p$ , density  $\rho_p$ , orbital velocity  $v_p$  and superficial gravity times the mass  $gm_p$  of Phoebe. Mean relative velocity of SDOs when entering the Hill's sphere of Saturn  $v(R)$ , when they intersect the orbit of Phoebe  $v_0$  and when collide with Phoebe  $v_i$ .

$R_p$ [km]	107.2
$\rho_p$ [gr cm <sup>-3</sup> ]	1.634
$v_p$ [km s <sup>-1</sup> ]	1.71
$gm_p$ [km s <sup>-2</sup> ]	0.5532
$v(R)$ [km s <sup>-1</sup> ]	4.06
$v_0$ [km s <sup>-1</sup> ]	4.65
$v_i$ [km s <sup>-1</sup> ]	4.96

where  $v_p$  is phoebe's orbital velocity and  $v_0$  is the mean relative velocity of SDOs when they cross the orbit of Phoebe. This velocity was computed in the same way as  $v(R)$  appropriately using Eq. (3) from  $v(q)$  registered in our outputs. All the mentioned velocities and the radius and orbital velocity of Phoebe are shown in Table 1.

Eq. (2) provides the number of collisions on Phoebe in relation to the number of encounters with Saturn that we had registered in our previous run.

There are many papers based on theoretical and observational work which argue that the initial mass of the trans-Neptunian region was  $\sim 100$  times greater than the present mass and decay to nearly its present value at most in 1 Gyr (see e.g. Morbidelli et al. 2008). The simulation of Di Sisto & Brunini (2007) studies the evolution of SDOs in the present configuration of the Solar System; that is, when the SD is expected to have roughly reached its present mass and dynamic state,  $\sim 3.5$  Gyrs ago. Then, we can estimate the total number of collisions on Phoebe during the last  $\sim 3.5$  Gyrs rescaling Eq. (2) in order to account for the total SDOs population.

Of the 1000 initial particles of our previous simulation (Di Sisto & Brunini 2007), 368 underwent 10257 encounters within Saturn's Hill sphere. Therefore, the total number of encounters with Saturn of the whole SDO population in the present configuration of the Solar System is estimated as

$$N_{er} = \left(\frac{368}{1000}\right) \left(\frac{10257}{368}\right) N, \quad (5)$$

where  $N$  is the number of different SDOs which have existed, in the last 3.5 Gyrs and which can be obtained from Eq. (1). Here we assume that the present number of SDOs is roughly the same as it was 3.5 Gyrs ago. Consequently, the total number of encounters with Saturn of the whole SDO population through the last 3.5 Gyrs depending on the diameter is given by:

$$N_{er}(d > d_0) = \left(\frac{10257}{1000}\right) N(d > d_0) \quad (6)$$

From this equation and Eq. (2) the total number of collisions of SDOs on Phoebe over the last 3.5 Gyrs depending on the SDO's diameter, is given by:

$$N_c(d > d_0) = \frac{v_i R_p^2}{v(R) R^2} N_{er}(d > d_0) \quad (7)$$

Table 2 shows some values of  $N_c$  for certain diameters of the impactors.

Depending on the values of  $s_2$ , the diameter of the largest SDO impactor onto Phoebe during the last 3.5 Gyrs has been calculated as ranging from 110 mts to 1.36 km.

**Table 2.** Number of collisions of SDO impactors on Phoebe with diameters  $d > d_0$  that produce craters with diameter  $D > D_0$  ( $N_c(D > D_0)$ ).

$d_0[km]$	$D_0[km]$	$N_c(D > D_0)$
0.081	1	2 - 1180
0.445	5	0 - 16
0.969	10	0 - 2
3.48	30	0

#### 4. Craters on Phoebe by SDOs

The estimation of the size of a crater produced by a particular impact has been extensively studied. Schmidt & Housen (1987) present a set of power-law scaling relations for the crater volume based on laboratory experiments that simulate crater formation and point-source solutions. Holsapple (1993) also describe the scaling law for impact processes in a review work. The derived scaling laws allow us to link impacts of different sizes, velocities and superficial gravity and then obtain the size of a crater produced by a collision on a solar system body. Holsapple & Housen (2007) present the updated scaling laws for cratering in a recent work dedicated to interpret the observations of the Deep Impact event. These cratering laws are used here to calculate the craters on Phoebe. Thus, the diameter  $D$  of a crater produced by an impactor of diameter  $d$  can be obtained from the general equation (Holsapple & Housen 2007):

$$D = K_1 \left[ \left( \frac{gd}{2v_i^2} \right) \left( \frac{\rho_t}{\rho_i} \right)^{\frac{2\nu}{\mu}} + \left( \frac{Y}{\rho_i v_i^2} \right)^{\frac{2+\mu}{2}} \left( \frac{\rho_t}{\rho_i} \right)^{\frac{\nu(2+\mu)}{\mu}} \right]^{-\frac{\mu}{2+\mu}} d \quad (8)$$

$\rho_t$  being the target density,  $g$  its superficial gravity,  $Y$  its strength,  $\rho_i$  the density of the impactor and  $v_i$  the collision velocity. This impact cratering scaling law depends on two exponents,  $\mu$  and  $\nu$ , and a constant,  $K_1$ , that characterize the different materials. The first term in the square brackets is a measure of the importance of gravity in the cratering event and the second is a measure of the importance of the target strength. Thus, if the first term dominates on the second term, the crater is under the gravity regime, and if the second term dominates we have the strength regime. The partition between the two size scales of impacts depends on the size of the event (Holsapple 1993). Eq. (8) is a convenient empirical smoothing function to span the transition between the gravity regime and the strength regime (Holsapple, 1993). Since Phoebe is a small satellite with a relatively low gravity, the strength regime can be important for the smaller craters.

As Phoebe's density ( $1.63 \text{ gr/cm}^3$ ) is similar to sand and lower compacting material is found on its surface, we adopt  $K_1 = 1.03$ ,  $\mu = 0.41$  and  $\nu = 0.4$  that correspond to sand or cohesive soil in Holsapple & Housen (2007). This value of  $\mu$  corresponds to materials with a porosity of  $\sim 30-35\%$  (Holsapple & Schmidt 1987) which is compatible with the ranges of Phoebe's predicted porosity. The value for dry soils from Holsapple (1993), i.e.  $Y = 0.18 \text{ mpa}$ , is used for the strength.

The calculated densities of TNOs vary considerably from  $\sim 0.5- \sim 3 \text{ gr/cm}$ . Although a dimension-density trend has been suggested (Sheppard et al. 2008, Perna et al. 2009), more data are required to confirm it. In addition; as crater experiments do not include variations in the impactor material, there is no data to precisely determine the dependence on the impactor density (Schmidt & Housen 1987, Housen & Holsapple 2003). Therefore, we assume  $\rho_i = \rho_t$ , which is also between the ranges of calculated densities in the trans-Neptunian region. By taking all this into account,


the diameter of a crater on Phoebe for a given impactor diameter can be calculated from Eq. (8) through:

$$D = 1.03 \left[ \left( \frac{gd}{2v_i^2} \right) + \left( \frac{Y}{\rho_i v_i^2} \right)^{1.205} \right]^{-0.17} d \quad (9)$$

This equation describes simple bowl-shape craters but, as mentioned in the introduction, Cassini images of Phoebe reveal basically simple crater shapes with the only exception of Jason with a diameter of  $\sim 100$  km (Giese et al. 2006). Hence we use Eq. (9) for calculating the diameters of all craters on Phoebe without any further correction for transient-to-final size. By combining Eq. (7) and (9) it is possible to calculate the number of craters on Phoebe according to the diameter of the crater. Fig. (2) shows the cumulative number of craters, with diameters greater than a given value for the two size distribution power laws for smaller SDOs on Phoebe. Note that the different slopes in the number of craters for each curve is due to the difference in both indexes  $s_2$  considered. As we mentioned before, there is a limit impactor diameter that accounts for the transition between the gravity regime and the strength regime. This diameter can be obtained equating the first and second terms of Eq. (9). This limit impactor diameter is  $d_l = 367$  mts, which produce a limit crater of  $D_l = 4.2$  km. Thus, for crater diameters  $D < D_l$ , the production of craters on Phoebe is under the strength regime and for  $D > D_l$ , the production of craters is under the gravity regime. Note that  $D_l$  depends strongly on the assumed value of strength which is actually unknown. Then considering other values of the strength for less cohesive soils as terrestrial dry desert alluvium of  $Y = 65$  kpa (Holsapple & Housen 2007) and Surface Lunar regolith  $Y = 10$  kpa (Holsapple 2011),  $D_l$  could take the values 1.5 km and 233 mts respectively. We will consider in the following that  $D_l = 4.2$  km but it must be taken into account that  $D_l$  can be as small as 233 mts.

Since in the strength regime the crater diameter depends linearly on the impactor diameter, the relation between the cumulative number of craters on Phoebe and the crater diameter follows the same power law relation as that followed by the number of SDOs. For  $D < 4.2$  km the cumulative number of craters on Phoebe follows a power law with a cumulative index of 1.5 and 2.5, according to the value of  $s_2 = 2.5$  or  $s_2 = 3.5$  respectively. For  $D > 4.2$  km, this is in the gravity regime, the crater diameter does not depend linearly on the impactor diameter. Therefore, we fit a power law for the cumulative number of craters on Phoebe depending on the crater diameter of index 2.8. Kirchoff & Schenk (2010) found that the crater size frequency distribution for Phoebe has a cumulative index of 2.348 for  $D = 0.15 - 1$  km and 1 for  $D = 1 - 4$  km. We can see that, for very small craters, this index is very similar to our value of  $s_2 = 3.5$  or cumulative index of 2.5. This is consistent with the fact that the size distribution of very small objects is expected to approach a Donhanyi size distribution ( $s_2 = 3.5$ ) and then the craters produced by those small projectiles (that are in the strength regime) have to follow the same power law size distribution. Besides Kirchoff & Schenk (2010) found that for  $D = 1 - 4$  km, Phoebe's crater distribution has a shallow slope and this implies that Phoebe has a deficiency of craters with  $D \sim 1.5$  km. This change of slope cannot be explained by our method and our proposed contribution of Centaurs from the SD to the craters on Phoebe, unless that the considered SDF of SDOs is different. Anyway more work in relation to another source of craters on Phoebe, as planetocentric objects, is needed and also its connection to the origin of the irregular satellite itself.

According to the differential size distribution index  $s_2$ , the largest crater on Phoebe produced by a Centaur from the SD has a diameter of between 1.4 km and 13.5 km. Table 2 shows the



**Fig. 2.** Cumulative number of craters with diameters greater than  $D_0$  in the last 3.5 Gyrs, produced by Centaurs from SDOs on Phoebe. Filled line corresponds to the differential power law index  $s_2 = 3.5$  and the dotted line to  $s_2 = 2.5$


cumulative number of craters on Phoebe greater than certain diameters produced by Centaurs from the SD in the current configuration of the Solar System during the last 3.5 Gyrs. Since the largest crater on Phoebe has a diameter of  $\sim 100$  km, it is unlikely that it was produced by a recent collision of an SDO. This will be discussed in a following section.

## 5. Rate of SDO collisions on Phoebe

From our outputs we can calculate the number of encounters within the Hill's sphere of Saturn as a function of time. In Fig. 3 is plotted the normalized cumulative number of encounters as a function of time. The whole plot can be fitted by a log-function given by  $f(t) = a + b \log t$ , where  $a = -3.24$  and  $b = 0.19$ . The total cumulative number of encounters with Saturn for all diameters for each time can be obtained from the plot and Eq. (6). We have calculated the number of collisions on Phoebe from the number of encounters with Saturn from our outputs (Eq. 7), and this number can be calculated for each time in the integration. Then the cumulative number of collisions (or cumulative number of craters) on Phoebe for all diameters as a function of time can be obtained by multiplying the fraction of encounters obtained from Fig. 3 by the number of collisions  $N_c(d > d_0)$ .

As we can see from Fig. 3 the rate of encounters and then the rate of collisions on Phoebe was high at the beginning but it has been decreasing up to the present. The first Myrs the shape of the curve is purely arbitrary due to initial conditions but then it will be significant and is stabilizing. In the last  $\sim 3.5$  Gyrs the rate has been almost constant. In fact, it is possible to fit a linear relation to the last 3.5 Gyrs of Fig. 3 given by  $g(t) = \hat{F}t + c$ , where  $\hat{F} = 7.1 \times 10^{-11}$  and  $c = 0.69$ . This linear approximation allows us to calculate a present rate of craters for a given diameter. The slope





**Fig. 3.** Fraction of encounters of SDOs with Saturn. The linear fit to the data the last 3.5 Gyrs is also shown.

of this linear function  $\dot{F} = 7.1 \times 10^{-11}$  is the present normalized rate of encounters of SDOs with Saturn per year. To obtain the present rate of encounters with Saturn for each diameter we must multiply  $\dot{F}$  by  $N_{et}(d > d_0)$  (obtained from Eq (6)). And then the present rate of collisions on Phoebe for each diameter can be obtained multiplying  $\dot{F}$  by  $N_c(d > d_0)$  (obtained from Eq (7)). Similarly, the current rate of cratering on Phoebe for craters greater than a given diameter can be obtained from the current rate of collisions and the relation (9) between the diameter of the impactor and the diameter of the crater. Thus, for example the current cratering rate on Phoebe of Centaurs from SDOs that produces craters with  $D > 1$  km is between  $1.4 \times 10^{-10}$  and  $8.3 \times 10^{-8}$  craters per year (depending on the  $s_2$  value); this is at least  $\sim 80$  craters with  $D > 1$  km in the last Gy. The current cratering rate of craters with  $D > 5$  km is at least  $1.14 \times 10^{-9}$  craters per year, this is  $\sim 1$  crater with  $D > 5$  km in the last Gy.

Zhanle et al. (2003) calculate cratering rates in the satellites of the outer planets. They obtained a cratering rate on Phoebe for craters with  $D > 10$  km of  $8.6 \times 10^{-11} \text{ year}^{-1}$ . In this study we obtain a cratering rate on Phoebe of craters with  $D > 10$  km of between  $2.7 \times 10^{-12}$  and  $1.4 \times 10^{-10}$  craters per year, depending on  $s_2$ . Our value for  $s_2 = 3.5$  is very similar to the previous calculation of Zhanle et al. (2003).

## 6. The contribution of escaped Plutinos to the craters on Phoebe

Plutinos could be another source of craters on Phoebe. In a recent paper, Di Sisto et al. (2010) study the post escape evolution of Plutinos when they escaped from the 3:2 mean motion resonance with Neptune, and in particular their contribution to the population of Centaurs. In that work, we perform two sets of numerical simulations in order first to identify the plutinos that have recently

escaped from the resonance and second to follow their evolution under the influence of the Sun and the four giant planets. This numerical simulation considers the evolution of plutinos in the present configuration of the Solar System, this is as we did for SDOs, when the trans-Neptunian region is expected to have reached roughly its present mass and dynamical state,  $\sim 3.5$  Gyrs ago. Following the same analysis that we made for SDOs described in Sects. 3 and 4, we calculate the number of craters on Phoebe from escaped plutinos the last 3.5 Gyrs, and also the greater impactor and crater. In the numerical simulations by Di Sisto et al. (2010), we start with 20 000 initial massless particles in the 3:2 mean motion resonance, 671 of them undergone 20 459 encounters within Saturn's Hill sphere during the integration. We obtained that the mean relative encounter velocity of plutinos when entering the Hill's sphere of Saturn is  $v(R) = 4.57 \text{ km/s}$ , the mean relative velocity of plutinos when they intersect the orbit of Phoebe is  $v_0 = 5.12 \text{ km/s}$  and the collision velocity of plutinos on Phoebe is  $v_i = 5.4 \text{ km/s}$ . We take the present number of plutinos from de Elía et al. (2008), but consider that the size distribution breaks at  $d \sim 60 \text{ km}$  with the two limit power law indexes  $s_2 = 3.5$  and 2.5 as we adopted for SDOs to be consistent (see Sect. 2). Those indexes give the highest and lowest number of SDOs and then the highest and lowest production of craters on Phoebe. Then the present cumulative number of plutinos is given by:

$$\begin{aligned} N(> D) &= C \left( \frac{1 \text{ km}}{D} \right)^p && \text{for } D \leq 60 \text{ km}, \\ N(> D) &= 7.9 \times 10^9 \left( \frac{1 \text{ km}}{D} \right)^3 && \text{for } D > 60 \text{ km}, \end{aligned} \quad (10)$$

where  $C = 7.9 \times 10^9 (60)^{p-3}$  by continuity for  $D = 60 \text{ km}$  and the cumulative power law index  $p$  adopt the values 2.5 and 1.5 ( $p = s - 1$ ). This law is plotted in Fig 1.

Considering all this and our method described above, we obtained that the greater escaped plutino impactor on Phoebe has a diameter between 1.5 mts and 102 mts that produces a crater between 19.3 mts and 1.3 km respectively, depending on the power index  $p$  of the size distribution of plutinos. Also, we can obtain the number of craters on Phoebe from escaped plutinos. We have at least two craters greater than 1 km on Phoebe from plutinos. Comparing this with the values obtained for the contribution of SDOs, it can be stated that the number of craters produced by escaped plutinos on Phoebe is negligible with respect to the contribution of SDOs. Also it can be stated that the greater craters are signed by the contribution of SDOs.

## 7. Discussion

In the previous sections, we have calculated the production of craters on Phoebe considering the present population in the SD and Plutinos. However - as mentioned - there are many papers based on theoretical and observational work that argue that the initial mass of the TN region was  $\sim 100$  times greater than the present mass (see e.g. Morbidelli et al. 2008). Observations predict a current mass of the Kuiper Belt that is very small with respect to that required for models to grow the objects that we see. The mass depletion due to a strong dynamical excitation of the Kuiper Belt is thought to be the scenario for this "mass deficit problem". There were several models that try to describe the mass depletion; the last model that described this mechanism is the "Nice model" where the Kuiper Belt had to be significantly depleted before the time of the LHB (Levison et al. 2008). The "Nice Model" assumes the giant planets initially in a more compacted region from  $\sim 5.5$  to  $\sim 14 \text{ AU}$  and a planetesimal disk of a total mass of  $\sim 35 M_T$  that extends beyond the orbits

of the giant planets up to  $\sim 34$  AU. The interaction between the planets and planetesimals makes the giant planets migrate for a long time removing particles from the system. After a time ranging from 350 My to 1.1 Gy, Jupiter and Saturn cross their mutual 1:2 mean motion resonance. Then, the eccentricities of Uranus and Neptune drives up and those planets penetrate into the planetesimal disk. This destabilizes the full disk and the planetesimals are scattered all over the Solar System.

Beyond the model and the mechanism responsible for the mass depletion of the trans-Neptunian zone, we can consider that primitive SDOs (that were 100 times more numerous than the present ones) follow the same dynamical evolution than the present population when they enter the planetary zone as Centaurs. Then we can calculate in the same way as we did in the previous sections and with the same model the cratering on Phoebe assuming an initial population of SDOs 100 times the present population.

This is an estimation since we have to know the real initial scenario of formation of the Solar System and in particular of SDOs. However when a SDO enter the Centaur zone, inside the orbit of Neptune, its dynamical evolution is governed by the giant planets and then the particular initial scenario can be considered secondary for the present study.

Doing that, we obtain that the greater impactor on Phoebe during the age of the Solar System has a diameter between 2.4 to 8.6 km and produces a crater of 21.6 to 64.2 km. The value corresponding to  $s_2 = 3.5$  (64.2 km) is in agreement (within the expected errors and statistical fluctuations) with the observation of the great crater “Jason” on Phoebe with a diameter of  $\sim 100$  km. The number of craters greater than a given diameter can be obtained increasing 100 times the values obtained in Sect. 4

Recently, Cassini images of Phoebe have allowed to study its surface and the craters in it. Kirchoff & Schenk (2010) obtain, a cumulative crater density for  $D \geq 5$  km of  $2233 \pm 1117$  based on crater counting from Cassini images. From our model, assuming an initial number of SDOs 100 times the present population, we obtain  $N_c(D > 5km) = 12 - 1640$ , again in a good agreement with the values obtained by Kirchoff & Schenk (2010)  $s_2 = 3.5$ .

## 8. Conclusion

We have studied the production of craters on Phoebe from SDOs and escaped plutinos that have reach the Saturn zone in the present configuration of the Solar System. We have obtained that the contribution of escaped plutinos is negligible with respect to the contribution of SDOs. We have obtained that both the number of craters and the greater crater on Phoebe produced by SDOs cannot account to the observations. But if we take into account that the initial mass of the trans-Neptunian region was 100 times the present one, we match the craters produced by SDOs on Phoebe with the observed characteristics of the satellite if  $s_2 = 3.5$ .

Those considerations imply that the main cratering features of Phoebe must be acquired when the SD had being depleted at the early times of evolution of the Solar System. More than that, if the “Nice model” describes correctly the scenario of the origin of the Solar System, the scattering inward of planetesimals by Neptune and Uranus in that model must be similar to the present scattering in our model, and the TNOs have to lose memory when they arrive to Saturn.

If this is what happened and the main crater characteristics on Phoebe were produced when Phoebe was a satellite of Saturn, the great agreement of our model with the observations constrain

the time when Phoebe had to be captured, very early in the evolution of our Solar System. This was also suggested by Levison et al. (2008).

We have obtained that the present normalized rate of encounters of SDOs with Saturn is:  $\dot{F} = 7.1 \times 10^{-11}$  per year. From this number we could obtain the present cratering rate on Phoebe for each crater diameter.

We have compared the size crater distribution on Phoebe obtained from our model with the observations of craters by Kirchoff & Schenk (2010). Our crater size frequency distribution agree with that obtained by Kirchoff & Schenk for very small impactors that produce craters with  $D = 0.15 - 1$  km. This distribution follow a power law with a cumulative index of 2.5 consistent with a Donhanyi size distribution. For craters of  $D = 1 - 4$  km Kirchoff & Schenk (2010) found a shallow slope and a deficiency of craters with  $D \sim 1.5$  km. This change of slope cannot be explained by our method and the contribution of Centaurs from the SD. More work in relation to another source of craters on Phoebe, as planetocentric objects is needed, and also its connection to the origin of the irregular satellite itself.

**Acknowledgments:** We thank Gonzalo de Elía for valuable discussion on this paper. We also acknowledge an anonymous referee that made valuable comments and suggestions that helped us to improve the manuscript.

## References

- Bernstein, G. M., Trilling, D. E., Allen, R. L., et al. 2004, *AJ*, 128, 1364.
- Buratti, B.J., Wegryn, E., Dalle Ore, C.M., et al, 2008. *Icarus*, 193, 334-343.
- Cruikshank, D.P., Wegryn, E., Soderlund, K. et al, 2008. *Icarus*, 193, 309-322.
- de Elía, G. C., Brunini, A. & R. P. Di Sisto. 2008, *A&A*, 490, 835
- Di Sisto, R. P. & Brunini, A. 2007, *Icarus*, 190, 224.
- Di Sisto, R. P., Brunini, A. & de Elía, G.C., 2010. *A&A*, 519, A112.
- Fernández, J. A., Gallardo, T & Brunini, A., 2004. *Icarus*172, 372-381.
- Fraser, W.C., Kavelaars, J. J., Holman, M. J. et al., 2008 *Icarus*195, 827-843.
- Fraser, W.C. & Kavelaars, J. J., 2009. *AJ*137, 72-82 .
- Fuentes, C.I. & Holman, M. J., 2008. *AJ*136, 83-97 .
- Fuentes, C.I., George, M.R. & Holman, M.J., 2009. *ApJ*696, 91-95 .
- Giese, B. Neukum, G., Roatsch, T. 2006, *Planet. Space Sci.*, 54, 1156-1166.
- Gil-Hutton, R., Licandro, J., Pinilla-Alonso, N. et al. 2009, *A&A*500, 909 - 916.
- Hendrix, A.R. & Hansen, C.J., 2008. *Icarus*, 193, 323-333.
- Holsapple, K.A. 1993, *Annu. Rev. Earth Planet Sci.*, 21, 333-373.
- Housen, K.R. & Holsapple, K.A., 2003. *Icarus*163, 102-119.
- Holsapple, K.A. & Housen, K.R., 2007. *Icarus*187, 345-356.
- Holsapple, K.A., 2011. Web page at <http://keith.aa.washington.edu/craterdata/scaling/index.htm>. Accessed August, 9 2011.
- Holsapple, K.A., Schmidt, R.M. 1987, *J. Geophys. Res.* 92, No B7, 6350-6376.
- Johnson, T.V. & Lunine, J. 2005. *Nature* 435, 69-71.
- Kirchoff, M.R., Schenk, P. 2010. *Icarus*206, 485-497.
- Levison, H. & Duncan, M. 1997. *Icarus*127, 13-32.
- Levison, H.F., Morbidelli, A., VanLaerhoven, C. et al. 2008. *Icarus*, 196, 258-273.
- Morbidelli A. 1997, *Icarus*, 127, 1
- Morbidelli, A., Levison, H.F., Gomes, R. 2008. In *The Solar System Beyond Neptune*, ed. M. A. Barucci, et al. (Tucson, USA: Univ. of Arizona Press), 275.
- Nesvorný, D., Alvarelos, J.L.A., Dones, L., Levison, H.F., 2003. *AJ*126, 398-429.

- Parker A.H. & Kavelaars, J.J., 2010(a) *PASP*, 122, 549-559.
- Parker A.H. & Kavelaars, J.J., 2010(b) *Icarus*, 209, 766-770.
- Perna, D., Dotto, E., Barucci, M.A. et al., 2009. *A&A*508, 451-455.
- Petit, J.-M., Holman, M.J., Gladman, B.J. et al., 2006. *MNRAS*, 365, 429-438.
- Pollack, J.B. Burns, J.A. Tauber, M.E. 1979, *Icarus*37, 587-611.
- Porco, C.C., Baker, E., Barbara, J. et al. 2005. *Science* 307, 1237-1242.
- Schmidt, R.M., Housen, K.R. 1987. *Int. J. Impact Eng.*, 5, 543-560.
- Sheppard, S.S., Lacerda, P. & Ortiz, J.L. 2008. in *The Solar System Beyond Neptune*, (Barucci, M.A., et al. Eds.), Univ. of Arizona Press, Tucson, USA, 129.
- Zahnle, K., Schenk, P. Levison, H. & Dones, L., 2003. *Icarus*, 163, 263-289.