Error free Quantum Reading by Quasi Bell State of Entangled Coherent States

Osamu Hirota*

Quantum ICT Research Institute, Tamagawa University, 6-1-1, Tamagawa Gakuen, Machida 194-8610 Tokyo, JAPAN (Dated: May 25, 2022)

Nonclassical states of light field have been exploited to provide marvellous results in quantum information science. Effectiveness of nonclassical states depends on whether physical parameter as signal is continuous or digital. Here we present an investigation on the potential of quasi Bell state of entangled coherent states in quantum reading of the classical digital memory which was pioneered by Pirandola. This is a typical example of discrete quantum discrimination. We show that the quasi Bell state gives the error free performance in the quantum reading that cannot be obtained by any classical state.

PACS numbers: 03.67.-a, 03.65.Ud, 42.50.-p, 89.20.Ff

I. INTRODUCTION

In estimating an unknown parameter of quantum system, one prepares a probe light with several quantum states. To reduce the uncertainty, one can take N identical measures and average the results of those measures [1]. These strategies are very attractive in quantum metrology, and many applications to the modern information technology have been examined [2].

The signals in almost all of the information devices are digital and the decision problem is a major subject. However, when one considers a quantum measurement of the classical signal of quantum systems, one needs to take into account the quantum noise effect in the measurement of the observable. In general, limitations in the precision or error to discriminate the quantum states are imposed by such quantum noise. The nonclassical effect may become effective in digital signal rather than continuous signal as in the classical information science. Recently, a basic model of reading scheme was discussed in the context of quantum information [3,4], employing the theory of the binary quantum states discrimination so called Helstrom bound [5].

It may be a logical consequence to discuss the above application of quantum information science. In this context, one of key properties which make a state nonclassical is entanglement. The quasi Bell state based on entangled coherent states has a potential of the perfect entanglement[6], and its applications to quantum teleportation and computation have been discussed[7,8,9,10]. Furthermore, the feasibility and experimental demonstration [11] of such states have been reported.

Thus, we are concerned with applications of the quasi Bell state of entangled coherent states to quantum reading where various new technologies can emerge. In this Letter, we show the fact that one of properties of the quasi Bell state may provide the error free quantum reading of classical digital memory.

II. THEORETICAL EVALUATION OF PHYSICAL LIMITATION AND QUASI BELL STATES

A. Ultimate limitation in quantum measurement

A general theory evaluating the performance of the measurement in classical and quantum systems is "communication theory" including estimation and detection.

We can divide the issue of the theoretical limit on the ultimate sensitive measurement into signals of continuous parameter and discrete parameter. The former clarifies the precision of the measurement observable like the phase, and the latter clarifies the minimum error performance of the decision for signals in the radar or weak external force.

The quantum Cramer-Rao inequality has been formulated[5,12,13,14], in which the bound is asymptotically achieved by the maximum likelihood estimator as well as the classical estimation theory. Here let $\rho(\theta)$ be the density operator of the system. The estimation bound is given as follows: First, the following operator equation is defined as

$$\frac{\partial \rho(\theta)}{\partial \theta} = \frac{1}{2} [A\rho(\theta) + \rho(\theta)A] \tag{1}$$

where A is called symmetric logarithmic derivative, which is self-adjoint operator. Then the bound is given by

$$(\delta\theta)^2 \ge \frac{1}{F_Q = Tr(\rho(\theta)A^2)} \tag{2}$$

 F_Q is also called quantum Fisher information. The above inequality defines the principally smallest possible uncertainty in estimation of the value of parameter.

On the other hand, in the radar detection or weak force detection, the problem is whether signal exists or not. The formulation of the ultimate detection performance is called quantum detection theory[5,12]. Especially, for the binary quantum state discrimination, the limitation can be evaluated as follows[5]:

$$P_{e} = \xi_{0} T r \rho_{0} \Pi_{1} + \xi_{1} T r \rho_{1} \Pi_{0} \xi_{0}$$

^{*}Electronic address: hirota@lab.tamagawa.ac.jp

where $\{\xi_i\}$ is a priori probability of quantum states $\{\rho_i\}$ of the system, $\{\Pi_i\}$ is the detection operator, respectively. In the case of pure states, the optimum solution is

$$P_e = \frac{1}{2} \left[1 - \sqrt{1 - 4\xi_0 \xi_1} \right] < \psi_0 |\psi_1 > |^2 \right]$$
(4)

B. Quasi Bell state of entangled coherent states

In general, the Bell states consist of four orthogonal states. However, the four entangled states based on nonorthogonal states are called quasi Bell state. The specific example is the following states based on coherent state:

$$\begin{cases} |\Psi_{1}\rangle = h_{1}(|\alpha\rangle_{A}|\alpha\rangle_{B} + |-\alpha\rangle_{A}| - \alpha\rangle_{B}) \\ |\Psi_{2}\rangle = h_{2}(|\alpha\rangle_{A}|\alpha\rangle_{B} - |-\alpha\rangle_{A}| - \alpha\rangle_{B}) \\ |\Psi_{3}\rangle = h_{3}(|\alpha\rangle_{A}| - \alpha\rangle_{B} + |-\alpha\rangle_{A}|\alpha\rangle_{B}) \\ |\Psi_{4}\rangle = h_{4}(|\alpha\rangle_{A}| - \alpha\rangle_{B} - |-\alpha\rangle_{A}|\alpha\rangle_{B}) \end{cases}$$
(5)

where $\{h_i\}$ are normalized constant: $h_1 = h_3 = 1/\sqrt{2(1+\kappa^2)}$, $h_2 = h_4 = 1/\sqrt{2(1-\kappa^2)}$, and where $\langle \alpha | -\alpha \rangle = \kappa$ and $\langle -\alpha | \alpha \rangle = \kappa^*$.

Some of these quasi Bell states are not orthogonal each other. Here, if $\kappa = \kappa^*$, then the Gram matrix of them becomes very simple as follows:

$$G = \begin{pmatrix} 1 & 0 & D & 0 \\ 0 & 1 & 0 & 0 \\ D & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$
(6)

where $D = \frac{2\kappa}{1+\kappa^2}$.

The degrees of entanglement for quasi Bell state are well known as follows:

$$E(|\Psi_1\rangle) = E(|\Psi_3\rangle)$$

$$= -\frac{1+C_{13}}{2}\log\frac{1+C_{13}}{2} - \frac{1-C_{13}}{2}\log\frac{1-C_{13}}{2}$$
(7)

where $C_{ij} = |\langle \Psi_i | \Psi_j \rangle|$, and we have the special property such as $E(|\Psi_2\rangle) = E(|\Psi_4\rangle) = 1$ [6].

Thus $|\Psi_2\rangle$ and $|\Psi_4\rangle$ have the perfect entanglement.

III. SPECIFIC MODEL OF QUANTUM READING

There are two types of reading scheme of classical digital memory. These are "Amplitude shift keying (ASK)" and "Phase shift keying(PSK)", respectively.

The problem in the current proposal of the quantum reading is to discriminate quantum channels or quantum states affected by the different reflections r_0 and r_1 to one of two modes [3,4]. This corresponds to the model of ASK scheme. If the reflection coefficients are $r_0 = 1$

and $r_1 << 1$, it imposes great energy loss effect. The typical systems in quantum information science suffer the degradation from the energy loss effect. If one employs the nonclassical state in the above model, the loss effect has to be taken into account.

On the other hand, PSK scheme to realize the classical digital memory does not have a loss effect. We will employ PSK scheme such that the memory on the classical disk consists of the flat and concave which correspond to "0" and "1", respectively. This type of memory has been invented in the beginning of the 1980's, employing a Gas laser with high coherence as the light source. Although, later, its scheme was replaced with ASK by employing a laser diode with low coherence, recently PSK scheme is revived by laser diode with high coherence.

The difference of two schemes is not important in the case of classical state, but they make great difference in the case of nonclassical states. Here, desirable scheme is PSK scheme, because one can describe the phase shift by an unitary operator as follows:

$$U(\theta) = exp(-\theta a^{\dagger}a) \tag{8}$$

where a and a^{\dagger} are the annihilation and creation operator in bosonic system, respectively.

The reading method in the current PSK scheme is to illuminate the laser light (coherent state) on a disk, and to read the phase difference. That is, the signal states of light to discriminate are

$$|\alpha(0)\rangle = I|\alpha\rangle \tag{9}$$

$$|\alpha(1)\rangle = U(\theta)|\alpha\rangle \tag{10}$$

where I is the identity operator.

|e|

IV. ERROR FREE QUANTUM READING

Let us consider the detection problem of the above model. Here we restrict the problem to the binary detection. So detection targets are two quantum states. Since the phase difference will be π , the problem is to read the phase shift π from the steady state or input state.

The coherent states are prepared in the current classical memory, and the target signal model becomes as follows:

$$\alpha(0)\rangle = I|\alpha\rangle = |\alpha\rangle \tag{11}$$

$$\alpha(1)\rangle = U(\theta = \pi)|\alpha\rangle = |-\alpha\rangle \tag{12}$$

If one employs the conventional homodyne receiver to discriminate the above quantum states, the limitation is imposed by so called quantum shot noise limit. So this is the standard quantum limit in the modern information technology. If one employs the injection back effect [15] to a laser diode to detect the phase delay of the reflection from a disk as the conventional system, the sensitivity is inferior to a homodyne receiver, though it is a typical scheme. However, in general, one can enjoy the Helstrom bound to overcome the standard quantum measurement which is achieved by a homodyne receiver. From Eq(4), the limitation depends on the inner product between two quantum states. For example, the inner product of two coherent states is

$$\langle \alpha | -\alpha \rangle = exp(-2|\alpha|^2) \tag{13}$$

Let us assume that $|\Psi_2\rangle$ of quasi Bell state in Eq(5) is employed as the light source, and the *B* mode is illuminated to a memory disk. The reflection effect $U_B(\theta)$ operates on *B* mode, so the channel model is

$$\epsilon_{A\otimes B} = I_A \otimes U_B(\pi) \tag{14}$$

Then, the target states become as follows:

$$\begin{aligned} |\Psi_{2}(0)\rangle &= h_{2}(|\alpha\rangle_{A}|\alpha\rangle_{B} - |-\alpha\rangle_{A}| - \alpha\rangle_{B}) \\ |\Psi_{2}(1)\rangle &= \epsilon_{A\otimes B}h_{2}(|\alpha\rangle_{A}|\alpha\rangle_{B} - |-\alpha\rangle_{A}| - \alpha\rangle_{B}) \\ &= h_{2}(|\alpha\rangle_{A}| - \alpha\rangle_{B} - |-\alpha\rangle_{A}|\alpha\rangle_{B}) \end{aligned} (15)$$

Thus, the input state $|\Psi_2(0)\rangle = |\Psi_2\rangle$ is changed to $|\Psi_2(1)\rangle = |\Psi_4\rangle$. The inner product between the above two entangled coherent states in quasi Bell state is

$$\langle \Psi_2(0) | \Psi_2(1) \rangle = 0 \tag{16}$$

That is, the inner product becomes zero, and it is independent of the energy of light source. This is a property of the quasi Bell state. To check this special property, one can examine the different phase shift as follows:

$$|\langle \Psi_2(0)|I_A \otimes U_B(\theta)|\Psi_2(0)\rangle| > 0 \tag{17}$$

where $\theta \neq \pi$.

Finally, let us employ the quantum optimum receiver for the binary pure state of two modes. The ultimate detection performances of systems with the coherent state and quasi Bell state of entangled coherent states are given, respectively, as follows:

$$P_e(C) = \frac{1}{2} [1 - \sqrt{1 - 4\xi_0 \xi_1 exp(-4|\alpha|^2)}]$$
$$P_e(ECS) = 0$$
(18)

Thus, one can see that the property of the quasi Bell state provides an attractive improvement, and this property can be obtained only by the combination of the nonclassical state and the quantum optimum receiver. Although there are many nonclassical states, almost all is not changed from the input state to the orthogonal state to the original input state just by reflection. So we interpret that such an effectiveness of the quasi Bell state comes from the special phenomena on entanglement based on non-orthogonal state, which is a feature of quasi Bell state.

V. CONCLUSION

In the quantum metrology, utilizing an entangled coherent state could be applicable to many super sensitive measurements. But, there is an attractive improvement especially in the digital quantum state discrimination rather than continuous quantum state[16]. We have examined a potential of the entangled coherent states with a perfect entanglement in quasi Bell state for a well known quantum reading issue, and shown that it has the potential to realize the error free quantum reading under the finite energy of light source.

Acknowledgments

The author is grateful S.J.van Enk, Bill Munro, K.Kato, and T.Usuda for helpful discussions.

- C.M.Caves, Quantum mechanical noise in an interferometer, Phys. Rev., D-23, 1693, (1981).
- [2] V.Giovannetti, S.Lloyd, and L.Maccone, Quantum metrology, Phys. Rev.Lett., 96, 010401, (2006).
- [3] S.Pirandola, Quantum reading of a classical digital memory, Phys. Rev. Lett., 106, 090504, (2011).
- [4] R.Nair, Discriminating quantum optical beam-splitter channels with number diagonal signal state, arXiv:quantph/1105.4063,(2011).
- [5] C.W.Helstrom, Quantum detection and estimation theory, (Academic, New York, 1976).
- [6] O.Hirota and M.Sasaki, Entangled state based on nonorthogonal state, Proceedings of QCMC-2000, Ed by P.Tombesi and O.Hirota, (Kluwer Academic/Plenum, 2001), arXiv:quant-ph/0101018,(2001).
 Also, O.Hirota, S.J.van Enk, K.Nakamura, M.Sohma, and K.Kato, arXiv:quant-ph/0101096,(2001).
- [7] S.J.van Enk and O.Hirota, Entangled coherent states:

Teleportation and decoherence, Phys. Rev., A-64, 022313 (2001)

- [8] X.Wang, Quantum teleportation of entangled coherent state, Phys. Rev., 64, 022302, (2001).
- [9] H.Jeong, M.S.Kim, and J.Lee, Quantum information processing for a coherent superposition state via a mixed entangled coherent channel, Phys. Rev., A-64, 052308 (2001).
- [10] T.C.Ralph, A.Gilchrist, G.J.Miburn, W.J.Munro, and S.Glancy, Quantum comutation with optical coherent states, Phys. Rev., 68, 042319, (2003).
- [11] A.Ourjoumtsev, F.Ferreyrol, R.Tualle-Brouri, and P.Grangier, Preparation of non-local superposition of quasi-classical light state, Nature Physics, 5, 189 (2009).
- [12] A.S.Holevo, Probabilistic and statistical aspect of quantum theory, (North Holland, Amsterdam, 1982).
- [13] H.P.Yuen and M.Lax, Multiple-parameters quantum estimation and measurement of non-selfadjoint observable,

IEEE. Transaction of Information Theory, **IT-21**, 125, (1973).

- [14] S.L.Braunstein and C.M.Caves, Statistical distance and the geometry of quantum states, Phys. Rev. Lett., 72, 3439, (1994).
- [15] O.Hirota, and Y.Suematsu, Noise properties of injection

lasers due to reflected waves, IEEE. Journal of Quantum Electronics, **QE-15**, 142, (1979).

[16] O.Hirota, K.Kato and D.Murakami, Effectiveness of entangled coherent state in quantum metrology, arXiv:quant-ph/1108.1517V3, 2011.