

On generalized Schrödinger semigroups

Batu Güneysu*

Institut für Mathematik
Humboldt-Universität zu Berlin

March 22, 2019

Abstract

We extend the Feynman-Kac formula for Schrödinger type operators on vector bundles over noncompact Riemannian manifolds to possibly very singular potentials that appear in hydrogen like quantum mechanical problems and that need not be bounded from below or locally square integrable. This path integral formula is then used to prove several L^p -type results, like bounds on the ground state energy and $L^2 \rightsquigarrow L^p$ smoothing properties of the corresponding Schrödinger semigroups. As another main result, we will prove that with a little control on the Riemannian structure, the latter semigroups are also $L^2 \rightsquigarrow \{\text{bounded continuous}\}$ smoothing for Kato decomposable potentials. These results in particular apply to a very general class of magnetic Schrödinger operators on Riemannian manifolds.

1 Setting and some notation

Let M be a smooth Riemannian manifold (connected and without boundary), equipped with the Riemannian volume measure $\text{vol}(\bullet)$. We set $m := \dim M$ and denote the minimal heat kernel of M with $p_t(x, y)$. For $x, y \in M$, the number $d(x, y)$ will stand for the geodesic distance of $x, y \in M$ and $K_r(x)$ for the open geodesic ball with radius r around x .

Let $E \rightarrow M$ be a smooth d -dimensional complex vector bundle with a fixed Hermitian structure $(\bullet, \bullet)_x$ and a fixed Hermitian covariant derivative ∇ . The symbol $\|\bullet\|_x$ stands for the corresponding norm and operator norm of each fiber E_x .

*E-Mail: guneysu@math.hu-berlin.de

For our probabilistic considerations, we will assume that the underlying filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_*, \mathbb{P})$ satisfies the usual hypothesis and that it carries a Brownian motion W in the Euclidean \mathbb{R}^l , where $l \geq m$ is large enough. We will also assume $\mathcal{F}_* = \mathcal{F}_*(W)$. If M is stochastically complete, then one can use this setting to construct a Brownian motion $B(x)$ on M , starting from x with an infinite lifetime, as the maximally defined solution of a Stratonovic¹ differential equation of the form

$$dB(x) = \sum_{j=1}^l A_j(B(x)) \underline{d}W^j, \quad B_0(x) = x, \quad (1)$$

where $A_1, \dots, A_l \in \Gamma_{C^\infty}(M, TM)$ are such that $\sum_{j=1}^l A_j^2$ equals the Laplace-Beltrami operator. If $\pi : P(E) \rightarrow M$ denotes the $U(d)$ -principal bundle of unitary frames in E , then the stochastic ∇ -horizontal lift of $B(x)$ from a \mathcal{F}_0 -random variable $u : \Omega \rightarrow P(E)$ with $\pi(u) = x$ \mathbb{P} -a.s. is given as the maximally defined solution of

$$dU(u) = \sum_{j=1}^l A_j^*(U(u)) \underline{d}W^j, \quad U_0(u) = u,$$

where $A_j^* \in \Gamma_{C^\infty}(P(E), TP(E))$ is the ∇ -lift of A_j . The corresponding stochastic parallel transport will be written as a pathwise isometry $//_t^x := U_t(u)u^{-1} : E_x \rightarrow E_{B_t(x)}$, with some $U(u)$ as above. As the notation indicates, the process $//^x$ does not depend on the particular choice of u ([11], proposition 2.17). The reader may find the details of these constructions for example in [13] [11] [9] and the references therein.

2 Main results

The operator $\nabla^* \nabla / 2$ with domain of definition $\Gamma_{C_0^\infty}(M, E)$ is a nonnegative symmetric operator in $\Gamma_{L^2}(M, E)$ and the corresponding Friedrichs extension will be denoted with $H(0)$. Since there won't be any danger of confusion, we will denote the Friedrichs realization of 1/2-times the Laplace Beltrami operator in $L^2(M)$ with $-\Delta/2$. The corresponding quadratic forms in $\Gamma_{L^2}(M, E)$ and in $L^2(M)$, respectively, will be written as $q_{H(0)}$ and $q_{-\Delta/2}$.

For a potential² $V : M \rightarrow \text{End}(E)$, the corresponding nonnegative potentials

¹We will write \underline{d} for Stratonovic differentials, whereas Itô differentials will be written as d .

²With "potential" we mean a measurable section V in $\text{End}(E)$ such that $V(x)$ is Hermitian for almost everywhere (a.e.) $x \in M$.

V^+ and V^- can be defined with the spectral calculus of the fibers of E , and one has $V = V^+ - V^-$. Note that V defines a quadratic form q_V in $\Gamma_{L^2}(M, E)$ as follows:

$$\begin{aligned} D(q_V) &= \left\{ f \mid f, (V^+)^{\frac{1}{2}} f, (V^-)^{\frac{1}{2}} f \in \Gamma_{L^2}(M, E) \right\}, \\ q_V(f) &= \int_M (V(x)f(x), f(x))_x \operatorname{vol}(dx). \end{aligned} \quad (2)$$

We will usually work under a global Kato assumption on the negative part of the potentials under consideration:

Definition 2.1 *A measurable section V in $\operatorname{End}(E)$ is said to be in the Kato class $\Gamma_{\mathcal{K}}(M, \operatorname{End}(E))$, if*

$$\limsup_{t \searrow 0} \int_0^t \int_M p_s(x, y) \|V(y)\|_y \operatorname{vol}(dy) ds = 0, \quad (3)$$

and V is said to be in the local Kato class $\Gamma_{\mathcal{K}_{\text{loc}}}(M, \operatorname{End}(E))$, if $1_K V$ is in $\Gamma_{\mathcal{K}}(M, \operatorname{End}(E))$ for all compact $K \subset M$.

As usual, we will write $\mathcal{K}(M)$ for the Kato class corresponding to functions on M (considered as sections in the trivial bundle $M \times \mathbb{C}$ with the canonical Hermitian structure), so that $\Gamma_{\mathcal{K}}(M, \operatorname{End}(E))$ is nothing but the complex linear space of measurable sections V in $\operatorname{End}(E)$ such that $|V| \in \mathcal{K}(M)$. The space $\mathcal{K}_{\text{loc}}(M)$ is defined analogously.

Remark 2.2 Clearly, $\Gamma_{\mathcal{K}}(M, \operatorname{End}(E))$ generally depends on the Riemannian structure of M and the Hermitian structure $(\bullet, \bullet)_x$ of E , and the space $\Gamma_{\mathcal{K}_{\text{loc}}}(M, \operatorname{End}(E))$ does not depend on $(\bullet, \bullet)_x$, but may still depend on the Riemannian structure of M .

Analogously to definition 2.1, in the following, $\Gamma_{\mathcal{C}}(M, \operatorname{End}(E))$ will always be understood with respect to the fiberwise operator norm of $\operatorname{End}(E)$, where \mathcal{C} is a class like L^p . By the general properties of $p_t(x, y)$, one easily gets [10] the general inclusions

$$\begin{aligned} \Gamma_{L^\infty}(M, \operatorname{End}(E)) &\subset \Gamma_{\mathcal{K}}(M, \operatorname{End}(E)), \\ \Gamma_{\mathcal{K}_{\text{loc}}}(M, \operatorname{End}(E)) &\subset \Gamma_{L^1_{\text{loc}}}(M, \operatorname{End}(E)), \end{aligned} \quad (4)$$

and with some control on the Riemannian structure of M , one can produce a large class of (local) Kato potentials:

Theorem 2.3 *Let M be geodesically complete with Ricci curvature bounded from below and a positive injectivity radius. Then for any p such that $p \geq 1$ if $m = 1$, and $p > m/2$ if $m \geq 2$, one has*

$$\Gamma_{L^p}(M, \text{End}(E)) + \Gamma_{L^\infty}(M, \text{End}(E)) \subset \Gamma_{\mathcal{K}}(M, \text{End}(E)). \quad (5)$$

In particular, for such p one has

$$\Gamma_{L^p_{\text{loc}}}(M, \text{End}(E)) \subset \Gamma_{\mathcal{K}_{\text{loc}}}(M, \text{End}(E)). \quad (6)$$

Proof. The first inclusion may be found in [10]. It essentially follows from one of the main results of [25]. The second inclusion is a trivial consequence of the first one. ■

We will frequently make use of the following two compatibility results of Kato potentials with Brownian motion:

Lemma 2.4 *Let M be stochastically complete and let $v \in \mathcal{K}_{\text{loc}}(M)$. Then for any $x \in M$ one has ³*

$$\mathbb{P} \{v(B_\bullet(x)) \in L^1_{\text{loc}}[0, \infty)\} = 1. \quad (7)$$

Proof. The arguments of [3], p.45, can be adapted to our situation. We use the fact that $B(x)$ is nonexplosive by assumption: For any $t \geq 0$ and $n \in \mathbb{N}$ one has

$$\begin{aligned} & \mathbb{P} \left\{ \int_0^t |v(B_s(x))| \, ds = \infty \right\} \\ &= \mathbb{P} \left\{ \int_0^t \left| \left(1_{K_n(x)}(B_s(x)) + 1_{M \setminus K_n(x)}(B_s(x)) \right) v(B_s(x)) \right| \, ds = \infty \right\} \\ &\leq \mathbb{P} \left\{ \int_0^t |(1_{K_n(x)}v)(B_s(x))| \, ds = \infty \right\} \\ &\quad + \mathbb{P} \left\{ d(x, B_s(x)) \geq n \text{ for all } s \in [0, t] \right\}. \end{aligned}$$

As $n \rightarrow \infty$, the last term tends to $\mathbb{P}\{B(x) \text{ explodes before } t\} = 0$, so the lemma is proved, if we can show that for any $t \geq 0$ and $w \in \mathcal{K}(M)$ one has

$$\mathbb{E} \left[\int_0^t |w(B_s(x))| \, ds \right] < \infty.$$

³Note that in any case one has $\{v(B_\bullet(x)) \in L^1_{\text{loc}}[0, \infty)\} \in \mathcal{F}$, since this set is nothing but $\bigcap_{n \in \mathbb{N}} \left\{ \int_0^n |v(B_s(x))| \, ds < \infty \right\}$.

The last assertion is trivial for small t , and the general case then follows from the Markoff property of $B(x)$ (see for example [24]).

■

Proposition 2.5 *For any $v \in \mathcal{K}(M)$ there is a $C(v) > 0$ such that for all $t \geq 0$,*

$$\sup_{x \in M} \mathbb{E} \left[e^{\int_0^t |v(B_s(x))| ds} \right] \leq 2e^{tC(v)}.$$

Proof. We set

$$C(v, s) := \sup_{x \in M} \mathbb{E} \left[\int_0^s |v(B_r(x))| dr \right]$$

and choose a $t_0(v) > 0$ with $C(v, t_0(v)) < 1/2$. Then using Khas'minskii's lemma and the Markoff property of $B(x)$ one easily gets (see for example p.9 in [34] for the arguments) the first inequality in

$$\begin{aligned} \mathbb{E} \left[e^{\int_0^t |v(B_s(x))| ds} \right] &\leq \frac{1}{1 - C(v, t_0(v))} e^{\frac{t}{t_0(v)} \log\left(\frac{1}{1 - C(v, t_0(v))}\right)} \\ &< 2 e^{\frac{t}{t_0(v)} \log\left(\frac{1}{1 - C(v, t_0(v))}\right)}. \end{aligned}$$

This proves the claim.

■

We refer the reader to [10] and the references therein for other facts about Kato potentials on Riemannian manifolds.

The following theorem follows from the main result of [10]:

Theorem 2.6 *Let V be a potential such that*

$$V^+ \in \Gamma_{L^1_{\text{loc}}}(M, \text{End}(E)) \quad \text{and} \quad V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E)).$$

Then the quadratic form $q_{H(0)} + q_V$ with

$$D(q_{H(0)} + q_V) = D(q_{H(0)}) \cap D(q_{V^+})$$

is densely defined, closed and bounded from below.

It follows from theorem 2.6 that the form $H(0) \dot{+} V$ is a well-defined self-adjoint bounded from below operator in $\Gamma_{L^2}(M, E)$ which will be denoted with $H(V)$. Generalizing the situation considered in [31], we will call

$$(e^{-tH(V)})_{t \geq 0} \subset \mathcal{L}(\Gamma_{L^2}(M, E))$$

the *Schrödinger semigroup* corresponding to $H(V)$.

Remark 2.7 1. We use the following notation for scalar operators on functions: If $\beta \in \Omega_{\mathbb{R}}^1(M)$ and if $v : M \rightarrow \mathbb{R}$ is such that $v^+ \in L_{\text{loc}}^1(M)$ and $v^- \in \mathcal{K}(M)$, then the self-adjoint bounded from below operator in $L^2(M)$ corresponding to $(d + i\beta)^*(d + i\beta)/2 + v$ in the sense of theorem 2.6 (applied to $\nabla = d + i\beta$) will be written as $H_{\beta}(v)$, with the convention $H_0(0) = -\Delta/2$. Operators of the form $H_{\beta}(v)$ describe the energy of charged nonrelativistic quantum mechanical particles with spin 0, which live on M under the influence of the electrical potential v and the magnetic potential β .

2. The above smoothness assumption on the magnetic potential β is satisfactory from the physics point of view, since, at least, this is a local assumption. The above class of electrical potentials v , on the other hand, is certainly big enough to deal with most physically relevant situation. This claim is motivated by (5), which implies that the Kato class is big enough to deal with Coulomb type singularities $-1/|x|_{\mathbb{R}^3}$ in the Euclidean \mathbb{R}^3 , which appear naturally in the quantum mechanical hydrogen problem.

We will first prove the following scalar Feynman-Kac formula for Schrödinger operators of the form $H_0(v)$. This formula will be used with $H_0(\underline{V})$ for certain existence and convergence arguments in the proof of a Feynman-Kac formula for operators of the type $H(V)$ (theorem 2.11). This “scalarization procedure” reflects the fact that operators of the form $H(V)$ always dominate scalar operators of the form $H_0(\underline{V})$, a statement which can be made precise by means of a Kato type inequality [16][2][10]. We will derive and use several aspects of this domination in this paper.

Theorem 2.8 *Let M be geodesically and stochastically complete and let $v : M \rightarrow \mathbb{R}$ be such that $v^+ \in L_{\text{loc}}^1(M)$ and $v^- \in \mathcal{K}(M)$. Then for a.e. $x \in M$ one has*

$$\mathbb{P} \{v(B_{\bullet}(x)) \in L_{\text{loc}}^1[0, \infty)\} = 1, \quad (8)$$

and the following formula holds for any $f \in L^2(M)$, $t \geq 0$ and a.e. $x \in M$,

$$e^{-tH_0(v)} f(x) = \mathbb{E} \left[e^{-\int_0^t v(B_s(x)) ds} f(B_t(x)) \right]. \quad (9)$$

Our proof of theorem 2.8 follows the strategy of the proof of theorem 6.2 in [32] (see also [20]). The proof shows that it is actually possible to derive (9) for v 's with v^- even more general than in $\mathcal{K}(M)$. However, (8) (which we will use in order to prove the existence of $\mathcal{V}^{(x)}$ in theorem 2.11 below) need not be true then and one has to define $\int_0^t v(B_s(x)) ds$ carefully.

Proof of theorem 2.8. We first remark that by writing $f = f_1 - f_2 + if_3 - if_4$ with $f_j \geq 0$, we can and we will assume $f \geq 0$. We divide the proof into two parts:

I) (8) and (9) hold under the additional assumption $v \geq C$.

Proof: We can assume $C = 0$, so $v \geq 0$. In this situation, the a priori pathwise definition of $\int_0^t v(B_s(x))ds$ does not cause any problems so that the right-hand side of (9) is well-defined as a number in $[0, \infty]$. Indeed, we will actually prove (9) first, and then deduce (8) from this formula.

Let us define a sequence of potentials $(v_n) \subset L^\infty(M)$ by $v_n := \min\{n, v\}$. Then we have

$$D(q_{H_0(v)}) = D(q_{-\Delta/2}) \cap \left\{ \psi \mid \sqrt{v}\psi \in L^2(M) \right\}, \quad (10)$$

$$q_{H_0(v)}(\psi) = q_{-\Delta/2}(\psi) + \int_M v(x)|\psi(x)|^2 \text{vol}(dx), \quad (11)$$

and for any n ,

$$\begin{aligned} D(q_{H_0(v_n)}) &= D(q_{-\Delta/2}), \\ q_{H_0(v_n)}(\psi) &= q_{-\Delta/2}(\psi) + \int_M v_n(x)|\psi(x)|^2 \text{vol}(dx), \end{aligned} \quad (12)$$

and it follows from $0 \leq v_n \leq v_{n+1} \leq v$, $v_n \rightarrow v$ a.e. in M as $n \rightarrow \infty$ and monotone convergence of integrals and convergence of monotonely increasing quadratic forms (see for example theorem 12.2.2 in [20]), that we may assume

$$\lim_{n \rightarrow \infty} e^{-tH_0(v_n)} f(x) = e^{-tH_0(v)} f(x) \quad \text{for a.e. } x \in M. \quad (13)$$

Corollary 1.5 in [9] (a Feynman-Kac formula for nonnegative locally square integrable potentials) implies

$$e^{-tH_0(v_n)} f(x) = \mathbb{E} \left[e^{-\int_0^t v_n(B_s(x))ds} f(B_t(x)) \right] \quad \text{for a.e. } x, \quad (14)$$

so that

$$e^{-tH_0(v)} f(x) = \lim_{n \rightarrow \infty} \mathbb{E} \left[e^{-\int_0^t v_n(B_s(x))ds} f(B_t(x)) \right] \quad \text{for a.e. } x. \quad (15)$$

Next, one gets from combining $e^{-\int_0^t v_1(B_s(x))ds} f(B_t(x)) \in L^1(\mathbb{P})$ with

$$e^{-\int_0^t v_1(B_s(x))ds} f(B_t(x)) \geq e^{-\int_0^t v_n(B_s(x))ds} f(B_t(x)) \geq e^{-\int_0^t v(B_s(x))ds} f(B_t(x))$$

and an generalized abstract convergence theorem for integrals (theorem 12.2.6 in [20]), that the right-hand side of (15) is equal to

$$\mathbb{E} \left[\exp \left(- \lim_{n \rightarrow \infty} \int_0^t v_n(B_s(x))ds \right) f(B_t(x)) \right] \quad \text{for all } x,$$

and $\int_0^t v_n(B_s(x))ds \rightarrow \int_0^t v(B_s(x))ds$ \mathbb{P} -a.s. as $n \rightarrow \infty$ follows from monotone convergence. Finally, (8) follows from applying (9) to some negative function in $L^2(M)$.

II) (8) and (9) hold in the general case.

Proof: In this case, (8) follows from $|v| \leq v^+ + v^-$, the validity of (8) for nonnegative locally integrable potentials (step I)) and lemma 2.4. It remains to prove (9). To this end, we define $v_n := \max\{-n, v\}$, so that each v_n is a bounded from below potential with $v_n^+ = v^+$ locally integrable and $v_n \geq v_{n+1} \geq v$, $v_n \rightarrow v$ a.e. in M as $n \rightarrow \infty$. One has

$$D(q_{H_0(v_n)}) = D(q_{H_0(v)}) = D(q_{-\Delta/2}) \cap \left\{ \psi \mid \sqrt{v^+} \psi \in L^2(M) \right\}, \quad (16)$$

$$q_{H_0(v)}(\psi) = q_{-\Delta/2}(\psi) + \int_M v(x) |\psi(x)|^2 \text{vol}(dx), \quad (17)$$

$$q_{H_0(v_n)}(\psi) = q_{-\Delta/2}(\psi) + \int_M v_n(x) |\psi(x)|^2 \text{vol}(dx). \quad (18)$$

Furthermore, one can use the above cited generalized convergence theorem for integrals and convergence of monotonely decreasing quadratic forms (the latter by subtracting $\min \sigma(H_0(v))$ if necessary) to see that we may assume (13) again. By I), we also have (14) now. It remains to prove

$$\mathbb{E} \left[e^{-\int_0^t v_n(B_s(x))ds} f(B_t(x)) \right] \rightarrow \mathbb{E} \left[e^{-\int_0^t v(B_s(x))ds} f(B_t(x)) \right] \quad (19)$$

as $n \rightarrow \infty$ for a.e. x . Noting that for a.e. x one has

$$e^{-\int_0^t v_1(B_s(x))ds} f(B_t(x)) \in L^1(\mathbb{P}), \quad \int_0^t |v_1(B_s(x))| ds < \infty \quad \mathbb{P}\text{-a.s.}$$

by I), we can use theorem 12.2.6 in [20] twice to see that (19) holds. ■

We will use the notation

$$\mathcal{E}_H := \min \sigma(H)$$

for the ground state energy of a self-adjoint semi-bounded from below operator H . It follows from theorem 2.8 that the operator $e^{-tH_0(v)}$ is positivity preserving for all $t \geq 0$. Using this fact, we get the following facts for $\mathcal{E}_{H_0(v)}$ directly from abstract results on self-adjoint semi-bounded from below operators on measure spaces:

Corollary 2.9 *Fix the assumptions of theorem 2.8.*

a) *If $\mathcal{E}_{H_0(v)}$ is an eigenvalue of $H_0(v)$, then $\mathcal{E}_{H_0(v)}$ is simple and the corresponding ground state eigenvector can be chosen strictly positive.*

b) Let $f_1, f_2 \in L^2(M)$ with $f_1, f_2 > 0$. Then the following formula holds,

$$\mathcal{E}_{H_0(v)} = - \lim_{t \rightarrow \infty} t^{-1} \log \mathbb{E} \left[\int_M f_1(x) e^{-\int_0^t v(B_s(x)) ds} f_2(B_t(x)) \text{vol}(dx) \right]. \quad (20)$$

Proof. a) This follows directly from the fact that $e^{-tH_0(v)}$ is positivity preserving for all t . See for example theorem XIII.44 in [29].

b) Using again that $e^{-tH_0(v)}$ is positivity preserving for all t , lemma 2.1 in [26] directly implies

$$\mathcal{E}_{H_0(v)} = - \lim_{t \rightarrow \infty} \frac{\log \langle f_1, e^{-tH_0(v)} f_2 \rangle}{t}. \quad (21)$$

Now (20) follows from (21) by the Feynman-Kac formula and Fubini's theorem. ■

We turn to the general vector valued setting again. If V is a appropriate measurable section in $\text{End}(E)$ and $x \in M$ is appropriate, then the process

$$\mathcal{V}^{(x)} : [0, \infty) \times \Omega \longrightarrow \text{End}(E)_x$$

will stand for the unique pathwise weak solution of

$$d\mathcal{V}_t^{(x)} = -\mathcal{V}_t^{(x)} \left(//_{/t}^{x,-1} V(B_t(x)) //_{/t}^x \right) dt, \quad \mathcal{V}_0^{(x)} = \mathbf{1}. \quad (22)$$

The process $\mathcal{V}^{(x)}$ is pathwise invertible and

$$\mathcal{V}^{(x),-1} : [0, \infty) \times \Omega \longrightarrow \text{End}(E)_x$$

is uniquely determined by

$$d\mathcal{V}_t^{(x),-1} = \left(//_{/t}^{x,-1} V(B_t(x)) //_{/t}^x \right) \mathcal{V}_t^{(x),-1} dt, \quad \mathcal{V}_0^{(x),-1} = \mathbf{1}. \quad (23)$$

We will use the notation

$$|\Psi|(x) := \|\Psi(x)\|_x \quad \text{for any section } \Psi \text{ in } E \text{ or in } \text{End}(E).$$

The following simple observation will be used throughout in the following:

Remark 2.10 If for any potential $V : M \rightarrow \text{End}(E)$ we define

$$\underline{V} : M \longrightarrow \mathbb{R}, \quad \underline{V}(x) := \min \sigma(V(x)),$$

then we have $\underline{V}^\pm(\bullet) = \min \sigma(V^\pm(\bullet))$ and $|\underline{V}^\pm(\bullet)| \leq |V^\pm(\bullet)|$, which is implied by the fiberwise spectral calculus of E .

The following Feynman-Kac type formula for sections in E will be the main tool of this paper. It is a generalization of theorem 1.3 in [9] to potentials that are not necessarily locally square integrable or bounded from below:

Theorem 2.11 *Let M be geodesically and stochastically complete and let V be a potential with*

$$V^+ \in \Gamma_{\text{loc}}^1(M, \text{End}(E)) \quad \text{and} \quad V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E)).$$

Then for a.e. $x \in M$, there is a unique process

$$\mathcal{Y}^{(x)} : [0, \infty) \times \Omega \longrightarrow \text{End}(E)_x$$

which satisfies (22) pathwise in the weak sense, and for any $f \in \Gamma_{L^2}(M, E)$, $t \geq 0$, a.e. $x \in M$ one has

$$e^{-tH(V)} f(x) = \mathbb{E} \left[\mathcal{Y}_t^{(x)} //_{t^{x,-1}} f(B_t(x)) \right]. \quad (24)$$

Proof. Firstly, we remark that since parallel transport is an isometric operation, the asserted existence of $\mathcal{Y}^{(x)}$ will follow from the Banach fixed point theorem, if we can show that for a.e. $x \in M$ one has

$$\mathbb{P} \left\{ \|V(B_{\bullet}(x))\|_{B_{\bullet}(x)} \in L_{\text{loc}}^1[0, \infty) \right\} = 1. \quad (25)$$

But since $|V| \leq |V^+| + |V^-|$, the integrability (25) follows from the assumptions on V by applying (8) with $v := |V^+|$ and lemma 2.4 with $v := |V^-|$. The remaining proof follows the strategy of the proof of theorem 2.8, with modifications that are necessary due to the vector valued character. We divide the proof into two parts again:

I) (24) holds under the additional assumption $V \geq C$.

Proof: We may assume $V \geq 0$. Using the the spectral calculus of the fibers of E we define $V_n := \min(n, V)$ for any $n \in \mathbb{N}$. Then each V_n is a potential in $\Gamma_{L^\infty}(M, \text{End}(E))$ and one has

$$0 \leq V_n \leq V_{n+1} \leq V, \quad V_n \rightarrow V \quad \text{a.e. in } M \text{ as } n \rightarrow \infty. \quad (26)$$

Using monotone convergence of quadratic forms as in the first part of the proof of theorem 2.8 shows that we may assume

$$\lim_{n \rightarrow \infty} e^{-tH(V_n)} f(x) = e^{-tH(V)} f(x) \quad \text{for a.e. } x. \quad (27)$$

With an obvious notation, theorem 1.1 in [9] implies

$$e^{-tH(V)} f(x) = \lim_{n \rightarrow \infty} \mathbb{E} \left[\mathcal{Y}_{n,t}^{(x)} //_{t^{x,-1}} f(B_t(x)) \right] \quad \text{for a.e. } x. \quad (28)$$

Let x be such that (25) holds from now on. In view of (25) and (26), proposition 4.2 implies⁴

$$\begin{aligned} & \left\| \mathcal{V}_{n,t}^{(x)} - \mathcal{V}_t^{(x)} \right\|_x \\ & \leq e^{3 \int_0^t \|V(B_s(x))\|_{B_s(x)} ds} \int_0^t \|V(B_s(x)) - V_n(B_s(x))\|_{B_s(x)} ds \quad \mathbb{P}\text{-a.s.}, \end{aligned} \quad (29)$$

so using (25) and (26) again, we get from dominated convergence that

$$\lim_{n \rightarrow \infty} \left\| \mathcal{V}_{n,t}^{(x)} - \mathcal{V}_t^{(x)} \right\|_x = 0 \quad \mathbb{P}\text{-a.s.} \quad (30)$$

Finally, we may use (30) and $\left\| \mathcal{V}_{n,t}^{(x)} \right\|_x \leq 1$ \mathbb{P} -a.s. (the latter follows from $V_n \geq 0$ and proposition 4.1 c)), to deduce (24) from (28) and dominated convergence.

II) (24) holds in the general case.

Proof: Now we define $V_n := \max(-n, V)$ for any n . Then each V_n is a bounded from below potential with $V_n^+ = V^+$ locally integrable and one has

$$V_n \geq V_{n+1} \geq V, \quad V_n \rightarrow V \text{ a.e. in } M \text{ as } n \rightarrow \infty, \quad (31)$$

so that one can use convergence of monotonely decreasing quadratic forms as in the second part of the proof of theorem 2.8 to see that we can assume (27). By I), we also have (28) now, and so it remains to prove that the limit may be put into the expectation value in (28) for a.e. x , which will be proved with a dominated convergence argument. To this end, note that we again have (29) and that (31) implies⁵

$$0 \leq |V_n - V| \leq |V_1 - V|. \quad (32)$$

As a consequence, we may use theorem 12.2.6 in [20] to deduce (30). Next, the inequality⁶ $-\underline{V}_n \leq -\underline{V}$ and proposition 4.1 c) give

$$\left\| \mathcal{V}_{n,t}^{(x)} \right\|_x \leq e^{-\int_0^t \underline{V}(B_s(x)) ds} \quad \mathbb{P}\text{-a.s.}$$

in particular,

$$\left\| \mathcal{V}_{n,t}^{(x)} / t^{x,-1} f(B_t(x)) \right\|_x \leq e^{-\int_0^t \underline{V}(B_s(x)) ds} \|f(B_t(x))\|_{B_t(x)} \quad \mathbb{P}\text{-a.s.}$$

⁴Note that $|V_n|(\bullet) \leq |V|(\bullet)$, which follows from $|V_n|(\bullet) = \max \sigma(V_n(\bullet))$, $|V|(\bullet) = \max \sigma(V(\bullet))$ and (126).

⁵To see this inequality, just note $|V_n - V|(\bullet) = \max \sigma(V_n(\bullet) - V(\bullet))$ and use (126).

⁶This follows directly from (126).

These arguments are valid for any x such that (25) holds. Finally, we have

$$\begin{aligned} & \mathbb{E} \left[e^{-\int_0^t \underline{V}(B_s(x)) ds} \| f(B_t(x)) \|_{B_t(x)} \right] \\ &= e^{-tH_0(\underline{V})} |f|(x) < \infty \quad \text{for a.e. } x, \end{aligned} \quad (33)$$

since the scalar potential $\underline{V} = \underline{V}^+ - \underline{V}^-$ satisfies the assumptions of theorem 2.8 by remark 2.10, so that formula (24) indeed follows from dominated convergence. ■

Using proposition 2.6 in [9], one can immediately derive a very general Feynman-Kac-Itô formula for magnetic Schrödinger operators on Riemannian manifolds from formula (24):

Corollary 2.12 *Let M be geodesically and stochastically complete, let $v : M \rightarrow \mathbb{R}$ be such that $v^+ \in L^1_{\text{loc}}(M)$ and $v^- \in \mathcal{K}(M)$, and let $\beta \in \Omega^1_{\mathbb{R}}(M)$. Then the following formula holds for any $f \in L^2(M)$, $t \geq 0$ and a.e. $x \in M$,*

$$e^{-tH_\beta(v)} f(x) = \mathbb{E} \left[e^{-\int_0^t v(B_s(x)) ds + i \int_0^t \beta(\underline{d}B_s(x))} f(B_t(x)) \right], \quad (34)$$

where $\int \beta(\underline{d}B(x))$ stands for the Stratonovic integral of β along $B(x)$.

Formula (34) generalizes a Feynman-Kac-Itô formula from [9].

Now we would like present some applications of (22). We first come to some L^p -type results. A key observation is the following semigroup domination. We refer the reader to [1] and [15] for an abstract formulation of semigroup domination and applications.

Theorem 2.13 *Under the assumptions of theorem 2.11, let $v : M \rightarrow \mathbb{R}$ be such that $v^+ \in L^1_{\text{loc}}(M)$, $v^- \in \mathcal{K}(M)$ and $V \geq v\mathbf{1}$. Then the following inequality holds for any $f \in \Gamma_{L^2}(M, E)$, $t \geq 0$ and a.e. $x \in M$,*

$$\| e^{-tH(V)} f(x) \|_x \leq e^{-tH_0(v)} |f|(x). \quad (35)$$

In particular, the following assertions hold:

i) *For any $f \in \Gamma_{L^2}(M, E)$, $t \geq 0$,*

$$\langle e^{-tH(V)} f, f \rangle \leq \langle e^{-tH_0(v)} |f|, |f| \rangle. \quad (36)$$

ii) *One has $|f| \in D(q_{H_0(v)})$ with $q_{H(V)}(f) \geq q_{H_0(v)}(|f|)$ for any $f \in D(q_{H(V)})$.*

iii) One has $\mathcal{E}_{H(V)} \geq \mathcal{E}_{H_0(v)}$.

iv) For any $f \in \Gamma_{L^2}(M, E)$, any $\lambda \in \mathbb{C}$ such that $\operatorname{Re}(\lambda) > \mathcal{E}_{H(V)}$ and a.e. $x \in M$,

$$\|(H(V) + \lambda)^{-1} f(x)\|_x \leq (H_0(v) + \lambda)^{-1} |f|(x). \quad (37)$$

Proof. It follows from proposition 4.1 c) that for a.e. $x \in M$ one has

$$\left\| \mathcal{Y}_t^{(x)} \right\|_x \leq e^{-\int_0^t v(B_s(x)) ds} \quad \mathbb{P}\text{-a.s.}, \quad (38)$$

which directly implies (35) and (36).

For ii) and iii), we can assume that $H(V)$ and $H_0(v)$ are nonnegative (otherwise we can consider $H(V + C)$ and $H_0(v + C)$ with $C \in \mathbb{R}$ small enough and use (128) and (129). Under this assumption, ii) is implied by i), (130), and iii) follows from ii), (128).

Finally, (37) follows from (35) by taking Laplace transforms

$$(H(V) + \lambda)^{-1} f(x) = \int_0^\infty e^{-t\lambda} e^{-tH(V)} f(x) dt$$

and

$$(H_0(v) + \lambda)^{-1} |f|(x) = \int_0^\infty e^{-t\lambda} e^{-tH_0(v)} |f|(x) dt,$$

and the proof is complete. ■

Remark 2.14 1. A canonical choice for v in theorem 2.13 is given by $v := \underline{V}$. This follows from remark 2.10.

2. If $V \geq 0$, then theorem 2.13 iii) implies

$$\mathcal{E}_{H(V)} \geq \mathcal{E}_{-\Delta/2},$$

a bound on the ground state energy of $H(V)$ which is purely ‘‘Riemann geometric’’ in the sense that it does not depend on the interaction V or any data corresponding to the underlying vector bundle E .

Combining theorem 2.13 iii) with remark 2.7 leads to an important consequence for quantum mechanics on manifolds which is well-known for quantum mechanics in Euclidean space:

Corollary 2.15 *Let M be geodesically and stochastically complete and let $v : M \rightarrow \mathbb{R}$ be an electrical potential with $v^+ \in L^1_{\text{loc}}(M)$, $v^- \in \mathcal{K}(M)$. Then the presence of a magnetic potential $\beta \in \Omega^1_{\mathbb{R}}(M)$ leads to an increase of the ground state energy of charged nonrelativistic spin 0 particles, which live on M under the influence of (β, v) .*

Proof. Mathematically, the assertion just means that

$$\mathcal{E}_{H_\beta(v)} \geq \mathcal{E}_{H_0(v)}, \quad (39)$$

and this inequality follows directly from theorem 2.13 iii).

However, we find it instructive to remark that it is almost trivial to deduce (39) directly from the Feynman-Kac-Itô formula: For (34) and the triangle inequality implies

$$\langle e^{-tH_\beta(v)} f, f \rangle \leq \langle e^{-tH_0(v)} |f|, |f| \rangle \quad (40)$$

for any $f \in L^2(M)$, from which (39) follows directly from combining (126) with (129). ■

Before we continue with our main results, we remark the following heat kernel estimate for future reference:

Proposition 2.16 *Let M be geodesically complete with Ricci curvature bounded from below and a positive injectivity radius. Then for any $t > 0$ there are $c(t), d(t) > 0$, which depend on the Riemannian structure of M , such that for all $0 < s \leq t$ and all $x, y \in M$ one has*

$$p_s(x, y) \leq \frac{c(t) e^{-d(t) \frac{d(x,y)^2}{s}}}{s^{m/2}}. \quad (41)$$

In particular,

$$C(t) := \sup_{x,y \in M} p_t(x, y) < \infty \quad \text{for all } t > 0. \quad (42)$$

Proof. The estimate (41) follow easily from the considerations of p.110 in [25] and a simple rescaling argument for the Riemannian structure of M . ■

For the next proposition we consider $e^{-tH(V)} f$ and $e^{-tH_0(v)} h$, where $f \in \Gamma_{L^2}(M, E)$, $h \in L^2(M)$, as an equivalence class of measurable sections in E and, respectively, as an equivalence class of measurable functions on M . In this sense, both $e^{-tH(V)} f$ and $e^{-tH_0(v)} h$ are given by the corresponding

Feynman-Kac formula. For any $p, q \in [1, \infty]$ let $\|\bullet\|_q$ denote the norm in $\Gamma_{L^q}(M, E)$ and let $\|\bullet\|_{p,q}$ denote the norm corresponding to

the Banach space $\mathcal{L}(\Gamma_{L^p}(M, E), \Gamma_{L^q}(M, E))$,

with the conventions $\|\bullet\| = \|\bullet\|_2$ and $\|\bullet\| = \|\bullet\|_{2,2}$ and analogous notations for functions. The following theorem 2.17 proves the $L^2 \rightsquigarrow L^q$ smoothing of the Schrödinger semigroup $(e^{-tH(V)})_{t \geq 0}$.

Theorem 2.17 *Fix the assumptions of theorem 2.13.*

a) *Let $q \in [1, \infty]$ and $t > 0$. Then one has the implication*

$$e^{-tH_0(v)} \in \mathcal{L}(L^2(M), L^q(M)) \Rightarrow e^{-tH(V)} \in \mathcal{L}\left(\Gamma_{L^2}(M, E), \Gamma_{L^q}(M, E)\right),$$

and it holds that

$$\|e^{-tH(V)}\|_{2,q} \leq \|e^{-tH_0(v)}\|_{2,q}. \quad (43)$$

b) *Let M be geodesically and stochastically complete and assume that*

$$C(t) := \sup_{x,y \in M} p_t(x, y) < \infty \quad \text{for all } t > 0. \quad (44)$$

Then for any $q \in [2, \infty]$, $t > 0$, one has

$$e^{-tH(V)} \in \mathcal{L}\left(\Gamma_{L^2}(M, E), \Gamma_{L^q}(M, E)\right). \quad (45)$$

In particular, any eigensection of $H(V)$ lies in $\Gamma_{L^q}(M, E)$ for all $q \in [2, \infty]$. More precisely, for any $q \in [2, \infty]$, $t > 0$, one has

$$\|e^{-tH(V)}\|_{2,q} \leq 2C(t)^{\frac{1}{2} - \frac{1}{q}} e^{tD(V^-)}, \quad (46)$$

where $D(V^-) := C(2\underline{V}^-)/2$, and where $C(2\underline{V}^-)$ is chosen as in proposition 2.5.

Remark 2.18 1. Combining part a) and part b) of theorem 2.17 shows the following surprising fact: If M is geodesically and stochastically complete and V is a potential with

$$V^+ \in \Gamma_{L^1_{\text{loc}}}(M, \text{End}(E)) \quad \text{and} \quad V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E)),$$

then the validity of

$$e^{-tH(V)} \in \mathcal{L}\left(\Gamma_{L^2}(M, E), \Gamma_{L^q}(M, E)\right)$$

can be achieved by *only* requiring additional control, namely (44), on the Riemannian structure of M .

2. If M is geodesically complete with Ricci curvature bounded from below and a positive injectivity radius, then the assumptions on the Riemannian structure of M are satisfied. This follows from proposition 2.16 and the well-known fact that geodesic completeness and a lower bound on the Ricci curvature imply stochastic completeness. Under these assumptions on M , the estimate (46) combined with (41) actually shows the following estimate for all $q \in [2, \infty]$, $0 \leq t \leq 1$ in the situation of part b) of theorem 2.17: One has

$$\|e^{-tH(V)}\|_{2,q} \leq \frac{C(q)}{t^{\frac{m}{2}(\frac{1}{2}-\frac{1}{q})}} e^{tD(V^-)}, \quad (47)$$

where $C(q) > 0$ is a constant which only depends on q and the Riemannian structure of M .

3. The inclusion (45) is contained in [14] for scalar operators of the form $H_0(v)$ in the Euclidean $L^2(\mathbb{R}^m)$. On the other hand, it seems as if (45) does not appear in the literature in this form even for operators of the form $H_\beta(v)$ in $L^2(\mathbb{R}^m)$. In the latter case, however, this result is included in [3] under the slightly stronger assumption $v^+ \in \mathcal{K}_{\text{loc}}(\mathbb{R}^m)$, but with β 's more general than smooth.

We will need the following proposition 2.19 for the proof of part b) of theorem 2.17. Although we will use the result only with $p = 2$, it does not cause much extra work to consider the general $L^p \rightsquigarrow L^q$ situation:

Proposition 2.19 *Assume that*

$$C(t) := \sup_{x,y \in M} p_t(x,y) < \infty \quad \text{for all } t > 0. \quad (48)$$

*Then the assignment*⁷

$$P_t h(x) := \int_M p_t(x,y) h(y) \text{vol}(dy)$$

defines an element of $\mathcal{L}(L^p(M), L^q(M))$ for all $1 \leq p \leq q \leq \infty$, $t > 0$, and one has

$$\|P_t\|_{p,q} \leq C(t)^{\frac{1}{p}-\frac{1}{q}}. \quad (49)$$

⁷Note that $P_t h = e^{\frac{t}{2}\Delta} h$ for $h \in L^2(M)$.

The proof of proposition 2.19 will be given in section 3. A short look at the proof shows that proposition 2.19 actually has a natural generalization to symmetric essentially bounded integral kernels on σ -finite measure spaces. Now we can prove theorem 2.17:

Proof of theorem 2.17. a) This follows from (35) and the Feynman-Kac formulae.

b) We will use part a) for the proof: Setting $v := \underline{V}$, we have to show that

$$e^{-tH_0(v)} \in \mathcal{L}(L^2(M), L^q(M)).$$

Note that using $-v \leq v^-$, proposition 2.5 implies

$$\operatorname{ess\,sup}_{x \in M} \mathbb{E} \left[e^{-\beta \int_0^t v(B_s(x)) ds} \right] \leq 2e^{tC(\beta v^-)} \quad \text{for all } \beta \geq 0. \quad (50)$$

Throughout, let $h \in L^2(M)$.

Case $q = \infty$: One has

$$\begin{aligned} \|e^{-tH_0(v)} h\|_\infty &\leq \operatorname{ess\,sup}_{x \in M} \mathbb{E} \left[e^{-\int_0^t v(B_s(x)) ds} |h(B_t(x))| \right] \\ &\leq \operatorname{ess\,sup}_{x \in M} \mathbb{E} \left[e^{-2 \int_0^t v(B_s(x)) ds} \right]^{\frac{1}{2}} \operatorname{ess\,sup}_{x \in M} \mathbb{E} [|h(B_t(x))|^2]^{\frac{1}{2}} \\ &\leq 2e^{\frac{tC(2v^-)}{2}} \|e^{t\Delta} h\|_\infty \\ &\leq 2e^{\frac{tC(2v^-)}{p}} C(t)^{\frac{1}{2}} \|h\|_2, \end{aligned} \quad (51)$$

where we have used Cauchy-Schwarz for the second step and (49) for the last step.

Case $q < \infty$: We set $l := q/2$. Then

$$\begin{aligned} \|e^{-tH_0(v)} h\|_q^q &\leq \int_M \mathbb{E} \left[e^{-2 \int_0^t v(B_s(x)) ds} |f(B_t(x))|^q \right] \operatorname{vol}(dx) \\ &\leq \int_M \mathbb{E} \left[e^{-2 \int_0^t v(B_s(x)) ds} \right]^l \mathbb{E} [|h(B_t(x))|^2]^l \operatorname{vol}(dx) \\ &\leq 2e^{ltC(2v^-)} \int_M \mathbb{E} [|h(B_t(x))|^2]^l \operatorname{vol}(dx) \end{aligned}$$

follows again from Cauchy-Schwarz. Setting $f := |h|^2$ we get from (49) the inequalities

$$\begin{aligned} \|e^{-tH_0(v)} h\|_q^2 &\leq e^{\frac{tC(2v^-)}{2}} \|e^{t\Delta}\|_{1,l}^{\frac{l}{q}} \|f\|_1^{\frac{l}{q}} = 2e^{\frac{tC(2v^-)}{2}} \|e^{t\Delta}\|_{1,l}^{\frac{l}{q}} \|h\|_2 \\ &\leq 2e^{\frac{tC(2v^-)}{2}} C(t)^{\frac{1}{2} - \frac{1}{q}} \|h\|_2. \end{aligned} \quad (52)$$

This completes the proof. ■

As a next goal, we want to prove that under very general assumptions on M and V (which should still include practically all physically relevant situations), the operator $e^{-tH(V)}$ has a

$$L^2 \rightsquigarrow \{\text{bounded continuous}\}$$

smoothing property for all $t > 0$. In detail, this is:

Theorem 2.20 *Let M be geodesically complete with Ricci curvature bounded from below and*

$$\sup_{x,y \in M} p_t(x,y) < \infty \quad \text{for all } t > 0.$$

Assume furthermore that V is a potential with

$$V^+ \in \Gamma_{\mathcal{K}_{\text{loc}}}(M, \text{End}(E)), \quad V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E)). \quad (53)$$

Then for any $t > 0$, $f \in \Gamma_{L^2}(M, E)$, the section

$$M \longrightarrow E, \quad x \longmapsto \mathbb{E} \left[\mathcal{V}_t^{(x)} //_{t^{x,-1}} f(B_t(x)) \right] \in E_x \quad (54)$$

is well-defined, continuous and bounded. In particular, $e^{-tH(V)}f$ has a continuous bounded representative for any $t > 0$, $f \in \Gamma_{L^2}(M, E)$, and each eigensection of $H(V)$ can be chosen continuous and bounded.

Potentials with the property (53) are usually called *Kato decomposable* in the mathematical physics literature.

Remark 2.21 1. If M is geodesically complete with Ricci curvature bounded from below and a positive injectivity radius, then the assumptions on the Riemannian structure of M are satisfied.

2. Theorem 2.20 generalizes one of the main results of [12], where we have considered the case $V_- = 0$ on Euclidean vector bundles of the form $\mathbb{R}^m \times \mathbb{C}^d$. It also generalizes one of the main results of [4]: In the latter paper the authors have considered scalar operators of the form $H_\beta(v)$ on manifolds with a bounded geometry.

3. Let us explain our approach for proving theorem 2.20: In the situation of theorem 2.20, let

$$Q_t^V f(x) := \mathbb{E} \left[\mathcal{V}_t^{(x)} //_{t^{x,-1}} f(B_t(x)) \right]. \quad (55)$$

Firstly, we remark that under our assumptions on M and V , the right-hand side of (55) will indeed turn out to be well-defined for *all* $x \in M$ (see proposition 2.22). We will use semigroup domination and theorem 2.16 to prove that $Q_t^V f$ is bounded. Furthermore, one can prove that $Q_\bullet^V f(x)$ satisfies a semigroup property for all $x \in M$ (a priori, this is only clear for *a.e.* $x \in M$, and the proof that it remains true for *all* $x \in M$ is actually quite technical). From these considerations, it is clear that we may assume

$$f \in \Gamma_{L^\infty}(M, E) \cap \Gamma_{L^2}(M, E).$$

Next, we will use local elliptic regularity to prove that $Q_t^0 \tilde{f}$ is C^∞ for any $t > 0$ and any essentially bounded square integrable \tilde{f} , so that the continuity of $Q_t^V f$ will follow, if we can locally uniformly approximate $Q_t^V f$ as $s \searrow 0$ by $Q_s^0 Q_{t-s}^V f$. This will in fact follow from the perturbation formula (64) below and the convergence (106). The latter of which strongly relies on the assumption that the potential is in the local Kato class. These techniques extend the corresponding ones from [5] (see also [3]) for usual scalar operators to our setting, where we remark that the proofs of assertions like proposition 2.23, proposition 2.24 or proposition 2.27 are almost trivial in the setting of [5].

The following five propositions will help us to turn the considerations of remark 2.21.2 into a full proof. Firstly, we shall prove the asserted well-definedness of the right-hand side of the Feynman-Kac formula. We will actually need a slightly more general result:

Proposition 2.22 *Under the assumptions of theorem 2.20, the process $\mathcal{V}^{(x)}$ exists for all $x \in M$, and for any $0 \leq s \leq t$ one has*

$$\mathbb{E} \left[\left\| \mathcal{V}_s^{(x), -1} \mathcal{V}_t^{(x)} \right\|_x \left\| f(B_t(x)) \right\|_x \right] \leq \sqrt{2} e^{\frac{C(2|V^-|)t}{2}} \sqrt{e^{\frac{t}{2}\Delta} |f|^2(x)} \quad (56)$$

$$\leq \sqrt{2} e^{\frac{C(2|V^-|)t}{2}} \left(\sup_{y, z \in M} p_t(y, z) \right)^{\frac{1}{2}} \|f\|. \quad (57)$$

Proof. Clearly, lemma 2.4 and the Banach fixed point theorem imply the existence of $\mathcal{V}^{(x)}$ for all x .

Noting

$$\left\| \mathcal{V}_s^{(x), -1} \mathcal{V}_t^{(x)} \right\|_x \leq e^{\int_0^t \|V^-(B_u(x))\|_{B_u(x)} du} \quad \mathbb{P}\text{-a.s.}, \quad (58)$$

which follows from combining proposition 4.1 d) with

$$-V \leq -\underline{V}\mathbf{1}, \quad -\underline{V} \leq \underline{V}^- \leq |V^-|, \quad (59)$$

we can use proposition 2.5 to estimate

$$\mathbb{E} \left[\left\| \mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} \right\|_x^2 \right] \leq 2e^{C(2|V^-)t}. \quad (60)$$

Using Cauchy-Schwarz, (60) implies

$$\begin{aligned} & \mathbb{E} \left[\left\| \mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} \right\|_x \|f(B_t(x))\|_x \right] \\ & \leq \mathbb{E} \left[\left\| \mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} \right\|_x^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\|f(B_t(x))\|_x^2 \right]^{\frac{1}{2}} \\ & = \mathbb{E} \left[\left\| \mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} \right\|_x^2 \right]^{\frac{1}{2}} \left(\int_M p_t(x, y) \|f(y)\|_y^2 \text{vol}(dy) \right)^{\frac{1}{2}} \\ & \leq \sqrt{2} e^{\frac{C(2|V^-)t}{2}} \left(\sup_{y, z \in M} p_t(y, z) \right)^{\frac{1}{2}} \|f\|, \end{aligned} \quad (61)$$

and the proof is complete. ■

Next, we prove the asserted semigroup property and the perturbation formula, respectively:

Proposition 2.23 *Under the assumptions of theorem 2.20, let*

$$Q_t^V f(x) := \mathbb{E} \left[\mathcal{V}_t^{(x)} //_{t,x}^{-1} f(B_t(x)) \right] \quad \text{for any } t \geq 0, x \in M. \quad (62)$$

a) $Q_{\bullet}^V f$ satisfies a pointwise semigroup identity,

$$Q_{s+t}^V f(x) = Q_s^V Q_t^V f(x) \quad \text{for any } s, t \geq 0, x \in M. \quad (63)$$

b) One has the following perturbation formula for any $t \geq s \geq 0, x \in M$,

$$Q_s^0 Q_{t-s}^V f(x) = \mathbb{E} \left[\mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} //_{t,x}^{-1} f(B_t(x)) \right]. \quad (64)$$

Proof. Note first that all terms in (62) - (64) are indeed pointwise well-defined, which is implied by proposition 2.22 and proposition 2.19. The proposition will now be proved in four steps.

I) (63) and (64) hold under the additional assumptions $V \in \Gamma_{L^\infty}(M, \text{End}(E))$ and $f \in \Gamma_{L^\infty}(M, E)$.

Proof: We have to introduce some notation first: In view of (1), for any starting time $a \geq 0$ and any appropriate \mathcal{F}_a -measurable $h : \Omega \rightarrow M$, we define the processes $B^{a,h}$, $//_{a,h}$ and $\mathcal{V}^{a,h}$ as follows:

$$B^{a,h} : [a, \infty) \times \Omega \longrightarrow M$$

is defined as the maximal solution of

$$dB^{a,h} = \sum_{j=1}^l A_j(B^{a,h}) \underline{d}W^j, \quad B_a^{a,h} = h,$$

$//^{a,h}$ is defined as the stochastic parallel transport corresponding to $B^{a,h}$, so that

$$//_b^{a,h} : E_h \longrightarrow E_{B_b^{a,h}} \quad \text{for any } b \geq a,$$

and, finally, for \mathbb{P} -a.e. $\omega \in \Omega$, the map

$$\mathcal{V}_{\bullet}^{a,h}(\omega) : [0, \infty) \longrightarrow \text{End}(E)_{h(\omega)}$$

is defined as the weak solution of

$$d\mathcal{V}_t^{a,h}(\omega) = -\mathcal{V}_t^{a,h}(\omega) \left(//_{a+t}^{a,h,-1} V(B_{a+t}^{a,h}) //_{a+t}^{a,h} \right) (\omega) dt, \quad \mathcal{V}_0^{a,h}(\omega) = \mathbf{1}.$$

Note that $\mathcal{V}_t^{a,h}(\omega)$ can be expanded as in (123) and that our usual notation implies

$$(B^{0,x}, //^{0,x}, \mathcal{V}^{0,x}) = (B(x), //^x, \mathcal{V}^x).$$

Proof of (63): Let U^x be a lift of $B(x)$ and let $U^{s,B_s(x)}$ be the lift of $B^{s,B_s(x)}$ from U_s^x . Then we have $//^x = U^x U_0^{x,-1}$ and $//^{s,B_s(x)} = U^{s,B_s(x)} U_s^{x,-1}$, so that the flow property of the solutions of

$$dU = \sum_{j=1}^l A_j^*(U) \underline{d}W^j \tag{65}$$

gives

$$//_{s+t}^x = //_{s+t}^{s,B_s(x)} //_s^x \quad \mathbb{P}\text{-a.s.} \tag{66}$$

Using (66) and the flow property of the solutions of

$$dB = \sum_{j=1}^l A_j(B) \underline{d}W^j, \tag{67}$$

one easily checks that for fixed s , the processes

$$\mathcal{V}_{s+\bullet}^{(x)} \quad \text{and} \quad \mathcal{V}_s^{(x)} //_s^{x,-1} \mathcal{V}_{\bullet}^{s,B_s(x)} //_s^x$$

both solve the same $\text{End}(E)_x$ -valued initial value problem, so that by uniqueness and (66) we get the multiplicative property

$$\mathcal{V}_{s+t}^{(x)} //_{s+t}^{x,-1} = \mathcal{V}_s^{(x)} //_s^{x,-1} \mathcal{V}_t^{s,B_s(x)} //_{s+t}^{s,B_s(x),-1} \quad \mathbb{P}\text{-a.s.} \tag{68}$$

With $\mathbb{E}^{\mathcal{F}_s}[\bullet] := \mathbb{E}[\bullet | \mathcal{F}_s]$ the last identity implies

$$\begin{aligned} Q_{s+t}^V f(x) &= \mathbb{E} \left[\mathcal{V}_s^x //_{s,x,-1} \mathcal{V}_t^{s,B_s(x)} //_{s+t}^{s,B_s(x),-1} f(B_{s+t}(x)) \right] \\ &= \mathbb{E} \left[\mathcal{V}_s^x //_{s,x,-1} \mathbb{E}^{\mathcal{F}_s} \left[\mathcal{V}_t^{s,B_s(x)} //_{s+t}^{s,B_s(x),-1} f \left(B_{s+t}^{s,B_s(x)} \right) \right] \right]. \end{aligned} \quad (69)$$

Since \mathcal{F}_s is independent from \mathcal{F}_{s+t} and since by its definition $\mathcal{V}_t^{s,B_s(x)}$ clearly is an \mathcal{F}_{s+t} -random variable, we can use lemma 6.3.1 in [13] to conclude

$$\begin{aligned} &\mathbb{E} \left[\mathcal{V}_s^x //_{s,x,-1} \mathbb{E}^{\mathcal{F}_s} \left[\mathcal{V}_t^{s,B_s(x)} //_{s+t}^{s,B_s(x),-1} f \left(B_{s+t}^{s,B_s(x)} \right) \right] \right] \\ &= \int_{\Omega} \mathcal{V}_s^x(\omega) //_{s,x,-1}(\omega) \int_{\Omega} Z_t^{s,B_s(x)(\omega)}(\tilde{\omega}) \mathbb{P}(d\tilde{\omega}) \mathbb{P}(d\omega), \end{aligned} \quad (70)$$

with

$$Z_t^{a,y} := \mathcal{V}_t^{a,y} //_{a+t}^{a,y} f(B_{a+t}^{a,y}) \quad \text{for any } a \geq 0, y \in M.$$

It follows that it is sufficient to prove

$$\mathbb{E} [Z_t^{s,y}] = \mathbb{E} [Z_t^{0,y}] \quad \text{for any } y \in M. \quad (71)$$

Let $\pi : P(E) \rightarrow M$ denote the principal bundle projection, let U^y be a lift of $B(y)$ and let $U^{s,y}$ be the lift of $B^{s,y}$ from U_0^y . Since parallel transport does not depend on the particular choice of the initial frame, we have $//^{s,y} = U^{s,y} U_0^{y,-1}$, and clearly we have $B^{s,y} = \pi(U^{s,y})$, $//^y = U^y U_0^{y,-1}$, $B(y) = \pi(U^y)$. For any $y \in M$ and $n \in \mathbb{N}$ we define a function $\mathcal{A}_n^{t,y}$ by setting

$$\begin{aligned} \mathcal{A}_n^{t,y} &: C([0, \infty), P(E)) \longrightarrow E_y, \\ \mathcal{A}_n^{t,y}(\gamma[\bullet]) &:= \left\{ \prod_{1 \leq j \leq n}^{\rightarrow} \left(\mathbf{1} + \frac{t}{n} U_0^y \gamma[(tj)/n]^{-1} V \left(\pi(\gamma[(tj)/n]) \right) \gamma[(tj)/n] U_0^{y,-1} \right) \right\} \\ &\quad \times U_0^y \gamma[t]^{-1} f \left(\pi(\gamma[t]) \right). \end{aligned} \quad (72)$$

Then we have the following inequalities,

$$\left\| \mathcal{A}_n^{t,y}(U_{s+\bullet}^{s,y}) \right\|_y \leq e^{t\|V\|_{\infty}} \|f(B_{s+t}^{s,y})\|_{B_{s+t}^{s,y}} \quad \mathbb{P}\text{-a.s.} \quad (73)$$

and

$$\left\| \mathcal{A}_n^{t,y}(U^y) \right\|_y \leq e^{t\|V\|_{\infty}} \|f(B_t(y))\|_{B_t(y)} \quad \mathbb{P}\text{-a.s.} \quad (74)$$

Since $\mathcal{V}^{s,y}$ and $\mathcal{V}^{(y)}$ can be represented as product integrals (this follows from applying theorem 7.1 in [6] with $z \mapsto 1 + z$ together with the corresponding remarks on page 56), one has

$$\lim_{n \rightarrow \infty} \mathcal{A}_n^{t,y}(U_{s+\bullet}^{s,y}) = Z_t^{s,y} \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathcal{A}_n^{t,y}(U^y) = Z_t^{0,y} \quad \mathbb{P}\text{-a.s.}$$

Now we note that $U_{s+\bullet}^{s,y}$ and U^y have the same law⁸, so that we can use dominated convergence (in view of (73) and (74)) to deduce

$$\mathbb{E}[Z_t^{s,y}] = \lim_{n \rightarrow \infty} \mathbb{E}[\mathcal{A}_n^{t,y}(U_{s+\bullet}^{s,y})] = \lim_{n \rightarrow \infty} \mathbb{E}[\mathcal{A}_n^{t,y}(U^y)] = \mathbb{E}[Z_t^{0,y}].$$

Proof of (64): We calculate

$$\begin{aligned} Q_s^0 Q_{t-s}^V f(x) &= \int_{\Omega} //_{s,x,-1}(\omega) \int_{\Omega} \mathcal{V}_{t-s}^{(B_s(x)(\omega))}(\tilde{\omega}) //_{t-s}^{B_s(x)(\omega)}(\tilde{\omega}) \\ &\quad \times f\left(B_{t-s}\left(B_s(x)(\omega)\right)(\tilde{\omega})\right) \mathbb{P}(d\tilde{\omega}) \mathbb{P}(d\omega) \\ &= \int_{\Omega} //_{s,x,-1}(\omega) \int_{\Omega} \mathcal{V}_{t-s}^{s,B_s(x)(\omega)}(\tilde{\omega}) //_{t-s}^{s,B_s(x)(\omega)}(\tilde{\omega}) \\ &\quad \times f\left(B_t^{s,B_s(x)(\omega)}(\tilde{\omega})\right) \mathbb{P}(d\tilde{\omega}) \mathbb{P}(d\omega) \\ &= \mathbb{E}\left[//_{s,x,-1} \mathbb{E}^{\mathcal{F}_s} \left[//_{s,x} \mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} //_{t,x,-1} f\left(B_t^{s,B_s(x)}\right)\right]\right] \\ &= \mathbb{E}\left[\mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} //_{t,x,-1} f\left(B_t(x)\right)\right], \end{aligned} \quad (75)$$

where we have used (71) for the second equality, lemma 6.3.1 in [13] together with (68) for the third equality, and the flow property of (67) for the last equality.

II) (63) and (64) hold under the additional assumptions $V \geq C$ and $f \in \Gamma_{L^\infty}(M, E)$.

Proof of (63): We can assume $V \geq 0$ and we define $V_n := \min(n, V)$ for any $n \in \mathbb{N}$. Then each V_n is a potential in $\Gamma_{L^\infty}(M, \text{End}(E))$ so that by applying II) we have for all n ,

$$Q_{s+t}^{V_n} f(x) = Q_s^{V_n} Q_t^{V_n} f(x). \quad (76)$$

Furthermore, the following two identities are included in the first part of the proof of theorem 2.11: For all $a \geq 0$, $y \in M$,

$$\lim_{n \rightarrow \infty} \mathcal{V}_{n,a}^{(y)} = \mathcal{V}_a^{(y)} \quad \mathbb{P}\text{-a.s.} \quad (77)$$

(with an obvious notation), and

$$\lim_{n \rightarrow \infty} Q_a^{V_n} f(y) = Q_a^V f(y). \quad (78)$$

⁸To see this, note first that the smoothness of the vector fields A_j^* implies the uniqueness in law for (65) (this follows from theorem 1.1.10 in [17] and the Whitney embedding theorem). Now one can use the same arguments as in the proof of corollary 1 to Satz 6.40 in [13] to deduce that $U_{s+\bullet}^{s,y}$ and U^y are equal in law.

Thus it remains to prove

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\mathcal{V}_{n,s}^{(x)} //_{s}^{x,-1} Q_t^{V_n} f(B_s(x)) \right] = \mathbb{E} \left[\lim_{n \rightarrow \infty} \mathcal{V}_{n,s}^{(x)} //_{s}^{x,-1} Q_t^{V_n} f(B_s(x)) \right]. \quad (79)$$

To this end, we remark

$$\left\| \mathcal{V}_{n,a}^{(y)} \right\|_y \leq 1 \quad \mathbb{P}\text{-a.s. for all } y \in M, n \in \mathbb{N}, a \geq 0 \quad (80)$$

(this is also included in the first part of the proof of theorem 2.11), so that

$$\left\| \mathcal{V}_{n,s}^{(x)} //_{s}^{x,-1} Q^{V_n} f(B_s(x)) \right\|_x \leq \|f\|_{\infty} \quad \mathbb{P}\text{-a.s.}, \quad (81)$$

and (79) follows from dominated convergence.

Proof of (64): Again, we may assume $V \geq 0$ and we define $V_n := \min(n, V)$. Then by II) we have for all n ,

$$Q_s^0 Q_{t-s}^{V_n} f(x) = \mathbb{E} \left[\mathcal{V}_{n,s}^{(x),-1} \mathcal{V}_{n,t}^{(x)} //_{t}^{x,-1} f(B_t(x)) \right]. \quad (82)$$

Furthermore, (77) implies

$$\lim_{n \rightarrow \infty} \mathcal{V}_{n,t}^{(x)} = \mathcal{V}_t^{(x)}, \quad \lim_{n \rightarrow \infty} \mathcal{V}_{n,s}^{(x),-1} = \mathcal{V}_s^{(x),-1} \quad \mathbb{P}\text{-a.s.}, \quad (83)$$

and we can use

$$\left\| \mathcal{V}_{n,s}^{(x),-1} \mathcal{V}_{n,t}^{(x)} //_{t}^{x,-1} f(B_t(x)) \right\|_x \leq \|f\|_{\infty} \quad \mathbb{P}\text{-a.s.} \quad (84)$$

(which follows from $-V_n \leq 0$ and proposition 4.1 d)) to conclude

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\mathcal{V}_{n,s}^{(x),-1} \mathcal{V}_{n,t}^{(x)} //_{t}^{x,-1} f(B_t(x)) \right] = \mathbb{E} \left[\mathcal{V}_s^{(x),-1} \mathcal{V}_t^{(x)} //_{t}^{x,-1} f(B_t(x)) \right]. \quad (85)$$

It remains to prove

$$\lim_{n \rightarrow \infty} Q_s^0 Q_{t-s}^{V_n} f(x) = Q_s^0 Q_{t-s}^V f(x). \quad (86)$$

To this end, we just note that by (80) we have

$$\left\| //_{s}^{x,-1} Q_{t-s}^{V_n} f(B_s(x)) \right\|_x \leq \|f\|_{\infty} \quad \mathbb{P}\text{-a.s.}, \quad (87)$$

so that (86) follows from dominated convergence and (78).

III) (63) and (64) hold under the additional assumption $f \in \Gamma_{L^\infty}(M, E)$.

Proof of (63): We define $V_n := \max(-n, V)$ for any $n \in \mathbb{N}$. Then each V_n is a bounded from below potential with $V_n^+ = V^+$ locally Kato. By III) we

have (76) and it follows from the second part of the proof of theorem 2.11 that one also has (77) again. For the proof of (78) note that for all $y \in M$, $n \in \mathbb{N}$, $a \geq 0$ one has

$$\|\mathcal{Y}_{n,a}^{(y)}\|_y \leq e^{-\int_0^a \underline{V}(B_u(y))du} \leq e^{\int_0^a \|V^-(B_u(y))\|_{B_u(y)} du} \quad (88)$$

(only the first inequality nontrivial; but this is included in the second part of the proof of theorem 2.11), so

$$\|\mathcal{Y}_{n,a}^{(y)} //_{a}^{x,-1} f(B_a(y))\|_x \leq \|f\|_\infty e^{\int_0^a \|V^-(B_u(x))\|_{B_u(x)} du} \quad \mathbb{P}\text{-a.s.} \quad (89)$$

and the last term is in $L^1(\mathbb{P})$ by proposition 2.5, so that (78) follows from (77) and dominated convergence. It remains to prove (79). But in view of (88) we have

$$\begin{aligned} & \|\mathcal{Y}_{n,s}^{(x)} //_{s}^{x,-1} Q_t^{V_n} f(B_s(x))\|_x \\ & \leq \|f\|_\infty e^{\int_0^s \|V^-(B_u(x))\|_{B_u(x)} du} \mathbb{E} \left[\|\mathcal{Y}_{n,t}^{(y)}\|_y \Big|_{y=B_s(x)} \right] \\ & \leq \|f\|_\infty e^{\int_0^s \|V^-(B_u(x))\|_{B_u(x)} du} \mathbb{E} \left[e^{\int_0^t \|V^-(B_u(y))\|_{B_u(y)} du} \Big|_{y=B_s(x)} \right] \\ & \leq 2 \|f\|_\infty e^{tC(\|V^-\|)} e^{\int_0^s \|V^-(B_u(x))\|_{B_u(x)} du} \quad \mathbb{P}\text{-a.s.}, \end{aligned} \quad (90)$$

and the last term is in $L^1(\mathbb{P})$ by proposition 2.5. Now (79) follows from dominated convergence.

Proof of (64): Again, let V_n be given by $\max(-n, V)$. Then by (77) we have (83), and one furthermore has

$$\left\| \mathcal{Y}_{n,s}^{(x),-1} \mathcal{Y}_{n,t}^{(x)} //_{t}^{x,-1} f(B_t(x)) \right\|_x \leq \|f\|_\infty e^{\int_0^t \|V^-(B_u(x))\|_{B_u(x)} du} \in L^1(\mathbb{P}) \quad (91)$$

(which follows from proposition 4.1 d) and $-\underline{V}_n \leq -\underline{V}$; the latter inequality is included in the second part of the proof of theorem 2.11) so that we have (85) by dominated convergence. It remains to prove (86). To this end, we can use (78) and

$$\begin{aligned} & \left\| //_{s}^{x,-1} Q_{t-s}^{V_n} f(B_s(x)) \right\|_x \leq \|f\|_\infty \mathbb{E} \left[\left\| \mathcal{Y}_{n,t-s}^{(y)} \right\|_y \Big|_{y=B_s(x)} \right] \\ & \leq \|f\|_\infty \mathbb{E} \left[e^{\int_0^{t-s} \|V^-(B_u(y))\|_{B_u(y)} du} \Big|_{y=B_s(x)} \right] \\ & \leq 2 \|f\|_\infty e^{(t-s)C(\|V^-\|)} \quad \mathbb{P}\text{-a.s.}, \end{aligned} \quad (92)$$

which follows from (88) and proposition 2.5, to deduce (86) with dominated convergence again.

IV) (63) and (64) hold in the general situation.

Proof: It remains to remove the condition that f is essentially bounded. To this end, one can consider $f_n := 1_{K_n(\mathcal{O})}f \in \Gamma_{L^\infty}(M, E)$ for some fixed reference point \mathcal{O} , apply III) to the f_n 's and take $n \rightarrow \infty$ to deduce this assertion with dominated convergence. ■

Next, we will prove:

Proposition 2.24 *Let M be geodesically complete and stochastically complete. Then for any $t > 0$ and any*

$$f \in \Gamma_{L^\infty}(M, E) \cap \Gamma_{L^2}(M, E),$$

the section given by

$$M \longrightarrow E, \quad x \longmapsto \mathbb{E} \left[//_{t}^{x, -1} f(B_t(x)) \right] \in E_x \quad (93)$$

is the C^∞ -representative of $e^{-tH(0)}f$.

Proof. By local elliptic regularity, $e^{-sH(0)}f$ has a C^∞ -representative which we denote with $f_s(\bullet)$ for all $s > 0$. Furthermore, the map $(s, y) \mapsto f_s(y)$ is C^∞ and one has

$$\partial_s f_s(y) = -\frac{1}{2} \nabla^* \nabla f_s(y) \quad \text{for all } s > 0, y \in M. \quad (94)$$

We fix arbitrary $x \in M$, $t > 0$ and $\epsilon > 0$ now. Then the time dependent version of formula (12) from [9] combined with the above (94) gives

$$\begin{aligned} & d_s \left(//_{s}^{x, -1} f_{t-s+\epsilon}(B_s(x)) \right) \\ &= //_{s}^{x, -1} \sum_{j=1}^l (\nabla_{A_j} f_{t-s+\epsilon})(B_s(x)) dW_s^j - \frac{1}{2} //_{s}^{x, -1} \nabla^* \nabla f_{t-s+\epsilon}(B_s(x)) ds \\ & \quad + //_{s}^{x, -1} \partial_s f_{t-s+\epsilon}(B_s(x)) ds \\ &= //_{s}^{x, -1} \sum_{j=1}^l (\nabla_{A_j} f_{t-s+\epsilon})(B_s(x)) dW_s^j \end{aligned} \quad (95)$$

for all $0 \leq s \leq t$, which implies that the process

$$N(x, t, \epsilon) : [0, t] \times \Omega \longrightarrow E_x, \quad N_s(x, t, \epsilon) := //_{s}^{x, -1} f_{t-s+\epsilon}(B_s(x))$$

is a continuous local martingale. It is in fact a martingale: For any $0 \leq s \leq t$ the following inequalities hold \mathbb{P} -a.s.,

$$\begin{aligned} \|N_s(x, t, \epsilon)\|_x &\leq \|f_{t-s+\epsilon}(B_s(x))\|_{B_s(x)} \\ &\leq \int_M p_{t-s+\epsilon}(B_s(x), y) \|f(y)\|_y \text{vol}(dy) \\ &\leq \|f\|_\infty \int_M p_{t-s+\epsilon}(B_s(x), y) \text{vol}(dy) = \|f\|_\infty, \end{aligned} \quad (96)$$

where we have used (35) with $V = 0$, $v = 0$ for the second inequality⁹, so that the martingale property of $N(x, t, \epsilon)$ follows from a standard criterion (see for example p.129 in [30]). This shows

$$f_{t+\epsilon}(x) = \mathbb{E}[N_0(x, t, \epsilon)] = \mathbb{E}[N_t(x, t, \epsilon)] = \mathbb{E}[\int_t^{x,-1} f_\epsilon(B_t(x))]. \quad (97)$$

Since $f_s = e^{-sH(0)} f \rightarrow f$ as $s \searrow 0$ with respect to $\Gamma_{L^2}(M, E)$, there exists a sequence $(\epsilon_n) \subset (0, \infty)$ with $\epsilon_n \rightarrow 0$ and $f_{\epsilon_n}(y) \rightarrow f(y)$ as $n \rightarrow \infty$ for a.e. $y \in M$, so that (in view of (96)) we can use dominated convergence and (97) with $\epsilon = \epsilon_n$ to conclude

$$\mathbb{E}[\int_t^{x,-1} f(B_t(x))] = \lim_{n \rightarrow \infty} \mathbb{E}[\int_t^{x,-1} f_{\epsilon_n}(B_t(x))] = \lim_{n \rightarrow \infty} f_{t+\epsilon_n}(x) = f_t(x),$$

and the proof is complete. ■

The next proposition is concerned with the first exit time of $B(x)$ from open geodesic balls, where x runs through a compact set. Although the arguments of the proof that we are going to present are certainly well-known from proofs of stochastic completeness, the result itself has not yet appeared in the literature, as far as we know.

Proposition 2.25 *Let M be geodesically complete with Ricci curvature bounded from below, fix some origin $\mathcal{O} \in M$, and for any $t > 0$, $x \in M$, $r > 0$ let*

$$\chi(r, t, x) := 1_{\{t < \zeta(r, x)\}},$$

where $\zeta(r, x)$ stands for the first exit time of $B(x)$ from $K_r(\mathcal{O})$. If $K \subset M$ is compact, then one has

$$\limsup_{t \searrow 0} \sup_{x \in K} \mathbb{E}[1 - \chi(r, t, x)] = 0 \quad \text{for any } r > \max_{x \in K} d(\mathcal{O}, x). \quad (98)$$

⁹Note that (35) is true for *all* x in the C^∞ case.

Remark 2.26 1. Note that (98) is nothing but

$$\liminf_{t \searrow 0} \inf_{x \in K} \mathbb{P}\{t < \zeta(r, x)\} = 1 \quad \text{for any } r > \max_{x \in K} d(\mathcal{O}, x). \quad (99)$$

2. By using the techniques of [18], it should be possible to relax the assumption $\text{Ric} \geq C$ considerably.

Proof of proposition 2.25. Since $B(x)$ is continuous, we can assume $K \neq \{\mathcal{O}\}$. Let $R(x) := d(\mathcal{O}, x)$. Then R is a smooth function on the open set $M \setminus (\text{Cut}(\mathcal{O}) \cup \{\mathcal{O}\})$. If $C > 0$ is such that the Ricci curvature of M is bounded from below by $-C$, then inequality (2.3) of [28] implies

$$\Delta R(x) \leq h(R(x)) \quad \text{for all } x \in M \setminus (\text{Cut}(\mathcal{O}) \cup \{\mathcal{O}\}), \quad (100)$$

where

$$h : (0, \infty) \longrightarrow (0, \infty), \quad h(r) := \frac{m-1}{r} + \frac{C}{3}r.$$

Furthermore, although $R \notin C^\infty(M)$, the process $R(B(x))$ is a continuous semi-martingale [23] which satisfies

$$R(B_t(x)) - R(x) = Z_t^x + \int_0^t \Delta R(B_s(x)) ds - L_t^x \quad \mathbb{P}\text{-a.s.} \quad (101)$$

for any $t \geq 0$, $x \in M$, where Z^x is a Brownian motion which starts in 0, L^x is a continuous nondecreasing process which starts in 0, and where the integral can be defined since $B(x)$ does not spend time in $\text{Cut}(\mathcal{O}) \cup \{\mathcal{O}\}$ (this follows from the well-known fact that $\text{Cut}(\mathcal{O}) \cup \{\mathcal{O}\}$ has measure zero; see p.527 in [13] for details). For any $x \in M$ let $Y^x : [0, \infty) \times \Omega \rightarrow (0, \infty)$ be the uniquely determined maximal solution of

$$dY^x = dZ^x + h(Y^x)dt, \quad Y_0^x = \max_{x \in K} R(x) (> 0), \quad (102)$$

where we remark that the Feller explosion test as formulated in proposition 4.2.2 in [17] can be checked with elementary estimates to prove that Y^x is indeed nonexplosive. Furthermore, (100), (101) and a classical comparison theorem for stochastic differential equations (theorem 1.1 in [19]) imply

$$R(B_t(x)) \leq Y_t^x \quad \mathbb{P}\text{-a.s. for any } t \geq 0, x \in K. \quad (103)$$

Now (103) shows the following uniform estimate in x : For any $t \geq 0$ and $r > 0$,

$$\begin{aligned} \inf_{x \in K} \mathbb{P}\{t < \zeta(r, x)\} &= \inf_{x \in K} \mathbb{P}\{R(B_s(x)) < r \text{ for all } s \in [0, t]\} \\ &\geq \inf_{x \in K} \mathbb{P}\{Y_s^x < r \text{ for all } s \in [0, t]\} \\ &= \mathbb{P}\{Y_s^{x^*} < r \text{ for all } s \in [0, t]\}, \end{aligned} \quad (104)$$

where x^* is an arbitrary point in M and where we have used uniqueness in law for the pair $(1, h)$. Finally, (99) follows from the fact that if $r > \max_{x \in K} R(x)$, then (by the continuity of Y^{x^*}) the last term in (104) tends to 1 as $t \searrow 0$. ■

We will use proposition 2.25 to prove part b) of:

Proposition 2.27 a) *Let M be stochastically complete and let*

$$V \in \Gamma_{\mathcal{K}}(M, \text{End}(E)).$$

Then one has

$$\limsup_{t \searrow 0} \sup_{x \in M} \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] = 0. \quad (105)$$

b) *Let M be geodesically complete with Ricci curvature bounded from below and let V be a potential with*

$$V^+ \in \Gamma_{\mathcal{K}_{\text{loc}}}(M, \text{End}(E)), \quad V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E)).$$

Then for all compact $K \subset M$ one has

$$\limsup_{t \searrow 0} \sup_{x \in K} \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] = 0. \quad (106)$$

Proof. a) By corollary 4.3 we have \mathbb{P} -a.s.

$$\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x \leq \left(\int_0^t \|V(B_s(x))\|_{B_s(x)} ds \right)^{\frac{1}{4}} e^{\int_0^t \|V(B_s(x))\|_{B_s(x)} ds}, \quad (107)$$

so that Cauchy-Schwarz implies

$$\mathbb{E} \left[\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] \leq \mathbb{E} \left[\int_0^t \|V(B_s(x))\|_{B_s(x)} ds \right]^{\frac{1}{2}} \mathbb{E} \left[e^{4 \int_0^t \|V(B_s(x))\|_{B_s(x)} ds} \right]^{\frac{1}{2}}. \quad (108)$$

Using proposition 2.5, (108) gives

$$\sup_{x \in M} \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] \leq \sqrt{2} e^{\frac{C(4|V|)t}{2}} \sup_{x \in M} \mathbb{E} \left[\int_0^t \|V(B_s(x))\|_{B_s(x)} ds \right]^{\frac{1}{2}}, \quad (109)$$

which tends to zero as $t \searrow 0$ by the definition of the Kato class.

b) Let $\chi(r, t, x)$ be as in proposition 2.25. Then we have

$$\begin{aligned}
& \sup_{x \in K} \mathbb{E} \left[\left(1 - \chi(r, t, x) + \chi(r, t, x) \right) \left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] \\
& \leq \sup_{x \in K} \mathbb{E} [1 - \chi(r, t, x)]^{\frac{1}{2}} \sup_{x \in K} \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right]^{\frac{1}{2}} \\
& \quad + \sup_{x \in K} \mathbb{E} \left[\chi(r, t, x) \left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] \\
& \leq \sqrt{2 + 4e^{C(2|V^-|)t}} \sup_{x \in K} \mathbb{E} [1 - \chi(r, t, x)]^{\frac{1}{2}} + \sup_{x \in K} \mathbb{E} \left[\chi(r, t, x) \left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right],
\end{aligned} \tag{110}$$

where we have used Cauchy-Schwarz and

$$1 - \chi(r, t, x) = (1 - \chi(r, t, x))^2$$

for the first step and (60) with $s = 0$ for the second step. In view of (98), it follows from (110) that it is sufficient to prove that for (some) $r > \max_{x \in K} d(\mathcal{O}, x)$ one has

$$\limsup_{t \searrow 0} \sup_{x \in K} \mathbb{E} \left[\chi(r, t, x) \left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] = 0. \tag{111}$$

To this end, let $t > 0$, $r > \max_{x \in K} d(\mathcal{O}, x)$ and take a $\Psi \in C_0^\infty(M)$ such that $\Psi = 1$ in $K_r(\mathcal{O})$. We denote with $\mathcal{Y}^{\Psi, (x)}$ the pathwise weak solution of (22) with V replaced with ΨV and remark that

$$\Psi V \in \Gamma_{\mathcal{K}}(M, \text{End}(E)).$$

Since in $\{\chi(r, t, x) \neq 0\}$ one \mathbb{P} -a.s. has

$$//_s^{x, -1} V(B_s(x)) //_s^x = //_s^{x, -1} \Psi(B_s(x)) V(B_s(x)) //_s^x \quad \text{for any } 0 \leq s \leq t,$$

expanding $\mathcal{Y}^{(x)}$ and $\mathcal{Y}^{\Psi, (x)}$ into path ordered exponentials as in (123) shows

$$\mathbb{E} \left[\chi(r, t, x) \left\| \mathbf{1} - \mathcal{Y}_t^{(x)} \right\|_x^2 \right] = \mathbb{E} \left[\chi(r, t, x) \left\| \mathbf{1} - \mathcal{Y}_t^{\Psi, (x)} \right\|_x^2 \right],$$

and (111) follows from part a). ■

Now we are prepared to prove theorem 2.20.

Proof of theorem 2.20. The asserted *boundedness* follows from setting $s = 0$ in (57).

Continuity: It follows from remark 2.21 that it is sufficient to prove that for any compact $K \subset M$ and any $f \in \Gamma_{L^\infty}(M, E) \cap \Gamma_{L^2}(M, E)$ one has

$$\limsup_{s \searrow 0} \sup_{x \in K} \|Q_s^0 Q_{t-s}^V f(x) - Q_t^V f(x)\|_x = 0. \quad (112)$$

By (64),

$$\begin{aligned} & \|Q_s^0 Q_{t-s}^V f(x) - Q_t^V f(x)\|_x \\ &= \left\| \mathbb{E} \left[\left(\mathcal{V}_s^{(x), -1} \mathcal{V}_t^{(x)} - \mathcal{V}_t^{(x)} \right) //_{t,x}^{-1} f(B_t(x)) \right] \right\|_x \\ &= \left\| \mathbb{E} \left[\left(\mathbf{1} - \mathcal{V}_s^{(x)} \right) \mathcal{V}_s^{(x), -1} \mathcal{V}_t^{(x)} //_{t,x}^{-1} f(B_t(x)) \right] \right\|_x \\ &\leq \|f\|_\infty \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{V}_s^{(x)} \right\|_x \left\| \mathcal{V}_s^{(x), -1} \mathcal{V}_t^{(x)} \right\|_x \right], \end{aligned} \quad (113)$$

so that using Cauchy-Schwarz with (60) and proposition 2.27 b) we get

$$\begin{aligned} & \sup_{x \in K} \|Q_s^0 Q_{t-s}^V f(x) - Q_t^V f(x)\|_x \\ &\leq 2 \|f\|_\infty e^{\frac{C(2|V^-|)t}{2}} \sup_{x \in K} \mathbb{E} \left[\left\| \mathbf{1} - \mathcal{V}_s^{(x)} \right\|_x^2 \right]^{1/2} \rightarrow 0, \text{ as } s \searrow 0. \end{aligned}$$

This completes the proof. ■

We finally remark the following corollary to theorem 2.20, which is important for geometric applications:

Corollary 2.28 *Let M be geodesically complete with Ricci curvature bounded from below and*

$$\sup_{x, y \in M} p_t(x, y) < \infty \text{ for all } t > 0.$$

Assume furthermore that $V \in \Gamma_{C^\infty}(M, \text{End}(E))$ is a potential with $V^- \in \Gamma_{\mathcal{K}}(M, \text{End}(E))$. Then for any $t > 0$, $f \in \Gamma_{L^2}(M, E)$, the section

$$M \longrightarrow E, \quad x \longmapsto \mathbb{E} \left[\mathcal{V}_t^{(x)} //_{t,x}^{-1} f(B_t(x)) \right] \in E_x \quad (114)$$

is C^∞ and bounded. In particular, $e^{-tH(V)} f$ has a bounded C^∞ -representative which is given by (114).

Proof. Let $Q_t^V f(x)$ be defined by the right hand side of (114). By local elliptic regularity, $e^{-tH(V)} f$ has a C^∞ -representative f_t^V . The Feynman-Kac formula implies $Q_t^V f(x) = f_t^V(x)$ for a.e. x , but since $x \mapsto Q_t^V f(x)$ is continuous (and bounded) by theorem 2.20, we actually have $Q_t^V f(x) = f_t^V(x)$ for *all* x . ■

3 Proof of proposition 2.19

Let us first note the following simple fact: For any $r \geq 1$ and $x \in M$ one has

$$\|p_t(x, \bullet)\|_r \leq C(t, r) := C(t)^{1-\frac{1}{r}}. \quad (115)$$

Throughout, let $h \in L^p(M)$.

Case $1 < p < q < \infty$: Let r be given as $1 - 1/r = 1/p - 1/q$. Applying Hölder's inequality with the exponents

$$q_1 = q, \quad q_2 = \frac{r}{1 - \frac{r}{q}}, \quad q_3 = \frac{p}{1 - \frac{p}{q}}$$

gives

$$\begin{aligned} & \|P_t h\|_q^q \\ & \leq \int_M \left(\int_M (p_t(x, y)^r |h(y)|^p)^{\frac{1}{q}} p_t(x, y)^{1-\frac{r}{q}} |h(y)|^{1-\frac{p}{q}} \text{vol}(dy) \right)^q \text{vol}(dx) \\ & \leq \int_M \left(\int_M p_t(x, y)^r |h(y)|^p \text{vol}(dy) \right) \left(\int_M p_t(x, y)^r \text{vol}(dy) \right)^{\frac{q}{r}(1-\frac{r}{q})} \\ & \quad \times \left(\int_M |h(y)|^p \text{vol}(dy) \right)^{\frac{q}{p}(1-\frac{p}{q})} \text{vol}(dx), \end{aligned}$$

so that using Fubini's theorem and (115),

$$\begin{aligned} \|P_t h\|_q^q & \leq C(t, r)^{q(1-\frac{r}{q})} \|h\|_p^{q(1-\frac{p}{q})} \int_M |h(y)|^p \int_M p_t(x, y)^r \text{vol}(dx) \text{vol}(dy) \\ & \leq C(t, r)^q \|h\|_p^q = C(t)^{q(\frac{1}{p}-\frac{1}{q})} \|h\|_p^q. \end{aligned} \quad (116)$$

Case $1 < p = q < \infty$: One has

$$\begin{aligned} \|P_t h\|_p^p &\leq \int_M \left(\int_M p_t(x, y) |h(y)| \text{vol}(dy) \right)^p \text{vol}(dx) \\ &\leq \int_M \int_M |h(y)|^p p_t(x, y) \text{vol}(dy) \text{vol}(dx) \end{aligned} \quad (117)$$

$$\begin{aligned} &= \int_M \int_M p_t(x, y) \text{vol}(dx) |h(y)|^p \text{vol}(dy) \\ &\leq \|h\|_p^p, \end{aligned} \quad (118)$$

where we have applied the Hölder inequality to the finite measure $\mu(dy) = p_t(x, y) \text{vol}(dy)$ for the second inequality.

Case $1 < p < q = \infty$: This works with the same argument that has been used for the inequality (117).

Case $1 = p < q < \infty$: One has

$$\|P_t h\|_q^q \leq \int_M \left(\int_M (p_t(x, y)^q |h(y)|)^{\frac{1}{q}} |h(y)|^{1-\frac{1}{q}} \text{vol}(dy) \right)^q \text{vol}(dx). \quad (119)$$

Applying the Hölder inequality with the exponents

$$q_1 = q, \quad q_2 = \frac{1}{1 - \frac{1}{q}}$$

gives

$$\|P_t h\|_q^q \leq \|h\|_1^{q-1} \int_M \int_M p_t(x, y)^q |h(y)| \text{vol}(dy) \text{vol}(dx), \quad (120)$$

so that the Fubini theorem and (115) imply

$$\|P_t h\|_q^q \leq C(t)^{q(1-\frac{1}{q})} \|h\|_1^q.$$

The cases $p = q = \infty$ and $p = q = 1$ and $p = 1, q = \infty$ are trivial. ■

4 Some inequalities

Let \mathcal{H} be a d -dimensional, $d < \infty$, complex or real Hilbert space with scalar product $\langle \bullet, \bullet \rangle$ and the corresponding norm $\|\bullet\|$. The induced operator norm will be denoted with the same symbol. If $a \geq 0$, $F \in L_{\text{loc}}^1([a, \infty), \mathcal{L}(\mathcal{H}))$,

then a standard use of the Banach fixed point theorem shows that there is a unique weak (= locally absolutely continuous) solution $Y : [a, \infty) \rightarrow \mathcal{L}(\mathcal{H})$ of the ordinary initial value problem

$$\frac{d}{ds}Y(s) = Y(s)F(s), \quad Y(a) = \mathbf{1}. \quad (121)$$

It is easily seen that Y is invertible with

$$\frac{d}{ds}Y^{-1}(s) = -F(s)Y^{-1}(s), \quad Y^{-1}(a) = \mathbf{1}.$$

Proposition 4.1 *Let F and Y be as above.*

a) *For any $t \geq a$,*

$$\|Y(t)\| \leq e^{\int_a^t \|F(s)\| ds}.$$

b) *Let $a = 0$. For any $t \geq 0$,*

$$\|Y(t) - \mathbf{1}\| \leq e^{\int_0^t \|F(s)\| ds}. \quad (122)$$

c) *Let $t \geq a$, assume that $F(s)$ is Hermitian for a.e. $s \in [a, t]$ and that there exists a real-valued function $c \in L^1[a, t]$ such that for all $v \in \mathcal{H}$ one has*

$$\langle F(s)v, v \rangle \leq c(s) \|v\|^2 \quad \text{for a.e. } s \in [a, t].$$

Then one has

$$\|Y(t)\| \leq e^{\int_a^t c(s) ds}.$$

d) *Let $a = 0$, $0 \leq t_1 \leq t_2$ and assume that $F(s)$ is Hermitian for a.e. $s \in [0, t_2]$ and that there exists a real-valued function $c \in L^1[0, t_2]$ such that for all $v \in \mathcal{H}$ one has*

$$\langle F(s)v, v \rangle \leq c(s) \|v\|^2 \quad \text{for a.e. } s \in [0, t_2].$$

Then one has

$$\|Y^{-1}(t_1)Y(t_2)\| \leq e^{\int_{t_1}^{t_2} c(s) ds}.$$

Proof. a) This is proposition B.1.(a) in [9].

b) This follows easily from expanding Y into the path ordered exponential

$$Y(t) = \mathbf{1} + \int_{0 \leq s_1 \leq \dots \leq s_k \leq t} F(s_1) \dots F(s_k) ds_1 \dots ds_k. \quad (123)$$

c) This is proposition B.1.(b) in [9].

d) This follows from part c), by noting that for fixed t_1 , the function

$$Y^{-1}(t_1)Y(\bullet) : [t_1, \infty) \longrightarrow \mathcal{L}(\mathcal{H})$$

is the solution of (121) with $a = t_1$. ■

Proposition 4.2 Let $F_1, F_2 \in L^1_{\text{loc}}([a, \infty), \mathcal{L}(\mathcal{H}))$ and let

$$Y_1, Y_2 : [a, \infty) \longrightarrow \mathcal{L}(\mathcal{H})$$

be the unique solutions of the ordinary initial value problems

$$\frac{d}{ds} Y_j(s) = Y_j(s) F_j(s), \quad Y_j(a) = \mathbf{1} \quad \text{for } j = 1, 2.$$

The following inequality holds for all $t \geq a$,

$$\|Y_1(t) - Y_2(t)\| \leq e^{2 \int_a^t \|F_1(s)\| ds + \int_a^t \|F_2(s)\| ds} \int_a^t \|F_1(s) - F_2(s)\| ds. \quad (124)$$

Proof. This is proposition B.2 in [9]. ■

Corollary 4.3 Let F and Y be as above and let $a = 0$. Then for any $t \geq 0$ and any $p \geq 1$ one has

$$\|Y(t) - \mathbf{1}\| \leq \left(\int_0^t \|F(s)\| ds \right)^{\frac{1}{p}} e^{\int_0^t \|F(s)\| ds}. \quad (125)$$

Proof. If $\int_0^t \|F(s)\| ds \leq 1$, then (125) follows from applying (124) with $F_1 = 0$. If $\int_0^t \|F(s)\| ds > 1$, then (125) follows from (122). ■

5 Some Hilbert space facts

We collect some well-known Hilbert space facts in the following theorem. The assertions are certainly well-known (see for example [35] [22] [36] for (126)-(129) and [10] for (130)). ■

Theorem 5.1 Let $\mathcal{H} = (\mathcal{H}, \langle \bullet, \bullet \rangle_{\mathcal{H}})$ be a Hilbert space.

a) Let $B \in \mathcal{L}(\mathcal{H})$ be self-adjoint. Then one has

$$\max \sigma(B) = \sup \left\{ \langle B\Psi, \Psi \rangle_{\mathcal{H}} \mid \Psi \in \mathcal{H}, \|\Psi\|_{\mathcal{H}} = 1 \right\}. \quad (126)$$

b) Let $H \geq c_1$ be self-adjoint and let q_H be the quadratic form corresponding to H . Then for any number $c_2 \leq c_1$ one has

$$D(q_H) = D\left((H - c_2)^{\frac{1}{2}}\right), \quad q_H(f) = \left\| (H - c_2)^{\frac{1}{2}} f \right\|_{\mathcal{H}}^2 + c_2 \|f\|_{\mathcal{H}}^2. \quad (127)$$

Furthermore, with $\mathcal{E}_H := \min \sigma(H)$ it holds that

$$\mathcal{E}_H = \inf_{f \in D(q_H)} q_H(f) \quad (128)$$

and

$$\max \sigma(e^{-H}) = e^{-\mathcal{E}_H}. \quad (129)$$

Finally, if $c_1 = 0$, that is $H \geq 0$, then

$$\begin{aligned} D(q_H) &= \left\{ f \mid f \in \mathcal{H}, \lim_{t \searrow 0} \left\langle \frac{f - e^{-tH} f}{t}, f \right\rangle_{\mathcal{H}} < \infty \right\}, \\ q_H(f) &= \lim_{t \searrow 0} \left\langle \frac{f - e^{-tH} f}{t}, f \right\rangle_{\mathcal{H}}. \end{aligned} \quad (130)$$

Acknowledgements. The author would like to thank A. Thalmaier, K. Kuwada and R. Philipowski for helpful discussions on proposition 2.24 and proposition 2.25. The reasearch has been financially supported by the BIGS and the SFB 647.

References

- [1] Bérard, P.H.: *Spectral Geometry: Direct and Inverse Problems*. Lecture Notes in Mathematics, Vol. 1207, Springer, 1986.
- [2] Braverman, M. & Milatovich, O. & Shubin, M.: *Essential self-adjointness of Schrödinger-type operators on manifolds*. Russian Math. Surveys 57 (2002), no. 4, 641–692.
- [3] Broderix, K. & Hundertmark, D. & Leschke, H.: *Continuity properties of Schrödinger semigroups with magnetic fields*. Reviews in Mathematical Physics 12 (2000), 181–225.
- [4] Brüning, J. & Geyler, V. & Pankrashkin, K.: *Continuity properties of integral kernels associated with Schrödinger operators on manifolds*. Ann. Henri Poincaré 8 (2007), no. 4, 781–816.

- [5] Chung, K.L. & Xin, Z.Z.: *From Brownian motion to Schrödinger's equation*. Grundlehren der Mathematischen Wissenschaften, 312. Springer-Verlag, Berlin, 1995.
- [6] Dollard, J.D. & Friedman, C.N.: *Product Integration*. Addison-Wesley, 1979.
- [7] Driver, B.K. & Thalmaier, A.: *Heat equation derivative formulas for vector bundles*. J. Funct. Anal. 183 (2001), no. 1, 42–108.
- [8] Emery, M.: *Stochastic calculus in manifolds*. Springer-Verlag, Berlin, 1989.
- [9] Güneysu, B.: *The Feynman-Kac formula for Schrödinger operators on vector bundles over complete manifolds*. J. Geom. Phys. 60 (2010) 1997–2010.
- [10] Güneysu, B.: *Kato's inequality and form boundedness of Kato potentials on arbitrary Riemannian manifolds*. Preprint.
- [11] Güneysu, B.: *On the Feynman Kac formula for Schrödinger semigroups on vector bundles*. PhD thesis, Bonn (2011).
- [12] Güneysu, B.: *Multiplicative matrix-valued functionals and continuity properties of semigroups corresponding to differential operators with matrix coefficients*. J. Math. Anal. Appl. 380 (2011), 709–725.
- [13] Hackenbroch, W. & Thalmaier, A.: *Stochastische Analysis*. B. G. Teubner, 1994.
- [14] Hempel, R. & Voigt, J.: *The spectrum of a Schrödinger operator in $L_p(\mathbb{R}^\nu)$ is p -independent*. Comm. Math. Phys. 104 (1986), no. 2, 243–250.
- [15] Hess, H. & Schrader, R. & Uhlenbrock, D.A.: *Domination of semigroups and generalization of Kato's inequality*. Duke Math. J. 44, Number 4 (1977), 893–904.
- [16] Hess, H. & Schrader, R. & Uhlenbrock, D.A.: *Kato's inequality and the spectral distribution of Laplacians on compact Riemannian manifolds*. J. Differential Geom. Volume 15, Number 1 (1980), 27–37.
- [17] Hsu, E.: *Stochastic Analysis on Manifolds*. AMS, 2002.
- [18] Hsu, E.: *Heat semigroup on a complete Riemannian manifold*. Annals of Probability 17, Number 3 (1989), 1248–1254.

- [19] Ikeda, N. & Watanabe, S.: *A comparison theorem for solutions of stochastic differential equations and its applications*. Osaka J. Math. 14 (1977), no. 3, 619–633.
- [20] Johnson, G.W. & Lapidus, M. L.: *The Feynman integral and Feynman's operational calculus*. The Clarendon Press, Oxford University Press, 2000.
- [21] Karatzas, I. & Shreve, S. E. *Brownian motion and stochastic calculus*. Graduate Texts in Mathematics, 113. Springer-Verlag, New York, 1988.
- [22] Kato, T.: *Perturbation theory for linear operators*. Die Grundlehren der mathematischen Wissenschaften, Band 132 Springer-Verlag New York, Inc., New York 1966.
- [23] Kendall, W.: *The radial part of Brownian motion on a manifold: a semimartingale property*. Ann. Probab. 15 (1987), no. 4, 1491–1500.
- [24] Kuwae, K. & Takahashi, M.: *Kato class functions of Markov processes under ultracontractivity*. Potential theory in Matsue, 193–202, Adv. Stud. Pure Math., 44, Math. Soc. Japan, Tokyo, 2006.
- [25] Kuwae, K. & Takahashi, M.: *Kato class measures of symmetric Markov processes under heat kernel estimates*. J. Funct. Anal. 250 (2007), no. 1, 86–113.
- [26] Lenz, D. & Keller, M. & Vogt, H. & Wojciechowski, R.: *Note on basic features of large time behaviour of heat kernels*. Preprint.
- [27] Milatovic, O.: *Localized self-adjointness of Schroedinger-type operators on Riemannian manifolds*. J. Math. Anal. Appl. 283 (2003), no. 1, 304–318.
- [28] Qian, Z.: *On conservation of probability and the strong Feller property*. Ann. Probab. 24, no. 1 (1996), 280–292.
- [29] Reed, M. & Simon, B.: *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press, Inc., 1978.
- [30] Revuz, D. & Yor, M.: *Continuous martingales and Brownian motion*. Grundlehren der Mathematischen Wissenschaften, 293. Springer-Verlag, Berlin, 1991.
- [31] Simon, B.: *Schrödinger semigroups*. Bull. Amer. Math. Soc. (N.S.) Volume 7, Number 3 (1982), 447–526.

- [32] Simon, B.: *Functional integration and quantum physics*. Academic Press. Inc., 1979.
- [33] Stroock, D.W.: *An introduction to the analysis of paths on a Riemannian manifold*. Mathematical Surveys and Monographs, 74. American Mathematical Society, Providence, RI, 2000.
- [34] Sznitman, A.S.: *Brownian motion, obstacles and random media*. Springer, Berlin, 1998.
- [35] Teschl, G.: *Mathematical methods in quantum mechanics. With applications to Schrödinger operators*. Graduate Studies in Mathematics, 99. American Mathematical Society, Providence, RI, 2009.
- [36] Weidmann, J.: *Lineare Operatoren in Hilberträumen. Teil 1*. B. G. Teubner, Stuttgart, 2000.