

Loose Laplacian spectra of random hypergraphs

Linyuan Lu ^{*} Xing Peng [†]

June 14, 2022

Abstract

Let $H = (V, E)$ be an r -uniform hypergraph with the vertex set V and the edge set E . For $1 \leq s \leq r/2$, we define a weighted graph $G^{(s)}$ on the vertex set $\binom{V}{s}$ as follows. Every pair of s -sets I and J is associated with a weight $w(I, J)$, which is the number of edges in H passing through I and J if $I \cap J = \emptyset$, and 0 if $I \cap J \neq \emptyset$. The s -th Laplacian $\mathcal{L}^{(s)}$ of H is defined to be the normalized Laplacian of $G^{(s)}$. The eigenvalues of $\mathcal{L}^{(s)}$ are listed as $\lambda_0^{(s)}, \lambda_1^{(s)}, \dots, \lambda_{\binom{n}{s}-1}^{(s)}$ in non-decreasing order. Let $\bar{\lambda}^{(s)}(H) = \max_{i \neq 0} \{1 - \lambda_i^{(s)}\}$. The parameters $\bar{\lambda}^{(s)}(H)$ and $\lambda_1^{(s)}(H)$, which were introduced in our previous paper, have a number of connections to the mixing rate of high-ordered random walks, the generalized distances/diameters, and the edge expansions.

For $0 < p < 1$, let $H^r(n, p)$ be a random r -uniform hypergraph over $[n] := \{1, 2, \dots, n\}$, where each r -set of $[n]$ has probability p to be an edge independently. For $1 \leq s \leq r/2$, $p(1-p) \gg \frac{\log^4 n}{n^{r-s}}$, and $1-p \gg \frac{\log n}{n^2}$, we prove that almost surely

$$\bar{\lambda}^{(s)}(H^r(n, p)) \leq \frac{s}{n-s} + (3 + o(1)) \sqrt{\frac{1-p}{\binom{n-s}{r-s}p}}.$$

We also prove that the empirical distribution of the eigenvalues of $\mathcal{L}^{(s)}$ for $H^r(n, p)$ follows the Semicircle Law if $p(1-p) \gg \frac{\log^{1/3} n}{n^{r-s}}$ and $1-p \gg \frac{\log n}{n^{2+2r-2s}}$.

1 Introduction

The spectrum of the adjacency matrix (and/or the Laplacian matrix) of a random graph was well-studied in the literature [1, 10, 11, 13, 14, 15, 17, 18, 21]. Given a graph G , let $\mu_1(G), \dots, \mu_n(G)$ be the eigenvalues of the adjacency matrix of G in the non-decreasing order, and $\lambda_0(G), \dots, \lambda_{n-1}(G)$ be the eigenvalues of (normalized) Laplacian matrix of G respectively. Let $G(n, p)$ be the Edős-Rényi random graph model. Füredi and Komlós [21] showed that if $np(1-p) \gg \log^6 n$ then almost surely $\mu_n = (1+o(1))np$ and $\max\{-\mu_1, \mu_{n-1}\} \leq (2+o(1))\sqrt{np(1-p)}$. The results are extended to sparse random graphs [17, 25] and general random matrices [15, 21]. Alon, Krivelevich, and Vu [1] proved the concentration of the s -th largest eigenvalue of a random symmetric matrix with independent random entries of absolute value at most 1. Friedman (in a series of papers [18, 19, 20]) proved that the second largest eigenvalue of random d -regular graphs is almost surely $(2+o(1))\sqrt{d-1}$ for any $d \geq 4$. Chung, Lu, and Vu [11] studied the Laplacian eigenvalues of random graphs with

^{*}University of South Carolina, Columbia, SC 29208, (lu@math.sc.edu). This author was supported in part by NSF grant DMS 1000475.

[†]University of South Carolina, Columbia, SC 29208, (pengx@mailbox.sc.edu). This author was supported in part by NSF grant DMS 1000475.

given expected degrees; their results were supplemented by Coja-Oghlan [13, 14] for much sparser random graphs.

In this paper, we study the spectra of the Laplacians of random hypergraphs. Laplacians for regular hypergraphs were first introduced by Chung [5] using the homology approach. Rodríguez [28, 29] treated a hypergraph as a multi-edge graph and then defined its Laplacian to be the Laplacian of the corresponding multi-edge graph. Inspired by these work, we [26] introduced the generalized Laplacian eigenvalues of hypergraphs through high-ordered random walks. Let $H = (V, E)$ be an r -uniform hypergraph on n vertices. We can associate $r - 1$ Laplacians $\mathcal{L}^{(s)}$ ($1 \leq s \leq r - 1$) to H ; roughly speaking, $\mathcal{L}^{(s)}$ captures the incidence relations between s -sets and edges in H . Our definition of the Laplacian at the special case $s = 1$ is the same as the Laplacian considered by Rodríguez [28, 29]. The s -th Laplacian is *loose* if $1 \leq s \leq r/2$, and is *tight* if $r/2 < s \leq r - 1$. Here we consider only the spectra of loose Laplacians.

For $1 \leq s \leq r/2$, we consider an auxiliary weighted graph $G^{(s)}$ defined as follows: the vertex set of $G^{(s)}$ is $\binom{V}{s}$ while the weight function $W: \binom{V}{s} \times \binom{V}{s} \rightarrow \mathbb{Z}$ is defined as

$$W(S, T) = \begin{cases} |\{F \in E(H): S \cup T \subset F\}| & \text{if } S \cap T = \emptyset; \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

The s -th Laplacian of H , denoted by $\mathcal{L}^{(s)}$, is the normalized Laplacian of $G^{(s)}$. For any s -set S , let d_S be the number of edges in H passing through S ; the degree of S in $G^{(s)}$ is $\binom{r-s}{s} d_S$. Let D be the diagonal matrix of the degrees $\{d_S\}$ and W be the weight matrix $\{w(S, T)\}$. Note that $T := \binom{r-s}{s} D$ is the diagonal matrix of degrees in $G^{(s)}$. We have

$$\mathcal{L}^{(s)} = I - T^{-1/2} W T^{-1/2}. \quad (2)$$

The eigenvalues of $\mathcal{L}^{(s)}$ are listed as $\lambda_0^{(s)}, \lambda_1^{(s)}, \dots, \lambda_{\binom{n}{s}-1}^{(s)}$ in non-decreasing order. We have

$$0 = \lambda_0^{(s)} \leq \lambda_1^{(s)} \leq \dots \leq \lambda_{\binom{n}{s}-1}^{(s)} \leq 2. \quad (3)$$

The first non-trivial eigenvalue $\lambda_1^{(s)} > 0$ if and only if $G^{(s)}$ is connected. When this occurs, we say H is *s-connected*. The diameter of $G^{(s)}$ is called the *s-th diameter* of H . The largest eigenvalue $\lambda_{\binom{n}{s}-1}^{(s)}$ is also denoted by $\lambda_{max}^{(s)}$. The (Laplacian) spectral radius, denoted by $\bar{\lambda}^{(s)}$, is the maximum of $1 - \lambda_1^{(s)}$ and $\lambda_{max}^{(s)} - 1$.

This definition differs slightly with the one in [26], where the vertex set of the auxiliary graph (denoted by $G^{(s)'}$) is the set of all distinct s -tuples instead. Note that $G^{(s)'}$ is the blow-up of $G^{(s)}$. Their Laplacian spectra differ only by the multiplicity of 1's. Therefore, two different definitions give the same values of $\lambda_1^{(s)}$, $\lambda_{max}^{(s)}$, and $\bar{\lambda}^{(s)}$.

For different s , the following inequalities were proved in [26].

$$\lambda_1^{(1)} \geq \lambda_1^{(2)} \geq \dots \geq \lambda_1^{(\lfloor r/2 \rfloor)}; \quad (4)$$

$$\lambda_{max}^{(1)} \leq \lambda_{max}^{(2)} \leq \dots \leq \lambda_{max}^{(\lfloor r/2 \rfloor)}. \quad (5)$$

The s -th Laplacian has a number of connections to the mixing rate of high-ordered random walks, the generalized distances/diameters, and the edge expansions. Here we list some applications, which are similar to results in [26], and results for graphs [4, 6, 7, 8, 9, 12].

Random s -Walks: The mixing rate of the random s -walk on H is at most $\bar{\lambda}^{(s)}$.

The s -Diameter: The s -diameter of H is at most

$$\left\lceil \frac{\log \frac{|E(H)| \binom{r}{s}}{\delta}}{\log \frac{\lambda_{\max}^{(s)} + \lambda_1^{(s)}}{\lambda_{\max}^{(s)} - \lambda_1^{(s)}}} \right\rceil.$$

Here $\delta = \min_{S \in \binom{V}{s}} d_S$ is the minimum degree among all s -sets.

Edge expansion: For $1 \leq t \leq s \leq \frac{r}{2}$, $\mathcal{S} \subset \binom{V}{t}$, and $\mathcal{T} \subset \binom{V}{t}$, define

$$E(\mathcal{S}, \mathcal{T}) = \{F \in E(H) : \exists S \in \mathcal{S}, \exists T \in \mathcal{T} \text{ such that } S \cap T = \emptyset, \text{ and } S \cup T \subset F\},$$

$$\begin{aligned} e(\mathcal{S}, \mathcal{T}) &= \frac{|E(\mathcal{S}, \mathcal{T})|}{\left|E\left(\binom{V}{s}, \binom{V}{t}\right)\right|}, \\ e(\mathcal{S}) &= \frac{\sum_{S \in \mathcal{S}} d_S}{\sum_{S \in \binom{V}{s}} d_S}, \\ e(\mathcal{T}) &= \frac{\sum_{T \in \mathcal{T}} d_T}{\sum_{T \in \binom{V}{t}} d_T}. \end{aligned}$$

Then we have

$$|e(\mathcal{S}, \mathcal{T}) - e(\mathcal{S})e(\mathcal{T})| \leq \bar{\lambda}^{(s)} \sqrt{e(\mathcal{S})e(\mathcal{T})e(\bar{\mathcal{S}})e(\bar{\mathcal{T}})}.$$

The proofs of these claims are very similar to those in [26] and are omitted here.

Our first result is the eigenvalues of the s -th Laplacian of the complete r -uniform hypergraph K_n^r .

Theorem 1 *Let K_n^r be the complete r -uniform hypergraph on n vertices. For $1 \leq s \leq r/2$, the eigenvalues of s -th Laplacian of K_n^r are given by*

$$1 - \frac{(-1)^i \binom{n-s-i}{s-i}}{\binom{n-s}{s}} \text{ with multiplicity } \binom{n}{i} - \binom{n}{i-1} \text{ for } 0 \leq i \leq s.$$

Here we point out an application of this theorem to the celebrated Erdős-Ko-Rado Theorem, which states ‘‘if the $n \geq 2s$, then the size of the maximum intersecting family of s -sets in $[n]$ is at most $\binom{n-1}{s-1}$.’’ (The theorem was originally proved by Erdős-Ko-Rado [16] for sufficiently large n ; the simplest proof was due to Katona [24].) Here we present a proof adapted from Calderbank-Frankl [2], where they use the eigenvalues of Kneser graph instead. (The relation between $\mathcal{L}^{(s)}(K_n^r)$ and the Laplacian of the Kneser graph is explained in section 2.)

It suffices to show for any intersecting family U of s -sets, $|U| \leq \binom{n-1}{s-1}$. Note that U is an independent set of $G^{(s)}(K_n^r)$. Restricting to U , $\mathcal{L}^{(s)}(K_n^r)$ becomes an identity matrix; whose eigenvalues are all equal to 1. By Cauchy’s interlace theorem, we have

$$\lambda_k^{(s)} \leq 1 \leq \lambda_{\binom{n}{s}-|U|+k}^{(s)} \tag{6}$$

for $0 \leq k \leq |U| - 1$. Let N^+ (or N^-) be the number of eigenvalues of $\mathcal{L}^{(s)}(K_n^r)$ which is ≥ 1 (or ≤ 1) respectively. Inequality (6) implies that $|U| \leq N^+$ and $|U| \leq N^-$. By Theorem 1, $N^+ = \sum_{i=0}^{\lfloor (s-1)/2 \rfloor} \left(\binom{n}{2i+1} - \binom{n}{2i} \right)$ and $N^- = \sum_{i=0}^{\lfloor s/2 \rfloor} \left(\binom{n}{2i} - \binom{n}{2i-1} \right)$. We have

$$|U| \leq \min\{N^+, N^-\} = \sum_{i=0}^{s-1} (-1)^{s-1-i} \binom{n}{i} = \binom{n-1}{s-1}.$$

For $0 < p < 1$, let $H^r(n, p)$ be a random r -uniform hypergraph over $[n] = \{1, 2, \dots, n\}$, where each r -set of $[n]$ has probability p to be an edge independently. We can estimate the Laplacian spectrum of $H^r(n, p)$ using the Laplacian spectrum of K_n^r as follows.

Theorem 2 *Let $H^r(n, p)$ be a random r -uniform hypergraph. For $1 \leq s \leq r/2$, if $p(1-p) \gg \frac{\log^4 n}{n^{r-s}}$ and $1-p \gg \frac{\log n}{n^2}$, then almost surely the s -th spectral radius $\bar{\lambda}^{(s)}(H^r(n, p))$ satisfies*

$$\bar{\lambda}^{(s)}(H^r(n, p)) \leq \frac{s}{n-s} + (3 + o(1)) \sqrt{\frac{1-p}{\binom{n-s}{r-s}p}}. \quad (7)$$

Moreover, for $1 \leq k \leq \binom{n}{s} - 1$, almost surely we have

$$|\lambda_k^{(s)}(H^r(n, p)) - \lambda_k^{(s)}(K_n^r)| \leq (3 + o(1)) \sqrt{\frac{1-p}{\binom{n-s}{r-s}p}}. \quad (8)$$

Note that $G(n, p)$ is a special case of $H^r(n, p)$ with $r = 2$. By choosing $s = 1$, Theorem 2 implies that

$$\bar{\lambda}(G(n, p)) \leq (3 + o(1)) \sqrt{\frac{1-p}{(n-1)p}} \quad \text{for } p(1-p) \gg \frac{\log^4 n}{n}. \quad (9)$$

Chung, Lu, and Vu's result[11], when restricted to $G(n, p)$, implies

$$\bar{\lambda}(G(n, p)) \leq (4 + o(1)) \frac{1}{\sqrt{np}} \quad \text{for } 1 - \epsilon \geq p \gg \frac{\log^6 n}{n}. \quad (10)$$

Inequality (9) has a smaller constant and works for a larger range of p than inequality (10).

Füredi and Komlós [21] proved the empirical distribution of the eigenvalues of $G(n, p)$ follows the Semicircle Law. Chung, Lu, and Vu [11] proved a similar result for the random graphs with given expected degrees. Here we prove a similar result for random hypergraphs.

Theorem 3 *For $1 \leq s \leq r/2$, if $p(1-p) \gg \frac{\log^{1/3} n}{n^{r-s}}$ and $1-p \gg \frac{\log n}{n^{2+2r-2s}}$, then almost surely the empirical distribution of eigenvalues of the s -th Laplacian of $H^r(n, p)$ follows the Semicircle Law centered at 1 and with radius $(2 + o(1)) \sqrt{\frac{1-p}{\binom{r-s}{s} \binom{n-s}{r-s} p}}$.*

Remark 1 *The proof of Theorem 3 actually implies the eigenvalues of $\mathcal{L}^{(s)}(H^r(n, p)) - \mathcal{L}^{(s)}(K_n^r)$ follows the Semicircle Law centered at 0 and with radius $(2 + o(1)) \sqrt{\frac{1-p}{\binom{r-s}{s} \binom{n-s}{r-s} p}}$. Thus we have*

$$\max_{1 \leq k \leq \binom{n}{s} - 1} |\lambda_k^{(s)}(H^r(n, p)) - \lambda_k^{(s)}(K_n^r)| \geq \left(\frac{2}{\sqrt{\binom{r-s}{s}}} + o(1) \right) \sqrt{\frac{1-p}{\binom{n-s}{r-s} p}}. \quad (11)$$

This shows that the upper bound of $|\lambda_k^{(s)}(H^r(n, p)) - \lambda_k^{(s)}(K_n^r)|$ in Theorem 2 is best up to a constant multiplicative factor.

The rest of the paper is organized as follows. In section 2, we introduce the notation and prove some basic lemmas. We will prove Theorem 1 in section 3 and Theorem 2 in section 4.

2 Notation and Lemmas

2.1 Laplacian eigenvalues of hypergraphs

Let $H = (V, E)$ be an r -uniform hypergraph. For any subset S ($|S| < r$), the degree of S , denoted by d_S , is the number of edges passing through S . For each $1 \leq s \leq r/2$, we associate a weighted graph $G^{(s)}$ on the vertex set $\binom{V}{s}$ to H as follows. Every pair of s -sets S and T is associated with a weight $w(S, T)$, which is given by

$$w(S, T) = \begin{cases} d_{S \cup T} & \text{if } S \cap T = \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

The s -th Laplacian $\mathcal{L}^{(s)}$ of H is defined to be the normalized Laplacian of $G^{(s)}$. The degree of S in $G^{(s)}$ is $\sum_T w(S, T) = \binom{r-s}{s} d_S$.

We assume that the s -sets in $\binom{V}{s}$ are ordered alphabetically. Let $N := \binom{n}{s}$; all square matrices considered in the paper have the dimension $N \times N$ and all vectors have dimension N . Let $W := (W(S, T))$ be the weight matrix, D be the diagonal matrix with diagonal entries $D(S, S) = d_S$, \mathbf{d} be the column vector with entries d_S at position $S \in \binom{V}{s}$, J be the square matrix of all 1's, and $\mathbf{1}$ be the column vector of all 1's. Let $T := \binom{r-s}{s} D$; here T is the diagonal matrix of degrees in $G^{(s)}$. Then, we have

$$\mathcal{L}^{(s)} = I - T^{-1/2} W T^{-1/2}.$$

We list the eigenvalues of $\mathcal{L}^{(s)}$ as

$$0 = \lambda_0^{(s)} \leq \lambda_1^{(s)}, \dots, \lambda_{\binom{n}{s}-1}^{(s)} \leq 2.$$

We aim to compute the spectral radius $\bar{\lambda}^{(s)}(H) = \max_{i \neq 0} |1 - \lambda_i^{(s)}|$. Let $\text{vol}^{(s)}(H) := \sum_{S \in \binom{V}{s}} d_S$ and $\phi_0 := \frac{1}{\sqrt{\text{vol}^{(s)}(H)}} D^{1/2} \mathbf{1}$. Note that ϕ_0 is the unit eigenvector corresponding to the trivial eigenvalue 0 of $\mathcal{L}^{(s)}$.

We are ready to prove theorem 1.

Proof of Theorem 1: We can express $\mathcal{L}^{(s)}(K_n^r)$ using the following notation. The Kneser graph $K(n, s)$ is a graph over the vertex set $\binom{[n]}{s}$; two s -sets S and T form an edge of $K(n, s)$ if and only if $S \cap T = \emptyset$. Let K be the adjacency matrix of $K(n, s)$; the eigenvalues of K are $(-1)^i \binom{n-s-i}{s-i}$ with multiplicity $\binom{n}{i} - \binom{n}{i-1}$ for $0 \leq i \leq s$ (see [22]). Note that $K(n, s)$ is a regular graph; so the Laplacian eigenvalues can be determined from the eigenvalues of its adjacency matrix. We observe that the associated weighted graph $G^{(s)}$ for the complete r -uniform hypergraph K_n^r is essentially the Kneser graph with each edge associated with a weight $\binom{n-2s}{r-2s}$. Note that the multiplicative factor $\binom{n-2s}{r-2s}$ is canceled after normalization. The $\mathcal{L}^{(s)}$ (for K_n^r) is exactly the Laplacian of Kneser graph. Hence,

$$\mathcal{L}^{(s)}(K_n^r) = I - \frac{1}{\binom{n-s}{s}} K.$$

Thus, the eigenvalues of s -th Laplacian of K_n^r are given by

$$1 - \frac{(-1)^i \binom{n-s-i}{s-i}}{\binom{n-s}{s}} \text{ with multiplicity } \binom{n}{i} - \binom{n}{i-1} \text{ for } 0 \leq i \leq s.$$

□

Remark 2 For $1 \leq s \leq r/2$, we have

$$\lambda_1^{(s)}(K_n^r) = 1 - \frac{s(s-1)}{(n-s)(n-s-1)}, \quad (12)$$

$$\lambda_{max}^{(s)}(K_n^r) = 1 + \frac{s}{n-s}, \quad (13)$$

$$\bar{\lambda}^{(s)}(K_n^r) = \frac{s}{n-s}. \quad (14)$$

2.2 Random hypergraphs

Let $H^r(n, p)$ be a random r -uniform hypergraph over the vertex set $V = [n]$ and each r -set has probability p to be an edge independently. We would like to bound the spectral radius of the s -th Laplacian of $H^r(n, p)$ for $1 \leq s \leq r/2$.

For any $F \in \binom{V}{r}$, let X_F be the random indicator variable for F being an edge in $H^r(n, p)$; all X_F 's are independent to each other. For any $S, T \in \binom{V}{s}$, we have

$$W(S, T) = \begin{cases} \sum_{\substack{F \in \binom{V}{r} \\ S \cup T \subset F}} X_F & \text{if } S \cap T = \emptyset; \\ 0 & \text{otherwise.} \end{cases}$$

Thus,

$$\mathbb{E}(W(S, T)) = \begin{cases} \binom{n-2s}{r-2s} p & \text{if } S \cap T = \emptyset; \\ 0 & \text{otherwise.} \end{cases} \quad (15)$$

The degree $d_S = \sum_{S \subset F \in \binom{V}{r}} X_F$; we have $\mathbb{E}(d_S) = \binom{n-s}{r-s} p$. For simplicity, let $d := \binom{n-s}{r-s} p$.

We use the following Lemma to compare the eigenvalues of two matrices.

Lemma 1 Given any two $(N \times N)$ -Hermitian matrices A and B , for $1 \leq k \leq N$, let $\mu_k(A)$ (or $\mu_k(B)$) be the k -th eigenvalues of A (or B) in the increasing order. We have

$$|\mu_k(A) - \mu_k(B)| \leq \|A - B\|.$$

Proof: By the Min-Max Theorem (see [27]), we have

$$\begin{aligned} \mu_k(A) &= \min_{S_k} \max_{x \in S_k, \|x\|=1} x' A x, \\ \mu_k(B) &= \min_{S_k} \max_{x \in S_k, \|x\|=1} x' B x. \end{aligned}$$

where the minimum is taken over all k -th dimensional subspace $S_k \subset \mathbb{R}^N$. We have

$$\begin{aligned} \mu_k(A) &= \min_{S_k} \max_{x \in S_k, \|x\|=1} x' A x \\ &= \min_{S_k} \max_{x \in S_k, \|x\|=1} (x' B x + x' (A - B) x) \\ &\leq \min_{S_k} \max_{x \in S_k, \|x\|=1} (x' B x + \|A - B\|) \\ &= \mu_k(B) + \|A - B\|. \end{aligned}$$

Similarly, we can show $\mu_k(A) \geq \mu_k(B) - \|A - B\|$. The proof of the Lemma is finished. \square

Our idea is to bound the spectral norm of the difference of $\mathcal{L}^{(s)}(H^r(n, p))$ and $\mathcal{L}^{(s)}(K_n^r)$. Let $M := \mathcal{L}^{(s)}(K_n^r) - \mathcal{L}^{(s)}(H^r(n, p)) = T^{-1/2} W T^{-1/2} - \frac{1}{\binom{n-s}{s}} K$. We write $M = M_1 + M_2 +$

$M_3 + M_4$, where

$$\begin{aligned} M_1 &= \frac{1}{\binom{r-s}{s}} \left(D^{-1/2}(W - \mathbf{E}(W))D^{-1/2} - d^{-1}(W - \mathbf{E}(W)) \right), \\ M_2 &= \frac{1}{\binom{r-s}{s}d} (W - \mathbf{E}(W)), \\ M_3 &= \frac{1}{\binom{r-s}{s}} D^{-1/2} \mathbf{E}(W) D^{-1/2} - \frac{d}{\binom{n}{s}} D^{-1/2} J D^{-1/2} - \frac{1}{\binom{n-s}{s}} K + \frac{1}{\binom{n}{s}} J, \\ M_4 &= \frac{1}{\binom{n}{s}} (dD^{-1/2} J D^{-1/2} - J). \end{aligned}$$

By the triangular inequality of matrix norms, we have

$$\|M\| \leq \|M_1\| + \|M_2\| + \|M_3\| + \|M_4\|.$$

Through this paper, the norm of any square matrix is the spectral norm. We would like to bound $\|M_i\|$ for $i = 1, 2, 3, 4$. We use the following Chernoff inequality.

Theorem 4 [3] *Let X_1, \dots, X_n be independent random variables with*

$$\Pr(X_i = 1) = p, \quad \Pr(X_i = 0) = 1 - p.$$

We consider the sum $X = \sum_{i=1}^n X_i$, with expectation $\mathbf{E}(X) = np$. Then we have

$$\begin{aligned} \text{(Lower tail)} \quad & \Pr(X \leq \mathbf{E}(X) - \lambda) \leq e^{-\lambda^2/2\mathbf{E}(X)}, \\ \text{(Upper tail)} \quad & \Pr(X \geq \mathbf{E}(X) + \lambda) \leq e^{-\frac{\lambda^2}{2(\mathbf{E}(X) + \lambda/3)}}. \end{aligned}$$

Lemma 2 *Suppose $d \geq \log N$. With probability at least $1 - \frac{1}{N^2}$, we have $d_S \in (d - 3\sqrt{d \log N}, d + 3\sqrt{d \log N})$ for all $S \in \binom{V}{s}$.*

Proof: Note $d_s = \sum_{F: S \subset F} X_F$ and $\mathbf{E}(d_s) = d$. Applying the lower tail of Chernoff's inequality with $\lambda = 3\sqrt{\mathbf{E}(X) \log N}$, we have

$$\Pr(X - \mathbf{E}(X) \leq -\lambda) \leq e^{-\lambda^2/2\mathbf{E}(X)} \leq \frac{1}{N^{9/2}}.$$

Applying the upper tail of Chernoff's inequality with $\lambda = 3\sqrt{\mathbf{E}(X) \log N}$, we have

$$\Pr(X - \mathbf{E}(X) \geq \lambda) \leq e^{-\frac{\lambda^2}{2(\mathbf{E}(X) + \lambda/3)}} \leq \frac{1}{N^{27/8}}.$$

The probability that $d_S \notin (d - 3\sqrt{d \log N}, d + 3\sqrt{d \log N})$ is at most $\frac{1}{N^3}$. Thus, with probability at least $1 - \frac{1}{N^2}$, we have $d_S \in (d - 3\sqrt{d \log N}, d + 3\sqrt{d \log N})$ for all $S \in \binom{V}{s}$. \square

For convenience, let $d_{\min} := d - 3\sqrt{d \log N}$, $d_{\max} := d + 3\sqrt{d \log N}$; almost surely we have $d_{\min} \leq d_S \leq d_{\max}$ for all S .

Lemma 3 *If $d \geq \log N$, then almost surely $\|M_3\| = O\left(\frac{\sqrt{\log N}}{n\sqrt{d}}\right)$.*

Proof: Note $\mathbf{E}(W) = \binom{n-2s}{r-2s} pK$, where K is the adjacency matrix of the Kneser graph $K(n, s)$. Let $M_0 := \frac{1}{\binom{n-s}{s}} K - \frac{1}{\binom{n}{s}} J$. We can rewrite M_3 as

$$M_3 = dD^{-1/2} M_0 D^{-1/2} - M_0.$$

Note $\|M_0\| = \bar{\lambda}^{(s)}(K_n^r) = \frac{s}{n-s}$. We have

$$\begin{aligned}
\|M_3\| &= \|dD^{-1/2}M_0D^{-1/2} - M_0\| \\
&\leq \|(dD^{-1/2} - d^{1/2}I)M_0D^{-1/2}\| + \|M_0(d^{1/2}D^{-1/2} - I)\| \\
&\leq \|(d^{1/2}I - dD^{-1/2})\| \|M_0\| \|D^{-1/2}\| + \|M_0\| \|(d^{1/2}D^{-1/2} - I)\| \\
&\leq \left| d^{1/2} - dd_{\min}^{-1/2} \right| \frac{s}{n-s} d_{\min}^{-1/2} + \frac{s}{n-s} \left| d^{1/2} d_{\min}^{-1/2} - 1 \right| \\
&= O\left(\frac{\sqrt{\log N}}{n\sqrt{d}}\right).
\end{aligned}$$

□

Lemma 4 *If $p(1-p) \gg \frac{\log n}{nr-s}$, then almost surely*

$$\sum_{S \in \binom{V}{s}} (d_S - d)^2 = (1 + o(1)) \binom{n}{s} d(1-p).$$

Proof: For $S \in \binom{V}{s}$, let $X_S = (d_S - d)^2$. We have

$$\mathbb{E}(X_S) = \mathbb{E}((d_S - d)^2) = \text{Var}(d_S) = \binom{n-s}{r-s} p(1-p) = d(1-p).$$

We use the second moment method to prove that $\sum_S X_S$ concentrates around its expectation $\binom{n}{s} d(1-p)$. For any $S, T \in \binom{V}{s}$, the covariance can be calculated as follows.

$$\begin{aligned}
\text{Cov}(X_S, X_T) &= \mathbb{E}(X_S X_T) - \mathbb{E}(X_S)\mathbb{E}(X_T) \\
&= \mathbb{E}((d_S - d)^2 (d_T - d)^2) - d^2(1-p)^2.
\end{aligned}$$

For $F \in \binom{V}{r}$, let $Y_F = X_F - \mathbb{E}(X_F)$. Then we have $d_S - d = \sum_{S \subset F} Y_F$.

$$\mathbb{E}((d_S - d)^2 (d_T - d)^2) = \sum_{\substack{F_1, F_2: S \subset F_1 \cap F_2 \\ F_3, F_4: T \subset F_3 \cap F_4}} \mathbb{E}(Y_{F_1} Y_{F_2} Y_{F_3} Y_{F_4}).$$

Since $\mathbb{E}(Y_{F_i}) = 0$, the non-zero terms occur only if

1. $F_1 = F_2 = F_3 = F_4$. In this case, we have

$$\mathbb{E}(Y_{F_1} Y_{F_2} Y_{F_3} Y_{F_4}) = \mathbb{E}(Y_{F_1}^4) = (1-p)^4 p + (-p)^4 (1-p) = p(1-p)(1-3p+3p^2).$$

The number of choices is $\binom{n-|S \cup T|}{r-|S \cup T|}$.

2. $F_1 = F_2 \neq F_3 = F_4$. In this case, we have

$$\mathbb{E}(Y_{F_1} Y_{F_2} Y_{F_3} Y_{F_4}) = \mathbb{E}(Y_{F_1}^2) \mathbb{E}(Y_{F_3}^2) = p^2(1-p)^2.$$

The number of choices is $\binom{n-s}{r-|S|} \binom{n-s}{r-|T|} - \binom{n-|S \cup T|}{r-|S \cup T|}$.

3. $F_1 = F_3 \neq F_2 = F_4$. In this case, we have

$$\mathbb{E}(Y_{F_1} Y_{F_2} Y_{F_3} Y_{F_4}) = \mathbb{E}(Y_{F_1}^2) \mathbb{E}(Y_{F_2}^2) = p^2(1-p)^2.$$

The number of choices is $\binom{n-|S \cup T|}{r-|S \cup T|}^2 - \binom{n-|S \cup T|}{r-|S \cup T|}$.

4. $F_1 = F_4 \neq F_2 = F_3$. This is the same as item 3.

Thus, we have

$$\begin{aligned}
\mathbb{E}(X_S X_T) &= \binom{n - |S \cup T|}{r - |S \cup T|} p(1-p)(1 - 3p + 3p^2) \\
&\quad + \left(\binom{n-s}{r-s}^2 + 2 \binom{n - |S \cup T|}{r - |S \cup T|}^2 - 3 \binom{n - |S \cup T|}{r - |S \cup T|} \right) p^2(1-p)^2 \\
&= \binom{n - |S \cup T|}{r - |S \cup T|} p(1-p)(1 - 6p + 6p^2) + \left(\binom{n-s}{r-s}^2 + 2 \binom{n - |S \cup T|}{r - |S \cup T|}^2 \right) p^2(1-p)^2.
\end{aligned}$$

This expression on the right depends only on the size of $S \cup T$. Putting together, we get

$$\begin{aligned}
\text{Var} \left(\sum_{S \in \binom{V}{s}} X_S \right) &= \sum_{S, T \in \binom{V}{s}} \text{Cov}(X_S, X_T) \\
&= \sum_{S, T \in \binom{V}{s}} (\mathbb{E}(X_S X_T) - d^2(1-p)^2) \\
&= \sum_{S, T \in \binom{V}{s}} \left(\mathbb{E}(X_S X_T) - \binom{n-s}{r-s}^2 p^2(1-p)^2 \right) \\
&= \sum_{i=s}^{2s} \sum_{|S \cup T|=i} \left(\binom{n-i}{r-i} p(1-p)(1 - 6p + 6p^2) + 2 \binom{n-i}{r-i}^2 p^2(1-p)^2 \right) \\
&\leq \sum_{i=s}^{2s} \sum_{|S \cup T|=i} \binom{n-i}{r-i} p(1-p) \left(1 - 6p + 6p^2 + 2 \binom{n-s}{r-s} p(1-p) \right) \\
&\leq \sum_{i=s}^{2s} \sum_{|S \cup T|=i} \binom{n-i}{r-i} 3dp(1-p)^2 \\
&= \binom{n}{r} 3dp(1-p)^2 \sum_{i=s}^{2s} \frac{r!}{(i-s)!2(2s-i)!(r-i)!} \\
&< 3 \cdot 4^r \binom{n}{r} dp(1-p)^2 \\
&= O \left(\binom{n}{s} d^2(1-p)^2 \right).
\end{aligned}$$

Let $X = \sum_S X_S$. We have $\mathbb{E}[X] = \binom{n}{s} d(1-p)$ and $\text{Var}(X) = O \left(\binom{n}{s} d^2(1-p)^2 \right)$. Applying Chebyshev's inequality to $X = \sum_{S \in \binom{V}{s}} X_S$, we have

$$\Pr \left(|X - \mathbb{E}(X)| \geq \log n \sqrt{\text{Var}(X)} \right) \leq \frac{1}{\log^2 n}.$$

Thus, almost surely $X = \mathbb{E}(X) + O(\log n \sqrt{\text{Var}(X)}) = (1 + o(1)) \binom{n}{s} d(1-p)$. \square

Lemma 5 *If $p(1-p) \gg \frac{\log n}{n^{r-s}}$, then almost surely $\|M_4\| \leq (1 + o(1)) \sqrt{\frac{1-p}{d}}$.*

Proof: We can rewrite M_4 as

$$\begin{aligned}
M_4 &= \frac{1}{\binom{n}{s}}(dD^{-1/2}JD^{-1/2} - J) \\
&= \frac{1}{\binom{n}{s}} \left((d^{1/2}D^{-1/2} - I) JD^{-1/2}d^{1/2} + J (d^{1/2}D^{-1/2} - I) \right) \\
&= \frac{1}{\binom{n}{s}} \left(\alpha \mathbf{1}' D^{-1/2} d^{1/2} + \mathbf{1} \alpha' \right).
\end{aligned}$$

Here $\alpha := d^{1/2}D^{-1/2}\mathbf{1} - \mathbf{1}$. Note that the spectral norm of a vector is the same as the L_2 -norm. We have

$$\begin{aligned}
\|\alpha\| &= \|d^{1/2}D^{-1/2}\mathbf{1} - \mathbf{1}\| \\
&= \sqrt{\sum_{S \in \binom{[n]}{s}} \left(\frac{\sqrt{d}}{\sqrt{d_S}} - 1 \right)^2} \\
&= \sqrt{\sum_{S \in \binom{[n]}{s}} \frac{(d_S - d)^2}{d_S(\sqrt{d} + \sqrt{d_S})^2}} \\
&\leq \frac{\sqrt{\sum_{S \in \binom{[n]}{s}} (d_S - d)^2}}{\sqrt{d_{\min}}(\sqrt{d} + \sqrt{d_{\min}})} \\
&= \left(\frac{1}{2} + o(1) \right) \sqrt{\frac{(1-p)\binom{n}{s}}{d}}.
\end{aligned}$$

In the last step, we applied Lemma 4. Therefore, we have

$$\begin{aligned}
\|M_4\| &= \left\| \frac{1}{\binom{n}{s}} \left(\alpha \mathbf{1}' D^{-1/2} d^{1/2} + \mathbf{1} \alpha' \right) \right\| \\
&= \frac{1}{\binom{n}{s}} \left(\left\| \alpha \mathbf{1}' D^{-1/2} d^{1/2} \right\| + \|\mathbf{1} \alpha'\| \right) \\
&\leq \frac{1}{\binom{n}{s}} \|\alpha\| \left(\|\mathbf{1}' D^{-1/2} d^{1/2}\| + \|\mathbf{1}\| \right) \\
&= \frac{1}{\binom{n}{s}} \|\alpha\| \left(\sqrt{\sum_{S \in \binom{[n]}{s}} \frac{d}{d_S}} + \sqrt{\binom{n}{s}} \right) \\
&\leq \frac{1}{\binom{n}{s}} \left(\frac{1}{2} + o(1) \right) \sqrt{\frac{(1-p)\binom{n}{s}}{d}} (2 + o(1)) \left(\sqrt{\binom{n}{s}} \right) \\
&= (1 + o(1)) \sqrt{\frac{1-p}{d}}.
\end{aligned}$$

3 Proof of Theorem 2

To estimate the spectral norm of M_1 and M_2 , we need consider the matrix $C := W - \mathbb{E}(W)$. We estimate the expectation of the trace of C^t as follows.

Lemma 6 For any k satisfying $k \ll \sqrt[4]{n^{r-s}p(1-p)}$, we have

$$\mathbb{E}(\text{Trace}(C^{2k})) \leq (1 + o(1)) \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k, \quad (16)$$

$$\mathbb{E}(\text{Trace}(C^{2k+1})) = O\left(\frac{2k^2 n^{s+k(r-s)} \binom{r-s}{s}^{k+1}}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k\right). \quad (17)$$

If further $k = o(\log(n^{r-s}p(1-p)))$, then we have

$$\mathbb{E}(\text{Trace}(C^{2k})) = (1 + o(1)) \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k. \quad (18)$$

The proof of this technical Lemma is quite long. We will delay its proof until the end of this section.

Lemma 7 If $p(1-p) \gg \frac{\log^4 n}{n^{r-s}}$, then we have $\|C\| \leq (2\binom{r-s}{s} + o(1)) \sqrt{d(1-p)}$ almost surely.

Proof: By Lemma 6, we have $\mathbb{E}(\text{Trace}(C^{2k})) \leq (1 + o(1)) \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k$. As $\mathbb{E}(\|C\|^{2k}) \leq \mathbb{E}(\text{Trace}(C^{2k}))$, we have

$$\mathbb{E}(\|C\|^{2k}) \leq (1 + o(1)) \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k.$$

Let $U := \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k$. By Markov's inequality,

$$\begin{aligned} \Pr\left(\|C\| \geq (1 + \epsilon) \sqrt[2k]{U}\right) &= \Pr\left(\|C\|^{2k} \geq (1 + \epsilon)^{2k} U\right) \\ &\leq \frac{\mathbb{E}(\|C\|^{2k})}{(1 + \epsilon)^{2k} U} \\ &\leq \frac{(1 + o(1))U}{(1 + \epsilon)^{2k} U} \\ &= \frac{1 + o(1)}{(1 + \epsilon)^{2k}}. \end{aligned}$$

Let $g(n)$ be a slowly growing function such that $g(n) \rightarrow \infty$ as n approaches the infinity and $g(n) \ll \frac{(n^{r-s}p(1-p))^{1/4}}{s \log n}$. This is possible because $n^{r-s}p(1-p) \gg \log^4 n$. Choose $k = sg(n) \log n$ and $\epsilon = 1/g(n)$. We have $k \ll (n^{r-s}p(1-p))^{1/4}$ and $\epsilon \rightarrow 0$. Then we have $(1 + o(1))/(1 + \epsilon)^{2k} = O(n^{-s})$, which implies that almost surely

$$\begin{aligned} \|C\| &\leq (1 + o(1)) \sqrt[2k]{U} \\ &= (1 + o(1)) \left(\frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k \right)^{\frac{1}{2k}} \\ &< n^{\frac{s}{2k}} 2 \sqrt{\frac{n^{r-s} \binom{r-s}{s} p(1-p)}{s!(r-2s)!}} \\ &= \left(2 \binom{r-s}{s} + o(1) \right) \sqrt{d(1-p)}. \end{aligned}$$

□

Recall $M_2 = \frac{1}{\binom{r-s}{s}d} C$. We have

Lemma 8 If $p(1-p) \gg \frac{\log^4 n}{n^{r-s}}$, then we have $\|M_2\| \leq (2+o(1))\sqrt{\frac{1-p}{d}}$ almost surely.

Lemma 9 If $p(1-p) \gg \frac{\log^4 n}{n^{r-s}}$, then we have $\|M_1\| = O\left(\frac{\sqrt{(1-p)\log N}}{d}\right)$ almost surely.

Proof: We have

$$\begin{aligned} M_1 &= \frac{1}{\binom{r-s}{s}} \left(D^{-1/2} C D^{-1/2} - d^{-1} C \right) \\ &= \frac{1}{\binom{r-s}{s}} \left((D^{-1/2} - d^{-1/2} I) C D^{-1/2} + d^{-1/2} C (D^{-1/2} - d^{-1/2} I) \right). \end{aligned}$$

Note $\|D^{-1/2} - d^{-1/2} I\| \leq |d_{\min}^{-1/2} - d^{-1/2}| = O\left(\frac{\sqrt{\log N}}{d}\right)$, $\|D^{-1/2}\| \leq d_{\min}^{-1/2} = (1+o(1))d^{-1/2}$, and $\|C\| = (2\binom{r-s}{s} + o(1))\sqrt{d(1-p)}$. We have

$$\begin{aligned} \|M_1\| &= \frac{1}{\binom{r-s}{s}} \left\| (D^{-1/2} - d^{-1/2} I) C D^{-1/2} + d^{-1/2} C (D^{-1/2} - d^{-1/2} I) \right\| \\ &= O\left(\frac{\sqrt{(1-p)\log N}}{d}\right). \end{aligned}$$

□

Proof of Theorem 2: Combining Lemmas 3, 5, 8, and 9, we have

$$\begin{aligned} \|M\| &= \|M_1 + M_2 + M_3 + M_4\| \\ &\leq \|M_1\| + \|M_2\| + \|M_3\| + \|M_4\| \\ &\leq O\left(\frac{\sqrt{(1-p)\log N}}{d}\right) + \frac{(2+o(1))\sqrt{1-p}}{\sqrt{d}} + O\left(\frac{\sqrt{\log N}}{n\sqrt{d}}\right) + (1+o(1))\sqrt{\frac{1-p}{d}} \\ &= (3+o(1))\sqrt{\frac{1-p}{d}}. \end{aligned}$$

In the last step, we use the fact $\frac{\sqrt{\log N}}{n\sqrt{d}} = o\left(\sqrt{\frac{1-p}{d}}\right)$ since $1-p \gg \frac{\log n}{n^2}$.

By Lemma 1, for $1 \leq k \leq \binom{n}{s} - 1$, we have

$$|\lambda_k^{(s)}(H^r(n, p)) - \lambda_k^{(s)}(K_n^r)| \leq \|M\| \leq (3+o(1))\sqrt{\frac{1-p}{d}}.$$

□

Recall that X_F is the random indicator variable for F being an edge in $H^r(n, p)$. For any fixed positive integer t , the terms in $\text{Trace}(C^t)$ are of the form

$$c_{S_1 S_2} c_{S_2 S_3} \cdots c_{S_t S_1}.$$

Here $c_{ST} = W(S, T) - \mathbb{E}(W(S, T)) = \sum_{\substack{F \in \binom{V}{r} \\ S \cup T \subset F}} (X_F - \mathbb{E}(X_F))$ if $S \cap T = \emptyset$; $c_{ST} = 0$ otherwise.

Note $c_{S_i S_j} = 0$ if $S_i \cap S_j \neq \emptyset$. Thus we need only to consider the sequence $S_1 S_2 \dots S_t S_1$ such that $S_i \cap S_{i+1} = \emptyset$ for each $1 \leq i \leq t$, here $t+1 = 1$.

For $F \in \binom{V}{r}$ and $S, T \in \binom{V}{s}$, we define a random variable c_{ST}^F as follows.

$$c_{ST}^F = \begin{cases} X_F - \mathbb{E}(X_F) & \text{if } S \cap T = \emptyset \text{ and } S \cup T \subseteq F; \\ 0 & \text{otherwise.} \end{cases}$$

The sequence $w := S_1 F_1 S_2 F_2 S_3 \dots S_t F_t S_1$ is called a closed s -walk of length t if

1. $S_1, \dots, S_t \in \binom{V}{s}$,
2. $F_1, \dots, F_t \in \binom{V}{r}$,
3. $S_i \cap S_{i+1} = \emptyset$, for $i = 1, 2, \dots, t$,
4. $S_i \cup S_{i+1} \subset F_i$, for $i = 1, 2, \dots, t$.

Here we use the convention $S_{t+1} = S_1$. Those r -sets F_i 's are referred as edges while those s -sets S_i 's are referred as stops. For $1 \leq i \leq t$, we say w walks from S_i to S_{i+1} at step i via the edge F_i .

Using the notation above, we rewrite the trace as

$$\text{Trace}(C^t) = \sum_{\text{closed } s\text{-walks}} c_{S_1 S_2}^{F_1} c_{S_2 S_3}^{F_2} \dots c_{S_t S_1}^{F_t},$$

where the summation is over all possible closed s -walks of length t .

Taking the expectation on both sides, we get

$$\mathbb{E}(\text{Trace}(C^t)) = \sum_{\text{closed } s\text{-walks}} \mathbb{E}(c_{S_1 S_2}^{F_1} c_{S_2 S_3}^{F_2} \dots c_{S_t S_1}^{F_t}).$$

The terms in the product above can be regrouped according to the values of F_i 's; those terms with distinct F 's are independent to each other. Since $\mathbb{E}(c_{S,T}^F) = 0$, the contribution of a closed walk is 0 if some F appears only once. Thus we need only to consider the set of closed walks where each edge appears at least twice or do not occur; we call these closed walks as *good* closed walks. A good closed walk can contain at most $\lfloor \frac{t}{2} \rfloor$ distinct edges.

Let \mathcal{G}_i be the set of good closed walks of length t with i distinct edges. For $1 \leq i \leq \lfloor \frac{t}{2} \rfloor$, let \mathcal{G}_i^j be the set of good closed walks with exactly i distinct edges and j distinct vertices; we have $\mathcal{G}_i := \cup_j \mathcal{G}_i^j$.

We consider a good closed walk in \mathcal{G}_i . When a new edge comes in the walk, it can bring in at most $(r-s)$ new vertices. Thus such a good closed walk covers at most $m_i := s + i(r-s)$ vertices. Any walk contains at least one edge. Hence, the number of vertices in a walk from \mathcal{G}_i is in the interval $[r, m_i]$.

We have

$$\mathbb{E}(\text{Trace}(C^t)) = \sum_{i=1}^{\lfloor \frac{t}{2} \rfloor} \sum_{S_1 F_1 S_2 \dots S_t S_1 \in \mathcal{G}_i} \mathbb{E}(c_{S_1 S_2}^{F_1} c_{S_2 S_3}^{F_2} \dots c_{S_t S_1}^{F_t}). \quad (19)$$

Assume that an edge F occurs q times in a good closed walk and $T := \{i : 1 \leq i \leq t \text{ and } F_i = F\}$. We have $\Pr\left(\prod_{i \in T} c_{S_i S_{i+1}}^F = (1-p)^q\right) = p$ and $\Pr\left(\prod_{i \in T} c_{S_i S_{i+1}}^F = (-p)^q\right) = 1-p$. Thus, for each positive integer $l \geq 2$, we have

$$\mathbb{E}\left(\prod_{i \in T} c_{S_i S_{i+1}}^F\right) = (1-p)^q p + (-p)^q (1-p) \leq p(1-p).$$

The equality holds for $q = 2$.

Pick a good closed walk $w := S_1 F_1 S_2 F_2 S_3 \dots S_t F_t S_1$ in \mathcal{G}_i . Let F^1, \dots, F^i be the list of distinct edges in the order as they appear in w .

For each $1 \leq l \leq i$, let $T_l := \{1 \leq j \leq t : F_j = F^l\}$; then $\sum_{l=1}^i |T_l| = t$. We have

$$\mathbb{E}(c_{S_1 S_2}^{F_1} c_{S_2 S_3}^{F_2} \dots c_{S_t S_1}^{F_t}) = \prod_{l=1}^i \prod_{j \in T_l} \mathbb{E}(c_{S_j S_{j+1}}^{F^l}) \leq \prod_{l=1}^i p(1-p) = p^i (1-p)^i.$$

This implies

$$\sum_{S_1 F_1 S_2 \dots S_t S_1 \in \mathcal{G}_i} \mathbb{E}(c_{S_1 S_2}^{F_1} c_{S_2 S_3}^{F_2} \dots c_{S_t S_1}^{F_t}) \leq |\mathcal{G}_i| p^i (1-p)^i \quad (20)$$

for all $1 \leq i \leq \lfloor \frac{t}{2} \rfloor$. In particular, the equality holds when $t = 2i$. Combining equation (19) and inequality (20), we get

$$\mathbb{E}(\text{Trace}(C^t)) \leq \sum_{i=1}^{\lfloor \frac{t}{2} \rfloor} |\mathcal{G}_i| p^i (1-p)^i. \quad (21)$$

Now we estimate the value of $|\mathcal{G}_i^j|$, the number of good closed walks of length t on i edges and j vertices. Let w be a good closed walk in \mathcal{G}_i^j . For $2 \leq k \leq i$, let $\dots S F^k S' \dots$ be a piece of sequence in w where the edge F^k occurs first time; S is called the in-stop of F^k and S' is called the out-stop of F^k .

The following lemma will state the hypergraph structure of these i edges; it is independent of the walk w . We will use the following notation. Let $\mathcal{S} = \cup_{l=1}^i \binom{F^l}{s}$. For any s -set $S \in \mathcal{S}$, the degree of S , denoted by d_S , is the number of edges in $\{F^1, F^2, \dots, F^i\}$ containing S .

Define

$$d'_S = \begin{cases} d_S - 1 & \text{if there exists a unique } k \text{ such that } S = F_k \cap (\cup_{l=1}^{k-1} F_l), \\ d_S & \text{otherwise.} \end{cases}$$

Lemma 10 *Assume that F^1, \dots, F^i is the list of distinct edges in the order as they appear in $w \in \mathcal{G}_i^j$. Then we have*

$$\sum_{S \in \mathcal{S}} (d'_S - 1) \leq \left(1 + \frac{2}{s} \binom{r}{s-1}\right) (m_i - j).$$

Proof: For $2 \leq k \leq i$, let $x_k = |F^k \setminus (\cup_{l=1}^{k-1} F^l)|$; we have

$$0 \leq x_k \leq r - s.$$

Thus,

$$j = r + x_2 + x_3 + \dots + x_i \leq r + (i-1)(r-s) = m_i.$$

Since a new edge F_k can contribute at most $\binom{r-x_k}{s}$ to $\sum_{S \in \mathcal{S}} (d_S - 1)$, we have

$$\sum_{S \in \mathcal{S}} (d_S - 1) \leq \sum_{k=2}^i \binom{r-x_k}{s}.$$

Let $K := \{k : x_k = r-s, 2 \leq k \leq i\}$ and $\overline{K} = \{2, \dots, i\} \setminus K$. The edges in the set $\{F_k : k \in K\}$ are called *forward* edges while the edges in the set $\{F_k : k \in \overline{K}\}$ are called *backward* edges. Note each backward edge contribute at least one to $m_i - j$; thus

$$m_i - j \geq |\overline{K}|.$$

Note for each $k \in K$, $\binom{r-x_k}{s} = 1$. We have

$$\begin{aligned}
\sum_{S \in \mathcal{S}} (d_S - 1) &\leq \sum_{k=2}^i \binom{r-x_k}{s} \\
&= |K| + \sum_{k \in \overline{K}} \binom{r-x_k}{s} \\
&= |K| + \sum_{k \in \overline{K}} \frac{r-x_k-s+1}{s} \binom{r-x_k}{s-1} \\
&\leq |K| + \sum_{k \in \overline{K}} \frac{2}{s} \binom{r}{s-1} (r-x_k-s) \\
&= |K| + \frac{2}{s} \binom{r}{s-1} (m_i - j).
\end{aligned}$$

For any $k \in K$, let $S(F_k) := F_k \cap (\cup_{l=1}^{k-1} F_l)$ be the starting stop of F_k when F_k first occurs in w . List the elements in K as $k_1, k_2, \dots, k_{|K|}$ in an increasing order. Consider the sequence of stops $S(F_{k_1}), S(F_{k_2}), \dots, S(F_{k_{|K|}})$ (not necessarily distinct). Let z be the number of distinct stops in the sequence. If $S(F_{k_i})$ does not appear the first time in the sequence above, then we consider the partial walk $S_{k_{l-1}} F_{k_{l-1}} \dots S_{k_i} F_{k_i}$. Since $F_{k_{l-1}}$ is a forward edge, there exists at least one backward edge $F^{l'}$ for some $l' \in (k_{l-1}, k_l)$. Thus,

$$|K| \leq z + |\overline{K}| \leq z + m_i - j.$$

Hence,

$$\begin{aligned}
\sum_{S \in \mathcal{S}} (d'_S - 1) &= \sum_{S \in \mathcal{S}} (d_S - 1) - z \\
&\leq |K| + \frac{2}{s} \binom{r}{s-1} (m_i - j) - z \\
&\leq \frac{2}{s} \binom{r}{s-1} (m_i - j) + m_i - j \\
&= \left(1 + \frac{2}{s} \binom{r}{s-1}\right) (m_i - j).
\end{aligned}$$

The proof of this Lemma is finished. \square

Lemma 11 For $1 \leq i \leq \lfloor \frac{t}{2} \rfloor$ and $r \leq j \leq m_i$, we have

$$|\mathcal{G}_i^j| \leq \binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^{m_i}}{(s!)^{i+1} ((r-2s)!)^i} \left(\frac{C_1 i C_2}{n}\right)^{m_i-j}.$$

Here C_1 and C_2 depend only on r and s , independent of i , j , and n .

Corollary 1 For $1 \leq i \leq \lfloor \frac{t}{2} \rfloor$ and $n \gg i^{C_2}$, we have

$$|\mathcal{G}_i| \leq (1 + o(1)) \binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^{m_i}}{(s!)^{i+1} ((r-2s)!)^i}. \quad (22)$$

Proof: We can associate a walk $w \in \mathcal{G}_i^j$ with a code of length t consisting of three symbols: ‘(’, ‘)’, and ‘*’. We scan the edges of the walk w from left to right; if an edge appears first time, then we assign the code ‘(’; if an edge appears second time, then we assign the code ‘)’; otherwise, we assign the code ‘*’.

For example, consider the following good walk with $i = 3$, $j = 8$, and $t = 8$:

$$w = S_1 F_1 S_2 F_2 S_3 F_3 S_4 F_1 S_5 F_1 S_4 F_3 S_3 F_2 S_2 F_1 S_1$$

Here edges are: $F_1 = (1, 2, 3, 4, 5)$, $F_2 = (4, 5, 6, 7, 8)$, $F_3 = (7, 8, 5, 3, 4)$. Stops are: $S_1 = (1, 2)$, $S_2 = (4, 5)$, $S_3 = (7, 8)$, $S_4 = (3, 4)$, $S_5 = (2, 5)$. The code for this walk is $((()*)*)$.

Since w has i distinct edges, there are i ‘(’, i ‘)’s, and $(t - 2i)$ ‘*’s. Note that the number of ‘(’ is always greater than or equal to the number of ‘)’ at any point when the sequence is read from left to right; each ‘(’ has a matched ‘)’ in the sequence. The symbol ‘*’ starts at position three and up. There are $\binom{t-2}{t-2i}$ ways to choose the ‘*’-positions and $\frac{1}{i+1} \binom{2i}{i}$ ways to choose i matched parentheses (the Catalan number). The number of such codes is

$$\binom{t-2}{t-2i} \frac{1}{i+1} \binom{2i}{i}.$$

To construct a walk from a given code, we scan the symbols from left to right. The first symbol is always ‘(’. There are $\binom{n}{s}$ ways to choose the first stop S_1 and $\binom{n-s}{r-s}$ ways to choose the rest of vertices in the first edge F^1 . Suppose that we already build a partial walk and need to decide the next stop and the next edge. There are at most $\binom{r-s}{s}$ ways to choose the next stop S . The choices of selecting the next edge depends on the next available symbol in the code sequence. Let $b_{(}$, $b_{)}$, and b_* be the product of the number of ways to choose the next edge at the ‘(’, ‘)’, and ‘*’ positions respectively. We have

$$|\mathcal{G}_i^j| \leq \binom{t-2}{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{n}{s} \binom{r-s}{s}^t b_{(} \cdot b_{)} \cdot b_{*}. \quad (23)$$

First we estimate $b_{(}$, the number of ways to choose new edges F^1, \dots, F^i given the first stop S_1 . Besides the s vertices selected at the first stop, there are $\binom{n-s}{j-s}$ ways to choose remaining $j - s$ vertices. Recall that F^1, \dots, F^i is the list of distinct edges in the order as they appear in w . For $2 \leq l \leq i$, let $\tilde{F}^l := F^l \setminus (\cup_{l'=1}^{l-1} F^{l'})$, $x_l := |\tilde{F}^l|$, and $y_l := r - s - x_l$. We also define $\tilde{F}^1 := F^1 \setminus S_1$; $x_1 := |\tilde{F}^1| = r - s$, and $y_1 = 0$. Note that $\cup_{l=1}^i \tilde{F}^l$ forms a partition of the remaining $(j - s)$ selected vertices. The number of ways to choose such a partition is

$$\frac{(j-s)!}{x_1! x_2! \cdots x_i!}.$$

To choose F_l , we need select x_l new vertices and y_l old vertices; each old vertex has at most j choices. We have

$$b_{(} \leq \sum_{x_2, \dots, x_i} \binom{n-s}{j-s} \frac{(j-s)!}{(r-s)! x_2! \cdots x_i!} j^{\sum_{l=2}^i y_l}.$$

Observe that $\sum_{l=2}^i y_l = m_i - j$ and

$$\frac{(j-s)!}{(r-s)! x_2! \cdots x_i!} \leq \binom{j-s}{r-s} \frac{(m_i - r)!}{((r-s)!)^{i-1}}.$$

The number of ways to choose x_2, \dots, x_i is the same as the number of ways to choose y_2, \dots, y_i , which is $\binom{m_i-j+i-2}{m_i-j} \leq (m_i-j+i-2)^{m_i-j}$. Therefore,

$$\begin{aligned}
b_{\langle} &\leq \sum_{x_2, \dots, x_i} \binom{n-s}{j-s} \frac{(j-s)!}{(r-s)!x_2! \cdots x_i!} j^{\sum_{i=2}^i y_i} \\
&\leq \binom{n-s}{j-s} \binom{j-s}{r-s} \frac{(m_i-r)!}{((r-s)!)^{i-1}} (m_i-j+i-2)^{m_i-j} j^{m_i-j} \\
&\leq \binom{n-s}{j-s} \binom{j-s}{r-s} \frac{(m_i-r)!}{((r-s)!)^{i-1}} \left(\frac{m_i+i-2}{2}\right)^{2(m_i-j)}. \tag{24}
\end{aligned}$$

There is at most i choices of edges at each ‘*’ position. Thus

$$b_* \leq i^{t-2i}. \tag{25}$$

It remains to bound b_{\rangle} . We first present an easy bound for b_{\rangle} . Edge F can be chosen at most one ‘)’-position. For any possible stop $S \in \mathcal{S}$, S can appear at the)-positions at most d_S times; each occurrence of S involves different edges since we are considering the second occurrence of edges. Thus,

$$b_{\rangle} \leq \prod_{S \in \mathcal{S}} d_S! \leq \prod_{S \in \mathcal{S}} d_S^{d_S-1} \leq i^{\sum_{S \in \mathcal{S}} (d_S-1)}.$$

We need a better upper bound for b_{\rangle} . Consider a stop S which is first chosen at a ‘)’-position. Let F be the edge on the walk right before the ‘)’-position; i.e., the walk w enters S through F . If this F occurred before, then the choices of edges at ‘)’-positions starting with S is at most

$$(d_S - 1)! \leq i^{d_S-2}.$$

If this F occurs first time and F is an forward edge, then there is only one choice for the next edge leaving S ; namely F itself. In this case, the choices of edges at ‘)’-positions starting with S is at most

$$(d_S - 1)! \leq i^{d_S-2}.$$

In the remaining case, F must be a backward edge. The number of backward edges is at most $m_i - j$. Since F contains at most $\binom{r}{s}$ stops, the number of such S is at most $\binom{r}{s} (m_i - j)$. A additional factor $i^{\binom{r}{s}(m_i-j)}$ is enough. We have

$$\begin{aligned}
b_{\rangle} &\leq i^{\binom{r}{s}(m_i-j)} \prod_{S: d_S \geq 2} i^{d_S-2} \\
&\leq i^{\binom{r}{s}(m_i-j)} \prod_S i^{d_S-1} \\
&\leq i^{\binom{r}{s}(m_i-j)} i^{(1+\frac{2}{s} \binom{r}{s-1})(m_i-j)} \\
&= i^{(\binom{r}{s}+1+\frac{2}{s} \binom{r}{s-1})(m_i-j)}. \tag{26}
\end{aligned}$$

Combining equations (23), (24), (25), and (26), we get

$$\begin{aligned}
|\mathcal{G}_i^j| &\leq \binom{t-2}{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{n}{s} \binom{r-s}{s}^t \binom{n-s}{j-s} \binom{j-s}{r-s} \\
&\quad \frac{(m_i-r)!}{((r-s)!)^{i-1}} \left(\frac{m_i+i-2}{2} \right)^{2(m_i-j)} i^{t-2i} i^{\binom{r}{s}+1+\frac{2}{s}\binom{r}{s-1}} (m_i-j) \\
&\leq \binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^j}{(s!)^{i+1} ((r-2s)!)^i} \\
&\quad \frac{(m_i-r)!}{(j-r)!} \left(\frac{m_i+i-2}{2} \right)^{2(m_i-j)} i^{\binom{r}{s}+1+\frac{2}{s}\binom{r}{s-1}} (m_i-j) \\
&\leq \binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^{m_i}}{(s!)^{i+1} ((r-2s)!)^i} \left(\frac{C_1 i^{C_2}}{n} \right)^{m_i-j}
\end{aligned}$$

Here we set $C_1 = 4(r-s)^3$ and $C_2 = \binom{r}{s} + 4 + \frac{2}{s} \binom{r}{s-1}$. \square

Lemma 12 *If $t = 2k$ is even, then we have*

$$|\mathcal{G}_k^{m_k}| = \binom{n}{m_k} \frac{m_k!}{(k+1)(s!)^{k+1} ((r-2s)!)^k} \binom{2k}{k}. \quad (27)$$

Proof: We will construct a bijection from $\mathcal{G}_k^{m_k}$ to a triple (U, \mathcal{P}, C) , where U is a set of m_k vertices, \mathcal{P} is a partition of U into $(k+1)$ s -sets and k $(r-2s)$ -sets, and C is a code consisting of k pairs valid parentheses.

For any good walk $w \in \mathcal{G}_k^{m_k}$, let U be the set of vertices covered by w . Note each edge appears exactly twice. We define a graph T , whose vertices are the stops in w . Two stops are connected if they belong to one edge. Observe that T is acyclic and connected; T must be a tree. Since T has exactly k edges, T must have $k+1$ vertices. Hence w has exactly $k+1$ stops; we list them as S^0, S^1, \dots, S^k . For $1 \leq i \leq k$, let E_i be the set of $(r-2s)$ vertices in F^i but not in any stops. We get a partition: $U = (\cup_{j=0}^i S^j) \cup (\cup_{j=1}^i E_j)$. A code consists of k ‘(’ and k ‘)’ is generated as follows. When we scan the walk from left to right, if an edge appears the first time, we append the code by a ‘(’; otherwise, we append the code by a ‘)’. The code is a valid sequence of k pairs of parentheses. (In this case, the number of ‘*’s is zero.) It suffices to recover a walk from a partition of $[m_k]$ and a sequence of valid parentheses.

Given a partition of U

$$(\cup_{j=0}^i S^j) \cup (\cup_{j=1}^i E_j)$$

and a sequence of k pairs valid parentheses, we first build a rooted tree T as follows. At each time, we maintain a tree T , a current stop S , a set of unused stops \mathcal{S} . Initially T contains nothing but the root stop S_0 , $S := S_0$, and $\mathcal{S} = \{S_1, S_2, \dots, S_k\}$. At each time, read a symbol from the sequence. If the symbol is an open parenthesis, then find an S_i in \mathcal{S} with index i as small as possible, delete S_i from \mathcal{S} , attach S_i to T as a child stop of S , and let $S := S_i$; if the symbol is “)””, then let S point to the the parent stop of the current S . Repeat this process until all symbols from the sequence are processed.

Since every closed parenthesis has a matching open parenthesis, this process never get stuck. When the process ends, a rooted tree T on the vertex set $\{S_0, \dots, S_k\}$ is created. For $1 \leq i \leq k$, let F_i be the union of E_i and two ends of i -th edge, which created in the process. For example, for $k = 3$, if the sequence is $(())()$, then the corresponding good closed walk is

$$S_1 F_1 S_2 F_2 S_3 F_2 S_2 F_1 S_1 F_3 S_4 F_3 S_1$$

where $F_1 = S_1 \cup S_2 \cup E_1$, $F_2 = S_2 \cup S_3 \cup E_2$, and $F_3 = S_4 \cup S_1 \cup E_3$.

Thus, this is a bijection from $\mathcal{G}_k^{m_k}$ to all triples $\{U, \mathcal{P}, C\}$. The number of ways to choose m_k vertices is $\binom{n}{m_k}$. The number of ways to choose these sets $S_0, S_1, \dots, S_k, E_1, \dots, E_k$ as a partition of U is

$$\binom{m_k}{s, \dots, s, r-2s, \dots, r-2s} = \frac{m_k!}{(s!)^{k+1}((r-2s)!)^k}.$$

The number of sequences of k pairs valid parentheses is the Catalan number $\frac{1}{k+1} \binom{2k}{k}$. By taking product of these three numbers, we get equation (27). \square

Proof of Lemma 6: By equations (21) and (22), we have

$$\mathbb{E}(\text{Trace}(C^t)) \leq \sum_{i=1}^{\lfloor \frac{t}{2} \rfloor} |\mathcal{G}_i| p^i (1-p)^i \leq (1+o(1)) \sum_{i=1}^{\lfloor \frac{t}{2} \rfloor} a_i.$$

Here $a_i := \binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^{m_i} p^i (1-p)^i}{(s!)^{i+1} ((r-2s)!)^i}$. We get

$$\begin{aligned} \frac{a_i}{a_{i+1}} &= \frac{\binom{t-2}{t-2i} i^{t-2i} \frac{1}{i+1} \binom{2i}{i} \binom{r-s}{s}^{t-i} \frac{n^{m_i} p^i (1-p)^i}{(s!)^{i+1} ((r-2s)!)^i}}{\binom{t-2}{t-2i-2} i^{t-2i-2} \frac{1}{i+2} \binom{2i+2}{i+1} \binom{r-s}{s}^{t-i-1} \frac{n^{m_{i+1}} p^{i+1} (1-p)^{i+1}}{(s!)^{i+2} ((r-2s)!)^{i+1}}} \\ &= \frac{i^3 (2i-1)(i+2)}{(2i+1)(t-2i)(t-2i-1)} \frac{(r-s)!}{n^{r-s} p (1-p)} \\ &< \frac{3i^4 (r-s)!}{n^{r-s} p (1-p)}. \end{aligned}$$

When $n^{r-s} p (1-p) \gg t^4$, we have $a_i = o(a_{i+1})$. Thus,

$$\mathbb{E}(\text{Trace}(C^t)) \leq (1+o(1)) a_{\lfloor \frac{t}{2} \rfloor}.$$

When $t = 2k$, we get

$$\mathbb{E}(\text{Trace}(C^{2k})) \leq (1+o(1)) \frac{n^{s+k(r-s)} \binom{r-s}{s}^k}{(k+1)(s!)^{k+1} ((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k.$$

For $t = 2k+1$, we have

$$\begin{aligned} \mathbb{E}(\text{Trace}(C^{2k+1})) &\leq (1+o(1)) a_k \\ &\leq (1+o(1)) \frac{2k^2 n^{s+k(r-s)} \binom{r-s}{s}^{k+1}}{(k+1)(s!)^{k+1} ((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k. \end{aligned}$$

Now we assume $k = o(\log(n^{r-s} p (1-p)))$. For $t = 2k$, let

$$b_k := |\mathcal{G}_k^{m_k}| p^k (1-p)^k = \binom{n}{m_k} \frac{m_k!}{(k+1)(s!)^{k+1} ((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k.$$

It is clear that $\mathbb{E}(\text{Trace}(C^{2k})) \geq b_k$. We also have

$$\begin{aligned} \mathbb{E}(\text{Trace}(C^{2k})) - b_k &\leq \sum_{i=1}^{k-1} |\mathcal{G}_i| p^i (1-p)^i + \sum_{j=r}^{m_k-1} |\mathcal{G}_k^j| p^k (1-p)^k \\ &= (1+o(1)) \sum_{i=1}^{k-1} a_i + a_k \sum_{j=r}^{m_k-1} \left(\frac{C_1 k C_2}{n} \right)^{m_k-j}. \end{aligned}$$

Note $a_k = (1 + o(1)) \binom{r-s}{s}^k b_k$ and $a_i = O\left(a_k \left(\frac{k^4}{n^{r-s}p(1-p)}\right)^{k-i}\right)$. We conclude

$$\mathbb{E}(\text{Trace}(C^{2k})) - b_k = O\left(b_k \binom{r-s}{s}^k \left(\frac{k^4}{n^{r-s}p(1-p)} + \frac{C_1 k^{C_2}}{n}\right)\right) = o(b_k).$$

Here we use the fact $\binom{r-s}{s}^k \left(\frac{k^4}{n^{r-s}p(1-p)} + \frac{C_1 k^{C_2}}{n}\right) = o(1)$ since $k = o(\log(n^{r-s}p(1-p)))$. \square

4 The semicircle law

Let us review the definition of the Semicircle Law. Let $F(x)$ be the continuous distribution function with density $f(x)$ such that $f(x) = (2/\pi)\sqrt{1-x^2}$ when $|x| \leq 1$ and $f(x) = 0$ when $|x| > 1$. Let A be a Hermitian matrix of dimension $N \times N$. The *empirical distribution* of the eigenvalues of A is

$$F(A, x) := \frac{1}{N} |\{\text{eigenvalues of } A \text{ less than } x\}|.$$

We say, the empirical distribution of the eigenvalues of A asymptotically follows the Semicircle Law centered at c with radius R if $F(\frac{1}{R}(A - cI), x)$ tends to $F(x)$ in probability as N goes to infinity. (In this case, we write $F(\frac{1}{R}(A - cI), x) \xrightarrow{P} F(x)$.) If c is the center of the Semicircle Law, then any $c' = c + o(R)$ is also the center of the Semicircle Law.

Theorem 5 *If $n^{r-s}p(1-p) \rightarrow \infty$, then the empirical distribution of the eigenvalues of $W - \mathbb{E}(W)$ follows the semicircle law centered at 0 with radius $2\sqrt{\binom{r-s}{s} \binom{n-s}{r-s} p(1-p)}$.*

Proof: Let $R := 2\sqrt{\binom{r-s}{s} \binom{n-s}{r-s} p(1-p)}$, $C := W - \mathbb{E}(W)$, and $C_{nor} := \frac{1}{R}C$.

To prove the theorem, we need to show that for any fixed t , the t -th moment of $F(C_{nor}, x)$ (with n goes to infinity) is asymptotically equal to the t -th moment of $F(x)$. We know the t -th moment of $F(C_{nor}, x)$ equals $\binom{n}{s}^{-1} \mathbb{E}(\text{Trace}(C_{nor}^t))$. For even $t = 2k$, the t -th moment of $F(x)$ is $(2k)!/2^{2k}k!(k+1)!$. For odd t , the t -th moment of $F(x)$ is 0.

In order to prove the theorem, we need to show for any fixed k ,

$$\frac{1}{\binom{n}{s}} \mathbb{E}(\text{Trace}(C_{nor}^{2k})) = (1 + o(1)) \frac{(2k)!}{2^{2k}k!(k+1)!}$$

and

$$\frac{1}{\binom{n}{s}} \mathbb{E}(\text{Trace}(C_{nor}^{2k+1})) = o(1).$$

We know

$$\mathbb{E}(\text{Trace}(C_{nor}^t)) = \frac{1}{R^t} \mathbb{E}(\text{Trace}(C^t))$$

for any t . By Lemma 6, we have

$$\mathbb{E}(\text{Trace}(C^{2k})) = (1 + o(1)) \frac{n^{s+k(r-s)}}{(k+1)(s!)^{k+1}((r-2s)!)^k} \binom{2k}{k} p^k (1-p)^k.$$

Then

$$\frac{1}{\binom{n}{s}} \mathbb{E}(\text{Trace}(C_{nor}^{2k})) = (1 + o(1)) \frac{(2k)!}{2^{2k}k!(k+1)!}$$

as desired.

By Lemma 6 again, we have

$$\mathbb{E}(\text{Trace}(C^{2k+1})) = O\left(\frac{2k^2 n^{s+k(r-s)} p^k (1-p)^k \binom{r-s}{s}^{k+1}}{(k+1)(s!)^{k+1} ((r-2s)!)^k} \binom{2k}{k}\right).$$

Thus

$$\frac{1}{\binom{n}{s}} \mathbb{E}(\text{Trace}(C_{nor}^{2k+1})) = O\left(\frac{2k^2 \binom{2k}{k} \binom{r-s}{s}^{k+1}}{2^{2k} (k+1) R}\right) = o(1).$$

Here k is any constant but $R \rightarrow \infty$. The theorem is proved. \square

The following Lemma is useful to derive the Semicircle Law from one matrix to the other.

Lemma 13 *Let A and B be two $(N \times N)$ -Hermitian matrices. Suppose that the empirical distribution of the eigenvalues of A follows the Semicircle Law centered at c with radius R . If either $\|B\| = o(R)$ or the rank of B is $o(N)$, then the empirical distribution of the eigenvalues of $A + B$ also follows the Semicircle Law centered at c with radius R .*

Proof: It suffices to show $F(\frac{1}{R}(A + B - cI), x) \xrightarrow{P} F(x)$. First we assume $\|B\| = o(R)$. By Lemma 1, for $1 \leq k \leq N$, we have

$$\left| \mu_k \left(\frac{1}{R}(A + B - cI) \right) - \mu_k \left(\frac{1}{R}(A - cI) \right) \right| \leq \frac{\|B\|}{R} = o(1).$$

Hence

$$F\left(\frac{1}{R}(A - cI), x - \frac{\|B\|}{R}\right) \leq F\left(\frac{1}{R}(A + B - cI), x\right) \leq F\left(\frac{1}{R}(A - cI), x + \frac{\|B\|}{R}\right).$$

Since $\|B\| = o(R)$, we have $F\left(\frac{1}{R}(A - cI), x - \frac{\|B\|}{R}\right) \xrightarrow{P} F(x)$ and $F\left(\frac{1}{R}(A - cI), x + \frac{\|B\|}{R}\right) \xrightarrow{P} F(x)$. By the Squeeze theorem, we have $F(\frac{1}{R}(A + B - cI), x) \xrightarrow{P} F(x)$.

Now we assume $\text{rank}(B) = o(N)$. Let U be the kernel of B (i.e. $B|_U = 0$); U has co-dimension $\text{rank}(B)$. Let $Z := \frac{1}{R}(A - cI)|_U = \frac{1}{R}(A + B - cI)|_U$. By Cauchy's interlace theorem [23], for $1 \leq j \leq N - \text{rank}(B)$, we have

$$\begin{aligned} \mu_j \left(\frac{1}{R}(A - cI) \right) &\leq \mu_j(Z) \leq \mu_{j+\text{rank}(B)} \left(\frac{1}{R}(A - cI) \right), \\ \mu_j \left(\frac{1}{R}(A + B - cI) \right) &\leq \mu_j(Z) \leq \mu_{j+\text{rank}(B)} \left(\frac{1}{R}(A + B - cI) \right). \end{aligned}$$

Thus, for $\text{rank}(B) + 1 \leq j \leq N - \text{rank}(B)$, we have

$$\mu_{j-\text{rank}(B)} \left(\frac{1}{R}(A - cI) \right) \leq \mu_j \left(\frac{1}{R}(A + B - cI) \right) \leq \mu_{j+\text{rank}(B)} \left(\frac{1}{R}(A - cI) \right).$$

It implies

$$F\left(\frac{1}{R}(A - cI), x\right) - \frac{\text{rank}(B)}{N} \leq F\left(\frac{1}{R}(A + B - cI), x\right) \leq F\left(\frac{1}{R}(A - cI), x\right) + \frac{\text{rank}(B)}{N}.$$

Since $\text{rank}(B) = o(N)$, we have $F\left(\frac{1}{R}(A - cI), x\right) \pm \frac{\text{rank}(B)}{N} \xrightarrow{P} F(x)$. By the Squeeze theorem, we have $F(\frac{1}{R}(A + B - cI), x) \xrightarrow{P} F(x)$. \square

Proof of Theorem 3: Recall

$$\mathcal{L}^{(s)}(K_n^r) - \mathcal{L}^{(s)}(H^r(n, p)) = M_1 + M_2 + M_2 + M_4.$$

We can write $\mathcal{L}^{(s)}(H^r(n, p))$ as $-M_2 + \left(1 - \frac{(-1)^s}{\binom{n}{s}}\right) I + B_1 - M_3 - M_4 - M_1$, where $B_1 = \mathcal{L}^{(s)}(K_n^r) - \left(1 - \frac{(-1)^s}{\binom{n}{s}}\right) I$.

By Theorem 5, the empirical distribution of the spectrum of $W - \mathbb{E}(W)$ follows the Semicircle Law centered at 0 with radius $(2 + o(1))\sqrt{\binom{r-s}{s}\binom{n-s}{r-s}p(1-p)}$. Since $M_2 = \frac{1}{\binom{r-s}{s}d}(W - \mathbb{E}(W))$, $\left(1 - \frac{(-1)^s}{\binom{n}{s}}\right) I - M_2$ follows the Semicircle Law centered at $c := 1 - \frac{(-1)^s}{\binom{n}{s}}$ with radius $R := (2 + o(1))\sqrt{\frac{1-p}{\binom{r-s}{s}\binom{n-s}{r-s}p}}$. Note $\frac{(-1)^s}{\binom{n}{s}} = o(R)$. We can change the center to 1.

By Theorem 1, $\mathcal{L}^{(s)}(K_n^r)$ has an eigenvalue $1 - (-1)^s \frac{\binom{n-s}{s}}{\binom{n}{s}}$ with multiplicity $\binom{n}{s} - \binom{n}{s-1}$. Thus B_1 has rank $\binom{n}{s-1} = o\left(\binom{n}{s}\right)$. We also observe that M_4 has rank at most 2, $\|M_1\| = O\left(\frac{\sqrt{(1-p)\log N}}{d}\right) = o(R)$, and $\|M_3\| = O\left(\frac{\sqrt{\log N}}{n\sqrt{d}}\right) = o(R)$. Here we notice $d \gg \log^{1/3} n$ and $1 - p \gg \frac{\log n}{n^2 d^2}$.

By Lemma 13, the matrices B_1 , M_1 , M_3 , and M_4 will not affect the Semicircle Law. The proof of this Lemma is finished. \square

References

- [1] N. Alon, M. Krivelevich, and V. H. Vu, Concentration of eigenvalues of random matrices *Israel Math. J.* **131**, (2002) 259–267.
- [2] A. R. Calderbank and P. Frankl, Improved upper bounds concerning the Erdős-Ko-Rado Theorem *Combinatorics, Probability and Computing* **1**, (1992) 115–122.
- [3] H. Chernoff, A note on an inequality involving the normal distribution, *Ann. Probab.*, **9**, (1981), 533–535.
- [4] F. Chung, Diameters and eigenvalues, *J. of the Amer. Math. Soc.*, **2** (1989), 187–196.
- [5] F. Chung, The Laplacian of a hypergraph, In J. Friedman (Ed.), *Expanding graphs (DIMACS series)*, **1993**, 21–36.
- [6] F. Chung, V. Faber, and T.A. Manteuffel, An upper bound on the diameter of a graph from eigenvalues associated with its Laplacian, *Siam. J. Disc. Math.*, **7-3** (1994), 443–457.
- [7] F. Chung, *Spectral graph theory*, AMS publications, 1997.
- [8] F. Chung, Laplacians and the Cheeger inequality for directed graphs, *Annals of Comb.*, **9** (2005), 1–19.
- [9] F. Chung, The diameter and Laplacian eigenvalues of directed graphs, *Electronic Journal of Combinatorics*, **13** (2006), N4.
- [10] F. Chung, L. Lu, and V. H. Vu, Eigenvalues of random power law graphs, *Annals of Combinatorics*, **7** (2003), 21–33.

- [11] F. Chung, L. Lu, and V. H. Vu, Spectra of random graphs with given expected degrees, *Proceedings of the National Academy of Sciences*, **100**(11) (2003), 6313-6318.
- [12] F. Chung and M. Radcliffe, On the spectra of general random graphs, *Preprint*.
- [13] A. Coja-Oghlan, On the Laplacian eigenvalues of $G(n, p)$, *Combinatorics, Probability and Computing*, **16**(6) (2007), 923-946.
- [14] A. Coja-Oghlan and A. Lanka, The spectral gap of random graphs with given expected degrees, *Electronic Journal of Combinatorics*, (2009) R138.
- [15] X. Ding and T. Jiang, Spectral distributions of adjacency and Laplacian matrices of random graphs, *The Annals of Applied Probability*, **20**(6) (2010), 2086-2117.
- [16] P. Erdős, C. Ko, R. Rado, Intersection theorems for systems of finite sets, *Quarterly Journal of Mathematics*, Oxford Series, series **2** 12: (1961) 313-320.
- [17] U. Feige and E. Ofek, Spectral techniques applied to sparse random graphs, *Random Structures and Algorithms*, **27**(2) (2005), 251-275.
- [18] J. Friedman, J. Kahn, and E. Szemerédi, On the second eigenvalue in random regular graphs, in Proc. 21st ACM Symposium on Theory of Computing, Association for Computing Machinery, New York, 1989, 587-598.
- [19] J. Friedman, On the second eigenvalue and random walks in random-regular graphs, *Combinatorica* **11**, Number 4, (1991) 331-362.
- [20] J. Friedman, *A Proof of Alon's Second Eigenvalue Conjecture and Related Problem*, Memoirs of the American Mathematical Society 2008; 100 pp.
- [21] Z. Füredi and J. Komlós. The eigenvalues of random symmetric matrices, *Combinatorica*, 1(3) 1981, 233-241.
- [22] C. Godsil and G. Royle, Algebraic Graph Theory, Springer-Verlag, New York, 2001.
- [23] S.-G. Hwang, Cauchy's interlace theorem for eigenvalues of Hermitian matrices, *Amer. Math. Monthly* **111** (2004), 157-159.
- [24] G. O. H. Katona, A simple proof of the Erdős-Chao Ko-Rado theorem, *Journal of Combinatorial Theory, Series B* **13**: (1972) 183-184.
- [25] M. Krivelevich and B. Sudakov, The largest eigenvalue of sparse random graphs, *Combinatorics, Probability and Computing*, **12** (2003), 61-72.
- [26] L. Lu and X. Peng, High-Ordered Random Walks and Generalized Laplacians on Hypergraphs, in A. Frieze, P. Horn, P. Pralat (Eds.): *Algorithms and Models for the Web Graph - 8th International Workshop, (WAW 2011) Proceedings*. Lecture Notes in Computer Science **6732** Springer 2011, 14-25.
- [27] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press, 1978.
- [28] J. A. Rodríguez, On the Laplacian Spectrum and Walk-regular Hypergraphs, *Linear and Multilinear Algebra*, **51** (3) (2003), 285-297.
- [29] J. A. Rodríguez, Laplacian eigenvalues and partition problems in hypergraphs, *Applied Mathematics Letters*, **22** (2009), 916-921.