ON THE DIAMETER OF PERMUTATION GROUPS

HARALD A. HELFGOTT AND ÁKOS SERESS

ABSTRACT. Given a finite group G and a set A of generators, the diameter $\operatorname{diam}(\Gamma(G,A))$ of the Cayley graph $\Gamma(G,A)$ is the smallest ℓ such that every element of G can be expressed as a word of length at most ℓ in $A \cup A^{-1}$. We are concerned with bounding $\operatorname{diam}(G) := \max_A \operatorname{diam}(\Gamma(G,A))$.

It has long been conjectured that the diameter of the symmetric group of degree n is polynomially bounded in n, but the best previously known upper bound was exponential in $\sqrt{n \log n}$. We give a quasipolynomial upper bound, namely,

$$\operatorname{diam}(G) = \exp\left(O((\log n)^4 \log \log n)\right) = \exp\left((\log \log |G|)^{O(1)}\right)$$

for G = Sym(n) or G = Alt(n), where the implied constants are absolute. This addresses a key open case of Babai's conjecture on diameters of simple groups. By a result of Babai and Seress (1992), our bound also implies a quasipolynomial upper bound on the diameter of all transitive permutation groups of degree n.

1. Introduction

1.1. **Groups and their diameters.** Let A be a set of generators for a group G. The (undirected) Cayley graph $\Gamma(G,A)$ is the graph whose set of vertices is V=G and whose set of edges is $E=\{\{g,ga\}:g\in G,a\in A\}$. The diameter diam(Γ) of a graph $\Gamma(V,E)$ is defined by

(1.1)
$$\operatorname{diam}(\Gamma) = \max_{v_1, v_2 \in V} \min_{\substack{P \text{ a path} \\ \text{from } v_1 \text{ to } v_2}} \operatorname{length}(P).$$

In particular, the diameter of a Cayley graph $\Gamma(G,A)$ is the maximum, for $g \in G$, of the length ℓ of the shortest expression $g = a_1^{\varepsilon_1} a_2^{\varepsilon_2} \cdots a_{\ell}^{\varepsilon_{\ell}}$ with $a_i \in A$ and $\varepsilon_i \in \{-1,1\}$ for each $i = 1, \ldots, \ell$. We may define the diameter diam(G) of a finite group to be the maximal diameter of the Cayley graphs $\Gamma(G,A)$ for all generating sets A of G.

Much recent work on group diameters has been motivated by the following conjecture:

Conjecture 1. (Babai, published as [BS92, Conj. 1.7]) For all finite simple groups G.

$$\operatorname{diam}(G) \le (\log |G|)^{O(1)},$$

where the implied constant is absolute.

Here and henceforth, |S| denotes the number of elements of a set S.

The first class of finite simple groups for which Conj. 1 was established was $PSL_2(\mathbb{Z}/p\mathbb{Z})$ with p prime, by Helfgott [Hel08]. The paper [Hel08] initiated a period

Ákos Seress passed away on February 13, 2013, after the paper's acceptance.

of intense activity [BG08a], [BG08b], [Din11], [BGS10], [Hel11], [GH11], [Var12], [BGS11], [PS], [BGT11], [GH], [GV12]¹ on the diameter problem and the related problem of expansion properties of Cayley graphs.

As far as work in this vein on the diameter of finite simple groups is concerned, the best results to date are those of Pyber, Szabó [PS] and Breuillard, Green, Tao [BGT11]. Their wide-ranging generalisation covers all simple groups of Lie type, but (just like [GH11]) the diameter estimates retain a strong dependence on the rank; thus, they prove Conj. 1 only for groups of bounded rank. The problem for the alternating groups remained wide open.²

These two issues are arguably related: product theorems (of the type $|A \cdot A \cdot A| \gg |A|^{1+\delta}$ familiar since [Hel08]) are false both in the unbounded-rank case and in the case of alternating groups, and the counterexamples described in both situations in [PPSS12], [PS] are based on similar principles.

In the present paper we address the case of alternating (and symmetric) groups. We expect that some of the combinatorial difficulties we overcome will also arise in the context of linear groups of large rank.

For G = Alt(n), Conj. 1 stipulates that diam(Alt(n)) = $n^{O(1)}$; [BS92] refers to this special case of Conj. 1 as a "folklore" conjecture. Indeed, this has long been a problem of interest in computer science (see [KMS84], [McK84], [BHK⁺90], [BBS04], [BH05]). On a more playful level, bounds on the diameter of permutation groups are relevant to every permutation puzzle (e.g., Rubik's cube).

The best previously known upper bound on diam(G) for G = Alt(n) or G = Sym(n) was more than two decades old:

(1.2)
$$\operatorname{diam}(G) \leq \exp((1+o(1))\sqrt{n\log n}) = \exp((1+o(1))\sqrt{\log |G|}),$$
 due to Babai and Seress [BS88]. (We write $\exp(x)$ for e^x .)

1.2. **Statement of results.** Recall that a function f(n) is called *quasipolynomial* if $\log(f(n))$ is a polynomial function of $\log n$. Our main result establishes a quasipolynomial upper bound for diam(Alt(n)) and diam(Sym(n)).

Main Theorem. Let
$$G = \operatorname{Sym}(n)$$
 or $\operatorname{Alt}(n)$. Then $\operatorname{diam}(G) \le \exp\left(O((\log n)^4 \log \log n)\right)$,

where the implied constant is absolute.

The quasipolynomial bound extends to a much broader class of permutation groups. Recall that a permutation group G acting on a set Ω is called *transitive* if

$$\forall \alpha, \beta \in \Omega \quad \exists g \in G \text{ such that } g \text{ takes } \alpha \text{ to } \beta.$$

The size $|\Omega|$ of the permutation domain is called the *degree* of G.

Kornhauser et al. [KMS84] and McKenzie [McK84] raised the question of what classes of permutation groups may have polynomial diameter bound in their degree.

¹This list is not meant to be exhaustive.

²See, e.g., I. Pak's remarks (made already before [PS], [BGT11]) on the relative difficulty of the work remaining to do in the linear case (to be finished "in the next 10 years") and of the problem on Alt(n), for which there was "much less hope" [Pak].

A weaker, quasipolynomial bound for all transitive groups was formally conjectured in [BS92]:

Conjecture 2. ([BS92, Conj. 1.6]) If G is a transitive permutation group of degree n then diam $(G) \le \exp((\log n)^{O(1)})$.

Babai and Seress [BS92] linked Conj. 2 to the diameter of alternating groups:

Theorem 1.1. ([BS92, Thm. 1.4]) If G is a transitive permutation group of degree n then

$$\operatorname{diam}(G) \le \exp\left(O(\log n)^3\right) \operatorname{diam}\left(\operatorname{Alt}(k)\right),$$

where Alt(k) is the largest alternating composition factor of G.

Combining our Main Theorem with Thm. 1.1, we immediately obtain

Corollary 1.2. Conjecture 2 is true; indeed the diameter of any transitive permutation group G of degree n is

$$\operatorname{diam}(G) \le \exp\left(O((\log n)^4 \log \log n)\right).$$

We note that Thm. 1.1 is not only used to prove Cor. 1.2 – it also comes into play as an inductive tool in the proof of the Main Theorem (see Lemma 6.3). Since Thm. 1.1 relies on the Classification of Finite Simple Groups, so does the Main Theorem.

It is well-known that, for any finite group G and any set A of generators of G, the eigenvalues $\lambda_0 \geq \lambda_1 \geq \lambda_2 \geq \ldots$ of the adjacency matrix of $\Gamma(G, A)$ satisfy

(1.3)
$$\lambda_0 - \lambda_1 \ge \frac{1}{\operatorname{diam}(\Gamma(G, A))^2}.$$

(See [DSC93, Cor. 1] or the references [Ald87], [Bab91], [Gan91], [Moh91] therein.) Because of (1.3), we obtain immediately that

$$\lambda_0 - \lambda_1 \ge \exp(-O((\log n)^4 \log \log n)),$$

with consequences on expansion and the mixing rate (see, e.g., [Lov96], [HLW06]). Finally, the Main Theorem and Cor. 1.2 extend to directed graphs. Given $G = \langle A \rangle$, the directed Cayley graph $\vec{\Gamma}(G, A)$ is the graph with vertex set G and edge set $\{(g, ga) : g \in G, a \in A\}$. The diameter of $\vec{\Gamma}(G, A)$ is defined by (1.1), where "path" should be read as "directed path"; $\overrightarrow{\text{diam}}(G)$ is the maximum of $\overrightarrow{\text{diam}}(\vec{\Gamma}(G, A))$ taken as A varies over all generating sets A of G. Thanks to Babai's bound $\overrightarrow{\text{diam}}(G) = O\left(\overrightarrow{\text{diam}}(G) \cdot (\log |G|)^2\right)$ [Bab06, Cor. 2.3], valid for all groups G, we obtain immediately from Cor. 1.2 that

Corollary 1.3. Let G be a transitive group on n elements. Then

$$\overrightarrow{\operatorname{diam}}(G) \le \exp(O((\log n)^4 \log \log n)).$$

1.3. **General approach.** An analogy underlies recent work on growth in groups: much³ of basic group theory carries over when, instead of subgroups, we study sets that grow slowly $(|A \cdot A \cdot A| \leq |A|^{1+\varepsilon})$. This realisation is clearer in [Hel11] than in [Hel08], and has become current since then. (The term "approximate group" [Tao08] actually first arose in a different context, namely, the generalisation of some arguments in classical additive combinatorics to the non-abelian case. (See also [Hel08, §2.3], [SSV05, Lem. 4.2].) The analogy between subgroups and slowly growing sets was also explored in a model-theoretic setting in later work by Hrushovski [Hru12].)

This analogy is more important than whether one works with approximate subgroups in Helfgott's sense $(|A \cdot A \cdot A| \leq |A|^{1+\varepsilon})$, or more generally $|A \cdot A \cdot A| \leq f(|A|)$ for some specified f) or Tao's sense [Tao08, Def. 3.7]; the two definitions are essentially equivalent, and we will actually work with neither. We could phrase part of our argument in terms of statements of the form $|A^k| \leq |A|^{1+\varepsilon}$, but k would sometimes be larger than n; applying the tripling lemma ([RT85], [Hel08, Lem. 2.2], [Tao08, Lem. 3.4]) to such statements would weaken them fatally.

There is another issue worth emphasising: the study of growth needs to be relative. We should not think simply in terms of a group acting on itself by multiplication – even if, in the last analysis, this is the only operation available to us. Rather, growth statements often need to be thought of in terms of the action of a group G on a set X, and the effect of this action on subsets $A \subseteq G$, $B \subseteq X$. (Here X may or may not be endowed with a structure of its own.) This was already clear in [Hel11, Prop. 3.1] and [GH], and is crucial here: a key step will involve the action of a normaliser $N_G(H)$ on a subgroup $H \subseteq G$ by conjugation.

1.4. **Relation to previous work.** Our debt to previous work on permutation groups is manifold. It is worthwhile to point out that some of our main techniques are adaptations to sets of classification-free arguments⁴ on the properties of subgroups of Sym(n) by Babai [Bab82], Pyber [Pyb93], Bochert [Boc89], and Liebeck [Lie83]. Of particular importance is Babai and Pyber's work on the order of 2-transitive groups [Bab82], [Pyb93].

We shall also utilise existing diameter bounds. Besides Thm 1.1, we shall use the main idea from [BS88] (see Lemma 3.19) and the following theorem by Babai, Beals, and Seress. For a permutation g of a set Ω , the support supp(g) is the subset of elements of Ω that are displaced by g.

Theorem 1.4. ([BBS04]) For every $\varepsilon < 1/3$ there exists $K(\varepsilon)$ such that, if $G = \operatorname{Alt}(n)$ or $\operatorname{Sym}(n)$ and A is a set of generators of G containing an element $x \in A$

³Or at least results on subgroups that rely on *grosso modo* quantitative arguments. (Crucially, the orbit-stabilizer theorem carries over (Lem. 3.1); Sylow theory, which is quantitative but relies on (necessarily delicate) congruences, does not.) As [BBS04, Lem. 2.1] (in retrospect) and Prop. 5.2 in the present work make clear, probabilistic arguments in combinatorics can also carry over, provided that the desired probability distribution on a set can be approximated quickly by the action of a random walk.

⁴Cf. the role of [LP11] (esp. Thm. 4.2, Thm. 6.2), which, in order to provide alternatives to the Classification of Finite Simple Groups, did (both more and less generally) for subgroups what [Hell1, §5] did for sets, and was later translated back to sets for use in [BGT11].

with $1 < |\operatorname{supp}(x)| \le \varepsilon n$, then

$$\operatorname{diam}(\Gamma(G, A)) \le K(\varepsilon)n^8.$$

We will use this theorem repeatedly in §6. As we shall make clear in §4, we also apply – crucially – one of the main methods involved in the proof of Thm. 1.4, namely, the use of short random walks to mimic a uniform distribution.

We note that until recently Theorem 1.4 gave the largest known explicit class of Cayley graphs of Sym(n) or Alt(n) that has polynomially bounded diameter. In late 2010, partly based on ideas from [BBS04], Bamberg et al. [BGH⁺] proved that if a set of generators of Sym(n) or Alt(n) contains an element of support size at most 0.63n then the diameter of the Cayley graph is bounded by a polynomial of n.

1.5. **Outline.** Let us begin *in medias res*, focusing on a crucial moment at which growth is achieved. Classical reasons aside, this will allow us to emphasize the link to [Hel08], [Hel11], [BGT11], [PS] and [GH], while repeating one of the main motifs: growth results from the action of a group on a set, often, as is the case here, by conjugation.

The setup for the crucial step will involve a set $A \subset \operatorname{Sym}([n])$ with $A = A^{-1}$ and a fairly large set $\Sigma \subset [n]$ ($[n] := \{1, 2, ..., n\}$) such that the pointwise stabilizer⁵ $A_{(\Sigma)}$ generates a group $\langle A_{(\Sigma)} \rangle$ with a large orbit $\Gamma \subset [n] \setminus \Sigma$. (Say, for concreteness, that $|\Sigma| \geq (\log n)^2$ and $|\Gamma| > 0.95n$.) The setwise stabilizer $\langle A_{\Sigma} \rangle$ acts on the pointwise stabilizer $\langle A_{(\Sigma)} \rangle$ by conjugation.

We can assume that $\langle A_{(\Sigma)} \rangle$ acts as the alternating or symmetric group on Γ , as otherwise we are done by a different argument (called *descent* in §6; we will discuss it later). It follows that we can find a set S of at most six elements of $(A_{(\Sigma)})^{\ell}$, ℓ fairly small, such that $\langle S \rangle$ is doubly transitive on Γ . (This implication is far from trivial; we prove a general result of this kind (Cor. 4.7) showing that, if a set A' generates $\operatorname{Sym}([m])$ or $\operatorname{Alt}([m])$, then there is a small set $S \subset (A')^{\ell}$, ℓ fairly small, such that $\langle S \rangle$ is k-transitive.)

Consider the action of the elements of A_{Σ} on the elements of S by conjugation. By an orbit-stabilizer principle, either (a) an element $g \neq e$ of A_{Σ} fixes (i.e., commutes with) every element of S, or (b) the orbit $\{gsg^{-1}:g\in A_{\Sigma}\}$ of some $s\in S$ is of size $\geq |A_{\Sigma}|^{1/6}$. In case (a), since $\langle S \rangle$ is doubly transitive, g fixes every point of Γ . We have thus constructed a non-identity element $g\in A$ with small support, and are done by Thm. 1.4. In case (b), we have constructed many $(\geq |A_{\Sigma}|^{1/6})$ distinct elements gsg^{-1} in the pointwise stabilizer $(A^3)_{(\Sigma)}$. This is what we call *creation* in §6.

The questions are now – how do we get to the point at which we began our narrative? And how do we use the conclusion we have just shown, namely, the creation of many elements in the pointwise stabilizer?

Let us start with the first question. For the conclusion to be strong, A_{Σ} should be large – for instance, large in comparison to $A_{(\Sigma)}$ or $(A^2)_{(\Sigma)}$. Now, A_{Σ} can be much

⁵Defined as in (2.1). The notation here follows Dixon and Mortimer [DM96] and Seress [Ser03] rather than Wielandt [Wie64]. Wielandt writes A_{Σ} for the pointwise stabilizer, which we denote by $A_{(\Sigma)}$; we write A_{Σ} for the setwise stabilizer.

larger than $(A^2)_{(\Sigma)}$ only if A occupies a large number R of cosets of $\mathrm{Sym}([n])_{(\Sigma)}$ in $\mathrm{Sym}([n])$. (By pigeonhole, $|(A^2)_{(\Sigma)}| \geq |A|/R$.) Our aim will be to find a large Σ such that R is larger than $(dn)^{|\Sigma|}$, where d > 1/2 is a constant.

This is also an intermediate aim in [Pyb93] (which treats subgroups, not sets). Much as there, we use this as follows: R is larger than $(dn)^{|\Sigma|}$, and so AA^{-1} intersects at least $d^{|\Sigma|}|\Sigma|!$ cosets of $(\operatorname{Sym}([n]))_{(\Sigma)}$ within $(\operatorname{Sym}([n]))_{\Sigma}$ (by pigeonhole); this means that the projection (by restriction) of $(AA^{-1})_{\Sigma}$ to $\operatorname{Sym}(\Sigma)$ has size at least $d^{|\Sigma|}|\Sigma|!$. At this point Pyber uses the fact (due to Liebeck [Lie83] and based on Bochert [Boc89]) that, if a subgroup H of $\operatorname{Sym}(\Sigma)$ is of size at least $s = d^{|\Sigma|}|\Sigma|!$, where d > 1/2, then there must be a large orbit $\Delta \subset \Sigma$ of H such that the restriction of H to Δ equals $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$. We will show (Prop. 3.15) that, even if $H \subset \operatorname{Sym}(\Sigma)$ is just a set, not a subgroup, the assumption that H is of size at least s implies that the restriction of H^{ℓ} to Δ equals all of $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$, where ℓ is relatively small. (This works because the proof of Bochert's nineteenth-century result is algorithmic.) The fact that we obtain all of $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$ is particularly important for what we called a "descent argument" (as in "infinite descent") in the above.

Now, as we said, we must find a large Σ such that A (or $A^{\ell'}$, ℓ' moderate) occupies a large number of cosets of $\mathrm{Sym}([n])_{(\Sigma)}$, i.e., sends (Σ) to many different tuples. Pyber shows this (for A a subgroup) by constructing $\Sigma = \{\alpha_1, \alpha_2, \ldots, \alpha_m\}$ so that

$$(1.4) |\alpha_i^{A_{(\alpha_1,\dots,\alpha_{i-1})}}| \ge dn$$

for every $1 \leq i \leq m$. (The use of stabilizer chains $A > A_{(\alpha_1)} > A_{(\alpha_1,\alpha_2)} > \dots$ goes back to the algorithmic work of Sims [Sim70], [Sim71], as does the use of the size of the orbits in (1.4); see [Ser03, §4.1].) This step also works when A is a subset (Lemma 3.17). The difficult part, of course, is to show that elements $\alpha_1, \alpha_2, \dots, \alpha_m$ satisfying (1.4) exist.

Here [Pyb93] uses Babai's splitting lemma [Bab82], which states that, if $H < \operatorname{Sym}([n])$ is a doubly transitive permutation group and $\Sigma \subset [n]$ is such that $H_{(\Sigma)}$ has no orbits of size $> (1-\epsilon)n$, then there is a set $\Sigma' \subset [n]$ with $|\Sigma'| \ll_{\epsilon} (\log n) |\Sigma|$ such that $H_{(\Sigma')}$ consists only of the identity. In fact, $\Sigma' = \Sigma^S = \{x^S : x \in \Sigma, s \in S\}$, where S is a subset of H of size $|S| \ll \log n$. Babai constructs S by choosing $O(\log n)$ elements randomly from H with the uniform distribution. A random element of H takes a pair (x,y) of distinct elements of [n] to any other such pair (x',y') with the same probability $((n(n-1)/2)^{-1})$ no matter what (x',y') is. Now, given any distinct $x,y \in [n]$, it is almost certain that they will be taken to elements x^g , y^g of different orbits of $H_{(\Sigma)}$ by some $g \in S \subset H$, simply because a positive proportion of all pairs (x',y') lie in different orbits (by the fact that there is no orbit of size $> (1-\epsilon)n$). Then, x and y belong to different orbits of $gH_{\Sigma}g^{-1} = H_{\Sigma}g^{-1}$, and thus to different orbits of $H_{\Sigma}s$. Summing probabilities over all x and y, we obtain that, with positive probability, every two distinct $x,y \in [n]$ belong to different orbits of $H_{\Sigma}s$. This implies that $H_{\Sigma}s$ is trivial.

We adapt this entire argument so as to hold for a set $A \subset \text{Sym}([n])$ instead of a subgroup H < Sym([n]); as usual, sometimes H is replaced by A and sometimes by

 AA^{-1} or A^{ℓ} , where ℓ is moderate ($\ell \ll n^{O(1)}$). The key here is that the outcome of a random walk of moderate length takes a pair (x,y) to any other pair (x',y') with almost uniform probability.

We apply the resulting generalization of the splitting lemma (Prop. 5.2) and point out that $(AA^{-1})_{(\Sigma')} = \{e\}$ implies $|\Sigma'| \gg \log_n |A|$ (by pigeonhole) and so $|\Sigma| \gg (\log |A|)/(\log n)^2$. In other words, we are guaranteed to be able to construct a stabilizer chain with long orbits as in (1.4) (for any d < 1) until m gets to size proportional to $(\log |A|)/(\log n)^2$. We call this the *organizing* step.

Now that we have the stabilizer chain, and thus the proper setup for the creation step, how do we use the outcome of the creation step? In [Hel08] and the work that followed, the main intermediate result stated that a generating set A always grew in size $(|A^3| \ge |A|^{1+\delta}$ [Hel08, Key Proposition]); to prove that the diameter $\Gamma(G, A)$ was small, one just had to apply this key proposition over and over $(|A^3| \ge |A|^{1+\delta}, |A^9| \ge |A|^{1+\delta} \ge |A|^{(1+\delta)^2}, \ldots)$. Here we will also prove our diameter bound by iteration; however, the quantity whose growth we will keep track of during iteration will not be the size of A^{ℓ} , but rather the length of the sequence $\alpha_1, \alpha_2, \ldots$ we have constructed satisfying (1.4) (for A^{ℓ} instead of A).

The iteration is conducted as follows. We actually construct the first $(\log n)^2$ elements of $\alpha_1, \alpha_2, \ldots$ by brute force, by raising A to an $n^{O((\log n)^2)}$ th power. (This works by Lemma 3.9.) Now we get to the main step that gets repeated (Prop. 6.4): given a sequence $\alpha_1, \ldots, \alpha_m$ satisfying (1.4) (for A^{ℓ} instead of A), we use the creation step to construct at least $(m!)^{1/6}$ elements of $(A^{\ell'})_{(\alpha_1,\ldots,\alpha_m)}$, where $\ell' \leq n^{O(\log n)}\ell$; then we use the organizing step to construct new elements $\alpha_{m+1},\ldots,\alpha_{m'}$ $(m' \geq m + cm(\log m)/(\log n)^2)$ so that (1.4) is satisfied for all $i = 1, 2, \ldots, m'$ (with $A^{\ell'}$ instead of A^{ℓ}). (We actually repeat the organizing step several times after each creation step; this helps us save a log in the final exponent.) Repeating this, we keep on lengthening the sequence $\alpha_1, \alpha_2, \ldots$ until it gets to be of length almost n, and then we are done easily.

* * *

Needless to say, in the above outline, we have left out details that will be treated in full in the body of the text. Let us discuss one more thing now – namely, what we have called the *descent* step. We reach it when we have constructed a set $\Sigma = \{\alpha_1, \alpha_2, \dots, \alpha_m\}$ such that (a) the restriction of A_{Σ} to Σ acts as $Alt(\Delta)$ or $Sym(\Delta)$ on a large subset $\Delta \subset \Sigma$, (b) the restriction of $\langle A_{\Sigma} \rangle$ to $[n] \setminus \Sigma$ does not act like Alt or Sym on any subset of $[n] \setminus \Sigma$ larger than 0.95n (say).

Now we can use Thm. 1.1 (Babai-Seress), and obtain from (b) that the diameter of $\langle A_{\Sigma} \rangle$ is bounded in terms of the diameter of $\mathrm{Alt}(k)$, k = [0.95n]. (It is here, and only here, that the Classification Theorem is needed, since Thm. 1.1 is based on it.) Now we can use, inductively, our own main theorem on the diameter of $\mathrm{Alt}(n)$, with k instead of n. This gives a bound on the diameter of $\langle A_{\Sigma} \rangle$. At this point we use Lemma 3.19 (which is [BS87, Lemma 3]; see also [BLS87]). This shows that (a) implies that $\langle A_{\Sigma} \rangle$ contains a non-identity element g of small support. We can now apply Thm. 1.4 (Babai-Beals-Seress) to bound the diameter of our group $G = \mathrm{Alt}(n)$ or $G = \mathrm{Sym}(n)$ with respect to A. Note that [BS87, Lemma 3] would be

prohibitively expensive if used as a constructive result; here we are using it to show the *existence* of an element, which we know can be constructed as a relatively short word thanks to the bound on the diameter of $\langle A_{\Sigma} \rangle$ we obtained through Thm. 1.1.

1.6. Acknowledgements. We are deeply grateful to both Pablo Spiga and Nick Gill for stimulating discussions and for their constant help. Gordon Royle organised the first author's visit to Australia; if it were not for him, our collaboration might not have happened. Thanks are also due to László Babai, Martin Kassabov, Igor Pak, Peter Sarnak and Andrzej Żuk for their advice. Detailed comments by two anonymous referees have certainly helped improve the paper.

Ákos Seress was supported in part by the NSF and by ARC Grant DP1096525. Travel was supported in part by H. A. Helfgott's Philip Leverhulme prize. We benefited from the kind hospitality of the University of Western Australia and the École Normale Supérieure during our visits to each other's institutions.

2. Notation

We write $[n] = \{1, 2, ..., n\}$. For a set Ω , $\operatorname{Sym}(\Omega)$ and $\operatorname{Alt}(\Omega)$ are the symmetric and alternating groups acting on Ω . As is customary, we often write $\operatorname{Alt}(n)$ and $\operatorname{Sym}(n)$ for $\operatorname{Alt}([n])$ and $\operatorname{Sym}([n])$ - particularly when we are thinking of these groups as abstract groups as opposed to their actions.

We write $H \leq G$ to mean that H is a subgroup of G and $H \triangleleft G$ to mean that H is a normal subgroup. We say that a group S is a *section* of a group G if there exist subgroups H and K of G with $K \triangleleft H$ and $H/K \cong S$. We denote the identity element of a group by e.

Let A be a subset of a group G. We write $A^{-1} = \{a^{-1} : a \in A\}$, $A^k = \{a_1 a_2 \cdots a_k : a_1, \ldots, a_k \in A\}$. In [Hel08], [Hel11], the first author wrote A_ℓ to mean $(A \cup A^{-1} \cup \{e\})^\ell$; this does not seem to have become standard, and would also not do here due to the potential confusion with alternating groups. (Recall that A_n is in common usage as a synonym for Alt(n).) We will often include $A = A^{-1}$, $e \in A$ explicitly in our assumptions so as to simplify notation. A set A with $A = A^{-1}$ is said to be symmetric.

We write |A| for the number of elements of a set A. (All of our sets and groups are finite.) Given a group G and a subgroup $H \leq G$, we write [G:H] for the index of H in G.

Let a group G act on a set X. As is customary in the study of permutation groups, given $g \in G$ and $\alpha \in X$, we write α^g for the image of α under the action of g. We speak of the *orbit* $\alpha^A = \{\alpha^g : g \in A\}$ of a point α under the action of a set A of permutations. Our actions are right actions by default: $(\alpha^g)^h = \alpha^{gh}$. In consequence, we also use right cosets by default, i.e., cosets Hg (and so G/H is the set of all such cosets). Clearly |G/H| = [G:H].

We define the commutator [g,h] by $[g,h] = g^{-1}h^{-1}gh$. Again, this choice is customary for permutation groups.

Define

(2.1)
$$A_{\Sigma} = \{ g \in A : \Sigma^g = \Sigma \}, \qquad \text{(the setwise stabilizer)}$$
$$A_{(\Sigma)} = \{ g \in A : \forall \alpha \in \Sigma \, (\alpha^g = \alpha) \}. \qquad \text{(the pointwise stabilizer)}$$

If $\Sigma = \{g_1, \dots, g_m\}$, the setwise stabilizer is denoted by $A_{\{g_1, \dots, g_m\}}$ and the pointwise stabilizer by $A_{\{g_1, \dots, g_m\}}$.

Given a permutation $g \in \operatorname{Sym}(\Omega)$, we define its $\operatorname{support} \operatorname{supp}(g)$ to be the set of elements of Ω moved by g: $\operatorname{supp}(g) = \{\alpha \in \Omega : \alpha^g \neq \alpha\}$. If a subset $\Delta \subseteq \Omega$ is invariant under g, i.e., Δ is a union of cycles of g, then we define $g|_{\Delta} \in \operatorname{Sym}(\Delta)$ as the restriction (natural projection) of g to Δ : the permutation $g|_{\Delta}$ acts on Δ as g does. If Δ is invariant under some $D \subseteq \operatorname{Sym}(\Omega)$ then $D|_{\Delta} = \{g|_{\Delta} : g \in D\}$.

A partition $\mathcal{B} = \{\Omega_1, \Omega_2, \dots, \Omega_k\}$ of a set Ω (Ω_i non-empty) is called a *system* of imprimitivity for a transitive group $G \leq \operatorname{Sym}(\Omega)$ if G permutes the sets Ω_i for $1 \leq i \leq k$. For $|\Omega| \geq 2$, a transitive group $G \leq \operatorname{Sym}(\Omega)$ is called *primitive* if there are only the two trivial systems of imprimitivity for G: the partition into one-element sets, and the partition consisting of one part $\Omega_1 = \Omega$.

We say that a graph (or a multigraph) is regular with degree or valency d if there are d edges adjoining every vertex; that is, "degree" and "valency" of a vertex mean the same thing. In a directed graph, the out-degree of a vertex x is the number of edges starting at x while the in-degree is the number of edges terminating at x. A directed graph is called strongly connected if for any two vertices x, y, there is a directed path from x to y.

By $f(n) \ll g(n)$, $g(n) \gg f(n)$ and f(n) = O(g(n)) we mean one and the same thing, namely, that there are N > 0, C > 0 such that $|f(n)| \leq C \cdot g(n)$ for all $n \geq N$. We write $\log_2 x$ to mean the logarithm base 2 of x (and not to mean $\log \log x$).

3. Preliminaries on sets, groups and growth

3.1. Orbits and stabilizers. The orbit-stabilizer theorem from elementary group theory carries over to sets. This is a fact whose importance to the area is difficult to overemphasise. It underlies already [Hel08] at a key point (Prop. 4.1); the action at stake there is that of a group G on itself by conjugation.

The setting for the theorem is the action of a group G on a set X. The *stabilizer* G_x of a point $x \in X$ is the set $\{g \in G : x^g = x\}$.

Lemma 3.1 (Orbit-stabilizer theorem for sets). Let G be a group acting on a set X. Let $x \in X$, and let $A \subseteq G$ be non-empty. Then

$$(3.1) |AA^{-1} \cap G_x| \ge \frac{|A|}{|x^A|}.$$

Moreover, for every $B \subseteq G$,

$$(3.2) |AB| \ge |A \cap G_x||x^B|.$$

The usual orbit-stabilizer theorem is the special case A = B = H, H a subgroup of G.

Proof. By the pigeonhole principle, there exists an image $x' \in x^A$ such that the set $S = \{a \in A : x^a = x'\}$ has at least $|A|/|x^A|$ elements. For any $a, a' \in S$, $x^{a(a')^{-1}} = (x')^{(a')^{-1}} = x$. Hence

$$|AA^{-1} \cap G_x| \ge |SS^{-1}| \ge |S| \ge \frac{|A|}{|x^A|}.$$

Let $b_1, b_2, \ldots, b_\ell \in B$, $\ell = |x^B|$, be elements with $x^{b_i} \neq x^{b_j}$ for $i \neq j$. Consider all products of the form ab_i , $a \in A \cap G_x$, $1 \leq i \leq \ell$. If two such products ab_i , $a'b_{i'}$ are equal, then $x^{b_i} = x^{ab_i} = x^{a'b_{i'}} = x^{b'_i}$. This implies $b_i = b_{i'}$. Since $ab_i = a'b_{i'}$, we conclude that a = a'. We have thus shown that all products ab_i , $a \in A \cap G_x$, $1 \leq i \leq \ell$, are in fact distinct. Hence

$$|AB| \ge |(A \cap G_x) \cdot \{b_i : 1 \le i \le \ell\}|$$

= $|A \cap G_x| \cdot \ell = |A \cap G_x| \cdot |x^B|$.

As the following corollaries show, the relation between the size of A, on the one hand, and the size of orbits and stabilizers, on the other, implies that growth in the size of either orbits or stabilizers induces growth in the size of A itself.

Corollary 3.2. Let G be a group acting on a set X. Let $x \in X$. Let $A \subseteq G$ be a non-empty set with $A = A^{-1}$. Then, for any k > 0,

$$(3.3) |A^{k+1}| \ge \frac{|A^k \cap G_x|}{|A^2 \cap G_x|} |A|.$$

Proof. By (3.2),

$$|A^{k+1}| \ge |A^k \cap G_x| |x^A| \ge \frac{|A^k \cap G_x|}{|A^2 \cap G_x|} |A^2 \cap G_x| |x^A|.$$

Since $|A^2 \cap G_x||x^A| \ge |A|$ (by (3.1)), we obtain (3.3).

Corollary 3.3. Let G be a group acting on a set X. Let $x \in X$. Let $A \subseteq G$ be a non-empty set with $A = A^{-1}$. Then, for any k > 0,

$$(3.4) |A^{k+2}| \ge \frac{|x^{A^k}|}{|x^A|} |A|.$$

Proof. By (3.2) and (3.1),

$$|A^{k+2}| \ge |A^2 \cap G_x||x^{A^k}| \ge \frac{|A|}{|x^A|}|x^{A^k}| = \frac{|x^{A^k}|}{|x^A|}|A|.$$

3.2. Lemmas on subgroups and quotients. We start by recapitulating some of the simple material in [Hel11, §7.1]. The first lemma guarantees that we can always find many elements of AA^{-1} in any subgroup of small enough index.

Lemma 3.4 ([Hel11, Lem. 7.2]). Let G be a group and H a subgroup thereof. Let $A \subseteq G$ be a non-empty set. Then

$$(3.5) |AA^{-1} \cap H| \ge \frac{|A|}{r},$$

where r is the number of cosets of H intersecting A. In particular,

$$|AA^{-1} \cap H| \ge \frac{|A|}{[G:H]}.$$

Proof. By the orbit-stabilizer principle (3.1) applied to the natural action of G on G/H by multiplication on the right.⁶ (Set x = He = H.)

The following two lemmas should be read as follows: growth in a subgroup gives growth in the group; growth in a quotient gives growth in the group.

Lemma 3.5 (essentially [Hel11, Lem. 7.3]). Let G be a group and H a subgroup thereof. Let $A \subseteq G$ be a non-empty set with $A = A^{-1}$. Then, for any k > 0,

$$(3.6) |A^{k+1}| \ge \frac{|A^k \cap H|}{|A^2 \cap H|} |A|.$$

Proof. By Cor. 3.2 applied to the action of G on G/H by multiplication on the right (with x = He = H).

For a group G and a subgroup $H \leq G$, we define the coset map $\pi_{G/H}: G \to G/H$ that maps each $g \in G$ to the right coset Hg containing g.

Lemma 3.6 (essentially [Hel11, Lem. 7.4]). Let $A \subseteq G$ be a non-empty set with $A = A^{-1}$. Then, for any k > 0,

$$|A^{k+2}| \ge \frac{|\pi_{G/H}(A^k)|}{|\pi_{G/H}(A)|} |A|.$$

Proof. By Cor. 3.3, applied with G acting on X := G/H by multiplication on the right and with x := H seen as an element of G/H.

The following lemma is a generalisation of Lemma 3.4.

Lemma 3.7. Let G be a group, let H, K be subgroups of G with $H \leq K$, and let $A \subseteq G$ be a non-empty set. Then

$$|\pi_{K/H}(AA^{-1} \cap K)| \ge \frac{|\pi_{G/H}(A)|}{|\pi_{G/K}(A)|} \ge \frac{|\pi_{G/H}(A)|}{[G:K]}.$$

In other words: if A intersects r[G:H] cosets of H in G, then AA^{-1} intersects at least r[G:H]/[G:K] = r[K:H] cosets of H in K. (As usual, all our cosets are right cosets.)

⁶Recall that we are following the convention that G/H is the set of right cosets Hq.

Proof. Since A intersects $|\pi_{G/H}(A)|$ cosets of H in G and $|\pi_{G/K}(A)|$ cosets of K in G, and every coset of K in G is a disjoint union of cosets of H in G, the pigeonhole principle implies that there exists a coset Kg of K such that A intersects at least $k = |\pi_{G/H}(A)|/|\pi_{G/K}(A)|$ cosets $Ha \subseteq Kg$. Let a_1, \ldots, a_k be elements of A in distinct cosets of H in Kg. Then $a_ia_1^{-1} \in AA^{-1} \cap K$ for each $i = 1, \ldots, k$. Finally, note that $Ha_1a_1^{-1}, \ldots, Ha_ka_1^{-1}$ are k distinct cosets of H.

The above lemmas fall into two types: either (a) they reduce the problem of proving growth in G to that of proving growth in a smaller structure (a subgroup in Lemma 3.5, a quotient in Lemma 3.6), or (b) they produce many elements in a smaller structure (a group in Lemma 3.4, a quotient in Lemma 3.7.

Lastly, a result of a somewhat different nature. It is a version of Schreier's lemma (rewritten slightly as in [GH, Lem. 2.10]). Usually, if a set A generates a group G, that does not mean that, for H a subgroup of G, the intersection $A \cap H$ will generate H. However, Lemma 3.8 tells us, if A projects onto G/H, then $A^3 \cap H$ does generate H. We will use Lemma 3.8 in the proof of Lemma 6.2 (for G a setwise stabilizer $(\operatorname{Sym}(n))_{\Delta}$ and H the corresponding pointwise stabilizer $(\operatorname{Sym}(n))_{(\Delta)}$).

Lemma 3.8 (Schreier). Let G be a group and H a subgroup thereof. Let $A \subseteq G$ with $A = A^{-1}$ and $e \in A$. Suppose A intersects each coset of H in G. Then $A^3 \cap H$ generates $\langle A \rangle \cap H$. Moreover, $\langle A \rangle = \langle A^3 \cap H \rangle A$.

Proof. Let $C \subseteq A$ be a full set of right coset representatives of H, with $e \in C$. We wish to show that $\langle A \rangle = \langle A^3 \cap H \rangle C$. (This immediately implies both $\langle A \rangle = \langle A^3 \cap H \rangle A$ and $\langle A \rangle \cap H = \langle A^3 \cap H \rangle$.)

Clearly $e \in \langle A^3 \cap H \rangle C$. It is thus enough to show that, if g = hc, where $h \in \langle A^3 \cap H \rangle$ and $c \in C$, and $a' \in A$, then ga' still lies in $\langle A^3 \cap H \rangle C$. This is easily seen: since C is a full set of coset representatives, there is a $c' \in C$ with c' = h'ca' for some $h' \in H$, and thus

$$ga' = hca' = h((h')^{-1})h'ca' = h((h')^{-1})c' \in \langle A^3 \cap H \rangle (A^3 \cap H)C = \langle A^3 \cap H \rangle C,$$
 where we use the fact that $h' = c'(a')^{-1}c^{-1} \in A^3$.

3.3. Actions and generators. The proofs of the next two lemmas share a rather simple idea. Indeed, both lemmas can be seen as consequences of the well-known fact that every connected graph has a spanning tree.⁷ The graph would be the union of the permutation graphs (with X as the vertex set) induced by the elements of the set A.

We give two brief proofs without graphs.

Lemma 3.9. Let G be a group acting transitively on a finite set X. Let $A \subseteq G$ with $A = A^{-1}$, $e \in A$ and $G = \langle A \rangle$. Then, for any $x \in X$,

$$x^{A^{\ell}} = X,$$

where $\ell = |X| - 1$.

⁷We thank an anonymous referee for this comment.

Proof. Consider the orbits $\{x\} \subseteq x^A \subseteq x^{A^2} \subseteq \cdots$. Let ℓ' be the smallest integer with $x^{A^{\ell'+1}} = x^{A^{\ell'}}$. As $x^{A^{\ell'+2}} = (x^{A^{\ell'+1}})^A = (x^{A^{\ell'}})^A = x^{A^{\ell'+1}} = x^{A^{\ell'}}$, we have $x^{A^{\ell'}} = x^{\langle A \rangle} = x^G = X$. Since

$$\{x\} \subsetneq x^A \subsetneq x^{A^2} \subsetneq \dots \subsetneq x^{A^{\ell'}} = X,$$

we have $\ell' \leq |X| - 1$.

Lemma 3.10. Let G be a group acting transitively on a finite set X. Let $A \subseteq G$ with $A = A^{-1}$ and $G = \langle A \rangle$. Then there is a subset $A' \subseteq A$, |A'| < |X|, such that $\langle A' \rangle$ acts transitively on X.

Proof. Let $x \in X$. Let $A_1 = \{g\}$, where g is any element of A such that $x^g \neq x$. For each $i \geq 1$, let A_{i+1} be $A_i \cup \{g_i\}$, where g_i is an element of A such that $x^{\langle A_i \cup \{g_i\} \rangle} \supseteq x^{\langle A_i \rangle}$. If no such element g_i exists, we can conclude that $x^{\langle A_i \rangle}$ is taken to itself by every $g_i \in A$. This implies that $x^{\langle A_i \rangle}$ is taken to itself by every product of elements of A, and thus $(x^{\langle A_i \rangle})^{\langle A \rangle} = x^{\langle A \rangle}$ equals $x^{\langle A_i \rangle}$.

Hence, we have a chain

$$\{x\} \subsetneq x^{\langle A_1 \rangle} \subsetneq x^{\langle A_2 \rangle} \subsetneq \cdots \subsetneq x^{\langle A_i \rangle} = x^{\langle A \rangle} = X.$$

Clearly $i \leq |X| - 1$, and so $|A_i| \leq |X| - 1$. Let $A' = A_i$.

3.4. Large subsets of Sym(n). Let us first prove a result on large subgroups of Sym(n).

Lemma 3.11. Let $n \ge 84$. Let $G \le \operatorname{Sym}(n)$ be transitive, with a section isomorphic to $\operatorname{Alt}(k)$ for some k > n/2. Then G is either $\operatorname{Alt}(n)$ or $\operatorname{Sym}(n)$.

Proof. Since $k \geq 5$, the group Alt(k) is simple. Hence some composition factor of G has a section isomorphic to Alt(k). Assume that G is imprimitive and let \mathcal{B} be a non-trivial system of imprimitivity for G. Write $b = |\mathcal{B}|$ and m = n/b and let K be the kernel of the action of G on \mathcal{B} . Since G/K is isomorphic to a subgroup of Sym(b), K is isomorphic to a subgroup of $Sym(m)^b$ and b, m < k, we obtain that G has no section isomorphic to Alt(k), a contradiction. This shows that G is primitive.

From [PS80], we obtain that either $G \ge \operatorname{Alt}(n)$ or $|G| \le 4^n$. Since $|G| \ge |\operatorname{Alt}(k)| = k!/2 \ge \lceil n/2 \rceil!/2$, a direct computation shows that the latter case arises only for n < 84.

Our aim for the rest of this subsection will be to show that, if $A \subset \operatorname{Sym}(n)$ is very large, then $A^{n^{O(1)}}$ contains a copy of $\operatorname{Alt}(\Delta)$, $|\Delta| > n/2$. The next lemma generalizes Bochert's theorem [Boc89], [DM96, Thm. 3.3B] to subsets. Recall that, for $g \in \operatorname{Sym}(\Omega)$, we define the *support* of g by $\operatorname{supp}(g) = \{\alpha \in \Omega : \alpha^g \neq \alpha\}$.

Lemma 3.12. Let $n \geq 5$. Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$, $e \in A$. If $\langle A \rangle$ is a primitive permutation group and $|A| > n!/(\lfloor n/2 \rfloor!)$, then A^{n^4} is either $\operatorname{Alt}([n])$ or $\operatorname{Sym}([n])$.

This is an example of how one can sometimes modify a proof of a result about subgroups to give a result about sets: the proof follows the lines of Bochert's essentially algorithmic proof, plus some bookkeeping.

Proof. Given $A \subseteq \operatorname{Sym}([n])$ as in the statement of the lemma, let k be the smallest integer such that there exists $\Delta \subseteq [n]$ with $|\Delta| = k$ and $(A^2)_{(\Delta)} = \{e\}$. Let Δ be one such set.

Suppose that $k \leq n/2$. Then $\operatorname{Sym}([n])_{(\Delta)}$ has n!/(n-k)! < |A| cosets in $\operatorname{Sym}([n])$. Thus, by the pigeonhole principle, there exist two distinct elements a and b of A in the same coset. Hence $ab^{-1} \in \operatorname{Sym}([n])_{(\Delta)}$, that is, $ab^{-1} \in (A^2)_{(\Delta)}$. This contradicts the definition of k. We conclude that k > n/2.

The set $\Omega = [n] \setminus \Delta$ has cardinality less than k, so by definition there exists $g \in (A^2)_{(\Omega)}$ with $g \neq e$. Let $\delta \in \Delta$ with $\delta^g \neq \delta$. As the set $\Delta \setminus \{\delta\}$ has cardinality less then k, by the definition of k, there exists $h \in (A^2)_{(\Delta \setminus \{\delta\})}$ with $h \neq e$. Then $\operatorname{supp}(h) \subset \Omega \cup \{\delta\}$. Necessarily, $\delta \in \operatorname{supp}(h)$, otherwise $(A^2)_{(\Delta)}$ contains the non-identity element h. Hence $\operatorname{supp}(g) \cap \operatorname{supp}(h) = \{\delta\}$ and so the commutator x = [g, h] is a 3-cycle. Note that $[g, h] \in A^8$.

Now, since $\langle A \rangle$ is primitive and contains a 3-cycle, by Jordan's theorem [DM96, Thm. 3.3A] we obtain that $\langle A \rangle \geq \operatorname{Alt}([n])$. In particular, $\langle A \rangle$ is 3-transitive, and thus its action by conjugation on the set X of all 3-cycles is transitive. By Lemma 3.9,

$$x^{A^{\ell}} = X$$
.

where $\ell=|X|=n(n-1)(n-2)/3$ and A^ℓ acts on x by conjugation. Thus $A^{n(n-1)(n-2)/3}[q,h]A^{n(n-1)(n-2)/3}$

contains all 3-cycles in $\mathrm{Alt}([n])$.

Since any element of Alt([n]) can be written as a product of at most $\lfloor n/2 \rfloor$ 3-cycles, we obtain that A^{n^4-1} contains Alt([n]). Also, if A contains an odd permutation, then $A^{n^4} = Sym([n])$.

What happens, however, if $\langle A \rangle$ is not transitive, let alone primitive? We shall see first that, if A is large, then $\langle A \rangle$ must have at least a large orbit. In the following two lemmas, we use the inequalities

$$\left(\frac{n}{e}\right)^n < n! < 3\sqrt{n}\left(\frac{n}{e}\right)^n$$

Lemma 3.13. Let $H < \operatorname{Sym}(n)$ with $|H| \ge d^n n!$, for some number d with 0.5 < d < 1. If n is greater than a bound depending only on d, then H has an orbit of length at least dn.

Proof. Let $k := \lfloor dn \rfloor$. Suppose that the longest orbit length of H is less than dn. Then, as is well-known, $|H| \le k!(n-k)!$. (The size of a direct product of symmetric groups $\operatorname{Sym}(\Omega_i)$ only goes up if we pass elements from the smaller sets Ω_i , $i \ge 2$, to the largest set Ω_1 .)

Now, by (3.7), we have the following inequalities:

$$\left(\frac{k}{n}\right)^{n} \left(\frac{n}{e}\right)^{n} < \left(\frac{k}{n}\right)^{n} n! \le d^{n} n! \le |A| \le |\langle A \rangle| \le k! (n-k)!$$

$$< 9\sqrt{k(n-k)} \left(\frac{k}{e}\right)^{k} \left(\frac{n-k}{e}\right)^{n-k} \le \frac{9}{2} n \frac{k^{k} (n-k)^{n-k}}{e^{n}}.$$

Simplifying the left-hand side together with the right-hand side, we obtain $k^{n-k} < \frac{9}{2}n(n-k)^{n-k}$, that is, $\left(\frac{k}{n-k}\right)^{n-k} < \frac{9}{2}n$.

We define $c := \left(\frac{d}{1-d}\right)^{1-d}$. As

$$\lim_{n \to \infty} \left(\frac{k}{n-k} \right)^{\frac{n-k}{n}} = c > 1,$$

for large enough n, depending only on d, we have $\left(\frac{k}{n-k}\right)^{n-k} > \left(\frac{1+c}{2}\right)^n$. However, $\left(\frac{1+c}{2}\right)^n < \frac{9}{2}n$ is false if n is greater than a bound depending only on d, proving our claim.

Using Bochert's theorem [Boc89], Liebeck derived a result ([Lie83, Lem. 1.1]; see [Jor70, pp. 68–75] for a classical result of the same kind) on large subgroups of $\operatorname{Sym}(n)$. It does not assume transitivity or primitivity. We will generalize it to sets (Prop. 3.15). In a somewhat strengthened version [DM96, Thm. 5.2B], the result from [Lie83] states the following, among other things: if H is a subgroup of $\operatorname{Sym}(n)$, $n \geq 9$, and

(3.9)
$$[\operatorname{Sym}(n): H] < \min\left(\frac{1}{2} \binom{n}{\lfloor n/2 \rfloor}, \binom{n}{m}\right)$$

for some $m \geq n/2$, then there is a set $\Delta \subset [n], |\Delta| > m$, such that

(3.10)
$$\operatorname{Alt}(n)_{([n]\setminus\Delta)} \le H \le \operatorname{Sym}(n)_{[n]\setminus\Delta}.$$

Here, of course, $\operatorname{Alt}(n)_{([n]\setminus\Delta)} \sim \operatorname{Alt}(\Delta)$ and $\operatorname{Sym}(n)_{[n]\setminus\Delta} = \operatorname{Sym}(n)_{\Delta}$; in particular, (3.10) implies that Δ is an orbit of [n]. It is easy to see that, if $|H| \geq d^n n!$, 0.5 < d < 1, then (3.9) is fulfilled for $m = \lceil dn \rceil$, provided that n is larger than a constant depending only on d: by Stirling's formula,

$$(3.11) \qquad \binom{n}{\lceil dn \rceil} \gg \frac{1}{\sqrt{n}} \frac{n^n}{\lceil dn \rceil^{\lceil dn \rceil} | (1-d)n |^{\lfloor (1-d)n \rfloor}} \gg \frac{1}{n^{3/2}} \left(\frac{1}{d^d (1-d)^{1-d}} \right)^n,$$

and, since $d^d(1-d)^{1-d} < d$ for $d \in (1/2,1)$, this is certainly greater than $(1/d)^n$ for n large enough. The inequality $\frac{1}{2}\binom{n}{[n/2]} \gg 2^n/\sqrt{n}$ implies $\frac{1}{2}\binom{n}{[n/2]} > (1/d)^n$ immediately for all large n. Thus (3.10) holds for some Δ with $|\Delta| > dn$.

We will show an analogue of (3.10) holds for a set A instead of a subgroup H (Prop. 3.15). This can be shown in two ways: we can use Liebeck's result (3.10) for groups, or we can give an elementary proof using only counting arguments. (Both [Lie83] and [DM96] do a detailed examination of the subgroup structure of $\operatorname{Sym}(n)$ in order to give a result valid for small n.)

Let us first give an elementary proof of a somewhat weaker statement.

Lemma 3.14. Let d be a number with 0.5 < d < 1. If $A \subseteq \operatorname{Sym}([n])$ (with $A = A^{-1}$) has cardinality $|A| \ge d^n n!$ and n is larger than a bound depending only on d, then there exists an orbit $\Delta \subseteq [n]$ of $\langle A \rangle$ such that $|\Delta| \ge dn$ and $(A^{n^4})|_{\Delta}$ is $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$.

Proof. By Lemma 3.13, for large enough n the group $\langle A \rangle$ has an orbit Δ of length $k \geq dn$. Write $\rho = k/n$ and note that $d \leq \rho \leq 1$. The group $G = B|_{\Delta}$ has order at least $d^n n!/(n-k)!$, so estimating k!(n-k)! from above as in (3.8) and estimating n! from below by (3.7), we obtain

$$[\operatorname{Sym}(\Delta):G] \le \frac{k!(n-k)!}{d^n n!} < \frac{9}{4} n \frac{k^k (n-k)^{n-k}}{d^n n^n} =$$

$$(3.12) \qquad \frac{9}{4} n \left(\frac{\rho^{\rho} (1-\rho)^{1-\rho}}{d}\right)^n = \frac{9}{4} n \left(2^{\frac{1}{\rho}} \rho (1-\rho)^{\frac{1-\rho}{\rho}}\right)^{\rho n} \left(\frac{1}{2d}\right)^n.$$

Next, we show that for large values of n the transitive group G cannot be imprimitive. Indeed, if G is imprimitive, then using (3.7) we have

$$(3.13) \qquad [\operatorname{Sym}(\Delta): G] \ge \frac{1}{2} \binom{k}{\lfloor k/2 \rfloor} > \frac{1}{2} \frac{(\frac{k}{e})^k}{9k(\frac{k}{2e})^k} > \frac{1}{18n} 2^{\rho n}.$$

A direct computation shows that the function $f(\rho) = 2^{1/\rho} \rho (1-\rho)^{(1-\rho)/\rho}$ is monotone increasing in the interval [1/2,1) with supremum 2. Hence, comparing the upper and lower bounds for $[\operatorname{Sym}(\Delta):G]$ deduced in (3.12) and (3.13), we obtain

(3.14)
$$\frac{9}{4}n2^{\rho n}\left(\frac{1}{2d}\right)^n > \frac{1}{18n}2^{\rho n}.$$

As d>1/2, for large enough n we have $(2d)^n>(18n)(\frac{9}{4}n)$ and therefore (3.14) cannot hold.

Hence G is primitive and $A|_{\Delta}$ is a set of size at least $d^n n!/(n-k)! \geq d^n k! > k!/(\lfloor k/2 \rfloor)!$ (where the last inequality holds for n greater than a lower bound depending only on d). Therefore, by Lemma 3.12, $(A|_{\Delta})^{n^4}$ is either $Alt(\Delta)$ or $Sym(\Delta)$, and hence so is $(A^{n^4})|_{\Delta} = (A|_{\Delta})^{n^4}$.

Now we get the full analogue of (3.10).

Proposition 3.15. Let d be a number with 0.5 < d < 1. Let $A \subseteq \operatorname{Sym}(n)$ with $A = A^{-1}$ and $e \in A$. If $|A| \ge d^n n!$ and n is larger than a bound depending only on d, then there exists an orbit $\Delta \subseteq [n]$ of $\langle A \rangle$ such that $|\Delta| \ge dn$ and $(A^{8n^5})_{([n]\setminus\Delta)}|_{\Delta}$ contains $\operatorname{Alt}(\Delta)$.

Proof. By Lemma 3.14, there is an orbit Δ of $\langle A \rangle$ such that $|\Delta| \geq dn$ and $(A^{n^4})|_{\Delta}$ is $\mathrm{Alt}(\Delta)$ or $\mathrm{Sym}(\Delta)$. Let $A' = A^{n^4}$.

It is clear that $|A'| \ge |\operatorname{Alt}(\Delta)| > |\operatorname{Sym}([n] \setminus \Delta)|$. Thus, by the pigeonhole principle, there are $h_1, h_2 \in A'$, $h_1 \ne h_2$, such that $h_1|_{[n] \setminus \Delta} = h_2|_{[n] \setminus \Delta}$, and so $g = h_1 h_2^{-1}$ fixes $[n] \setminus \Delta$ pointwise.

We show that $((A')^{14})_{([n]\setminus\Delta)}$ contains an element g' such that $g'|_{\Delta}$ is a 3-cycle. If $g|_{\Delta}$ has at least two fixed points then there exists an element $h \in A'$ such that $h|_{\Delta}$ is a 3-cycle, with $\operatorname{supp}(h|_{\Delta})$ intersecting $\operatorname{supp}(g|_{\Delta})$ in exactly one point. Then $g' = [g, h] \in (A')^{2+1+2+1} = (A')^6$ fixes $[n] \setminus \Delta$ pointwise and $g'|_{\Delta}$ is a 3-cycle. If g contains a cycle $(\alpha\beta\gamma\delta\dots)$ of length at least 4, then we choose an element $h \in A'$ with $h|_{\Delta} = (\alpha\beta\gamma)$ and let $g' = [g, h] \in (A')^6$. Then g' fixes $[n] \setminus \Delta$ pointwise and $g'|_{\Delta}$ is the 3-cycle $(\alpha\beta\delta)$.

In all other cases, $|\operatorname{supp}(g|_{\Delta})| \ge |\Delta| - 1 \ge 6$ (assuming $n \ge 13$, which implies $|\Delta| \ge 7$) and all nontrivial cycles of g have length 2 or 3. Hence $g|_{\Delta}$ contains at least two 3-cycles, or at least two 2-cycles.

If $g|_{\Delta}$ contains the cycles $(\alpha\beta\gamma)$ and $(\delta\eta\nu)$ then we choose an element $h \in A'$ with $h|_{\Delta} = (\alpha\eta)(\beta\delta\gamma\nu)$. A little computation shows that g' = [g,h] fixes $[n] \setminus \Delta$ pointwise and $g'|_{\Delta}$ is the 3-cycle $(\delta\eta\nu)$.

Finally, suppose g contains the 2-cycles $(\alpha\beta)$ and $(\gamma\delta)$. We choose again an element $h \in A'$ with $h|_{\Delta} = (\alpha\beta\gamma)$; then $\operatorname{supp}([g,h]) = \{\alpha,\beta,\gamma,\delta\}$ and [g,h] fixes $[n] \setminus \Delta$ pointwise. Since $[g,h] \in (A')^6$ also fixes at least two points of Δ , we deduce as in the very first case of our analysis that the commutator g' = [[g,h],h'] with an appropriate $h' \in A'$ is a 3-cycle. Note that $g' \in (A')^{6+1+6+1} = (A')^{14}$.

Given any 3-cycle s in $\operatorname{Sym}(\Delta)$, we can conjugate g' by an appropriate element of A' to get an element of $((A')^{16})_{([n]\setminus\Delta)}$ whose restriction to Δ equals s. Now, every element of $\operatorname{Alt}(\Delta)$ is the product of at most $\lfloor |\Delta|/2 \rfloor$ 3-cycles. Hence $((A')^{16\lfloor n/2 \rfloor})_{(|n|\setminus\Delta)}|_{\Delta}$ contains $\operatorname{Alt}(\Delta)$.

An anonymous referee kindly provides the following argument, showing that Prop. 3.15, which is a generalization of (3.10), can be proven using (3.10).

Second proof of Prop. 3.15. (This proof gives Prop. 3.15 with $A^{2(n^4+1)n^4}$ instead of A^{8n^5} .) By (3.10) applied to $H = \langle A \rangle$, there is a set Δ with $|\Delta| > dn$ such that (a) H is contained in $\operatorname{Sym}(n)_{\Delta}$ and (b) H contains the subgroup $D = \operatorname{Alt}(n)_{([n]\setminus\Delta)}$, i.e., $H|_{\Delta}$ contains $\operatorname{Alt}(\Delta)$. Let $B = A^2 \cap D$. By Lemma 3.4,

$$|B| = |A^2 \cap D| \ge \frac{|A|}{[\operatorname{Sym}(n)_{\Delta} : D]} \ge \frac{d^n n!}{2(n - |\Delta|)!} > \frac{d^n |\Delta|!}{2} \ge \frac{|\Delta|!}{2^{n+1}} \ge \frac{|\Delta|!}{2^{2|\Delta|+1}}.$$

For n sufficiently large (and hence $|\Delta|$ sufficiently large), $2^{2|\Delta|+1} < \lfloor |\Delta|/2 \rfloor!$, and so we obtain that $|B| > |\Delta|!/|\Delta|/2|!$.

Since $\langle A|_{\Delta}\rangle=H|_{\Delta}$ contains $\mathrm{Alt}(\Delta),\ \langle (A\cup B)|_{\Delta}\rangle$ is $\mathrm{Alt}(\Delta)$ or $\mathrm{Sym}(\Delta)$ – and, in particular, it is primitive. Hence, a first application of Lemma 3.12 (with Δ instead of [n]) implies that $((A\cup B)|_{\Delta})^{n^4}$ is $\mathrm{Alt}(\Delta)$ or $\mathrm{Sym}(\Delta)$.

The set $S = \{gbg^{-1} : g \in (A \cup B)^{n^4}, b \in B\}$ is in D; moreover, $\langle S|_{\Delta}\rangle$ is normal in $Alt(\Delta)$. Since S_{Δ} is non-trivial (by |B| > 1), we conclude that $\langle S|_{\Delta}\rangle = Alt(\Delta)$. Now we apply Lemma 3.12 (again with Δ instead of [n]) and obtain that $(S|_{\Delta})^{n^4} = Alt(\Delta)$. Since $S^{n^4} \subset A^{(2n^4+2)n^4}$, we are done.

3.5. Bases and stabilizer chains. Given a permutation group G on a set Ω , a subset Σ of Ω is called a *base* if $G_{(\Sigma)} = \{e\}$. This definition goes back to Sims [Sim70]. If, instead of G, we consider a subset A of $\operatorname{Sym}(\Omega)$, then, as the following lemma suggests, it makes sense to see whether $(AA^{-1})_{(\Sigma)}$ (rather than $A_{(\Sigma)}$) equals $\{e\}$.

Lemma 3.16. Let $A \subseteq \operatorname{Sym}(\Omega)$, $|\Omega| = n$. If $\Sigma \subseteq \Omega$ is such that $(AA^{-1})_{(\Sigma)} = \{e\}$, then $|\Sigma| \ge \log_n |A|$.

Proof. Notice first that $[\operatorname{Sym}(\Omega):(\operatorname{Sym}(\Omega))_{(\Sigma)}] \leq n^{|\Sigma|}$. By the pigeonhole principle, if $|A| > n^{|\Sigma|}$, then there exists a right coset of $(\operatorname{Sym}(\Omega))_{(\Sigma)}$ containing more than one element of A, and thus

$$|(AA^{-1})_{(\Sigma)}| = |AA^{-1} \cap (\operatorname{Sym}(\Omega))_{(\Sigma)}| > 1.$$

Hence, if
$$(AA^{-1})_{(\Sigma)} = \{e\}$$
, then we have $|A| \leq n^{|\Sigma|}$, i.e., $|\Sigma| \geq \log_n |A|$.

The use of stabilizer chains $H > H_{\alpha_1} > H_{(\alpha_1,\alpha_2)} > \cdots$ is very common in computational group theory (starting, again, with the work of Sims; see references in [Ser03, §4.1]). We may study a similar chain $A > A_{\alpha_1} > A_{(\alpha_1,\alpha_2)} > \cdots$ when A is merely a set.

Lemma 3.17. Let $\Sigma = \{\alpha_1, \ldots, \alpha_m\} \subseteq [n]$ and $A \subseteq \operatorname{Sym}([n])$. Suppose that

$$\left| \alpha_i^{A_{(\alpha_1, \dots, \alpha_{i-1})}} \right| \ge r_i$$

for all i = 1, 2, ..., m. Then A^m intersects at least $\prod_{i=1}^m r_i$ cosets of $\operatorname{Sym}([n])_{(\Sigma)}$.

Proof. For each $1 \leq i \leq m$, write $\Delta_i = \alpha_i^{A_{(\alpha_1, \dots, \alpha_{i-1})}}$; thus $|\Delta_i| \geq r_i$. For each $\delta \in \Delta_i$, pick $g_\delta \in A_{(\alpha_1, \dots, \alpha_{i-1})}$ with $\alpha_i^{g_\delta} = \delta$ and write $S_i = \{g_\delta : \delta \in \Delta_i\}$. Clearly, $|S_i| = |\Delta_i|$ and $S_i \subseteq A$. We show that for every two distinct tuples

$$(s_1, s_2, \dots, s_m), (s'_1, s'_2, \dots, s'_m) \in S_1 \times \dots \times S_m$$

the products $P = s_m s_{m-1} \cdots s_1$ and $P' = s'_m s'_{m-1} \cdots s'_1$ belong to two distinct cosets of $\operatorname{Sym}([n])_{(\Sigma)}$. From this it follows that A^m intersects at least $|S_1| \cdots |S_m| = |\Delta_1| \cdots |\Delta_m| \geq \prod_{i=1}^m r_i$ cosets of $\operatorname{Sym}([n])_{(\Sigma)}$.

We argue by contradiction, that is, we assume that P and P' map $(\alpha_1, \ldots, \alpha_m)$ to the same m-tuple. Let j be the smallest index such that $s_j \neq s'_j$. Then $Q = Ps_1^{-1} \cdots s_{j-1}^{-1}$ and $Q' = P's_1^{-1} \cdots s_{j-1}^{-1} = P's_1'^{-1} \cdots s_{j-1}'^{-1}$ also map $(\alpha_1, \ldots, \alpha_m)$ to the same m-tuple. Note that for all $k \leq m$, s_k and s'_k fix $(\alpha_1, \ldots, \alpha_{k-1})$ pointwise. Thus

$$\alpha_j^Q = \alpha_j^{s_j} \neq \alpha_j^{s_j'} = \alpha_j^{Q'},$$

contradicting our assumption.

We thus see that, if we choose $\alpha_1, \alpha_2, \ldots$ so that the orbits $\alpha_i^{A_{(\alpha_1, \ldots, \alpha_{i-1})}}$ are large, we get to occupy many cosets of $(\operatorname{Sym}([n]))_{(\Sigma)}$. By Lemma 3.7, this will enable us to occupy many cosets of $(\operatorname{Sym}([n]))_{(\Sigma)}$ in the setwise stabilizer $(\operatorname{Sym}([n]))_{\Sigma}$. We will then be able to apply Prop. 3.15 to build a large alternating group within $\operatorname{Sym}(\Sigma) \cong (\operatorname{Sym}([n]))_{\Sigma}/(\operatorname{Sym}([n]))_{(\Sigma)}$. This procedure is already implicit in [Pyb93, Lem. 3]; indeed, what amounts to this is signalled by Pyber as the main new element in his refinement [Pyb93, Thm. A] of Babai's theorem on the order of doubly transitive groups [Bab82]. The main difference is that we have to work, of course, with sets rather than groups; we also obtain a somewhat stronger conclusion due to our using Prop. 3.15 rather than invoking Liebeck's lemma directly.

Lemma 3.18. Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$ and $e \in A$. Let $\Sigma = \{\alpha_1, \ldots, \alpha_m\} \subseteq [n]$ be such that

$$\left|\alpha_{i}^{A_{(\alpha_{1},...,\alpha_{i-1})}}\right| \geq dn$$

for all i = 1, 2, ..., m, where d > 0.5. Then, provided that m is larger than a bound C(d) depending only on d, there exists $\Delta \subseteq \Sigma$ with $|\Delta| \ge d|\Sigma|$ and

$$Alt(\Delta) \subseteq ((A^{16m^6})_{\Sigma})_{(\Sigma \setminus \Delta)}|_{\Delta}.$$

Proof. By (3.15) and Lemma 3.17, A^m intersects at least $(dn)^m$ cosets of $\operatorname{Sym}([n])_{(\Sigma)}$ in $\operatorname{Sym}([n])$. Since

$$[\operatorname{Sym}([n]) : \operatorname{Sym}([n])_{\Sigma}] = \frac{[\operatorname{Sym}([n]) : \operatorname{Sym}([n])_{(\Sigma)}]}{[\operatorname{Sym}([n])_{\Sigma} : \operatorname{Sym}([n])_{(\Sigma)}]} \le \frac{n^m}{m!},$$

Lemma 3.7 implies (with $G = \operatorname{Sym}([n])$, $K = \operatorname{Sym}([n])_{\Sigma}$, $H = \operatorname{Sym}([n])_{(\Sigma)}$, and A^m instead of A) that

$$|\pi_{K/H}(A^{2m} \cap K)| \ge \frac{|\pi_{G/H}(A^m)|}{n^m/m!} \ge \frac{(dn)^m}{n^m/m!} = d^m m!.$$

Note that $|\pi_{K/H}(A^{2m} \cap K)| = |(A^{2m})_{\Sigma}|_{\Sigma}|$. We can thus apply Prop. 3.15 (with m instead of n, and $A' = (A^{2m})_{\Sigma}|_{\Sigma}$ instead of A) and obtain that there is a set $\Delta \subseteq \Sigma$ such that $|\Delta| \ge dm$ and $((A')^{8m^5})_{(\Sigma \setminus \Delta)}|_{\Delta}$ contains $Alt(\Delta)$.

3.6. Existence of elements of small support. The following lemma is essentially [BS87, Lemma 3] (or [BS88, Lemma 1]; see also [BLS87]).

Lemma 3.19. Let $\Delta \subseteq [n]$, $|\Delta| \ge c(\log n)^2$, c > 0. Let $H \le (\operatorname{Sym}(n))_{\Delta}$. Assume $H|_{\Delta}$ is $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$.

Let Γ be any orbit of H. Then, if n is larger than a bound depending only on c, H contains an element g with $g|_{\Delta} \neq 1$ and $|\operatorname{supp}(g|_{\Gamma})| < |\Gamma|/4$.

Proof. Let $p_1 = 2$, $p_2 = 3, \ldots, p_k$ be the sequence of the first k primes, where k is the least integer such that $p_1p_2\cdots p_k > n^4$. Much as in [BS87], we remark that, by elementary bounds towards the prime number theorem,

$$(3.16) 2p_1 + p_2 + \dots + p_k < c(\log n)^2,$$

provided that n be larger than a bound depending only on c. Thus H contains an element h such that $h|_{\Delta}$ consists of $|\Delta| - (2p_1 + p_2 + \cdots + p_k)$ fixed points and cycles of length $p_1, p_1, p_2, p_3, \ldots, p_k$. (We need two cycles of length $p_1 = 2$ because we want an even permutation on Δ .)

We can now reason as in [BS87, Lemma 3] or [BS88, Lemma 1]. For every $\gamma \in \Gamma$, denote by κ_{γ} the length (possibly 1) of the cycle of h containing γ and for $i \leq k$ define $\Gamma_i := \{ \gamma \in \Gamma : p_i \mid \kappa_{\gamma} \}$. Then

(3.17)
$$\sum_{\gamma \in \Gamma} \sum_{p_i \mid \kappa_{\gamma}} \log p_i < |\Gamma| \log n$$

because $\kappa_{\gamma} < n$ implies that for all γ the inner sum is less than $\log n$. Exchanging the order of summation,

$$\sum_{\gamma \in \Gamma} \sum_{p_i \mid \kappa_{\gamma}} \log p_i = \sum_{i=1}^k |\Gamma_i| \log p_i.$$

If $|\Gamma_i| \geq |\Gamma|/4$ for all $i \leq k$ then

$$\sum_{i=1}^{k} |\Gamma_i| \log p_i \ge \frac{|\Gamma|}{4} \log \left(\prod_{i=1}^{k} p_i \right) > \frac{|\Gamma|}{4} \log(n^4) = |\Gamma| \log n,$$

contradicting (3.17). Hence there is a prime $p \leq p_k$ such that $p|\kappa_{\gamma}$ for fewer than $|\Gamma|/4$ elements γ of Γ . Denoting the order of h by |h|, we define $g = h^{\ell}$ for $\ell := |h|/p$. We obtain that $|\sup(g|_{\Gamma})| < |\Gamma|/4$. We also have that g is non-trivial, since $g|_{\Delta}$ contains a p-cycle. Clearly $g \in H$, and so we are done.⁸

4. RANDOM WALKS AND GENERATION

4.1. Random walks. The aim of this subsection is to present some basic material on random walks. As stated in the outline, our later use of random walks to mimic the uniform distribution in combinatorial arguments is clearly influenced by [BBS04]; indeed, this subsection is very close to the first two thirds of [BBS04, §2].

Let Γ be a strongly connected directed multigraph with vertex set $V = V(\Gamma)$. For $x \in V(\Gamma)$, we denote by $\Gamma(x)$ the multiset of endpoints of the edges starting at x (counted with multiplicities in case of multiple edges). We are interested in the special case when Γ is regular of valency d (i.e., $|\Gamma(x)| = d$, for each $x \in V(\Gamma)$) and Γ is also symmetric in the sense that for all vertices $x, y \in V(\Gamma)$, the number of edges connecting x to y is the same as the number of edges connecting y to x. These two conditions imply that the adjacency matrix A of Γ is symmetric and all row and column sums are equal to d.

A lazy random walk on Γ is a stochastic process where a particle moves from vertex to vertex; if the particle is at vertex x such that $\Gamma(x) = \{y_1, \ldots, y_d\}$, then the particle

- stays at x with probability $\frac{1}{2}$;
- moves to vertex y_i with probability $\frac{1}{2d}$, for all $i = 1, \ldots, d$.

Here we are concerned with the asymptotic rate of convergence for the probability distribution of a particle in a lazy random walk on Γ . For $x,y \in V(\Gamma)$, write $p_k(x,y)$ for the probability that the particle is at vertex y after k steps of a lazy random walk starting at x. For a fixed $\varepsilon > 0$, the ℓ_{∞} -mixing time for ε is the minimum value of k such that

$$\frac{1}{|V(\Gamma)|}(1-\varepsilon) \le p_k(x,y) \le \frac{1}{|V(\Gamma)|}(1+\varepsilon)$$

for all $x, y \in V(\Gamma)$.

⁸Since we need only the existence of g for the moment, we are not concerned by the fact that l is very large. Compare this to the situation in [BS88], where the use of a large l causes diameter bounds much weaker than those in the present paper.

We can give a crude (and well-known; see, e.g., [BBS04, Fact 2.1]) upper bound on the ℓ_{∞} mixing time for regular symmetric multigraphs in terms of $N = |V(\Gamma)|$, ε and the valency d alone.

Lemma 4.1. Let Γ be a connected, regular and symmetric multigraph of valency d and with N vertices. Then the ℓ_{∞} mixing time for ε is at most $N^2 d \log(N/\varepsilon)$.

Proof. Let A be the adjacency matrix of Γ . Since A is symmetric, the eigenvalues of A are real; moreover, their modulus is clearly no more than d in magnitude. Let

$$d = \mu_1 \ge \mu_2 \ge \cdots \ge \mu_N \ge -d$$

be the eigenvalues of A and write P = I/2 + A/2d, where I is the $N \times N$ -identity matrix. The matrix P is the probability transition matrix for the Markov process described by a lazy random walk on Γ .

The sum of every row or column of P is 1, i.e., P is a doubly stochastic matrix. The eigenvalues of P are

$$1 = \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_N \ge 0$$

with $\lambda_i = 1/2 + \mu_i/2d$ for each i = 1, ..., N. It is well-known that the asymptotic rate of convergence to the uniform distribution of a lazy random walk is determined by λ_2 : since P is symmetric, there is a basis of \mathbb{R}^N consisting of orthogonal eigenvectors $v_1, v_2, ..., v_n$ of P with eigenvalues $\lambda_1, ..., \lambda_N$, where every eigenvector v_i has ℓ_2 -norm 1 with respect to (say) the counting measure; writing e_x for the probability distribution having value 1 at x and 0 elsewhere, we see that (by Cauchy-Schwarz and Plancherel)

$$\sum_{j=1}^{N} |\langle e_x, v_j \rangle| \, |\langle v_j, e_y \rangle| \le \sqrt{\sum_{j=1}^{N} |\langle e_x, v_j \rangle|^2} \sqrt{\sum_{j=1}^{N} |\langle v_j, e_y \rangle|^2} \le |e_x|_2 \cdot |e_y|_2 = 1,$$

and, since

$$p_k(x,y) = \langle P^k e_x, e_y \rangle = \langle \sum_{j=1}^N \langle e_x, v_j \rangle \cdot P^k v_j, e_y \rangle = \sum_{j=1}^N \langle e_x, v_j \rangle \cdot \lambda_j^k \langle v_j, e_y \rangle$$
$$= \frac{1}{\sqrt{N}} \cdot 1^k \cdot \frac{1}{\sqrt{N}} + \sum_{j=2}^N \langle e_x, v_j \rangle \cdot \lambda_j^k \langle v_j, e_y \rangle,$$

we see that

$$\left| p_k(x,y) - \frac{1}{N} \right| \le \lambda_2^k \sum_{j=2}^N |\langle e_x, v_j \rangle| \, |\langle v_j, e_y \rangle| \le \lambda_2^k.$$

By [Fie72, Lemma 2.4 and Theorem 3.4], we have

$$\lambda_2 \le 1 - 2(1 - \cos(\pi/N))\mu(P),$$

where $\mu(P) = \min_{\emptyset \neq M \subseteq V} \sum_{i \in M, j \notin M} p_{ij}$. As Γ is a connected regular graph of valency d, we have $\mu(P) \geq 1/2d$. Using the Taylor series for $\cos(x)$, we see that $(1-\cos(\pi/N)) \geq 1/N^2$. Thence $|p_k(x,y)-1/N| \leq (1-1/(N^2d))^k$. Since $1-x \leq e^{-x}$ for all x, we obtain $|p_k(x,y)-1/N| \leq \varepsilon/N$ for $k \geq N^2d\log(N/\varepsilon)$, as desired. \square

We will generally study regular symmetric multigraphs of the following type. (The following argument is already present in [BBS04, §2]; indeed, the only difference between Lemma 4.2 here and corresponding material in [BBS04, §2] is that Lemma 4.2 applies to ordered as opposed to unordered k-tuples.) Let G be a group and A be a subset of G with $A = A^{-1}$ and $e \in A$. Let G act on a set X. We take the elements of X as the vertices of our multigraph, and draw one edge from $x \in X$ to $x' \in X$ for every $a \in A$ such that $x^a = x'$. A walk on the graph then corresponds to the action of an element of A^{ℓ} on an element x of X, where ℓ is the length of the walk and x is the starting point of the walk.

Lemma 4.1 then gives us a lower bound on how large ℓ has to be for the action of A^{ℓ} on X to have a rather strong randomising effect.

Lemma 4.2. Let H be a k-transitive subgroup of $\operatorname{Sym}([n])$. Let A be a set of generators of H with $A = A^{-1}$ and $e \in A$. Then there is a subset $A' \subseteq A$ with $A' = (A')^{-1}$, such that, for every $\varepsilon > 0$, for any $\ell \geq 2n^{3k} \log(n^k/\varepsilon)$, and for any k-tuples $\overline{x} = (x_1, \ldots, x_k)$, $\overline{y} = (y_1, \ldots, y_k)$ of distinct elements of [n], the probability of the event

$$\overline{y} = \overline{x}^{g_1 g_2 \cdots g_\ell}$$

for $g_1, \ldots, g_\ell \in A'$ (chosen independently, with uniform distribution on $A' \setminus \{e\}$ and with the identity being assigned probability 1/2) is at least $(1-\varepsilon)\frac{(n-k)!}{n!}$ and at most $(1+\varepsilon)\frac{(n-k)!}{n!}$.

Proof. Let Δ be the set of k-tuples of distinct elements of [n]. Since H acts transitively on Δ and since $\langle A \rangle = H$, Lemma 3.10 gives us a subset A' of A with $\langle A' \rangle$ transitive on Δ and with $|A'| < |\Delta|$. Set $A_0 = A' \cup A'^{-1}$. Let Γ be the multigraph with vertex set Δ and with $\Gamma(\overline{x}) = \{\overline{x}^a \mid a \in A_0\}$ as the multiset of neighbours of \overline{x} for each $\overline{x} \in \Delta$. Clearly, Γ is a regular graph of valency $|A_0| \leq 2|\Delta|$ and with $|\Delta| \leq n^k$ vertices. Now the statement follows from Lemma 4.1 applied to Γ .

4.2. **Generators.** Given $A \subseteq \operatorname{Sym}([n])$ such that $\langle A \rangle$ is $\operatorname{Alt}([n])$ or $\operatorname{Sym}([n])$, how long can it take to construct a *small* set of generators for a *transitive* subgroup of $\langle A \rangle$? This subsection is devoted to answering that question. We start by proving two auxiliary lemmas.

Lemma 4.3. Let $A \subset \operatorname{Sym}([n])$, $e \in A$. Assume $\langle A \rangle$ is transitive. Then there is a $g \in A^n$ such that $|\operatorname{supp}(g)| \geq n/2$.

Proof. For each $i \in [n]$, let g_i be an element of A moving i. (If no such element existed, then $\langle A \rangle$ could not be transitive.) Let $g = g_1^{r_1} g_2^{r_2} \dots g_n^{r_n}$, where $r_1, r_2, \dots, r_n \in \{0, 1\}$ are independent random variables taking the values 0 and 1 with equal probability.⁹

Let $\alpha \in [n]$ be arbitrary. Let j be the largest integer such that g_j moves α . Then g moves α if and only if $g' = g_1^{r_1} \dots g_j^{r_j}$ moves α . Take r_1, r_2, \dots, r_{j-1} as given. If

⁹Such an element g is called a random subproduct of the sequence (g_i) . This notion was introduced by [BLS88] in the context of the analysis of algorithms on permutation groups. See, e.g., [Ser03, $\S 2.3$] for other applications.

 $\beta = \alpha^{g_1^{r_1} \dots g_{j-1}^{r_{j-1}}}$ equals α , then g' moves α if and only if $r_j = 1$; this happens with probability 1/2. If $\beta \neq \alpha$, then g' certainly moves α if $r_j = 0$, and thus moves α with probability at least 1/2. Thus g moves α with probability at least 1/2.

Summing over all α , we see that the expected value of the number of elements of [n] moved by g is at least n/2. In particular, there is a $g \in A^n$ moving at least n/2 elements of [n].

The following is the simplest sphere-packing lower bound, applied to the Hamming distance. (The *Hamming distance* on $\{0,1\}^k$ is $d(\vec{x},\vec{y}) = |\{1 \le j \le k : x_i \ne y_i\}|.$)

Lemma 4.4. Let n > 0, $k \ge 4.404 \log_2 n$, $\rho > 1$. Let $U = \{0,1\}^k$ the set of $\{0,1\}$ -sequences of length k. Then there exists $V \subseteq U$, |V| > n such that any two sequences in V differ in more than $\log_2 n$ coordinates.

Proof. In general, for U a metric space and $V \subseteq U$ maximal with respect to the property that the distance between any two points of V is greater than r, the closed balls of radius r around the points of V clearly cover U; hence, if the notion of volume is well-defined, |V| is at most $\operatorname{Vol}(U)$ divided by the volume of a closed ball of radius r. Applying this to the Hamming distance, we obtain that, for $V \subseteq U$ maximal,

$$|V| \ge \frac{2^k}{\sum_{j=0}^{\lfloor r \rfloor} \binom{k}{j}}.$$

By, e.g., [MS77, §10.11, Lem. 8],

$$\sum_{j=0}^{\lfloor r\rfloor} \binom{k}{j} \le 2^{kH(\lfloor r\rfloor/k)} \le 2^{kH(r/k)}$$

for $0 \le r \le k/2$, where $H(x) = -x \log_2 x - (1-x) \log_2 (1-x)$ is the binary entropy function. Let $r = \log_2 n$. It is easy to check that, for $0 \le \rho \le 1/4.404$, $1 - H(\rho) > \rho$. Hence

$$|V| \ge 2^{k(1-H(r/k))} > 2^{k \cdot r/k} = 2^r = n.$$

The following lemma is the main step toward answering the question raised at the beginning of the subsection. Most of the proof goes to show that, for some $g \in A^n$, $h \in A^{\ell}$ and a random $\beta \in [n]$, the orbit of β under $\langle g, h \rangle$ is rather large. The following is a brief sketch. If β were being acted upon by many random elements of $\mathrm{Sym}([n])$ in succession, it would indeed traverse many points. Now think of this obvious remark as being strengthened twice. First, let g have large support and let h be a random element of $\mathrm{Sym}([n])$. If we let h act on β and then let g act (or not) on β^h , and we let this happen over and over, the effect is a great deal as if β were

 $^{^{10}}$ This argument essentially appears in [BLS88, §6.2] (without proof). It appears again, with proof and in a much more general context, in [BCF⁺91]. Indeed, Lemma 4.3 here follows immediately from [BCF⁺91, Lem. 2.2] (with K equal to a point stabilizer), and the idea of the proof of Lemma 4.3 given here is exactly the same as that of [BCF⁺91, Lem. 2.2]. We thank an anonymous referee for this remark.

being acted upon by random elements in succession: if β has arrived at a point x where it has not been before, then the random element h acts on it in a way that, as far as we are concerned, is essentially random, in that it is almost independent of any of the parts of h we have seen so far. This makes the action of the fixed element g on x^h itself random. Here comes the second strengthening: it is actually enough for h to be the outcome of a random walk of moderate length $\ell \leq n^{O(\log n)}$: as we know (Lemma 4.2), such an h pretends to be a random element of $\operatorname{Sym}([n])$ very ably as far as its action on k tuples, $k \ll \log n$, is concerned; these are all the tuples that we have to deal with, since the above argument gives us large orbits after $O(\log n)$ steps.

The proof below is just a detailed and rigourous version of this sketch.

Lemma 4.5. Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. Then there are $g \in A^n$, $h \in A^{\lfloor n^{27 \log n} \rfloor}$ such that the action of $\langle g, h \rangle$ on [n] has at most $175(\log n)^2$ orbits, provided that n is larger than an absolute constant.

Proof. We will show that, for some $g \in A^n$, for $h \in A^{\ell}$ ($\ell \leq \lfloor n^{27 \log n} \rfloor$) taken randomly in a sense we will specify, and for any $\beta \in [n]$, the expected value of $1/|\beta^{\langle g,h\rangle}|$ is at most $175(\log n)^2/n$. (Here $\beta^{\langle g,h\rangle}$ denotes the orbit of β under the action of $\langle g,h\rangle \leq \operatorname{Sym}([n])$.) Now, $\sum_{\beta \in [n]} 1/|\beta^{\langle g,h\rangle}|$ is just the number of orbits of $\langle g,h\rangle$ (since each such orbit contributes $|\beta^{\langle g,h\rangle}| \cdot 1/|\beta^{\langle g,h\rangle}| = 1$ to the sum). Hence, by the additivity of expected values,

$$\mathbb{E}(\text{number of orbits of } \langle g, h \rangle) = \sum_{\beta \in [n]} \mathbb{E}\left(\frac{1}{|\beta^{\langle g, h \rangle}|}\right) \leq 175(\log n)^2.$$

In particular, this will imply that there exists an $h \in A^{\ell}$ such that the number of orbits of $\langle q, h \rangle$ is at most $175(\log n)^2$, and so we will be done.

Let $k = \lceil 4.404 \log_2 n \rceil$. By Lemma 4.3, there is an element $g \in A^n$ with

$$|\operatorname{supp}(g)| = \alpha n \ge n/2.$$

Let $\varepsilon = 1/n$ and $\ell = \lceil 2n^{6k} \log(n^{2k}/\varepsilon) \rceil$. (It is easy to check that, for n larger than an absolute constant, $\ell \leq \lfloor n^{27\log n} \rfloor$.) Let $h \in A^{\ell}$ be the outcome of a random walk of length ℓ as in Lemma 4.2.

Consider all words of the form

$$f(\vec{a}) = hg^{a_1}hg^{a_2}\dots hg^{a_k},$$

where $\vec{a} = (a_i : 1 \le i \le k)$ runs through all sequences in $U = \{0, 1\}^k$. For $\beta \in [n]$, we wish to estimate $|\beta^{\langle g, h \rangle}|$ from below by counting the number of different images $f_{\beta}(\vec{a}) := \beta^{f(\vec{a})}$, for $\vec{a} \in U$.

To this end, for fixed elements $\vec{a} = (a_1, \ldots, a_k)$ and $\vec{a'} = (a'_1, \ldots, a'_k)$ in U and $\beta \in [n]$, we wish to bound from above the probability that $f_{\beta}(\vec{a}) = f_{\beta}(\vec{a'})$. We will do this by examining all possible trajectories $(\beta_1, \ldots, \beta_k)$, $(\beta'_1, \ldots, \beta'_k)$, where

$$\beta_1 = \beta^{hg^{a_1}}, \ \beta_2 = \beta_1^{hg^{a_2}}, \ \dots, \ \beta_k = \beta_{k-1}^{hg^{a_k}} \quad \text{and} \quad \beta_1' = \beta^{hg^{a_1'}}, \ \dots, \ \beta_k' = \beta_{k-1}^{hg^{a_k'}},$$

counting how many satisfy $\beta_k \neq \beta'_k$, and then estimating the probability (for h chosen randomly in the manner described above) that such a pair of trajectories be traversed following $f(\vec{a})$ and $f(\vec{a'})$.

Let $R = \{1 \leq i \leq k : a_i \neq a_i'\}$; let the elements of R be $k_1 < k_2 < \cdots < k_r$, where r = |R|. Let $r_0 \leq r$ be fixed. Let $k' = k_{r_0}$. Consider all tuples $(\beta_1, \beta_2, \ldots, \beta_k, \beta'_{k'}, \ldots, \beta'_k) \in [n]^{(2k-k')+1}$ such that

- (a) $\beta_1, \beta_2, \dots, \beta_k, \beta'_{k'}, \dots, \beta'_k$ are distinct from each other and from β ,
- (b) $\beta_1^{g^{-a_1}}, \beta_2^{g^{-a_2}}, \dots, \beta_k^{g^{-a_k}}, (\beta'_{k'+1})^{g^{-a'_{k'+1}}}, \dots, (\beta'_k)^{g^{-a'_k}}$ are distinct from each other,
- (c) $\beta_{k_i} \notin \text{supp}(g)$ for every $j < r_0$, but $\beta_{k'} \in \text{supp}(g)$,
- (d) $(\beta'_{k'})^{g^{-a'_{k'}}} = (\beta_{k'})^{g^{-a_{k'}}}.$

The number of such tuples is at least

(4.1)
$$\left(\prod_{j=1}^{r_0-1} (n - |\operatorname{supp}(g)| - j) \right) \cdot (|\operatorname{supp}(g)| - 1) \cdot \prod_{j=(r_0+1)}^{2k-k'} (n - (2j-1)),$$

where we count tuples by choosing first $\beta_{k_j} \in [n] \setminus \text{supp}(g)$ for $1 \leq j < r_0$, then $\beta_{k'} \in \text{supp}(g)$, then the other β_i and β_i' . To justify the estimate on the number of choices at each stage, notice that at the j^{th} choice with $j \leq r_0 - 1$ we have to make selections from $[n] \setminus \text{supp}(g)$ so as to satisfy (c) while keeping them different from previous selections and from β (to satisfy (a)). Then $\beta_{k'}$ can be chosen as an arbitrary element of supp(g) different from β . At this point, (b) is still satisfied automatically. At later choices, if β_i or β_i' is selected at stage j then enforcing (a) eliminates j possibilities and enforcing (b) eliminates j-1, not necessarily different, possibilities. Note that (4.1) also gives a valid lower estimate (namely, 0) in the case when $r_0 - 1 \geq n - |\sup(g)| > 0$ (the negative terms in the first product in (4.1) are made harmless by a term equal to 0).

By Lemma 4.2 (with 2k - k' instead of k, and with properties (a), (b) as inputs), the probability that a random $h \in A^{\ell}$ satisfies (4.2)

$$(\beta, \beta_1, \dots, \beta_{k-1}, \beta'_{k'}, \dots, \beta'_{k-1})^h = (\beta_1^{g^{-a_1}}, \beta_2^{g^{-a_2}}, \beta_k^{g^{-a_k}}, (\beta'_{k'+1})^{g^{-a'_{k'+1}}}, \dots, (\beta'_k)^{g^{-a'_k}})$$
 is at least $(1 - \varepsilon) \frac{(n - (2k - k'))!}{n!} > (1 - \varepsilon) \frac{1}{n^{2k - k'}}$. If h satisfies (4.2) then $\beta^{hg^{a_1}} = \beta_1$, $\beta_1^{hg^{a_2}} = \beta_2, \dots, \beta_{k-1}^{hg^{a_k}} = \beta_k$. By properties (c) and (d), we also have $\beta^{hg^{a'_1}} = \beta_1$, $\beta_1^{hg^{a'_2}} = \beta_2, \dots, (\beta_{k'-1})^{hg^{a'_{k'}}} = (\beta_{k'})^{g^{-a_{k'}}g^{a'_{k'}}} = (\beta'_{k'})^{g^{-a'_{k'}}g^{a'_{k'}}} = \beta'_{k'}$; by (4.2), we also have $(\beta'_{k'})^{hg^{a'_{k'+1}}} = \beta'_{k'+1}, \dots, (\beta'_{k-1})^{hg^{a'_k}} = \beta'_k$. Thus, in particular, any two distinct tuples

$$(\beta_1, \beta_2, \dots, \beta_k, \beta'_{k'}, \dots, \beta'_k)$$

give us mutually exclusive events, even for different values of r_0 . Note also that, by property (a) and what we have just said, $f_{\beta}(\vec{a}) = \beta_k \neq \beta_k' = f_{\beta}(\vec{a'})$.

Hence the probability P that $f_{\beta}(\vec{a}) \neq f_{\beta}(\vec{a'})$ is at least

$$P \ge \sum_{r_0=1}^{r} \frac{1-\varepsilon}{n^{2k-k_{r_0}}} \cdot \left(\prod_{j=1}^{r_0-1} (n-\alpha n-j)\right) \cdot (\alpha n-1) \cdot \prod_{j=(r_0+1)}^{2k-k_{r_0}} (n-(2j-1))$$

$$(4.3) \qquad > \sum_{r_0=1}^{r} \left(1-\frac{1}{n}\right) \left(1-\frac{4k}{n}\right)^{2k} \left(\alpha-\frac{1}{n}\right) \cdot \prod_{j=1}^{r_0-1} \left(1-\alpha-\frac{j}{n}\right).$$

If $\alpha n = |\operatorname{supp}(g)| \ge n - k$ then we estimate P from below by the summand $r_0 = 1$ in (4.3), yielding

$$P > \left(1 - \frac{1}{n}\right) \left(1 - \frac{4k}{n}\right)^{2k} \left(\alpha - \frac{1}{n}\right) > 1 - \frac{1}{n} - \frac{8k^2}{n} - \frac{k+1}{n} \ge 1 - \frac{9k^2}{n},$$

with the last inequality valid for $n \geq 2$.

If $\alpha n = |\sup(g)| < n - k$ then, estimating the terms $(1 - \alpha - j/n)$ in the last product in (4.3) from below by $(1 - \alpha - k/n)$, we obtain

$$(4.4) P > \left(1 - \frac{1}{n}\right) \left(1 - \frac{4k}{n}\right)^{2k} \left(\alpha - \frac{1}{n}\right) \sum_{r_0 = 1}^r \left(1 - \alpha - \frac{k}{n}\right)^{r_0 - 1}$$

$$> \left(1 - \frac{4k}{n}\right)^{2k + 1} \left(\alpha - \frac{1}{n}\right) \frac{1 - (1 - \alpha - (k/n))^r}{(1 - (1 - \alpha - (k/n)))}$$

$$= \left(1 - \frac{4k}{n}\right)^{2k + 1} \frac{\alpha - (1/n)}{\alpha + (k/n)} \left(1 - (1 - \alpha - (k/n))^r\right).$$

Since $\alpha \geq 1/2$, we have $\frac{\alpha - (1/n)}{\alpha + (k/n)} \geq 1 - \frac{2(k+1)}{n}$ and $(1 - \alpha - (k/n))^r < (1/2)^r$, implying

$$P > 1 - \frac{4k(2k+1)}{n} - \frac{2(k+1)}{n} - \frac{1}{2^r} > 1 - \frac{9k^2}{n} - \frac{1}{2^r}$$

if $n \geq 3$ (since then $k \geq 7$).

We conclude that, for any two non-identical tuples

$$\vec{a} = (a_1, \dots, a_k) \in \{0, 1\}^k, \quad \vec{a'} = (a'_1, \dots, a'_k) \in \{0, 1\}^k$$

and for any $\beta \in [n]$,

$$\operatorname{Prob}(\beta^{hg^{a_1}hg^{a_2}...hg^{a_k}} = \beta^{hg^{a'_1}hg^{a'_2}...hg^{a'_k}}) < \frac{9k^2}{n} + \frac{1}{2^{d(\vec{a},\vec{a'})}},$$

where $d(\vec{a}, \vec{a'})$ is the Hamming distance between \vec{a} and $\vec{a'}$, i.e., the number of indices $1 \le j \le k$ for which $a_j \ne a'_j$.

By Lemma 4.4, there exists a set V of more than n tuples so that any two tuples differ in more than $\log_2 n$ coordinates. For fixed $\beta \in [n]$, writing $f_{\beta}(\vec{a}) = \beta^{hg^{a_1}hg^{a_2}...hg^{a_k}}$, $\vec{a} \in V$ for the random variable $\beta \mapsto f_{\beta}(\vec{a})$ defined using a random

 $h \in A^{\ell}$, we obtain that

$$\mathbb{E}(|\{(\vec{a}, \vec{a}') \in V^2 : f_{\beta}(\vec{a}) = f_{\beta}(\vec{a}')\}|) = \sum_{\vec{a}, \vec{a}' \in V} \text{Prob}(f_{\beta}(\vec{a}) = f_{\beta}(\vec{a}'))$$

$$\leq |V| + \left(\frac{9k^2}{n} + \frac{1}{2^{d(\vec{a}, \vec{a}')}}\right) |V|(|V| - 1) < \frac{|V|^2}{n} + \left(\frac{9k^2}{n} + \frac{1}{n}\right) |V|^2$$

$$< (9k^2 + 2) \frac{|V|^2}{n} < 175(\log n)^2 \frac{|V|^2}{n}$$

for n larger than an absolute constant.

Concerning the length of the orbit $\beta^{\langle g,h\rangle}$, we have

$$\mathbb{E}\left(\frac{1}{|\beta^{\langle g,h\rangle}|}\right) \leq \mathbb{E}\left(\frac{1}{|\{f_{\beta}(\vec{a}) : \vec{a} \in V\}|}\right)$$

$$\leq \mathbb{E}\left(\frac{|\{(\vec{a}, \vec{a}') \in V^2 : f_{\beta}(\vec{a}) = f_{\beta}(\vec{a}')\}|}{|V|^2}\right) \leq \frac{175(\log n)^2}{n},$$

where we use Cauchy-Schwarz in the second step for the numbers m_i that measure how many times a particular value γ_i occurs among the $f_{\beta}(\vec{a})$, for some $\vec{a} \in V$. \square

Proposition 4.6. Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. If n is larger than an absolute constant, then there are $g_1, g_2, g_3 \in A^{\lfloor n^{27 \log n} \rfloor}$ such that $\langle g_1, g_2, g_3 \rangle$ is transitive.

Proof. Let g, h be as in Lemma 4.5. Let $\varepsilon = 1/n^2$, $\ell = \lceil 2n^6 \log(n^2/\varepsilon) \rceil$. Let $g' \in A^{\ell}$ be the outcome of a random walk of length ℓ as in Lemma 4.2. Note that $\ell \leq \lfloor n^{27 \log n} \rfloor$ for n larger than an absolute constant.

Let Δ be the union of orbits of $\langle g, h \rangle$ of length less than \sqrt{n} . Since, by Lemma 4.5, there are at most $175(\log n)^2$ orbits of $\langle g, h \rangle$, we have $|\Delta| < 175\sqrt{n}(\log n)^2$. Let S be a set consisting of one element α of each orbit of length less than \sqrt{n} . Then, for each $\alpha \in S$, Lemma 4.2 implies that

$$\operatorname{Prob}\left(\alpha^{g'} \in \Delta\right) \le (1+\varepsilon)\frac{|\Delta|}{n} < \left(1+\frac{1}{n^2}\right)\frac{175(\log n)^2}{\sqrt{n}}$$

and so

(4.5)
$$\operatorname{Prob}\left((\exists \alpha \in S) \ (\alpha^{g'} \in \Delta)\right) < \left(1 + \frac{1}{n^2}\right) \frac{175^2(\log n)^4}{\sqrt{n}}.$$

Let κ be an orbit of $\langle g, h \rangle$ contained in $n \setminus \Delta$; by definition, $|\kappa| \geq \sqrt{n}$. Let κ_0 be the largest orbit; by the pigeonhole principle, $|\kappa_0| > n/(175(\log n)^2)$. Then

$$\mathbb{E}(|\kappa^{g'} \cap \kappa_0|) = \sum_{\alpha \in \kappa} \operatorname{Prob}(\alpha^{g'} \in \kappa_0) \ge \sum_{\alpha \in \kappa} (1 - \varepsilon) \frac{|\kappa_0|}{n} = (1 - \varepsilon) \frac{|\kappa||\kappa_0|}{n},$$

whereas

$$\mathbb{E}\left(|\kappa^{g'} \cap \kappa_{0}|^{2}\right) = \sum_{\alpha,\beta \in \kappa} \operatorname{Prob}\left(\alpha^{g'} \in \kappa_{0} \wedge \beta^{g'} \in \kappa_{0}\right)$$

$$= \sum_{\alpha \in \kappa} \operatorname{Prob}(\alpha^{g'} \in \kappa_{0}) + \sum_{\substack{\alpha,\beta \in \kappa \\ \alpha \neq \beta}} \sum_{\substack{\alpha',\beta' \in \kappa_{0} \\ \alpha' \neq \beta'}} \operatorname{Prob}((\alpha,\beta)^{g'} = (\alpha',\beta'))$$

$$\leq \sum_{\alpha \in \kappa} (1+\varepsilon) \frac{|\kappa_{0}|}{n} + \sum_{\substack{\alpha,\beta \in \kappa,\alpha \neq \beta \\ \alpha,\beta \in \kappa,\alpha \neq \beta}} (1+\varepsilon) \frac{|\kappa_{0}|(|\kappa_{0}|-1)}{n(n-1)}$$

$$\leq (1+\varepsilon) \left(\frac{|\kappa_{0}||\kappa|}{n} + \frac{|\kappa|(|\kappa|-1)|\kappa_{0}|(|\kappa_{0}|-1)}{n(n-1)}\right)$$

$$\leq (1+\varepsilon) \left(\frac{|\kappa_{0}||\kappa|}{n} + \frac{|\kappa|^{2}|\kappa_{0}|^{2}}{n^{2}}\right).$$

Thus

$$\operatorname{Var}(|\kappa^{g'} \cap \kappa_0|) = \mathbb{E}(|\kappa^{g'} \cap \kappa_0|^2) - \mathbb{E}(|\kappa^{g'} \cap \kappa_0|)^2$$

$$\leq (1+\varepsilon) \left(\frac{|\kappa_0||\kappa|}{n} + \frac{|\kappa_0|^2|\kappa|^2}{n^2}\right) - (1-\varepsilon)^2 \frac{|\kappa_0|^2|\kappa|^2}{n^2}$$

$$\leq 3\varepsilon \frac{|\kappa|^2|\kappa_0|^2}{n^2} + (1+\varepsilon) \frac{|\kappa_0||\kappa|}{n} < \left(1 + \frac{4}{n}\right) \frac{|\kappa_0||\kappa|}{n}.$$

By Chebyshev's inequality,

$$\operatorname{Prob}(\kappa^{g'} \cap \kappa_0 = \emptyset) \leq \frac{\operatorname{Var}(|\kappa^{g'} \cap \kappa_0|)}{\mathbb{E}(|\kappa^{g'} \cap \kappa_0|)^2}$$

$$\leq \frac{(|\kappa||\kappa_0|/n)(1 + 4/n)}{(1 - \varepsilon)^2 \frac{|\kappa|^2 |\kappa_0|^2}{n^2}} \leq \frac{12n}{|\kappa||\kappa_0|} < \frac{12 \cdot 175(\log n)^2}{\sqrt{n}}.$$

Hence

(4.6)
$$\operatorname{Prob}\left((\exists \kappa \subseteq ([n] \setminus \Delta)) \ (\kappa^{g'} \cap \kappa_0 = \emptyset)\right) < \frac{12 \cdot 175^2 (\log n)^4}{\sqrt{n}}.$$

Now, for n larger than a constant,

$$\left(1 + \frac{1}{n^2}\right) \frac{175^2 (\log n)^4}{\sqrt{n}} + \frac{12 \cdot 175^2 (\log n)^4}{\sqrt{n}} < 1.$$

Therefore, (4.5) and (4.6) imply that with positive probability, (a) $\kappa^{g'}$ intersects $[n] \setminus \Delta$ for every orbit κ not contained in $[n] \setminus \Delta$ and (b) $\kappa^{g'}$ intersects κ_0 for every orbit κ contained in $[n] \setminus \Delta$. In particular, this happens for some $g' \in A^{\ell}$. Properties (a) and (b) imply that $\langle g, h, g' \rangle$ is transitive. We set $g_1 = g$, $g_2 = h$, $g_3 = g'$ and are done.

We will later use¹¹ the following corollary with k = 2.

Corollary 4.7. Let $A \subseteq \operatorname{Sym}[(n])$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. Let $k \geq 1$. If n is larger than a constant depending only on k, then there is a set $S \subseteq A^{\lfloor n^{28 \log n} \rfloor}$ of size at most 3k such that $\langle S \rangle$ is k-transitive.

Proof. Let $\alpha_1 \in [n]$ be arbitrary. Since $\langle A \rangle$ is transitive, Lemma 3.9 implies that $\alpha_1^{A^n} = [n]$. Let $G = \operatorname{Sym}([n])$, $H = G_{\alpha_1}$, $A' = A^n$. Since $\alpha_1^{A'} = [n]$, A' intersects every coset of H in G. By Schreier's Lemma (Lem 3.8), it follows that $(A')^3 \cap H$ generates $\langle A \rangle \cap H$, which is either $\operatorname{Sym}([n] \setminus \{\alpha_1\})$ or $\operatorname{Alt}([n] \setminus \{\alpha_1\})$. Let $A_1 = (A')^3 \cap H$.

Iterating, we obtain a sequence of sets $A_0 = A, A_1, A_2, \ldots, A_{k-1} \subseteq \operatorname{Sym}([n])$ and a sequence of elements $\alpha_1, \alpha_2, \ldots, \alpha_{k-1} \in [n]$ such that $A_i \subseteq A_{i-1}^{3n}$ and $\langle A_i \rangle$ is $\operatorname{Sym}([n] \setminus \{\alpha_1, \ldots, \alpha_i\})$ or $\operatorname{Alt}([n] \setminus \{\alpha_1, \ldots, \alpha_i\})$.

Let $(g_1)_i$, $(g_2)_i$, $(g_3)_i$ be as in Prop. 4.6, applied with A_i instead of A. Then $(g_1)_i$, $(g_2)_i$, $(g_3)_i \in A^{(3n)^i \lfloor n^{27 \log n} \rfloor}$ and $\langle (g_1)_i, (g_2)_i, (g_3)_i \rangle \subseteq \operatorname{Sym}([n] \setminus \{\alpha_1, \ldots, \alpha_i\})$ is transitive on $[n] \setminus \{\alpha_1, \ldots, \alpha_i\}$ for $0 \leq i \leq k-1$. Thus, for $S = \bigcup_{i=0}^{k-1} A_i$, $\langle S \rangle$ is k-transitive on [n].

5. The splitting Lemma and its consequences

We will prove what is in effect an adaptation of Babai's splitting lemma (proven for groups in [Bab82, Lem. 3.1]) to the case of sets. This is a key point in this paper: the splitting lemma will allow us to construct long stabilizer chains with large orbits.

The following easy lemma will make an "unfolding" step possible.

Lemma 5.1. Let $A \subseteq \text{Sym}([n]), \Sigma \subseteq [n]$ and $g \in \text{Sym}([n])$. Then

$$gA_{(\Sigma^g)}g^{-1} = (gAg^{-1})_{(\Sigma)}.$$

Proof. We have $\operatorname{Sym}([n])_{(\Sigma^g)} = g^{-1} \operatorname{Sym}([n])_{(\Sigma)} g$. Therefore,

$$A_{(\Sigma^g)} = A \cap \text{Sym}([n])_{(\Sigma^g)} = A \cap g^{-1} \text{Sym}([n])_{(\Sigma)} g$$

= $g^{-1}(gAg^{-1} \cap \text{Sym}([n])_{(\Sigma)})g = g^{-1}(gAg^{-1})_{(\Sigma)}g.$

Notice a feature of the following statement – there is a high power of A in the assumptions, not just in the conclusion. We will "unfold" the high power of A in the course of the proof. (By Σ^S we mean the set $\Sigma^S = \{\alpha^g : \alpha \in \Sigma, g \in S\}$.)

Proposition 5.2 (Splitting Lemma). Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle$ 2-transitive. Let $\Sigma \subseteq [n]$. Assume that there are at least $\rho n(n-1)$ ordered pairs (α, β) of distinct elements of [n] such that there is no $g \in (A^{\lfloor 9n^6 \log n \rfloor})_{(\Sigma)}$ with $\alpha^g = \beta$. Then there is a subset S of $A^{\lfloor 5n^6 \log n \rfloor}$ with

$$(AA^{-1})_{(\Sigma^S)} = \{e\}$$

П

¹¹If we wished to, we could use it to obtain a set S of generators of Alt([n]) or Sym([n]) simply by setting k = 6: the Classification of Finite Simple Groups implies that a 6-transitive group must be either alternating or symmetric.

and

$$|S| \le \left\lceil \frac{2}{\log(3/(3-2\rho))} \cdot \log n \right\rceil.$$

Proof. Set $\ell = \lceil 2n^6 \log(n^2/(1/3)) \rceil$; note that $\ell \leq \lfloor 5n^6 \log n \rfloor$ and $2\ell+2 \leq \lfloor 9n^6 \log n \rfloor$ for $n \geq 5$. (For n < 5, the statement is trivial.) By Lemma 4.2 applied with k = 2 and $\varepsilon = 1/3$, we obtain that given any two distinct elements $\alpha, \beta \in [n]$ and $g \in A^{\ell}$, the pair (α^g, β^g) adopts any possible value (α', β') with probability at least (1 - 1/3)/(n(n - 1)), where we choose $g \in A^{\ell}$ with the distribution in Lemma 4.2 $(g = g_1g_2 \cdots g_{\ell}, g_i$ chosen independently from $A' \cup \{e\}$, where A' is a symmetric subset of A). Since this distribution is symmetric, this is the same as saying that $(\alpha^{g^{-1}}, \beta^{g^{-1}})$ adopts any possible value (α', β') with probability at least (1 - 1/3)/(n(n - 1)).

Now, given (α, β) and $g \in A^{\ell}$, we have $h \in (AA^{-1})_{(\Sigma^g)}$ and $\alpha^h = \beta$ if and only if $ghg^{-1} \in g(AA^{-1})_{(\Sigma^g)}g^{-1}$ and $(\alpha^{g^{-1}})^{ghg^{-1}} = \beta^{g^{-1}}$. By Lemma 5.1 applied to AA^{-1} , we have that $ghg^{-1} \in g(AA^{-1})_{(\Sigma^g)}g^{-1}$ only if $ghg^{-1} \in (gAA^{-1}g^{-1})_{(\Sigma)}$, which in turn can happen only if $ghg^{-1} \in (A^{2\ell+2})_{(\Sigma)}$. Thus, if there is no element $j \in (A^{2\ell+2})_{(\Sigma)}$ with $\alpha^{g^{-1}j} = \beta^{g^{-1}}$, then there is no element $h \in (AA^{-1})_{(\Sigma^g)}$ with $\alpha^h = \beta$. (This is the "unfolding" step we referred to before.)

Since by hypothesis there are at least $\rho n(n-1)$ ordered pairs (α', β') such that there is no element $j \in (A^{2\ell+2})_{(\Sigma)}$ with $\alpha'^j = \beta'$, and since $(\alpha^{g^{-1}}, \beta^{g^{-1}})$ equals any such pair with probability at least (2/3)/(n(n-1)), we see that the probability that there is no element $h \in (AA^{-1})_{(\Sigma^g)}$ with $\alpha^h = \beta$ is at least $2\rho/3$.

Let S be a set of r random $g \in A^{\ell}$ (chosen independently, with the distribution as above). The probability that for every $g \in S$ there is an element $h \in (AA^{-1})_{(\Sigma^g)}$ with $\alpha^h = \beta$ is at most $(1 - 2\rho/3)^r$. This must happen if there is an element $h \in (AA^{-1})_{\Sigma^S}$ such that $\alpha^h = \beta$. Thus, the probability that there is such an h is at most $(1 - 2\rho/3)^r$, and the probability that there is such an h for at least one of the n(n-1) pairs (α, β) is at most $n(n-1)(1-2\rho/3)^r$.

Setting $r = \lceil (\log n^2)/(\log 3/(3-2\rho)) \rceil$, we obtain that the probability that there is such an h for at least one pair is less than 1. Hence there is a set $S \subseteq A^{\ell}$ with at most r elements such that, for every pair (α, β) of distinct elements of [n], there is no $h \in (AA^{-1})_{(\Sigma^S)}$ with $\alpha^h = \beta$. This implies immediately that the only element of $(AA^{-1})_{(\Sigma^S)}$ is the identity.

Corollary 5.3. Let $A \subseteq \operatorname{Sym}([n])$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle$ 2-transitive. Let $A' = A^{\lfloor 9n^6 \log n \rfloor}$. Let $\Sigma \subseteq [n]$ be such that

$$|\alpha^{A'_{(\Sigma)}}| < (1 - \rho)n$$

for every $\alpha \in [n]$, where $\rho \in (0,1)$. Then

$$|\Sigma| > \frac{\log |A|}{\left\lceil \frac{2}{\log(3/(3-2\rho))} \cdot \log n \right\rceil \cdot \log n}.$$

In particular, if $\rho = 0.05$ then $|\Sigma| > (\log |A|)/(60(\log n)^2)$.

Proof. Since $|\alpha^{A'_{(\Sigma)}}| < (1-\rho)n$ for every $\alpha \in [n]$, there are at least $\rho n(n-1)$ pairs (α, β) such that there is no $g \in A'_{(\Sigma)}$ with $\alpha^g = \beta$. By Prop. 5.2, there is a set $S \subseteq \operatorname{Sym}([n])$ such that $(AA^{-1})_{(\Sigma^S)} = \{e\}$ and $|S| \le \left\lceil \frac{2}{\log(3/(3-2\rho))} \cdot \log n \right\rceil$. Since $(AA^{-1})_{(\Sigma^S)} = \{e\}$, we know, by Lemma 3.16, that $|\Sigma^S| \ge \log_n |A|$. Clearly $|\Sigma^S| \le |S||\Sigma|$. Hence

$$|\Sigma| \geq \frac{\log_n |A|}{|S|} \geq \frac{\log |A|}{\left\lceil \frac{2}{\log(3/(3-2\rho))} \cdot \log n \right\rceil \cdot \log n}.$$

A key idea in the proof of the Main Theorem is the following. For $A \subseteq \operatorname{Sym}([n])$, we can construct $A' = A^{\lfloor 5n^6 \log n \rfloor}$ and a set $\Sigma = \{\alpha_1, \alpha_2, \dots\} \subseteq [n]$ starting with an empty set and taking at each step α_i to be an element such that $|\alpha_i^{(A')}(\alpha_1,\dots,\alpha_{i-1})| \ge (1-\rho)n$ (say); if no such element exists, we stop the procedure. By Cor. 5.3, $|\Sigma|$ must be large.

An application of Lemma 3.18 will give that, for $A'' = (A')^{16n^6}$, the set $(A'')_{\Sigma}$ contains a copy of $Alt(\Delta)$, where $\Delta \subseteq \Sigma$ and $|\Delta| \ge (1 - \rho)|\Sigma|$. Such a large alternating group certainly looks like a valuable tool.

6. Proof of the main theorem

The core of this section is Proposition 6.4. It is a growth result, but not quite of type $|A \cdot A \cdot A| \ge |A|^{1+\varepsilon}$ or $|A^k| \ge |A|^{1+\varepsilon}$. What will grow by a factor at each step is not the number of elements |A| of A, but rather the length m of a sequence $\alpha_1, \ldots, \alpha_m$ such that the orbits

(6.1)
$$\alpha_1^A, \alpha_2^{A_{\alpha_1}}, \alpha_3^{A_{(\alpha_1,\alpha_2)}}, \dots, \alpha_m^{A_{(\alpha_1,\alpha_2,\dots,\alpha_{m-1})}}$$

are all large.

This growth result (Prop. 6.4) will be applied iteratively. There are two ways for the iteration to stop: (a) an element we construct could fix a large set pointwise (we call this the case of exit), or (b) a group we work with could fail to have a large alternating composition factor. In case (a), we obtain all of G = Alt([n]) in a few steps by Thm. 1.4. In case (b), we can descend to the problem of proving small diameter for n' smaller than n by a constant factor. (Here, as in "infinite descent", the term "descent" means the same as induction, seen backwards.)

* * *

Let us sketch briefly the proof of Prop. 6.4. First, we use (6.4) to construct many elements in the setwise stabilizer G_{Σ} , where $\Sigma = \{\alpha_1, \ldots, \alpha_m\}$; in fact we get an entire copy of a large alternating group in $(G_{\Sigma})|_{\Sigma}$ (Lemma 3.18). This is the setup. Then comes the creation step: we use the action by conjugation of G_{Σ} on the pointwise stabilizer $G_{(\Sigma)}$ to construct many elements of $G_{(\Sigma)}$ (Lemma 6.1). We organise these new elements (all in a power A' of A) as follows: we apply Cor. 5.3 (a consequence of the splitting lemma) to lengthen our stabilizer chain $A' \supseteq A'_{\alpha_1} \supseteq$

 $\ldots \supseteq A'_{(\alpha_1,\ldots,\alpha_m)} \supseteq \ldots$ up to $A'_{(\alpha_1,\ldots,\alpha_{m+\ell})}$ in such a way that the orbits (defined as in (6.1)) are still large. We repeat the *organiser* step about $\gg (\log n)/(\log m)$ times. There are only two ways for this procedure to stop prematurely, namely, exit and descent (cases (a) and (b) discussed above).

We start by proving the lemma containing the *creation* step: we give a way to construct many elements in a subgroup H^- of a group G. The basic idea is the application of the orbit-stabilizer principle to the action by conjugation of a subgroup $H^+ \leq N_G(H^-)$ on H^- , where $N_G(H^-)$ is the normaliser of H^- .

Lemma 6.1. Let $G = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$, $H^- \leq G$, $H^+ \leq N_G(H^-)$, Γ an orbit of both H^- and H^+ . Let $Y = \{y_1, \ldots, y_r\} \subseteq H^-$ be such that $\langle Y \rangle|_{\Gamma}$ is 2-transitive on Γ . Let $B \subseteq H^+$. Then either

- (a) there is a $b \in BB^{-1} \setminus \{e\}$ fixing Γ pointwise, or (b) $|B^{-1}YB \cap H^{-}| \geq |B|^{1/r}$.

Proof. Consider the action of B on $\vec{y} = (y_1, \dots, y_r)$ by conjugation: for $b \in B$, we define $\vec{y}^b := (y_1^b, \dots, y_r^b)$, where $y^b = b^{-1}yb$. Assume first that there are two distinct elements $b_1, b_2 \in B$ such that $\vec{y}^{b_1}|_{\Gamma} = \vec{y}^{b_2}|_{\Gamma}$. Then $b_1b_2^{-1}|_{\Gamma}$ centralizes $\vec{y}|_{\Gamma}$, implying that $b_1b_2^{-1}|_{\Gamma} \in C(\langle Y \rangle|_{\Gamma}) = \{e\}$. (As is well-known and can be easily seen, the centralizer of a doubly transitive group, such as $\langle Y \rangle|_{\Gamma} < \operatorname{Sym}(\Gamma)$, is trivial.) Hence $b_1b_2^{-1} \in B$ fixes Γ pointwise without being the identity, i.e., conclusion (a) holds.

Assume now that the restrictions $\vec{y}^b|_{\Gamma}$ are all distinct. Hence, by the pigeonhole principle, there exists an index $j \in \{1, \dots, r\}$ such that the set W of conjugates of y_i by B satisfies $|W|_{\Gamma}| \geq |B|^{1/r}$. Observe that all elements of W are in H^- , as $Y \subset H^{-} \text{ and } B \subset N_{G}(H^{-}). \text{ Hence } |B^{-1}YB \cap H^{-}| \geq |W| \geq |B|^{1/r}.$

The following useful lemma is in part an easy application of Schreier's lemma and in part a consequence of a trick based on the following trivial fact: one clearly cannot have two disjoint copies within [n] of an orbit of size greater than n/2.

Lemma 6.2. Let $\Delta \subseteq [n]$. Let $B^+ \subseteq (\operatorname{Sym}(n))_{\Delta}$ with $B^+ = (B^+)^{-1}$, $e \in B^+$. Assume $B^+|_{\Delta}$ is $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$. Let $B^- = ((B^+)^3)_{(\Delta)}$.

Then $\langle B^- \rangle = \langle B^+ \rangle_{(\Delta)} \triangleleft \langle B^+ \rangle$. Furthermore, if $\langle B^- \rangle$ has an orbit Γ of length greater than n/2, then Γ is also an orbit of $\langle B^+ \rangle$.

Proof. Since $B^+|_{\Delta}$ is a group $(Alt(\Delta) \text{ or } Sym(\Delta)), B^+|_{\Delta} = \langle B^+ \rangle|_{\Delta}$. Thus B^+ contains an element from every coset of $\langle B^+ \rangle_{(\Delta)}$ in $\langle B^+ \rangle$ and so, by Lemma 3.8, B^- contains a set of generators of $\langle B^+ \rangle_{(\Delta)}$. Hence $\langle B^- \rangle = \langle B^+ \rangle_{(\Delta)}$. In particular, $\langle B^- \rangle \triangleleft \langle B^+ \rangle$, as $\langle B^+ \rangle_{(\Delta)}$ is the kernel of the action of $\langle B^+ \rangle$ on Δ .

The orbits of the normal subgroup $\langle B^- \rangle \triangleleft \langle B^+ \rangle$ are blocks of imprimitivity for $\langle B^+ \rangle$. Since one cannot have two blocks of length greater than n/2, $\langle B^+ \rangle$ leaves Γ invariant as a set, and so Γ is an orbit of $\langle B^+ \rangle$.

The following lemma is also crucial to the descent step. In the proof of the lemma, we use Lemma 3.19 to guarantee the existence of an element that we then construct by other means.

Lemma 6.3. Let $G = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. Let $\Delta \subseteq [n]$, $|\Delta| \ge (\log n)^2$. Let $A \subseteq G$ with $A = A^{-1}$, $e \in A$ and $\langle A \rangle = G$. Let $B^+ \subseteq (A^l)_{\Delta}$, $l \ge 1$, with $B^+ = (B^+)^{-1}$, $e \in B^+$. Assume $B^+|_{\Delta}$ is $\operatorname{Alt}(\Delta)$ or $\operatorname{Sym}(\Delta)$. Let $B^- = ((B^+)^3)_{(\Delta)}$. Assume $\langle B^- \rangle$ has an orbit Γ of length at least ρn , for some $\rho > 8/9$.

If all alternating composition factors $\mathrm{Alt}(k)$ of $\langle B^- \rangle$ satisfy $k \leq \delta n$, where $\delta > 0$, and

(6.2)
$$\max_{k < \delta n} \operatorname{diam}(\operatorname{Alt}(k)) \le D_{\delta},$$

for some $D_{\delta} > 0$, and n is larger than an absolute constant, then

$$A^{\lfloor le^{c(\log n)^3}\cdot D_\delta\rfloor}\supseteq \mathrm{Alt}([n]),$$

where $c = c(\rho)$ depends only on ρ .

Proof. The group $U := \langle B^- \rangle|_{\Gamma}$ is transitive. It is also isomorphic to a quotient of $\langle B^- \rangle$, so U also has no alternating composition factors $\mathrm{Alt}(k)$ with $k > \delta n$. By Thm. 1.1 and by (6.2), there exists an absolute constant C_1 such that for

(6.3)
$$u := |e^{C_1(\log n)^3} \cdot D_{\delta}|, \quad (B^-)^u|_{\Gamma} = U.$$

Let $H = \langle B^+ \rangle$. By Lemma 6.2, Γ is an orbit of H. If n is large enough that Lemma 3.19 applies then there exists a non-identity element $g \in H$ of support less than $|\Gamma|/4$ on Γ . Take $h \in B^+$ with $h|_{\Delta} = g|_{\Delta}$. Then $gh^{-1} \in \langle B^+ \rangle_{(\Delta)} = \langle B^- \rangle$ and so, by (6.3), there exists $b \in (B^-)^u$ with $gh^{-1}|_{\Gamma} = b|_{\Gamma}$. Therefore, $bh \in (B^+)^{3u+1}$ satisfies $bh|_{\Gamma} = g|_{\Gamma}$. Since g fixes at least $(3/4)|\Gamma| \geq (3/4) \cdot \rho n > (2/3)n$ points in Γ , we have $|\sup(bh)| \leq (1 - (3/4)\rho)n < n/3$. By Thm. 1.4, $(A \cup \{bh, (bh)^{-1}\})^{Kn^8}$ contains Alt([n]), where $K = K(\varepsilon)$ ($\varepsilon = 1 - (3/4)\rho < 1/3$) is the number defined in Thm. 1.4. Since $A \cup \{bh, (bh)^{-1}\} \subseteq A^{(3u+1)l}$, we are done.

We come to the key results in the paper. They will be given as two separate propositions, proved by a back-and-forth inductive process. For the sake of clarity, we will state them in terms of functions $F_1, F_2 : \mathbb{R}^+ \to \mathbb{R}^+$ obeying certain relations; we will later specify functions satisfying these relations.

Proposition 6.4. Let $G = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. Let $A \subset G$ with $A = A^{-1}$, $e \in A$, and $\langle A \rangle = G$. Let $\alpha_1, \alpha_2, \ldots, \alpha_{m+1} \in [n]$ be such that

(6.4)
$$\left| \alpha_i^{A_{(\alpha_1, \dots, \alpha_{i-1})}} \right| \ge \frac{9}{10} n$$

for every $i = 1, 2, \dots, m + 1$, where $m \ge (\log n)^2$.

There are absolute constants $n_0 \in \mathbb{Z}^+$ and $K, c_1, c_2, c_3 > 0$ such that the following holds. Assume $n \geq n_0$. Assume also that Proposition 6.5 holds for all smaller values of n with respect to some increasing function $F_2 : \mathbb{R}^+ \to \mathbb{R}^+$. Let $F_1 : \mathbb{R}^+ \to \mathbb{R}^+$ be such that, for all $n \in \mathbb{Z}^+$,

(6.5)
$$F_1(n) \ge \max\left(n^{c_3 \log n} e^{c_1(\log n)^3} F_2(0.95n), 2Kn^{c_3 \log n + 8}\right).$$

Then either

$$(6.6) A^{\lfloor F_1(n) \rfloor} \supseteq Alt([n])$$

or there are $\alpha_{m+2}, \alpha_{m+3}, \dots, \alpha_{m+l+1} \in [n], l \geq c_2(m \log m)/(\log n)$, such that

$$\left|\alpha_i^{A'_{(\alpha_1,\dots,\alpha_{i-1})}}\right| \ge \frac{9}{10}n$$

for $A' = A^{\lfloor n^{c_3 \log n} \rfloor}$ and every $i = 1, 2, \dots, m + l + 1$.

An easy application of Proposition 6.4 proves Proposition 6.5 (which is equivalent to our Main Theorem). Conversely, in order to prove Proposition 6.4, we will use Proposition 6.5 for smaller values of n in an inductive process. In the proofs of Prop. 6.4 and Prop. 6.5, we assume that n is greater than a well-defined (but not explicitly computed) absolute constant n_0 ; we take n_0 to be large enough to satisfy the assumptions made in the course of both proofs. In the statement of Prop. 6.4, the assumption is made explicitly; in the statement of Prop. 6.5, the assumption is allowed by (6.8), which implies that, when $n \leq n_0$, the bound diam($\Gamma(G, Y)$) $\leq F_2(n)$ is trivial and there is nothing to prove.

Proposition 6.5. Let $G = \operatorname{Sym}([n])$ or $\operatorname{Alt}([n])$. Let $Y \subseteq G$ with $Y = Y^{-1}$, $e \in Y$ and $G = \langle Y \rangle$.

Assume Prop. 6.4 holds for n with respect to some function $F_1: \mathbb{R}^+ \to \mathbb{R}^+$. Let c_2 and c_3 be the absolute constants in the statement of Prop. 6.4; let n_0 be at least as large as in Prop. 6.4. Let $F_2: \mathbb{R}^+ \to \mathbb{R}^+$ be such that

$$(6.8) F_2(n) \ge \max \left(e^{(\log n)^3 + 2\log n + c'c_3(\log n)^3 \log\log n} F_1(n) + 2, n_0! \right)$$

for some $c' > c_2$ and all $n \in \mathbb{Z}^+$. Then

$$\operatorname{diam}(\Gamma(G, Y)) \leq F_2(n),$$

provided that n_0 is larger than a constant depending only on c_2 and c'.

The proof consists just of a repeated use of Proposition 6.4, plus some accounting.

Proof. We can assume that n is large enough that $m_0 \leq 0.1n \leq n-3$ for $m_0 = \lfloor (\log n)^2 \rfloor + 1$ and so G acts transitively on the set X of all $(m_0 + 1)$ -tuples. Hence, by Lemma 3.9, the set $A_0 := Y^{n^{m_0+1}} \supseteq Y^{|X|}$ acts transitively on the set of all $(m_0 + 1)$ -tuples. Thus (6.4) holds with A_0 instead of A, m_0 instead of m and $\alpha_i = i$ for $i = 1, 2, \ldots, m_0 + 1$. We apply Proposition 6.4 with these parameters, assuming $n \geq n_0$, where n_0 is the absolute constant in the statement of Prop. 6.4. We obtain either (6.6) or (6.7).

In the latter case, we set $\ell_0 = \ell$, $m_1 = m_0 + \ell_0$, and iterate: we apply Proposition 6.4 to

$$A_1 = A_0^r$$
, $A_2 = A_1^r = A_0^{r^2}$, $A_3 = A_2^r = A_0^{r^3}$,...

where $r = \lfloor n^{c_3 \log n} \rfloor$. (After each step, we "save" the output ℓ to ℓ_i and set $m_{i+1} = m_i + \ell_i$.) We stop when we obtain (6.6); say this happens when we apply Proposition 6.4 with $A = A_k = A_0^{r^k}$.

It remains to estimate k. By Proposition 6.4,

(6.9)
$$m_{i+1} > (1 + (c_2 \log m_i)/(\log n)) \cdot m_i$$

We want to compute how many times we have to iterate (6.9) before we run into a contradiction with $m_i \leq n$.

For $1 \le j \le \log n$, let t_j be the largest index i between 0 and k such that $m_i < e^j$; if no such index exists, set $t_j = 1$. We have $m_0 \ge 3$ and so $t_1 = 1$. By (6.9) and $(1 + c_2 j/(\log n))^{\lfloor (\log n)/(c_2 j) \rfloor + 2} > e$, we have $t_{j+1} \le t_j + \lfloor (\log n)/(c_2 j) \rfloor + 3$. Thus

$$t_{\lfloor \log n \rfloor} + 1 \le t_1 + 1 + \sum_{j=1}^{\lfloor \log n \rfloor - 1} (t_{j+1} - t_j)$$

$$\le 2 + \sum_{1 \le j \le \log n} \left(\frac{\log n}{c_2 j} + 3 \right) \le c' \log n \log \log n$$

for any $c' > 1/c_2$, with the last inequality valid if n is larger than a constant depending only on c and c'. Since $t_{\lfloor \log n \rfloor} + 2 > k$ (because $m_k \leq n$), we get that $k \leq c' \log n \log \log n$.

Thus

$$A_k = A_0^{r^k} \subseteq Y^{n^{\lfloor (\log n)^2 \rfloor + 2} \cdot r^{\lfloor c' \log n \log \log n \rfloor}} \subseteq Y^{\lfloor e^{(\log n)^3 + 2 \log n + c' c_3 (\log n)^3 \log \log n} \rfloor}.$$

Then, by (6.6) (valid for $A = A_k$), we obtain

$$\operatorname{Alt}([n]) \subseteq (Y^{\lfloor e^{(\log n)^3 + 2\log n + c'c_3(\log n)^3\log\log n}\rfloor})^{\lfloor F_1(n)\rfloor} \subseteq Y^{\lfloor F_2(n)\rfloor - 1}$$

for n larger than a constant. If $Y \subseteq \text{Alt}([n])$, then $Y^{\lfloor F_2(n) \rfloor - 1} = \text{Alt}([n])$. If Y contains an odd permutation then $Y^{\lfloor F_2(n) \rfloor} = \text{Sym}([n])$.

We finally turn to the proof of Proposition 6.4.

Proof of Proposition 6.4. We can assume that n is large enough that $m \geq (\log n)^2 > C(0.9)$, where C(0.9) is as in Lemma 3.18. Apply Lemma 3.18 with d = 0.9 and $\Sigma = \{\alpha_1, \ldots, \alpha_m\}$. We obtain a set $\Delta \subseteq \Sigma$ such that $|\Delta| \geq 0.9|\Sigma|$ and $\left(\left(A^{16m^6}\right)_{\Sigma}\right)_{(\Sigma \setminus \Delta)} |_{\Delta}$ contains $\mathrm{Alt}(\Delta)$. Let

$$B^{+} = \left\{ g \in \left(\left(A^{16m^{6}} \right)_{\Sigma} \right)_{(\Sigma \setminus \Delta)} : g|_{\Delta} \in \operatorname{Alt}(\Delta) \right\}, \quad B^{-} = \left((B^{+})^{3} \right)_{(\Delta)}.$$

This is our initial setup: we have a large set B^+ in the setwise stabilizer G_{Σ} ; furthermore, we have constructed a large subset $\Delta \subseteq \Sigma$ such that $B^+ \subseteq (G_{\Sigma})_{(\Sigma \setminus \Delta)}$ and $B^+|_{\Delta} = \operatorname{Alt}(\Delta)$. We also have a set B^- in the pointwise stabilizer $G_{(\Sigma)}$. By (6.4) with i = m + 1, $\left|\alpha_{m+1}^{B^-}\right| \ge \frac{9}{10}n$, and so $\langle B^- \rangle$ has an orbit Γ of length at least 0.9n. By Lemma 6.2, Γ is also an orbit of $\langle B^+ \rangle$.

We would like $\langle B^- \rangle$ to act as an alternating or symmetric group on Γ ; let us show that, if this is not the case, we obtain *descent*. We are assuming that Proposition 6.5 holds for n' < n (inductive hypothesis). Hence, if $\langle B^- \rangle$ has no composition factor Alt(k) with k > 0.95n, then Lemma 6.3 (descent) gives us

$$A^{\lfloor 16m^6e^{c_1(\log n)^3}\cdot F_2(0.95n)\rfloor} \supseteq \operatorname{Alt}([n]),$$

for n larger than an absolute constant, where $c_1 = c(0.9)$ is from Lemma 6.3. By (6.5), we conclude that (6.6) holds and we are done. (We are assuming that n is larger than a constant, so that $16n^6 \le e^{c_3 \log n}$, where $c_3 > 0$ will be set later.)

Thus, we can suppose from now on that $\langle B^- \rangle$ does have a composition factor $\mathrm{Alt}(k)$ for some k>0.95n. The only orbit of $\langle B^- \rangle$ that can be of length at least k is Γ , so $\langle B^- \rangle|_{\Gamma} = \langle B^-|_{\Gamma} \rangle$ must contain $\mathrm{Alt}(k)$ as a section. Hence, by Lemma 3.11, $\langle B^-|_{\Gamma} \rangle \geq \mathrm{Alt}(\Gamma)$. (We can assume 0.95n>84, and thus Lemma 3.11 does apply.) Note we also get that $|\Gamma|>0.95n$.

Now that we know that $\langle B^-|_{\Gamma}\rangle \geq \mathrm{Alt}(\Gamma)$, Corollary 4.7 gives us a small set of elements $Y=\{y_1,y_2,\ldots,y_6\}\subseteq (B^-)^{\lfloor n^{28\log n}\rfloor}$ such that $\langle Y\rangle|_{\Gamma}$ is 2-transitive on Γ . We apply Lemma 6.1 (*creation*) with $H^-=\langle B^-\rangle$, $H^+=\langle B^+\rangle$, $B=B^+$ and r=6. (The condition $H^-\lhd H^+$ is fulfilled thanks to Lemma 6.2.)

If conclusion (a) in Lemma 6.1 holds, then there is a $b \in B^+(B^+)^{-1} \setminus \{e\}$ with $\operatorname{supp}(b) \leq 0.05n$. Thm. 1.4 thus gives us that $(A \cup \{b\})^{Kn^8} \supseteq \operatorname{Alt}([n])$, where $K = K(0.1) \ge K(0.05)$ is an absolute constant. (We set K = K(0.1), instead of K = K(0.05), because we are planning to use the same constant later.) By (6.5),

$$2 \cdot 48m^6 \cdot Kn^8 < 96Kn^{14} \le F_1(n),$$

and so (provided that n is larger than a constant) (6.6) holds and we are done. (This is what we call an *exit* from the procedure.)

We can thus assume that conclusion (b) in Lemma 6.1 holds, i.e., we have created a set $W = (B^+)^{-1}YB^+ \cap \langle B^- \rangle$ with $|W| \geq |B^+|^{1/6}$. Note that $(B^+)^{-1}YB^+ \subset A^{\lfloor n^{29\log n} \rfloor}$ (for n larger than a constant) and $|B^+| \geq |\operatorname{Alt}(\Delta)| = (1/2)|\Delta|! \geq m^{0.899m}$ (for m larger than a constant; recall that $|\Delta| \geq 0.9m$). Hence

$$\left| A^{\lfloor n^{29\log n} \rfloor} \cap \langle B^- \rangle \right| \ge m^{0.149m}.$$

Now that we have *created* many elements in the pointwise stabilizer of Σ , it is our task to *organise* them: we wish to produce $\alpha_{m+2}, \ldots, \alpha_{m+\ell+1}$ satisfying (6.7).

This can be done in two ways. One is short and simple, gives a bound of $l \gg m(\log m)/(\log n)^2$, and results in a bound of $O((\log n)^5(\log \log n))$ in the exponent of the final result. The other is longer, but gives the stronger bound of $l \gg m(\log m)/(\log n)$ promised in the statement of the proposition, and results in a bound of $O((\log n)^4 \log \log n)$ in the exponent of the final result. Let us go through both arguments for the sake of clarity.

In the first argument, we simply apply Corollary 5.3 with $\operatorname{Sym}(\Gamma)$ instead of $\operatorname{Sym}([n])$ and $A^{[n^{29\log n}]} \cap \langle B^- \rangle \supset B^-$ instead of A. We obtain that any maximal sequence of elements $\alpha_{m+2}, \ldots, \alpha_{m+\ell+1}$ satisfying (6.7) must be of length

$$\gg (\log |A^{[n^{29\log n}]} \cap \langle B^- \rangle|)/(\log n)^2 \gg \frac{\log m^{0.149m}}{(\log n)^2} \gg \frac{m(\log m)}{(\log n)^2}.$$

Thus $\ell \gg m(\log m)/(\log n)^2$.

Let us now carry out the second argument in detail. The basic idea is that the creation step has given us enough elements that we can apply the organiser step several times in succession.

For $i \geq 0$, we define recursively $A_i, B_i \subseteq \langle A \rangle$ and a sequence Σ_i of points in [n]. Let $A_0 = A^{\lfloor n^{29 \log n} \rfloor}$, $m_0 = m$, $\Sigma_0 = (\alpha_1, \dots, \alpha_{m_0+1})$, and $B_0 = (A_0)_{(\Sigma_0 \setminus \{\alpha_{m_0+1}\})}$.

If A_i, Σ_i, B_i are already defined then let $A'_{i+1} = A_i^{\lfloor 9n^6 \log n \rfloor}$ and let Σ_{i+1} be a maximal extension $\Sigma_{i+1} = (\alpha_1, \dots, \alpha_{m_{i+1}+1})$ of $\Sigma_i = (\alpha_1, \dots, \alpha_{m_i+1})$ such that

$$\left|\alpha_{j}^{(A'_{i+1})_{(\alpha_{1},\ldots,\alpha_{j-1})}}\right| \geq 0.9n,$$

for all $j = 1, 2, \ldots, m_{i+1} + 1$. Finally, let

$$A_{i+1} = (A'_{i+1})^{29n^6}$$
 and $B_{i+1} = (A_{i+1})_{(\Sigma_{i+1} \setminus {\alpha_{m_{i+1}+1}})}$.

Note that for all $i \geq 0$, $\langle B_i \rangle$ has an orbit Γ_i of length at least 0.9n because $\left| \alpha_{m_i+1}^{B_i} \right| \geq$ 0.9n. (We went up to i = m+1 in condition (6.4) and up to i = m+l+1 in conclusion (6.7) (rather than i = m and i = m + l, respectively) so that we could do this useful trick!)

We stop the recursion, and set w := i for the last i for which A_i is defined, if either

- (a) $|B_i|_{\Gamma_i}| < |B_i|$, i.e., there are two elements $b_1, b_2 \in B_i$ such that $b_1b_2^{-1}$ fixes Γ_i pointwise; or
- (b) $|\Gamma_i| \leq 0.95n$ or $\langle B_i|_{\Gamma_i} \rangle \not\supset \text{Alt}(\Gamma_i)$ or (c) $n^{m_i-m_0} > \sqrt{m^{0.149m}}$.

By (6.10), we have $|B_0| \ge m^{0.149m}$.

First, we estimate the differences $m_{i+1} - m_i$. If the recursion did not stop after the definition of A_i, B_i , and Σ_i then, in particular, the stopping criterion (c) is not fulfilled at step i. Lemma 3.4, applied with $\langle B_0 \rangle$ as G, $G_{(\Sigma_i \setminus \{\alpha_{m_i}+1\})}$ as H, and B_0 as A, then implies that

$$|B_i| \ge |B_0^2 \cap H| \ge \frac{|B_0|}{n^{m_i - m_0}} \ge \sqrt{m^{0.149m}}.$$

Also, by the criteria (a) and (b), we have $|B_i|_{\Gamma_i} = |B_i|$ and $\langle B_i \rangle|_{\Gamma_i}$ acts as Alt (Γ_i) or Sym(Γ_i) on Γ_i , where $|\Gamma_i| > 0.95n$.

Since $0.9n < 0.95 \cdot 0.95n \le 0.95 |\Gamma_i|$, we can apply Corollary 5.3 with $\rho = 0.05$, $B_i|_{\Gamma_i}$ instead of A, and Γ_i instead of [n], and obtain that, for $1 \leq i < w$,

(6.12)
$$m_{i+1} - m_i > \frac{\log |B_i|}{60(\log n)^2} \ge \frac{c_2 m \log m}{60(\log n)^2},$$

where we define $c_2 := 0.149/2 = 0.0745$. (This is what we have called an organiser step. It is ultimately based on the splitting lemma (Prop. 5.2), of which Cor. 5.3 is a corollary.)

At the same time, $n^{m_{w-1}-m_0} < \sqrt{m^{0.149m}}$ implies

$$m_{w-1} - m_0 \le \frac{c_2 \log m}{\log n} m.$$

Since $m_{w-1} - m_0 = \sum_{i=1}^{w-1} (m_i - m_{i-1})$, from (6.12) it follows that

$$\frac{c_2 \log m}{\log n} m > (w - 1) \frac{c_2 m \log m}{60(\log n)^2}$$

and we conclude that $w-1 < 60 \log n$. Hence

$$A_w = A_0^{\lfloor 9n^6 \log n \rfloor^w (48n^6)^w} \subseteq A^{\lfloor n^{29 \log n} \rfloor \cdot \lfloor 432n^{12} \log n \rfloor^w} \subseteq A^{\lfloor n^{c_3 \log n} \rfloor}$$

for $c_3 := 750 > 29 + 12 \cdot 60$, provided that n is larger than an absolute constant. If $n^{m_w - m_0} > \sqrt{m^{0.149m}}$ (stopping condition (c)), then

$$m_w - m_0 \ge \frac{c_2 \log m}{\log n} m,$$

and so, setting $\ell = m_w - m_0$, we obtain (6.7).

(In other words: as long as our *organizing* has consumed less than the square-root of the material we *created*, we are organizing rapidly; if our organizing has consumed at least the square-root of the said material, then we have already organized plenty.)

If we stopped because condition (a) holds then A_w^2 contains a non-trivial element $b_1b_2^{-1}$ with support less than 0.1n. By Theorem 1.4, $(A \cup \{b_1b_2^{-1}\})^{Kn^8} \supseteq \text{Alt}([n])$, where K = K(0.1) is an absolute constant. By (6.5),

$$2 \cdot |n^{c_3 \log n}| \cdot Kn^8 \le F_1(n),$$

and so we obtain (6.6). (This is an exit case.)

Finally, suppose we stopped in case (b), i.e., $\langle B_w|_{\Gamma_w}\rangle \not\supset \mathrm{Alt}(\Gamma_w)$ or $|\Gamma_w| \leq 0.95n$. As $|\Sigma_w| \geq m > C(0.9)$, we can apply Lemma 3.18 with $\Sigma_w \setminus \alpha_{m_w+1}$ as Σ and A'_w as A, to obtain $\Delta_w \subseteq \Sigma_w \setminus \alpha_{m_w+1}$, $|\Delta_w| \geq 0.9|\Sigma_w \setminus \alpha_{m_w+1}|$ such that

$$B_w^+ = (((A_w')^{16n^6})_{\Sigma_w \setminus \{\alpha_{m_w+1}\}})_{(\Sigma_w \setminus (\{\alpha_{m_w+1}\} \cup \Delta_w))}$$

satisfies $(B_w^+)|_{\Delta_w} = \text{Alt}(\Delta_w)$. (This is a fresh setup.) Also, by Lemma 6.2, $B_w^- = ((B_w^+)^3)_{(\Delta_w)}$ generates $\langle B_w^+ \rangle_{(\Delta_w)} \lhd \langle B_w^+ \rangle$. Note that $B_w^- \subseteq B_w$ and $\langle B_w^- \rangle$ has an orbit of length at least 0.9n, simply because B_w^- contains $(A_w')_{\Sigma_w \setminus \{\alpha_{mw+1}\}}$, and the orbit of α_{mw+1} under $(A_w')_{\Sigma_w \setminus \{\alpha_{mw+1}\}}$ is of length $\geq 0.9n$ by (6.11).

We are ready for another descent. The group $\langle B_w^- \rangle$ has no composition factor $\mathrm{Alt}(k)$ with k>0.95n, because such a factor would be a section of $\langle B_w \rangle$ and Lemma 3.11 would imply that $\langle B_w|_{\Gamma_w} \rangle$ is an alternating group on >0.95n elements, in contradiction with condition (b). Thus the hypotheses of Lemma 6.3 are satisfied with $\delta=0.95$ and $\rho=0.9$ and, by the assumption that Prop. 6.5 holds for $n'\leq 0.95n < n$ (inductive hypothesis), Lemma 6.3 gives us that

$$A^{\lfloor n^{c_3 \log n} e^{c(\log n)^3} \cdot F_2(0.95n) \rfloor} \supseteq \text{Alt}([n]),$$

where c = c(0.9). We apply (6.5), and conclude that (6.6) holds.

We now use Proposition 6.5 to prove both the Main Theorem and Cor. 1.3 (for Sym(n) and Alt(n)).

Theorem 6.6. Let G = Sym(n) or Alt(n). Then

(6.13)
$$\frac{\operatorname{diam}(G) = O(e^{c(\log n)^4 \log \log n}),}{\operatorname{diam}(G) = O(e^{(c+1)(\log n)^4 \log \log n}),}$$

for an absolute constant c > 0.

As we shall see, $c_1 = 49071$ is valid (and by no means optimal).

Proof. We must find functions F_1 , F_2 satisfying (6.5) and (6.8). We can set

$$F_2(n) = e^{(\log n)^3 + 2\log n + c'c_3(\log n)^3\log\log n}F_1(n) + 2$$

for $c' > c_2$ arbitrary. Now we must make sure that

(6.14)
$$F_1(n) \ge n^{c_3 \log n} e^{c_1 (\log n)^3} \cdot \left(e^{c'c_3 (\log 0.95n)^3 \log \log 0.95n + (\log 0.95n)^3 + 2\log 0.95n} F_1(0.95n) + 2 \right).$$

(Here we can assume n > 1, so that $\log \log n$ is well-defined.) Choose $c_4 > c'c_3$. Then, for n larger than a constant n'_0 depending only on c_1 , c_3 , c' and c_4 , (6.14) will hold provided that

(6.15)
$$F_1(n) \ge e^{c_4(\log n)^3 \log \log n} \max(F_1(0.95n), 1).$$

For any $c > c_4/(4|\log 0.95|)$ and any $C \ge 1$, (6.15) is fulfilled by

$$F_1(n) = Ce^{c(\log n)^4 \log \log n},$$

provided that n is larger than a constant n_0'' depending only on c and c_4 . We set $C = n_0'''!$, where $n_0''' = \max(n_0, n_0', n_0'', 2K)$. Then (6.5) holds for all $n \ge n_0'''$, and (6.8) holds with $n_0!$ replaced by $n_0'''!$. We now apply Proposition 6.5 for our n, with n_0 replaced by n_0''' ; it uses Proposition 6.4, which in turn uses Proposition 6.5 for smaller n, and so on. The recursion ends when $n \le \max(n_0''', 1)$, as then Proposition 6.5 is trivially true (due to the bound $F_2(n) \ge n_0'''!$ in (6.8)).

We obtain that

(6.16)
$$\operatorname{diam}(\Gamma(G, Y)) \le Ce^{c(\log n)^4 \log \log n}$$

for any set Y of generators of G with $Y = Y^{-1}$, $e \in Y$. A quick calculation shows that, since $c_2 = 0.0745$ and $c_3 = 750$ (see the proof of Prop. 6.4), we can set c' = 13.423 > 1/0.0745, $c_4 = 10068 > c'c_3$ and

$$c = \left\lfloor \frac{c_4}{4|\log 0.95|} \right\rfloor = 49071.$$

Let A be an arbitrary set of generators of G. Let $Y = A \cup A^{-1} \cup \{e\}$. The undirected Cayley graph $\Gamma(G, Y)$ is just the undirected Cayley graph $\Gamma(G, A)$ with a loop at every vertex; their diameters are the same. Thus, by (6.16),

$$\operatorname{diam}(\Gamma(G,A)) = \operatorname{diam}(\Gamma(G,Y)) \le Ce^{c(\log n)^4 \log \log n}.$$

By [Bab06, Cor. 2.3],

$$\operatorname{diam}(\vec{\Gamma}(G,A)) \le O\left(\operatorname{diam}(G)(n\log n)^2\right) \le O\left(e^{(c+1)(\log n)^4\log\log n}\right).$$

References

- [Ald87] D. Aldous. On the Markov chain simulation method for uniform combinatorial distributions and simulated annealing. *Prob. Engng. Info. Sci.*, 1(1):33–46, 1987.
- [Bab91] L. Babai. Local expansion of vertex transitive graphs and random generation in finite groups. In 23rd ACM Symposium on Theory of Computing, pages 164–174. ACM Press, New York, NY, 1991.
- [Bab06] L. Babai. On the diameter of Eulerian orientations of graphs. In Proceedings of the Seventeenth Annual ACM-SIAM Symposium on Discrete Algorithms, pages 822–831, New York, 2006. ACM.
- [Bab82] L. Babai. On the order of doubly transitive permutation groups. Invent. Math., 65(3):473-484, 1981/82.
- [BBS04] L. Babai, R. Beals, and Á. Seress. On the diameter of the symmetric group: polynomial bounds. In *Proceedings of the Fifteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1108–1112 (electronic), New York, 2004. ACM.
- [BCF⁺91] L. Babai, G. Cooperman, L. Finkelstein, E. M. Luks, and Á. Seress. Fast Monte-Carlo algorithms for permutation groups. In 23rd ACM Symposium on Theory of Computing, pages 90–100. ACM Press, New York, NY, 1991.
- [BG08a] J. Bourgain and A. Gamburd. On the spectral cap for finitely generated subgroups of SU(2). Invent. Math., 171:83-121, 2008.
- [BG08b] J. Bourgain and A. Gamburd. Uniform expansion bounds for Cayley graphs of $SL_2(\mathbb{F}_p)$. Ann. of Math. (2), 167(2):625–642, 2008.
- [BGH⁺] J. Bamberg, N. Gill, T. Hayes, H. A. Helfgott, G. Royle, Á. Seress, and P. Spiga. Bounds on the diameter of Cayley graphs of the symmetric group. Submitted. Available as arxiv.org:1205.1596 since 2012.
- [BGS10] J. Bourgain, A. Gamburd, and P. Sarnak. Affine linear sieve, expanders, and sumproduct. Invent. Math., 179(3):559-644, 2010.
- [BGS11] J. Bourgain, A. Gamburd, and P. Sarnak. Generalization of Selberg's $\frac{3}{16}$ theorem and affine sieve. *Acta Math.*, 207(2):255–290, 2011.
- [BGT11] E. Breuillard, B. Green, and T. Tao. Approximate subgroups of linear groups. *Geom. Funct. Anal.*, 21(4):774–819, 2011.
- [BH05] L. Babai and T. Hayes. Near-independence of permutations and an almost sure polynomial bound on the diameter of the symmetric group. In Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms, pages 1057–1066. ACM, New York, 2005.
- [BHK⁺90] L. Babai, G. Hetyei, W. M. Kantor, A. Lubotzky, and Á. Seress. On the diameter of finite groups. In 31st Annual Symposium on Foundations of Computer Science, Vol. I, II (St. Louis, MO, 1990), pages 857–865. IEEE Comput. Soc. Press, Los Alamitos, CA, 1990.
- [BLS87] L. Babai, E. M. Luks, and Á. Seress. Permutation groups in NC. In Proceedings of the Nineteenth Annual ACM Symposium on the Theory of Computing, pages 409–420, New York, 1987. ACM.
- [BLS88] L. Babai, E. M. Luks, and Á. Seress. Fast management of permutation groups. In Proceedings of the 29th IEEE Symposium on Foundations of Computer Science, pages 272–282, Singer Island, FL, 1988. IEEE Computer Society Press.
- [Boc89] A. Bochert. Ueber die Zahl der verschiedenen Werthe, die eine Function gegebener Buchstaben durch Vertauschung derselben erlangen kann. *Math. Ann.*, 33:584–590, 1889.
- [BS87] L. Babai and Á. Seress. On the degree of transitivity of permutation groups: a short proof. J. Combin. Theory Ser. A, 45(2):310–315, 1987.
- [BS88] L. Babai and A. Seress. On the diameter of Cayley graphs of the symmetric group. J. Combin. Theory Ser. A, 49(1):175–179, 1988.

- [BS92] L. Babai and Á. Seress. On the diameter of permutation groups. *European J. Combin.*, 13(4):231–243, 1992.
- [Din11] O. Dinai. Growth in SL₂ over finite fields. J. Group Theory, 14(2):273–297, 2011.
- [DM96] J. D. Dixon and B. Mortimer. Permutation Groups, volume 163 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1996.
- [DSC93] P. Diaconis and L. Saloff-Coste. Comparison techniques for random walk on finite groups. $Ann.\ Probab.,\ 21(4):2131-2156,\ 1993.$
- [Fie72] M. Fiedler. Bounds for eigenvalues of doubly stochastic matrices. Linear Algebra and Appl., 5:299–310, 1972.
- [Gan91] A. Gangolli. Convergence bounds for Markov chains and applications to sampling. PhD thesis, Dept. Computer Science, Stanford Univ., 1991.
- [GH] N. Gill and H. A. Helfgott. Growth in solvable subgroups of $GL_r(\mathbb{Z}/p\mathbb{Z})$. Submitted. Available as arxiv.org:1008.5264 since 2010.
- [GH11] N. Gill and H. A. Helfgott. Growth of small generating sets in $SL_n(\mathbb{Z}/p\mathbb{Z})$. Int. Math. Res. Not. IMRN, (18):4226-4251, 2011.
- [GV12] A. S. Golsefidy and P. P. Varjú. Expansion in perfect groups. *Geom. Funct. Anal.*, 22(6):1832–1891, 2012.
- [Hel08] H. A. Helfgott. Growth and generation in $SL_2(\mathbb{Z}/p\mathbb{Z})$. Ann. of Math. (2), 167(2):601–623, 2008.
- [Hell1] H. A. Helfgott. Growth in $SL_3(\mathbb{Z}/p\mathbb{Z})$. J. Eur. Math. Soc. (JEMS), 13(3):761–851, 2011.
- [HLW06] Sh. Hoory, N. Linial, and A. Wigderson. Expander graphs and their applications. *Bull. Amer. Math. Soc. (N.S.)*, 43(4):439–561 (electronic), 2006.
- [Hru12] E. Hrushovski. Stable group theory and approximate subgroups. J. Amer. Math. Soc., 25(1):189–243, 2012.
- [Jor70] C. Jordan. Traité des substitutions et des équations algébriques. Gauthier-Villars, Paris, 1870. Reprinted: 1957, Paris: Albert Blanchard.
- [KMS84] D. Kornhauser, G. Miller, and P. Spirakis. Coordinating pebble motion on graphs, the diameter of permutation groups, and applications. In *Proceedings of the 25th IEEE* Symposium on Foundations of Computer Science, pages 241–250, Singer Island, FL, 1984. IEEE Computer Society Press.
- [Lie83] M. W. Liebeck. On graphs whose full automorphism group is an alternating group or a finite classical group. Proc. London Math. Soc. (3), 47(2):337–362, 1983.
- [Lov96] L. Lovász. Random walks on graphs: a survey. In *Combinatorics, Paul Erdős is eighty, Vol. 2 (Keszthely, 1993)*, volume 2 of *Bolyai Soc. Math. Stud.*, pages 353–397. János Bolyai Math. Soc., Budapest, 1996.
- [LP11] M. J. Larsen and R. Pink. Finite subgroups of algebraic groups. J. Amer. Math. Soc., 24(4):1105–1158, 2011.
- [McK84] P. McKenzie. Permutations of bounded degree generate groups of polynomial diameter. Inform. Process. Lett., 19(5):253–254, 1984.
- [Moh91] B. Mohar. Eigenvalues, diameter, and mean distance in graphs. *Graphs Combin.*, 7(1):53–64, 1991.
- [MS77] F. J. MacWilliams and N. J. A. Sloane. The theory of error-correcting codes. North-Holland Publishing Co., Amsterdam, 1977. North-Holland Mathematical Library, Vol. 16.
- [Pak] I. Pak. Problems: new, old and unusual. Talk at National University of Ireland, Galway, Ireland, Dec 1, 2009. Available as http://larmor.nuigalway.ie/~detinko/Igor.pdf.
- [PPSS12] C. E. Praeger, L. Pyber, P. Spiga, and E. Szabó. Graphs with automorphism groups admitting composition factors of bounded rank. Proc. Amer. Math. Soc., 140(7):2307– 2318, 2012.
- [PS] L. Pyber and E. Szabó. Growth in finite simple groups of Lie type of bounded rank. Submitted. Available as arxiv.org:1005.1881 since 2010.
- [PS80] Ch. E. Praeger and J. Saxl. On the orders of primitive permutation groups. Bull. London Math. Soc., 12(4):303–307, 1980.

- [Pyb93] L. Pyber. On the orders of doubly transitive permutation groups, elementary estimates. J. Combin. Theory Ser. A, 62(2):361–366, 1993.
- [RT85] I. Z. Ruzsa and S. Turjányi. A note on additive bases of integers. Publ. Math. Debrecen, 32(1-2):101-104, 1985.
- [Ser03] Á. Seress. Permutation Group Algorithms, volume 152 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2003.
- [Sim70] Ch. C. Sims. Computational methods in the study of permutation groups. In *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, pages 169–183. Pergamon, Oxford, 1970.
- [Sim71] Ch. C. Sims. Computation with permutation groups. In Proc. Second Symposium on Symbolic and Algebraic Manipulation, pages 23–28. ACM Press, New York, NY, 1971.
- [SSV05] B. Sudakov, E. Szemerédi, and V. H. Vu. On a question of Erdős and Moser. Duke Math. J., 129(1):129-155, 2005.
- [Tao08] T. Tao. Product set estimates for non-commutative groups. Combinatorica, 28(5):547–594, 2008.
- [Var12] P. P. Varjú. Expansion in $SL_d(\mathcal{O}_K/I)$, I square-free. J. Eur. Math. Soc. (JEMS), 14(1):273–305, 2012.
- [Wie64] H. Wielandt. Finite Permutation Groups. Translated from the German by R. Bercov. Academic Press, New York, 1964.

HARALD A. HELFGOTT, ÉCOLE NORMALE SUPÉRIEURE, DÉPARTEMENT DE MATHÉMATIQUES, 45 RUE D'ULM, F-75230 PARIS, FRANCE

E-mail address: harald.helfgott@ens.fr

ÁKOS SERESS, CENTRE FOR THE MATHEMATICS OF SYMMETRY AND COMPUTATION, THE UNIVERSITY OF WESTERN AUSTRALIA, CRAWLEY, WA 6009 AUSTRALIA, AND DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, USA