

# SOME EXAMPLES OF SELF-SIMILAR SOLUTIONS AND TRANSLATING SOLITONS FOR LAGRANGIAN MEAN CURVATURE FLOW

Hiroshi Nakahara

Department of Mathematics  
Tokyo Institute of Technology  
2-21-1 O-okayama, Meguro  
Tokyo 152-8551 Japan

*E-mail address:* nakahara.h.ab@m.titech.ac.jp

November 27, 2024

## Abstract

We construct new examples of self-similar solutions and translating solitons for Lagrangian mean curvature flow by extending the method of Joyce, Lee and Tsui [3]. Those examples include examples in which the Lagrangian angle is arbitrarily small as the examples of Joyce, Lee and Tsui [3].

## 1 Introduction

In recent years the Lagrangian mean curvature flow has been extensively studied, as it is a key ingredient in the Strominger-Yau-Zaslow Conjecture and Thomas-Yau Conjecture. Strominger-Yau-Zaslow Conjecture explains Mirror Symmetry of Calabi-Yau 3-folds. In Joyce, Lee and Tsui [3], many examples of self-similar solutions and translating solitons for Lagrangian mean curvature flow are constructed. Those Lagrangian submanifold  $L$  are the total space of a 1-parameter family of quadrics  $Q_s$ ,  $s \in I$ , where  $I$  is an open interval in  $\mathbb{R}$ . In this paper, we construct new examples of those Lagrangian submanifolds that link up with the examples of Lagrangian submanifolds given in [1], [2], [3], [4] and so on. To do so we improve theorems in [3] by describing Lagrangian submanifolds of the forms of Ansatz 3.1 and Ansatz 3.3 in [3].

Let  $L$  be a Lagrangian submanifold in  $\mathbb{C}^n$ . Define a function  $c : L \rightarrow \mathbb{C}$  by the relation that

$$dz_1 \wedge \cdots \wedge dz_n|_L \equiv c \cdot \text{vol}_L$$

where  $\text{vol}_L$  is the volume form of  $L$ . Then  $|c| \equiv 1$  holds. So we can define *Lagrangian angle*  $\theta : L \rightarrow \mathbb{R}$  or  $\theta : L \rightarrow \mathbb{R}/2\pi\mathbb{Z}$  by the relation that

$$dz_1 \wedge \cdots \wedge dz_n|_L \equiv e^{i\theta} \text{vol}_L.$$

On a Lagrangian submanifold  $L$  in  $\mathbb{C}^n$ , the mean curvature vector  $H$  is given by

$$H = J\nabla\theta, \quad (1)$$

where  $\nabla$  is the gradient on  $L$  and  $J$  is the standard complex structure in  $\mathbb{C}^n$ . The proof of (1) is given in [5].

**Definition 1.1.** Let  $L \subset \mathbb{R}^N$  be a submanifold in  $\mathbb{R}^N$ .  $L$  is called a *self-similar solution* if  $H \equiv \alpha F^\perp$  on  $L$  for some constant  $\alpha \in \mathbb{R}$ , where  $F^\perp$  is the orthogonal projection of the position vector  $F$  in  $\mathbb{R}^N$  to the normal bundle of  $L$ , and  $H$  is the mean curvature vector of  $L$  in  $\mathbb{R}^N$ . It is called a *self-shrinker* if  $\alpha < 0$  and a *self-expander* if  $\alpha > 0$ . On the other hand  $L \subset \mathbb{R}^N$  is called a *translating soliton* if there exists a constant vector  $T$  in  $\mathbb{R}^N$  such that  $H \equiv T^\perp$ , where  $T^\perp$  is the orthogonal projection of the constant vector  $T$  in  $\mathbb{R}^N$  to the normal bundle of  $L$  and  $H$  is the mean curvature vector of  $L$  in  $\mathbb{R}^N$ . We call  $T$  a *translating vector*.

It is well-known that if  $F$  is a self-similar solution then  $F_t = \sqrt{2\alpha t}F$  is moved by the mean curvature flow, and if  $F$  is a translating soliton then  $F_t = F + tT$  is also moved by the mean curvature flow.

First we consider self-similar solutions.

**Theorem 1.2.** Let  $B, C, \lambda_1, \dots, \lambda_n \in \mathbb{R} - \{0\}$ ,  $E > 1$ ,  $a_1, \dots, a_n > 0$ , and  $\alpha, \psi_1, \dots, \psi_n \in \mathbb{R}$  be constants. Let  $I \subset \mathbb{R}$  be a connected open neighborhood of  $0 \in \mathbb{R}$  such that  $E\{\prod_{k=1}^n (1 + 2a_k B \lambda_k s)\}e^{2B\alpha s} - 1$  and  $1/a_j + 2\lambda_j B s$  are positive for any  $j \in \{1, \dots, n\}$  and any  $s \in I$ . Define  $r_1, \dots, r_n : I \rightarrow \mathbb{R}$  by

$$r_j = \sqrt{\frac{1}{a_j} + 2\lambda_j B s}.$$

Define  $\phi_1, \dots, \phi_n : I \rightarrow \mathbb{R}$  by

$$\phi_j = \psi_j + \int_0^s \frac{\lambda_j |B|}{\left(\frac{1}{a_j} + 2\lambda_j B t\right) \sqrt{E\{\prod_{k=1}^n (1 + 2a_k B \lambda_k t)\}e^{2B\alpha t} - 1}} dt.$$

Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1 r_1(s) e^{i\phi_1(s)}, \dots, x_n r_n(s) e^{i\phi_n(s)}) \mid \sum_{j=1}^n \lambda_j x_j^2 = C, x_j \in \mathbb{R}, s \in I\}$$

is an immersed Lagrangian submanifold, and its position vector  $F$  and mean curvature vector  $H$  satisfy  $CH \equiv \alpha F^\perp$ .

*Remark 1.2.1.* In the situation of Theorem 1.2, let  $I' \subset I$  be a subinterval such that  $r_j$  has positive lower and upper bounds. Put  $\alpha \neq 0, n \geq 2, \lambda_j > 0$  for  $1 \leq j \leq k < n$  and  $\lambda_j < 0$  for  $k < j \leq n$  where  $k$  is a positive integer less than  $n$ . Let  $t \in \mathbb{R}$  be a constant. Define

$$L_t = \{(x_1 r_1(s) e^{i\phi_1(s)}, \dots, x_n r_n(s) e^{i\phi_n(s)}) \mid \sum_{j=1}^n \lambda_j x_j^2 = 2\alpha t, x_j \in \mathbb{R}, s \in I'\}.$$

The fact that the varifold  $\cup_t L_t, -\infty < t < \infty$ , forms an eternal solution for Brakke flow without mass loss is proved similarly to Lee and Wang [4]. So see Lee and Wang [4]. By Theorem 1.2,  $L_t$  is a Lagrangian self-shrinker if  $t < 0$ , a Lagrangian self-expander if  $t > 0$ , and a Lagrangian cone if  $t = 0$ .

**Theorem 1.3.** Let  $a_j > 0, \psi_j \in \mathbb{R}, @E > 1$ , and  $\alpha \geq 0$  be constants. Define  $r_j(s) : \mathbb{R} \rightarrow \mathbb{R}$  by  $r_j(s) = \sqrt{\frac{1}{a_j} + s^2}$ . Define  $\phi_j(s) : \mathbb{R} \rightarrow \mathbb{R}$  by

$$\phi_j(s) = \psi_j + \int_0^s \frac{|t|}{\left(\frac{1}{a_j} + t^2\right) \sqrt{E \left\{ \prod_{k=1}^n (1 + a_k t^2) \right\} e^{\alpha t^2} - 1}} dt. \quad (2)$$

Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1 r_1(s) e^{i\phi_1(s)}, \dots, x_n r_n(s) e^{i\phi_n(s)}) \mid \sum_{j=1}^n x_j^2 = 1, x_j \in \mathbb{R}, s \in \mathbb{R}, s \neq 0\}$$

is an immersed Lagrangian diffeomorphic to  $(\mathbb{R} - \{0\}) \times S^{n-1}$ , and its position vector  $F$  and mean curvature vector  $H$  satisfy  $H \equiv \alpha F^\perp$ .

*Remark 1.3.1.* The manifold obtained as  $\lim_{E \rightarrow 1+0} L$  is the same as Theorem C in [3]. So the condition  $s \neq 0$  on the definition of  $L$  is not necessary if  $E = 1$ . If we put  $E = 1$  then changing  $0 \mapsto -\infty$  in the integral of (2) gives the example of [2].

*Remark 1.3.2.* Define  $\bar{\phi}_1, \dots, \bar{\phi}_n > 0$  by

$$\bar{\phi}_j = \int_0^\infty \frac{|t|}{\left(\frac{1}{a_j} + t^2\right) \sqrt{E \left\{ \prod_{k=1}^n (1 + a_k t^2) \right\} e^{\alpha t^2} - 1}} dt.$$

We put  $\alpha > 0$ . From the proof of Theorem 1.3 and (6) in § 2 the Lagrangian angle  $\theta$  satisfies

$$\begin{aligned} \theta &= \sum_j \phi_j + \arg\left(s + i \frac{|s|}{\sqrt{E \left\{ \prod_{k=1}^n (1 + a_k s^2) \right\} e^{\alpha s^2} - 1}}\right) \quad \text{and} \\ \dot{\theta} &= \frac{-\alpha |s|}{\sqrt{E \left\{ \prod_{k=1}^n (1 + a_k s^2) \right\} e^{\alpha s^2} - 1}}. \end{aligned} \quad (3)$$

It follows that  $\theta$  is strictly decreasing. We define the submanifold  $L_1 \subset L$  by restricting  $s > 0$  and  $L_2 \subset L$  by restricting  $s < 0$ . Therefore we have  $L = L_1 \cup L_2$ . We rewrite  $\theta_1, \theta_2$  as the Lagrangian angle of  $L_1, L_2$  respectively. Then  $\lim_{s \rightarrow +\infty} \theta_1(s) < \theta_1(s) < \lim_{s \rightarrow 0+0} \theta_1(s)$  and  $\lim_{s \rightarrow 0-0} \theta_2(s) < \theta_2(s) < \lim_{s \rightarrow -\infty} \theta_2(s)$ . So from the first equation of (3) we have

$$\sum_j \psi_j + \sum_j \bar{\phi}_j < \theta_1 < \sum_j \psi_j + \tan^{-1} \frac{1}{\sqrt{E-1}}$$

and

$$\sum_j \psi_j + \pi - \tan^{-1} \frac{1}{\sqrt{E-1}} < \theta_2 < \sum_j \psi_j + \pi - \sum_j \bar{\phi}_j.$$

Therefore by choosing  $\tan^{-1}(1/\sqrt{E-1})$  close to 0, that is, choosing  $E$  close to  $\infty$ , the oscillation of the Lagrangian angle of  $L_1, L_2$  can be made arbitrarily small. Furthermore  $\Phi : (a_1, \dots, a_n) \mapsto (\bar{\phi}_1, \dots, \bar{\phi}_n)$  gives a diffeomorphism

$$\Phi : (0, \infty)^n \rightarrow \{(\bar{\phi}_1, \dots, \bar{\phi}_n) \in (0, \tan^{-1} \frac{1}{\sqrt{E-1}})^n \mid 0 < \sum_j \bar{\phi}_j < \tan^{-1} \frac{1}{\sqrt{E-1}}\}.$$

We can prove that  $\Phi$  is a diffeomorphism similarly to the proof of Theorem D in [3]. So we omit the proof. Therefore by choosing  $\sum_j \bar{\phi}_j$  close to  $\tan^{-1}(1/\sqrt{E-1})$ , the oscillation of the Lagrangian angle of  $L_1, L_2$  can also be made arbitrarily small.

*Remark 1.3.3.* If we put  $B = 1/2, C = \lambda_1 = \dots = \lambda_n = 1, \alpha \geq 0$  in the situation of Theorem 1.2 then  $L$  is the same as  $L$  in Theorem 1.3 where  $s > 0$ .

Next we turn to translating solitons.

**Theorem 1.4.** *Let  $B, \lambda_1, \dots, \lambda_{n-1} \in \mathbb{R} - \{0\}, E > 1, a_1, \dots, a_{n-1} > 0$ , and  $\alpha, \psi_1, \dots, \psi_{n-1} \in \mathbb{R}, K \in \mathbb{C}$  be constants. Let  $I \subset \mathbb{R}$  be a connected open neighborhood of  $0 \in \mathbb{R}$  such that  $E\{\prod_{k=1}^{n-1} (1 + 2a_k B \lambda_k s)\} e^{2B\alpha s} - 1$  and  $1/a_j + 2\lambda_j B s$  are positive for any  $j \in \{1, \dots, n-1\}$  and any  $s \in I$ . Define  $r_1, \dots, r_{n-1} : I \rightarrow \mathbb{R}$  by*

$$r_j = \sqrt{\frac{1}{a_j} + 2\lambda_j B s}.$$

Define  $\phi_1, \dots, \phi_{n-1} : I \rightarrow \mathbb{R}$  by

$$\phi_j = \psi_j + \int_0^s \frac{\lambda_j |B|}{(\frac{1}{a_j} + 2\lambda_j B t) \sqrt{E\{\prod_{k=1}^{n-1} (1 + 2a_k B \lambda_k t)\} e^{2B\alpha t} - 1}} dt.$$

Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1 r_1(s) e^{i\phi_1(s)}, \dots, x_{n-1} r_{n-1}(s) e^{i\phi_{n-1}(s)}, -\frac{1}{2} \sum_{j=1}^{n-1} \lambda_j x_j^2 + Bs + i|B| \int_0^s \frac{dt}{\sqrt{E\{\prod_{k=1}^{n-1} (1 + 2a_k B \lambda_k t)\} e^{2B\alpha t} - 1}} + K) | x_1, \dots, x_{n-1} \in \mathbb{R}, s \in I\}$$

is an immersed Lagrangian submanifold, and its mean curvature vector  $H$  satisfy  $H \equiv T^\perp$  where  $T = (0, \dots, 0, \alpha) \in \mathbb{C}^n$ .

**Theorem 1.5.** Let  $a_1, \dots, a_{n-1} > 0$ ,  $\psi_1, \dots, \psi_{n-1} \in \mathbb{R}, E > 1$ , and  $\alpha \geq 0$  be constants. Define  $r_j(s) : \mathbb{R} \rightarrow \mathbb{R}$  by  $r_j(s) = \sqrt{\frac{1}{a_j} + s^2}$ . Define  $\phi_j(s) : \mathbb{R} \rightarrow \mathbb{R}$  by

$$\phi_j(s) = \psi_j + \int_0^s \frac{|t|}{(\frac{1}{a_j} + t^2) \sqrt{E\{\prod_{k=1}^{n-1} (1 + a_k t^2)\} e^{\alpha t^2} - 1}} dt.$$

Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1 r_1(s) e^{i\phi_1(s)}, \dots, x_{n-1} r_{n-1}(s) e^{i\phi_{n-1}(s)}, -\frac{1}{2} \sum_{j=1}^{n-1} x_j^2 + \frac{1}{2} s^2 + i \int_0^s \frac{|t| dt}{\sqrt{E\{\prod_{k=1}^{n-1} (1 + a_k t^2)\} e^{\alpha t^2} - 1}}) | x_1 \dots, x_{n-1} \in \mathbb{R}, s \in \mathbb{R}, s \neq 0\}$$

is an immersed Lagrangian diffeomorphic to  $(\mathbb{R} - \{0\}) \times \mathbb{R}^{n-1}$ , and its mean curvature vector  $H$  satisfy  $H \equiv T^\perp$  where  $T = (0, \dots, 0, \alpha) \in \mathbb{C}^n$ .

*Remark 1.5.1.* If we put  $\psi_1 = \dots = \psi_{n-1} = 0$  then the manifold obtained as  $\lim_{E \rightarrow 1+0} L$  is the same as Theorem G in [3]. So the condition  $s \neq 0$  on the definition of  $L$  is not necessary if  $E = 1$ .

*Remark 1.5.2.* We define the submanifold  $L_1 \subset L$  by restricting  $s > 0$  and  $L_2 \subset L$  by restricting  $s < 0$ . Similarly to Remark 1.3.2 if we fix  $\alpha > 0$  then by choosing the parameters  $a_1, \dots, a_n > 0$  or  $E > 1$ , the oscillation of the Lagrangian angle of  $L_1, L_2$  can be made arbitrarily small.

*Remark 1.5.3.* If we put  $B = 1/2$ ,  $\lambda_1 = \dots = \lambda_{n-1} = 1$ ,  $K = 0$ ,  $\alpha \geq 0$  in the situation of Theorem 1.4 then  $L$  is the same as  $L$  in Theorem 1.5 where  $s > 0$ .

## 2 Proofs for self-similar solutions

In order to prove Theorem 1.2 and Theorem 1.3 we use the following Theorem 2.1 and Theorem 2.3 that are a slight generalization of Theorem A and Theorem B in [3]. The following Theorem 2.1 sets up the ordinary differential equations for immersed Lagrangian submanifolds diffeomorphic to  $\mathcal{S}^1 \times \mathcal{S}^{m-1} \times \mathbb{R}^{n-m}$  or  $\mathbb{R} \times \mathcal{S}^{m-1} \times \mathbb{R}^{n-m}$  where  $1 \leq m \leq n$ .

**Theorem 2.1.** *Let  $I \subset \mathbb{R}$  be an open interval. Let  $\lambda_1, \dots, \lambda_n, C \in \mathbb{R} - \{0\}$  be constants, and  $\omega_1, \dots, \omega_n : I \rightarrow \mathbb{C} - \{0\}$  and  $f : I \rightarrow \mathbb{C} - \{0\}$  be smooth functions. Suppose that*

$$\frac{d\omega_j}{ds} = \frac{\lambda_j f}{\omega_j}, \quad j = 1, \dots, n \quad (4)$$

hold in  $I$ . Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1\omega_1(s), \dots, x_n\omega_n(s)) \mid \sum_{j=1}^n \lambda_j x_j^2 = C, x_j \in \mathbb{R}, s \in I\} \quad (5)$$

is a Lagrangian submanifold in  $\mathbb{C}^n$ , with Lagrangian angle

$$\theta(s) = \arg(\omega_1 \cdots \omega_n f) \quad (6)$$

at  $(x_1\omega_1(s), \dots, x_n\omega_n(s)) \in L$ , so  $\theta$  is a function depending only on  $s$ , not on  $x_1, \dots, x_n$ . Further we have

$$H = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right)$$

and

$$F^\perp = -\frac{C \operatorname{Im}(f)}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right).$$

*Remark 2.1.1.* A direct calculation shows that

$$\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle = \sum_{j=1}^n \frac{\lambda_j^2 x_j^2 |f|^2}{|\omega_j|^2}.$$

*Remark 2.1.2.* Let  $\tilde{\theta} : I \rightarrow \mathbb{R}$  be a function satisfying

$$\frac{d\tilde{\theta}}{ds} = -\alpha \operatorname{Im}(e^{i\tilde{\theta}} \overline{\omega_1 \cdots \omega_n}).$$

Put

$$f = e^{i\tilde{\theta}} \overline{\omega_1 \cdots \omega_n}.$$

Then it is proved in [3] that  $L$  is a Lagrangian submanifold and  $L$  is a self-similar solution. This is Theorem A in [3].

**Corollary 2.2.** *Let  $I \subset \mathbb{R}$  be an open interval. Let  $\lambda_1, \dots, \lambda_n, C \in \mathbb{R} - \{0\}$  be constants, and  $\omega_1, \dots, \omega_n : I \rightarrow \mathbb{C} - \{0\}$  and  $f : I \rightarrow \mathbb{C} - \{0\}$  be smooth functions. Suppose that*

$$\frac{d\omega_j}{ds} = \frac{\lambda_j f}{\bar{\omega}_j}, \quad j = 1, \dots, n$$

and

$$\frac{d}{ds} \arg(\omega_1 \cdots \omega_n f) = -\alpha \operatorname{Im}(f)$$

hold in  $I$ . Then the submanifold  $L_C$  in  $\mathbb{C}^n$  given by

$$L_C = \{(x_1 \omega_1(s), \dots, x_n \omega_n(s)) \mid \sum_{j=1}^n \lambda_j x_j^2 = C, x_j \in \mathbb{R}, s \in I\}$$

is a Lagrangian submanifold in  $\mathbb{C}^n$ , and its position vector  $F$  and mean curvature vector  $H$  satisfy  $CH \equiv \alpha F^\perp$ . Moreover, If  $|\omega_j|$  has positive lower and upper bounds and  $\alpha \in \mathbb{R} - \{0\}, n \geq 2, \lambda_j > 0$  for  $1 \leq j \leq k < n$  and  $\lambda_j < 0$  for  $k < j \leq n$  where  $k$  is a positive integer less than  $n$  then the varifold  $\cup_t L_{2\alpha t}, -\infty < t < \infty$ , forms an eternal solution for Brakke flow without mass loss. The fact is proved similarly to Lee and Wang [4]. Then  $L_{2\alpha t}$  is a Lagrangian self-shrinker if  $t < 0$ , a Lagrangian self-expander if  $t > 0$ , and a Lagrangian cone if  $t = 0$ .

The following Theorem 2.3 gives the solution to ordinary differential equations in Theorem 2.1.

**Theorem 2.3.** *In the situation of Theorem 2.1 write  $\omega_j := r_j e^{i\phi_j}$ , for functions  $r_1, \dots, r_n : I \rightarrow (0, \infty), \phi_1, \dots, \phi_n : I \rightarrow \mathbb{R}$ . Fix  $s_0 \in I$ . Define  $u : I \rightarrow \mathbb{R}$  by*

$$u(s) := 2 \int_{s_0}^s \operatorname{Re}(f(t)) dt.$$

Then we have

$$r_j^2 = \alpha_j + \lambda_j u, \quad \phi_j = \psi_j + \int_{s_0}^s \frac{\lambda_j \operatorname{Im}(f(t))}{\alpha_j + \lambda_j u(t)} dt,$$

with  $\alpha_j = r_j^2(s_0), \quad \psi_j = \phi_j(s_0)$ .

*Remark 2.3.1.* Thus if  $f$  is explicitly given, the ordinary differential equation (4) is solved by Theorem 2.3.

*Proof of Theorem 2.1.* If we set

$$x_n = \sqrt{\frac{1}{\lambda_n} (C - \lambda_1 x_1^2 - \dots - \lambda_{n-1} x_{n-1}^2)},$$

then  $(x_1, \dots, x_{n-1}, s)$  is a coordinate of  $L$ . With this coordinate we have

$$\begin{aligned}
& (dz_1 \wedge d\bar{z}_1 + \dots + dz_n \wedge d\bar{z}_n)|_L \\
&= (\omega_1 dx_1 + x_1 \dot{\omega}_1 ds) \wedge (\bar{\omega}_1 dx_1 + x_1 \bar{\dot{\omega}}_1 ds) + \dots + \\
& \quad (\omega_{n-1} dx_{n-1} + x_{n-1} \dot{\omega}_{n-1} ds) \wedge (\bar{\omega}_{n-1} dx_{n-1} + x_{n-1} \bar{\dot{\omega}}_{n-1} ds) + \\
& \quad \left(-\frac{\lambda_1 x_1}{\lambda_n x_n} \omega_n dx_1 - \dots - \frac{\lambda_{n-1} x_{n-1}}{\lambda_n x_n} \omega_n dx_{n-1} + x_n \dot{\omega}_n ds\right) \wedge \\
& \quad \left(-\frac{\lambda_1 x_1}{\lambda_n x_n} \bar{\omega}_n dx_1 - \dots - \frac{\lambda_{n-1} x_{n-1}}{\lambda_n x_n} \bar{\omega}_n dx_{n-1} + x_n \bar{\dot{\omega}}_n ds\right) \\
&= 2\text{Im}(x_1 \omega_1 \bar{\dot{\omega}}_1) dx_1 \wedge ds + \dots + 2\text{Im}(x_{n-1} \omega_{n-1} \bar{\dot{\omega}}_{n-1}) dx_{n-1} \wedge ds \\
& \quad + 2\text{Im}\left(-\frac{\lambda_1}{\lambda_n} x_1 \omega_n \bar{\dot{\omega}}_n\right) dx_1 \wedge ds + \dots + 2\text{Im}\left(-\frac{\lambda_{n-1}}{\lambda_n} x_{n-1} \omega_n \bar{\dot{\omega}}_n\right) dx_{n-1} \wedge ds \\
&= 0,
\end{aligned}$$

and

$$\begin{aligned}
& dz_1 \wedge \dots \wedge dz_n|_L \\
&= \begin{vmatrix} \omega_1 & \dots & \dots & x_1 \dot{\omega}_1 \\ & \ddots & 0 & \vdots \\ 0 & & \omega_{n-1} & x_{n-1} \dot{\omega}_{n-1} \\ -\frac{\lambda_1 x_1}{\lambda_n x_n} \omega_n & \dots & -\frac{\lambda_{n-1} x_{n-1}}{\lambda_n x_n} \omega_n & x_n \dot{\omega}_n \end{vmatrix} dx_1 \wedge \dots \wedge dx_{n-1} \wedge ds \\
&= \frac{\omega_1 \dots \omega_n}{\lambda_n x_n} \left( \sum_{j=1}^n \frac{\lambda_j x_j^2 \dot{\omega}_j}{\omega_j} \right) dx_1 \wedge \dots \wedge dx_{n-1} \wedge ds \\
&= \frac{\omega_1 \dots \omega_n f}{\lambda_n x_n} \sum_j \frac{\lambda_j^2 x_j^2}{|\omega_j|^2} dx_1 \wedge \dots \wedge dx_{n-1} \wedge ds.
\end{aligned}$$

It follows that  $L$  is a nonsingular immersed Lagrangian, with Lagrangian angle

$$\theta(s) = \arg(\omega_1 \dots \omega_n f)$$

at  $(x_1 \omega_1(s), \dots, x_n \omega_n(s))$ . Since

$$d\theta = \dot{\theta} ds,$$

and

$$\begin{aligned}
\left\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial x_j} \right\rangle &= \langle (x_1 \dot{\omega}_1(s), \dots, x_n \dot{\omega}_n(s)), (0, \dots, 0, \omega_j, 0, \dots, 0, -\frac{\lambda_j x_j}{\lambda_n x_n} \omega_n) \rangle \\
&= \text{Re}(x_j \dot{\omega}_j \bar{\omega}_j - \frac{\lambda_j}{\lambda_n} x_j \dot{\omega}_n \bar{\omega}_n) \\
&= x_j \lambda_j \text{Re}(f - f) \\
&= 0
\end{aligned}$$



for any  $1 \leq j \leq n-1$ , it follows that

$$\nabla\theta = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} \left( \frac{\partial}{\partial s} \right).$$

Therefore we obtain

$$H = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right).$$

The normal projection of the position vector  $F$  is computed by

$$\begin{aligned} \langle F, J \frac{\partial}{\partial x_j} \rangle &= \langle (x_1\omega_1, \dots, x_n\omega_n), i(0, \dots, 0, \omega_j, 0, \dots, 0, -\frac{\lambda_j x_j}{\lambda_n x_n} \omega_n) \rangle \\ &= 0, \end{aligned}$$

and

$$\begin{aligned} \langle F, J \frac{\partial}{\partial s} \rangle &= \langle (x_1\omega_1, \dots, x_n\omega_n), i(x_1\dot{\omega}_1(s), \dots, x_n\dot{\omega}_n(s)) \rangle \\ &= \operatorname{Re}(\sum x_j^2 \bar{\omega}_j \cdot i\dot{\omega}_j) \\ &= -C \operatorname{Im}(f). \end{aligned}$$

It follows that

$$F^\perp = -\frac{C \operatorname{Im}(f)}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right).$$

This completes the proof of Theorem 2.1. □

*Proof of Theorem 2.3.* Since we have

$$\frac{dr_j^2}{ds} = \frac{d}{ds}(\omega_j \bar{\omega}_j) = \lambda_j f + \lambda_j \bar{f} = 2\lambda_j \operatorname{Re}(f),$$

we obtain

$$r_j^2 = \alpha_j + \lambda_j u.$$

So we have

$$\dot{r}_j = \frac{\lambda_j \operatorname{Re} f}{r_j}.$$

By our assumption we have

$$\dot{\omega}_j = \frac{\lambda_j f}{\bar{\omega}_j}.$$

This is equivalent to

$$\dot{r}_j e^{i\phi_j} + r_j i \dot{\phi}_j e^{i\phi_j} = \frac{\lambda_j f}{r_j e^{-i\phi_j}}.$$

Then we have

$$\frac{\lambda_j \operatorname{Re}(f)}{r_j} + r_j i \dot{\phi}_j = \frac{\lambda_j f}{r_j}.$$

Then

$$\dot{\phi}_j = \frac{\lambda_j \operatorname{Im}(f)}{\alpha_j + \lambda_j u}.$$

Therefore

$$\phi_j = \psi_j + \int_{s_0}^s \frac{\lambda_j \operatorname{Im}(f(t))}{\alpha_j + \lambda_j u(t)} dt.$$

This completes the proof of Theorem 2.3.  $\square$

From Theorem 2.1 if  $\dot{\theta} = -\alpha \operatorname{Im}(f)$  then  $L$  is a self-similar solution. From Theorem 2.3 and (6),  $\dot{\theta} = -\alpha \operatorname{Im}(f)$  is equivalent to

$$\sum_j \frac{\lambda_j \operatorname{Im}(f)}{\alpha_j + 2\lambda_j \int_{s_0}^s \operatorname{Re}(f) dt} + \frac{d}{ds} \arg(f) = -\alpha \operatorname{Im}(f). \quad (7)$$

Therefore if (7) holds then  $L$  is a self-similar solution. So we can get self-similar solutions by getting solutions of (7). For example if we put  $f \equiv i$  then  $f$  is a solution of (7). In this case if we put  $\alpha_1 = \dots = \alpha_n = 0$ ,  $\lambda_1, \dots, \lambda_n \in \mathbb{Z} - \{0\}$  then the construction reduces to that of Lee and Wang [4].

*Proof of Theorem 1.2.* Define  $\omega_j : I \rightarrow \mathbb{C} - \{0\}$  by  $\omega_j = r_j e^{i\phi_j}$ . Define  $f : I \rightarrow \mathbb{C} - \{0\}$  by

$$f = \frac{\overline{\omega_j} \dot{\omega}_j}{\lambda_j}.$$

A direct calculation shows that

$$f = B + i \frac{|B|}{\sqrt{E\{\prod_{k=1}^n (1 + 2a_k B \lambda_k s)\} e^{2B\alpha s} - 1}}.$$

Then  $f \neq 0$  in  $I$ . Apply Theorem 2.1 to the data  $\omega_j, f$  above. Then we get a Lagrangian submanifold  $L$  defined by (5). A direct calculation shows that  $f$  satisfies (7). So this completes the proof.  $\square$

*Proof of Theorem 1.3.* Put  $I = \mathbb{R}_{>0}$  or  $I = \mathbb{R}_{<0}$ . Define  $\omega_j : I \rightarrow \mathbb{C} - \{0\}$  by  $\omega_j = r_j e^{i\phi_j}$ . Define  $f : I \rightarrow \mathbb{C} - \{0\}$  by

$$f = \overline{\omega_j} \dot{\omega}_j.$$

A direct calculation shows that

$$f = s + i \frac{|s|}{\sqrt{E\{\prod_{k=1}^n (1 + a_k s^2)\} e^{\alpha s^2} - 1}}.$$

Then  $f \neq 0$  in  $I$ . Apply Theorem 2.1 to the data  $\omega_j, f$  above and  $\lambda_1 = \dots = \lambda_n = C = 1$ . Then we get a Lagrangian submanifold  $L$  defined by (5). A direct calculation shows that  $f$  satisfies (7). So this completes the proof.  $\square$

### 3 Proofs for translating solitons

In order to prove Theorem 1.4 and Theorem 1.5 we use the following Theorem 3.1 and Corollary 3.2 that are a slight generalization of Theorem A and Theorem B in [3]. The following Theorem 3.1 sets up the ordinary differential equations for immersed Lagrangian submanifolds diffeomorphic to  $\mathbb{R}^n$ .

**Theorem 3.1.** *Fix  $n \geq 2$ . Let  $\lambda_1, \dots, \lambda_{n-1} \in \mathbb{R} \setminus \{0\}$  and  $\alpha \in \mathbb{R}$  be constants,  $I$  be an open interval in  $\mathbb{R}$ , and  $\omega_1, \dots, \omega_{n-1} : I \rightarrow \mathbb{C} \setminus \{0\}$  and  $\beta : I \rightarrow \mathbb{C}$  be smooth functions. Suppose that*

$$\frac{d\omega_j}{ds} = \frac{\lambda_j}{\omega_j} \cdot \frac{d\beta}{ds} \quad (8)$$

and

$$\frac{d\beta}{ds} \neq 0$$

hold in  $I$ . Then the submanifold  $L$  in  $\mathbb{C}^n$  given by

$$L = \{(x_1\omega_1(s), \dots, x_{n-1}\omega_{n-1}(s), -\frac{1}{2} \sum_{j=1}^{n-1} \lambda_j x_j^2 + \beta(s)) \mid x_1, \dots, x_{n-1} \in \mathbb{R}, s \in I\} \quad (9)$$

is an immersed Lagrangian submanifolds diffeomorphic to  $\mathbb{R}^n$ , with Lagrangian angle

$$\theta(s) = \arg(\omega_1 \cdots \omega_{n-1} \beta). \quad (10)$$

Further we have

$$H = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right)$$

and

$$T^\perp = \frac{-\alpha \operatorname{Im}(\beta)}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right),$$

where  $T = (0, \dots, 0, \alpha) \in \mathbb{C}^n$  is a constant vector.

*Remark 3.1.1.* A direct calculation shows that

$$\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle = \sum_{j=1}^{n-1} \frac{\lambda_j^2 x_j^2 |\dot{\beta}|^2}{|\omega_j|^2} + \left| -\frac{1}{2} \sum_{j=1}^{n-1} \lambda_j x_j^2 + \beta \right|^2.$$

*Remark 3.1.2.* Let  $\tilde{\theta} : I \rightarrow \mathbb{R}$  be a function satisfying

$$\frac{d\tilde{\theta}}{ds} = -\alpha \operatorname{Im}(e^{i\tilde{\theta}} \overline{\omega_1 \cdots \omega_{n-1}}).$$

Suppose that

$$\frac{d\beta}{ds} = e^{i\bar{\theta}} \overline{\omega_1 \cdots \omega_{n-1}}.$$

hold in  $I$ . Then it is proved in [3] that  $L$  is a Lagrangian submanifold and  $L$  is a translating soliton with translating vector  $(0, \dots, 0, \alpha) \in \mathbb{C}^n$ . This is Theorem G in [3].

The following Corollary 3.2 gives the solution to ordinary differential equations in Theorem 3.1.

**Corollary 3.2.** *In the situation of Theorem 3.1, write  $\omega_j := r_j e^{i\phi_j}$ , for functions  $r_1, \dots, r_{n-1} : I \rightarrow (0, \infty)$ ,  $\phi_1, \dots, \phi_{n-1} : I \rightarrow \mathbb{R}$ . Define  $u : I \rightarrow \mathbb{R}$  by*

$$u(s) := 2 \int_{s_0}^s \operatorname{Re}(\dot{\beta}(t)) dt.$$

Then we have

$$r_j^2 = \alpha_j + \lambda_j u, \quad \phi_j = \psi_j + \int_{s_0}^s \frac{\lambda_j \operatorname{Im}(\dot{\beta}(t))}{\alpha_j + \lambda_j u(t)} dt$$

with  $\alpha_j = r_j^2(s_0)$ ,  $\psi_j = \phi_j(s_0)$ .

*Remark 3.2.1.* Thus if  $\beta$  is explicitly given, the ordinary differential equation (8) is solved by Theorem 3.2.

*Proof of Theorem 3.1.* We consider the map

$$\iota : \mathbb{R}^{n-1} \times I \rightarrow \mathbb{C}^n$$

given by

$$\iota((x_1, \dots, x_{n-1}), s) = (x_1 \omega_1(s), \dots, x_{n-1} \omega_{n-1}(s), \frac{1}{2} \sum_{j=1}^{n-1} \lambda_j x_j^2 + \beta(s)).$$

Then we have  $L = \iota(\mathbb{R}^{n-1} \times I)$ . So  $(x_1, \dots, x_{n-1}, s)$  is a coordinate of  $L$ . With this coordinate we have

$$\begin{aligned} & (dz_1 \wedge d\bar{z}_1 + \cdots + dz_n \wedge d\bar{z}_n)|_L \\ &= (\omega_1 dx_1 + x_1 \dot{\omega}_1 ds) \wedge (\bar{\omega}_1 dx_1 + x_1 \overline{\dot{\omega}_1} ds) + \cdots + \\ & \quad (\omega_{n-1} dx_{n-1} + \dot{\omega}_{n-1} ds) \wedge (\bar{\omega}_{n-1} dx_{n-1} + x_{n-1} \overline{\dot{\omega}_{n-1}} ds) \\ & \quad + (-\lambda_1 x_1 dx_1 - \cdots - \lambda_{n-1} x_{n-1} dx_{n-1} + \dot{\beta}(s) ds) \\ & \quad \wedge (-\lambda_1 x_1 dx_1 - \cdots - \lambda_{n-1} x_{n-1} dx_{n-1} + \overline{\dot{\beta}(s)} ds) \\ &= 2i \operatorname{Im}(x_1 \omega_1 \overline{\dot{\omega}_1}) dx_1 \wedge ds + \cdots + 2i \operatorname{Im}(x_{n-1} \omega_{n-1} \overline{\dot{\omega}_{n-1}}) dx_{n-1} \wedge ds \\ & \quad - 2i \operatorname{Im}(\lambda_1 x_1 \overline{\dot{\beta}}) dx_1 \wedge ds - \cdots - 2i \operatorname{Im}(\lambda_{n-1} x_{n-1} \overline{\dot{\beta}}) dx_{n-1} \wedge ds \\ &= 0 \end{aligned}$$

and

$$\begin{aligned}
& dz_1 \wedge \cdots \wedge dz_n|_L \\
&= (\omega_1 dx_1 + x_1 \dot{\omega}_1 ds) \wedge \cdots \wedge (\omega_{n-1} dx_{n-1} + x_{n-1} \dot{\omega}_{n-1} ds) \wedge \\
&\quad (-\lambda_1 x_1 dx_1 - \cdots - \lambda_{n-1} x_{n-1} dx_{n-1} + \dot{\beta} ds) \\
&= \omega_1 \cdots \omega_{n-1} \dot{\beta} \left(1 + \sum_{j=1}^{n-1} \frac{\lambda_j^2 x_j^2}{|\omega_j|^2}\right) dx_1 \wedge \cdots \wedge dx_{n-1} \wedge ds.
\end{aligned}$$

It follows that  $L$  is a nonsingular immersed Lagrangian, with Lagrangian angle

$$\theta = \arg(\omega_1 \cdots \omega_{n-1} \dot{\beta}).$$

Since

$$\begin{aligned}
& \left\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial x_j} \right\rangle \\
&= \langle (x_1 \dot{\omega}_1(s), \cdots, x_{n-1} \dot{\omega}_{n-1}(s), \dot{\beta}), (0, \cdots, 0, \omega_j, 0, \cdots, 0, -\lambda_j x_j) \rangle \\
&= \operatorname{Re}(\overline{\omega_j} x_j \dot{\omega}_j - \lambda_j x_j \dot{\beta}) \\
&= 0,
\end{aligned}$$

and  $d\theta = \dot{\theta} ds$ , it follows that

$$\nabla \theta = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} \left( \frac{\partial}{\partial s} \right).$$

Therefore we obtain

$$H = \frac{\dot{\theta}}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right).$$

The normal projection of the position vector  $F$  is computed by

$$\begin{aligned}
& \left\langle T, J \frac{\partial}{\partial x_j} \right\rangle \\
&= \langle (0, \cdots, 0, \alpha), i(0, \cdots, 0, \omega_j, 0, \cdots, 0, -\lambda_j x_j) \rangle \\
&= 0
\end{aligned}$$

and

$$\begin{aligned}
\left\langle T, J \frac{\partial}{\partial s} \right\rangle &= \langle (0, \cdots, 0, \alpha), i(x_1 \dot{\omega}_1(s), \cdots, x_{n-1} \dot{\omega}_{n-1}(s), \dot{\beta}) \rangle \\
&= \operatorname{Re}(\alpha i \dot{\beta}) \\
&= -\alpha \operatorname{Im}(\dot{\beta}).
\end{aligned}$$

It follows that

$$T^\perp = -\frac{\alpha \operatorname{Im}(\dot{\beta})}{\langle \frac{\partial}{\partial s}, \frac{\partial}{\partial s} \rangle} J \left( \frac{\partial}{\partial s} \right).$$

This completes the proof of Theorem 3.1.  $\square$

As the proof of Theorem G in [3] there is another proof of Theorem 3.1 which obtains Theorem 3.1 from Theorem 2.1 by a limiting procedure. See Joyce, Lee and tsui [3].

*Proof of Corollary 3.2.* Changing  $f$  to  $\dot{\beta}$  in the proof of Theorem 2.3 gives the proof of Corollary 3.2.  $\square$

From Theorem 3.1 if  $\dot{\theta} = -\alpha \operatorname{Im}(\dot{\beta})$  then  $L$  is a translating soliton with translating vector  $T = (0, \dots, 0, \alpha) \in \mathbb{C}^n$ . From Theorem 3.2 and (10),  $\dot{\theta} = -\alpha \operatorname{Im}(\dot{\beta})$  is equivalent to

$$\sum_{j=1}^{n-1} \frac{\lambda_j \operatorname{Im}(\dot{\beta})}{\alpha_j + 2\lambda_j \int_{s_0}^s \operatorname{Re}(\dot{\beta}) dt} + \frac{d}{ds} \arg(\dot{\beta}) = -\alpha \operatorname{Im}(\dot{\beta}). \quad (11)$$

Therefore if (11) holds then  $L$  is a translating soliton with translating vector  $T = (0, \dots, 0, \alpha) \in \mathbb{C}^n$ .

*Proof of Theorem 1.4.* Define  $\omega_j : I \rightarrow \mathbb{C} - \{0\}$  by  $\omega_j = r_j e^{i\phi_j}$ . Define  $\beta : I \rightarrow \mathbb{C} - \{0\}$  by

$$\dot{\beta} = \frac{\overline{\omega_j} \dot{\omega}_j}{\lambda_j} \quad \text{and} \quad \beta(0) = -K.$$

A direct calculation shows that

$$\beta(s) = \int_0^s \left( B + i \frac{|B|}{\sqrt{E \{ \prod_{k=1}^{n-1} (1 + 2a_k B \lambda_k s) \}} e^{2B\alpha s} - 1} \right) ds + K.$$

Then  $\dot{\beta} \neq 0$  in  $I$ . Apply Theorem 3.1 to the data  $\omega_j, \beta$  above. Then we get a Lagrangian submanifold  $L$  defined by (9). A direct calculation shows that  $\beta$  satisfies (11). So this completes the proof.  $\square$

*Proof of Theorem 1.5.* Put  $I = \mathbb{R}_{>0}$  or  $I = \mathbb{R}_{<0}$ . Define  $\omega_j : I \rightarrow \mathbb{C} - \{0\}$  by  $\omega_j = r_j e^{i\phi_j}$ . Define  $\beta : I \rightarrow \mathbb{C} - \{0\}$  by

$$\dot{\beta} = \overline{\omega_j} \dot{\omega}_j \quad \text{and} \quad \beta(0) = 0.$$

A direct calculation shows that

$$\beta(s) = \int_0^s \left( s + i \frac{|s|}{\sqrt{E \{ \prod_{k=1}^{n-1} (1 + a_k s^2) \}} e^{\alpha s^2} - 1} \right) ds.$$

Then  $\dot{\beta} \neq 0$  in  $I$ . Apply Theorem 3.1 to the data  $\omega_j, \beta$  above and  $\lambda_1 = \dots = \lambda_{n-1} = 1$ . Then we get a Lagrangian submanifold  $L$  defined by (9). A direct calculation shows that  $\beta$  satisfies (11). So this completes the proof.  $\square$

## Acknowledgements

The author would like to thank the supervisor Akito Futaki. He also wishes to thank Masataka Shibata, Yuji Terashima, Mitutaka Murayama and Kota Hattori for useful conversations.

## References

- [1] H. Ancaux. Construction of Lagrangian self-similar solutions to the mean curvature flow in  $\mathbb{C}^n$ . *Geom. Dedicata*, 120:37–48, 2006.
- [2] D.D. Joyce. Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications. *J. Differential Geom.*, 63(2):279–347, 2003.
- [3] D.D. Joyce, Y.-I. Lee, and M.-P. Tsui. Self-similar solutions and translating solitons for Lagrangian mean curvature flow. *J. Differential Geom.*, 84(1):127–161, 2010.
- [4] Y.-I. Lee and M.-T. Wang. Hamiltonian stationary cones and self-similar solutions in higher dimension. *Trans. Amer. Math. Soc.*, 362(3):1491–1503, 2010.
- [5] R. P. Thomas and S.-T. Yau. Special Lagrangians, stable bundles and mean curvature flow. *Comm. Anal. Geom.*, 10(5):1075–1113, 2002.