

Width Distributions for Convex Regular Polyhedra

STEVEN R. FINCH

March 12, 2016

ABSTRACT. The mean width is a measure on three-dimensional convex bodies that enjoys equal status with volume and surface area [1]. As the phrase suggests, it is the mean of a probability density f . We verify formulas for mean widths of the regular tetrahedron and the cube. Higher-order moments of f_{tetra} and f_{cube} have not been examined until now. Assume that each polyhedron has edges of unit length. We deduce that the mean square width of the regular tetrahedron is $1/3 + (3 + \sqrt{3})/(3\pi)$ and the mean square width of the cube is $1 + 4/\pi$.

Let C be a convex body in \mathbb{R}^3 . A **width** is the distance between a pair of parallel C -supporting planes. Every unit vector $u \in \mathbb{R}^3$ determines a unique such pair of planes orthogonal to u and hence a width $w(u)$. Let u be uniformly distributed on the unit sphere $S^2 \subset \mathbb{R}^3$. Then w is a random variable and

$$E(w_{\text{tetra}}) = \frac{3}{2\pi} \arccos\left(-\frac{1}{3}\right)$$

for $C =$ the regular tetrahedron with edges of unit length and

$$E(w_{\text{cube}}) = \frac{3}{2}$$

for $C =$ the cube with edges of unit length [1, 2, 3]. The probability density of w is not known. Our humble contribution is to verify the preceding **mean width** results and to deduce the following **mean square width** results:

$$E(w_{\text{tetra}}^2) = \frac{1}{3} \left(1 + \frac{3 + \sqrt{3}}{\pi} \right),$$

$$E(w_{\text{cube}}^2) = 1 + \frac{4}{\pi}$$

which appear to be new.

We start with two-dimensional analogs of these results in Sections 1 and 2. The tetrahedral case (Section 3) is more difficult than the cubic case (Section 4); details of

⁰Copyright © 2011, 2016 by Steven R. Finch. All rights reserved.

the evaluation of a certain definite integral are relegated to Section 5. We note that the phrases **mean breadth** and **mean caliper diameter** are synonymous with mean width [2], and hope that this paper might inspire relevant computations for other convex bodies.

1. EQUILATERAL TRIANGLE

Given a convex region C in \mathbb{R}^2 , a width is the distance between a pair of parallel C -supporting lines. Every unit vector $u \in \mathbb{R}^2$ determines a unique such pair of lines orthogonal to u and hence a width $w(u)$. Let u be uniformly distributed on the unit circle $S^1 \subset \mathbb{R}^2$. We wish to study the distribution of the random variable w in the case $C =$ the equilateral triangle with sides of unit length.

For simplicity, let Δ be the equilateral triangle with vertices

$$v_1 = (0, 1), \quad v_2 = \left(\frac{\sqrt{3}}{2}, -\frac{1}{2}\right), \quad v_3 = \left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}\right).$$

At the end, it will be necessary to normalize by $\sqrt{3}$, the side-length of Δ .

Also let $\tilde{\Delta}$ be the union of three overlapping disks of radius $1/2$ centered at $v_1/2, v_2/2, v_3/2$. Clearly $\Delta \subset \tilde{\Delta}$ and $\tilde{\Delta}$ has centroid $(0, 0)$. A **diameter** of $\tilde{\Delta}$ is the length of the intersection between $\tilde{\Delta}$ and a line passing through the origin.

Computing all widths of Δ is equivalent to computing all diameters of $\tilde{\Delta}$. The latter is achieved as follows. Fix a point (a, b) on the unit circle. The line L passing through $(0, 0)$ and (a, b) has parametric representation

$$x = ta, \quad y = tb, \quad t \in \mathbb{R}$$

and hence $y = (b/a)x$ assuming $a \neq 0$. The nontrivial intersection between top circle and L satisfies

$$x^2 + \left(\frac{b}{a}x - \frac{1}{2}\right)^2 = \frac{1}{4}$$

thus $x_1 = ab$ since $a^2 + b^2 = 1$. The nontrivial intersection between right circle and L satisfies

$$\left(x - \frac{\sqrt{3}}{4}\right)^2 + \left(\frac{b}{a}x + \frac{1}{4}\right)^2 = \frac{1}{4}$$

thus $x_2 = a(\sqrt{3}a - b)/2$. The nontrivial intersection between left circle and L satisfies

$$\left(x + \frac{\sqrt{3}}{4}\right)^2 + \left(\frac{b}{a}x + \frac{1}{4}\right)^2 = \frac{1}{4}$$

thus $x_3 = -a(\sqrt{3}a + b)/2$.

We now examine all pairwise distances, squared, between the three intersection points:

$$(x_i - x_j)^2 + \left(\frac{b}{a}x_i - \frac{b}{a}x_j\right)^2 = \begin{cases} \frac{1}{4}(3 - 6\sqrt{3}ab + 6b^2) & \text{if } i = 1, j = 2 \\ \frac{3}{4}(1 + 2\sqrt{3}ab + 2b^2) & \text{if } i = 1, j = 3 \\ 3a^2 & \text{if } i = 2, j = 3 \end{cases}$$

and define

$$g(a, b) = \max \left\{ \frac{1}{4}(3 - 6\sqrt{3}ab + 6b^2), \frac{3}{4}(1 + 2\sqrt{3}ab + 2b^2), 3a^2 \right\}.$$

Therefore the mean width for C is

$$\frac{1}{\sqrt{3}} \frac{1}{2\pi} \int_0^{2\pi} \sqrt{g(\cos \theta, \sin \theta)} d\theta = \frac{3}{\pi} = \frac{\text{perimeter of } C}{\pi}$$

and the mean square width is

$$\frac{1}{3} \frac{1}{2\pi} \int_0^{2\pi} g(\cos \theta, \sin \theta) d\theta = \frac{1}{2} \left(1 + \frac{3\sqrt{3}}{2\pi} \right).$$

2. SQUARE

We turn to the case C = the square with sides of unit length. For simplicity, let \square be the square with vertices

$$v_1 = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right), \quad v_2 = \left(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right), \quad v_3 = \left(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right), \quad v_4 = \left(-\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right).$$

At the end, it will be necessary to normalize by $\sqrt{2}$, the side-length of \square .

Also let $\tilde{\square}$ be the union of four overlapping disks of radius $1/2$ centered at $v_1/2, v_2/2, v_3/2, v_4/2$. A diameter of $\tilde{\square}$ is the length of the intersection between $\tilde{\square}$ and a line passing through the origin.

Computing all widths of \square is equivalent to computing all diameters of $\tilde{\square}$. Fix a point (a, b) on the unit circle. The line L passing through $(0, 0)$ and (a, b) can be represented as $y = (b/a)x$ assuming $a \neq 0$. The nontrivial intersection between northeast circle and L satisfies

$$\left(x - \frac{\sqrt{2}}{4} \right)^2 + \left(\frac{b}{a}x - \frac{\sqrt{2}}{4} \right)^2 = \frac{1}{4}$$

thus $x_1 = \sqrt{2}a(a+b)/2$ since $a^2 + b^2 = 1$. The nontrivial intersection between southeast circle and L satisfies

$$\left(x - \frac{\sqrt{2}}{4} \right)^2 + \left(\frac{b}{a}x + \frac{\sqrt{2}}{4} \right)^2 = \frac{1}{4}$$

thus $x_2 = \sqrt{2}a(a-b)/2$. The nontrivial intersection between northwest circle and L satisfies

$$\left(x + \frac{\sqrt{2}}{4}\right)^2 + \left(\frac{b}{a}x - \frac{\sqrt{2}}{4}\right)^2 = \frac{1}{4}$$

thus $x_3 = -\sqrt{2}a(a-b)/2$. The nontrivial intersection between southwest circle and L satisfies

$$\left(x + \frac{\sqrt{2}}{4}\right)^2 + \left(\frac{b}{a}x + \frac{\sqrt{2}}{4}\right)^2 = \frac{1}{4}$$

thus $x_4 = -\sqrt{2}a(a+b)/2$.

We now examine all pairwise distances, squared, between the four intersection points:

$$(x_i - x_j)^2 + \left(\frac{b}{a}x_i - \frac{b}{a}x_j\right)^2 = \begin{cases} 2b^2 & \text{if } i = 1, j = 2 \\ 2a^2 & \text{if } i = 1, j = 3 \\ 2 + 4ab & \text{if } i = 1, j = 4 \\ 2 - 4ab & \text{if } i = 2, j = 3 \\ 2a^2 & \text{if } i = 2, j = 4 \\ 2b^2 & \text{if } i = 3, j = 4 \end{cases}$$

and define

$$\begin{aligned} g(a, b) &= \max \{2a^2, 2b^2, 2(1 + 2ab), 2(1 - 2ab)\} \\ &= \max \{2(1 + 2ab), 2(1 - 2ab)\}. \end{aligned}$$

Therefore the mean width for C is

$$\frac{1}{\sqrt{2}} \frac{1}{2\pi} \int_0^{2\pi} \sqrt{g(\cos \theta, \sin \theta)} d\theta = \frac{4}{\pi} = \frac{\text{perimeter of } C}{\pi}$$

and the mean square width is

$$\frac{1}{2} \frac{1}{2\pi} \int_0^{2\pi} g(\cos \theta, \sin \theta) d\theta = 1 + \frac{2}{\pi}.$$

3. REGULAR TETRAHEDRON

Returning to the three-dimensional setting of the introduction, let C = the regular tetrahedron with edges of unit length.

For simplicity, let Δ be the tetrahedron with vertices

$$\begin{aligned} v_1 &= (0, 0, 1), & v_2 &= \left(\frac{2\sqrt{2}}{3}, 0, -\frac{1}{3}\right), \\ v_3 &= \left(-\frac{\sqrt{2}}{3}, \sqrt{\frac{2}{3}}, -\frac{1}{3}\right), & v_4 &= \left(-\frac{\sqrt{2}}{3}, -\sqrt{\frac{2}{3}}, -\frac{1}{3}\right), \end{aligned}$$

At the end, it will be necessary to normalize by $2\sqrt{2/3}$, the edge-length of Δ .

Also let $\tilde{\Delta}$ be the union of four overlapping balls of radius $1/2$ centered at $v_1/2, v_2/2, v_3/2, v_4/2$. Clearly $\Delta \subset \tilde{\Delta}$ and $\tilde{\Delta}$ has centroid $(0,0,0)$. A diameter of $\tilde{\Delta}$ is the length of the intersection between $\tilde{\Delta}$ and a line passing through the origin.

Computing all widths of Δ is equivalent to computing all diameters of $\tilde{\Delta}$. The latter is achieved as follows. Fix a point (a, b, c) on the unit sphere. The line L passing through $(0, 0, 0)$ and (a, b, c) has parametric representation

$$x = ta, \quad y = tb, \quad z = tc, \quad t \in \mathbb{R}$$

and hence $y = (b/a)x, z = (c/a)x$ assuming $a \neq 0$. The nontrivial intersection between top sphere and L satisfies

$$x^2 + \left(\frac{b}{a}x\right)^2 + \left(\frac{c}{a}x - \frac{1}{2}\right)^2 = \frac{1}{4}$$

thus $x_1 = ac$ since $a^2 + b^2 + c^2 = 1$. The nontrivial intersection between front sphere and L satisfies

$$\left(x - \frac{\sqrt{2}}{3}\right)^2 + \left(\frac{b}{a}x\right)^2 + \left(\frac{c}{a}x + \frac{1}{6}\right)^2 = \frac{1}{4}$$

thus $x_2 = a(2\sqrt{2}a - c)/3$. The nontrivial intersection between left sphere and L satisfies

$$\left(x + \frac{\sqrt{2}}{6}\right)^2 + \left(\frac{b}{a}x + \frac{\sqrt{6}}{6}\right)^2 + \left(\frac{c}{a}x + \frac{1}{6}\right)^2 = \frac{1}{4}$$

thus $x_3 = -a(\sqrt{2}a + \sqrt{6}b + c)/3$. The nontrivial intersection between right sphere and L satisfies

$$\left(x + \frac{\sqrt{2}}{6}\right)^2 + \left(\frac{b}{a}x - \frac{\sqrt{6}}{6}\right)^2 + \left(\frac{c}{a}x + \frac{1}{6}\right)^2 = \frac{1}{4}$$

thus $x_4 = a(-\sqrt{2}a + \sqrt{6}b - c)/3$.

We now examine all pairwise distances, squared, between the four intersection points:

$$\begin{aligned} & (x_i - x_j)^2 + \left(\frac{b}{a}x_i - \frac{b}{a}x_j\right)^2 + \left(\frac{c}{a}x_i - \frac{c}{a}x_j\right)^2 \\ &= \begin{cases} \frac{8}{9}(a^2 - 2\sqrt{2}ac + 2c^2) & \text{if } i = 1, j = 2 \\ \frac{1}{9}(\sqrt{2}a + \sqrt{6}b + 4c)^2 & \text{if } i = 1, j = 3 \\ \frac{1}{9}(\sqrt{2}a - \sqrt{6}b + 4c)^2 & \text{if } i = 1, j = 4 \\ \frac{2}{3}(3a^2 + 2\sqrt{3}ab + b^2) & \text{if } i = 2, j = 3 \\ \frac{2}{3}(3a^2 - 2\sqrt{3}ab + b^2) & \text{if } i = 2, j = 4 \\ \frac{2}{3}b^2 & \text{if } i = 3, j = 4 \end{cases} \end{aligned}$$

and define

$$g(a, b) = \max \left\{ \frac{8}{9} (a^2 - 2\sqrt{2}ac + 2c^2), \frac{1}{9} (\sqrt{2}a + \sqrt{6}b + 4c)^2, \frac{1}{9} (\sqrt{2}a - \sqrt{6}b + 4c)^2, \right. \\ \left. \frac{2}{3} (3a^2 + 2\sqrt{3}ab + b^2), \frac{2}{3} (3a^2 - 2\sqrt{3}ab + b^2), \frac{8}{3}b^2 \right\}.$$

As for the equilateral triangle, no simplification of g seems possible. The mean width for C is

$$\frac{1}{2\sqrt{2/3}} \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi \sqrt{g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)} \sin \varphi \, d\varphi \, d\theta = \frac{3}{2\pi} \arccos \left(-\frac{1}{3} \right)$$

and the mean square width is

$$\frac{1}{8/3} \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi \, d\varphi \, d\theta = \frac{1}{3} \left(1 + \frac{3 + \sqrt{3}}{\pi} \right).$$

Details on the final integral are given in Section 5.

4. CUBE

We turn to the case C = the cube with edges of unit length. For simplicity, let \square be the cube with vertices

$$v_k = \left(\pm \frac{\sqrt{3}}{3}, \pm \frac{\sqrt{3}}{3}, \pm \frac{\sqrt{3}}{3} \right)$$

for $1 \leq k \leq 8$. At the end, it will be necessary to normalize by $2/\sqrt{3}$, the edge-length of \square .

Also let $\tilde{\square}$ be the union of eight overlapping balls of radius $1/2$ centered at $v_1/2, v_2/2, v_3/2, \dots, v_8/2$. A diameter of $\tilde{\square}$ is the length of the intersection between $\tilde{\square}$ and a line passing through the origin.

Computing all widths of \square is equivalent to computing all diameters of $\tilde{\square}$. Fix a point (a, b, c) on the unit sphere. The line L passing through $(0, 0, 0)$ and (a, b, c) can be represented as $y = (b/a)x, z = (c/a)x$ assuming $a \neq 0$. The nontrivial intersection between each of the eight spheres and L satisfies

$$\left(x \pm \frac{\sqrt{3}}{6} \right)^2 + \left(\frac{b}{a}x \pm \frac{\sqrt{3}}{6} \right)^2 + \left(\frac{c}{a}x \pm \frac{\sqrt{3}}{6} \right)^2 = \frac{1}{4}$$

and consequently

$$g(a, b) = \max \left\{ \frac{4}{3}a^2, \frac{4}{3}b^2, \frac{4}{3}c^2, \frac{4}{3} (1 + 2ab - c^2), \frac{4}{3} (1 + 2ac - b^2), \right. \\ \frac{4}{3} (1 - 2ab - c^2), \frac{4}{3} (1 - 2ac - b^2), \frac{4}{3} (b + c)^2, \frac{4}{3} (b - c)^2, \\ \frac{4}{3} (1 + 2ab + 2ac + 2bc), \frac{4}{3} (1 + 2ab - 2ac - 2bc), \\ \left. \frac{4}{3} (1 - 2ab - 2ac + 2bc), \frac{4}{3} (1 - 2ab + 2ac - 2bc) \right\}$$

after examining all 28 pairwise distances and extracting 13 distinct expressions. As for the square (in which g simplified to a maximum over two terms), here g simplifies to a maximum over four terms:

$$g(a, b) = \max \left\{ \frac{4}{3} (1 + 2ab + 2ac + 2bc), \frac{4}{3} (1 + 2ab - 2ac - 2bc), \right. \\ \left. \frac{4}{3} (1 - 2ab - 2ac + 2bc), \frac{4}{3} (1 - 2ab + 2ac - 2bc) \right\}.$$

The mean width for C is

$$\frac{1}{2/\sqrt{3}} \frac{1}{4\pi} \int_0^{2\pi} \int_0^{\pi} \sqrt{g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)} \sin \varphi \, d\varphi \, d\theta = \frac{3}{2}$$

and the mean square width is

$$\frac{1}{4/3} \frac{1}{4\pi} \int_0^{2\pi} \int_0^{\pi} g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi \, d\varphi \, d\theta = 1 + \frac{4}{\pi}.$$

5. A DEFINITE INTEGRAL

Considerable work is required to evaluate the definite integral at the end of Section 3. A plot of the surface

$$(\theta, \varphi) \mapsto \sqrt{\frac{g(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)}{8/3}}$$

appears in Figure 1, where $0 \leq \theta \leq 2\pi$ and $0 \leq \varphi \leq \pi$. Figure 2 contains the same surface, but viewed from above. Our focus will be on the part of the surface in the lower right corner, specifically $0 \leq \theta \leq \pi/3$ and $2 \lesssim \varphi \leq \pi$. The volume under this part is $1/24^{\text{th}}$ of the volume under the full surface.

We need to find the precise lower bound on φ as a function of θ . Recall the formula for g as a maximum over six terms in Section 3; let g_ℓ denote the ℓ^{th} term, where $1 \leq \ell \leq 6$. Then the lower bound on φ is found by solving the equation

$$g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) = g_4(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi)$$

for φ . We obtain $\varphi(\theta) = 2 \arctan(h(\theta))$, where

$$h(\theta) = \frac{\cos \theta + \sqrt{3} \sin \theta + \sqrt{10 - \cos(2\theta) + \sqrt{3} \sin(2\theta)}}{2\sqrt{2}}$$

and, in particular,

$$\varphi(0) = 2 \arctan(\sqrt{2}) \approx 1.9106,$$

$$\varphi(\pi/3) = 2 \arctan \left(\frac{1 + \sqrt{3}}{\sqrt{2}} \right) \approx 2.1862.$$

It follows that $g = g_1$ for $0 \leq \theta \leq \pi/3$ and $2 \arctan(h) \leq \varphi \leq \pi$. Now we have

$$\begin{aligned} & \frac{1}{8/3} \frac{1}{4\pi} \int g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi d\varphi \\ &= \frac{(-3 + \cos(2\theta)) \cos(3\varphi) - 3(7 + 3 \cos(2\theta)) \cos \varphi - 16\sqrt{2} \cos \theta \sin^3 \varphi}{288\pi} \end{aligned}$$

and

$$\begin{aligned} \cos(3\varphi)|_{2 \arctan(h)}^{\pi} &= -2 \frac{(1 - 3h^2)^2}{(1 + h^2)^3}, \\ \cos(\varphi)|_{2 \arctan(h)}^{\pi} &= -\frac{2}{1 + h^2}, \quad \sin(\varphi)|_{2 \arctan(h)}^{\pi} = -\frac{2h}{1 + h^2} \end{aligned}$$

therefore

$$\begin{aligned} & \frac{1}{8/3} \frac{1}{4\pi} \int_{2 \arctan(h)}^{\pi} g_1(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) \sin \varphi d\varphi \\ &= \frac{6h^4 + 8\sqrt{2}h^3 \cos(\theta) + 3h^2(1 + \cos(2\theta)) + (3 + \cos(2\theta))}{18\pi(1 + h^2)^3}. \end{aligned}$$

Integrating this expression from 0 to $\pi/3$ gives the desired formula for $E(w_{\text{tetra}}^2)$.

REFERENCES

- [1] G.-C. Rota, Mathematical snapshots, unpublished note (1997), <http://www-groups.dcs.st-and.ac.uk/~history/Extras/rota.pdf>.
- [2] S. R. Finch, Colliding dice probabilities, unpublished note (2011), <http://www.people.fas.harvard.edu/~sfinch/csolve/dce.pdf>.
- [3] D. A. Klain and G.-C. Rota, *Introduction to Geometric Probability*, Cambridge Univ. Press, 1997, pp. 139–140; MR1608265 (2001f:52009).
- [4] S. R. Finch, Simulations in R involving colliding dice and mean widths, <http://www.people.fas.harvard.edu/~sfinch/csolve/rsimul.html>.

Steven R. Finch
 Dept. of Statistics
 Harvard University
 Cambridge, MA, USA
steven_finch@harvard.edu




Figure 1: Surface plot of $\sqrt{3g/8}$, where $0 \leq \theta \leq 2\pi$ and $0 \leq \varphi \leq \pi$.




Figure 2: Another view of $\sqrt{3g/8}$, with contours of intersection.