## FACTORIAL CLUSTER ALGEBRAS

#### CHRISTOF GEISS, BERNARD LECLERC, AND JAN SCHRÖER

ABSTRACT. We show that cluster algebras do not contain non-trivial units and that all cluster variables are irreducible elements. Both statements follow from Fomin and Zelevinsky's Laurent phenomenon. As an application we give a criterion for a cluster algebra to be a factorial algebra. This can be used to construct cluster algebras, which are isomorphic to polynomial rings. We also study various kinds of upper bounds for cluster algebras, and we prove that factorial cluster algebras coincide with their upper bounds.

#### CONTENTS

1.	Introduction and main results	1
2.	Invertible elements in cluster algebras	6
3.	Irreducibility of cluster variables	7
4.	Factorial cluster algebras	8
5.	The divisibility group of a cluster algebra	11
6.	Examples of non-factorial cluster algebras	11
7.	Examples of factorial cluster algebras	14
8.	Applications	20
References		22

# 1. INTRODUCTION AND MAIN RESULTS

1.1. Introduction. The introduction of cluster algebras by Fomin and Zelevinsky [FZ1] triggered an extensive theory. Most results deal with the combinatorics of seed and quiver mutations, with various categorifications of cluster algebras, and with cluster phenomena occuring in various areas of mathematics, like representation theory of finite-dimensional algebras, quantum groups and Lie theory, Calabi-Yau categories, non-commutative Donaldson-Thomas invariants, Poisson geometry, discrete dynamical systems and algebraic combinatorics.

On the other hand, there are not many results on cluster algebras themselves. As a subalgebra of a field, any cluster algebra  $\mathcal{A}$  is obviously an integral domain. It is also easy to show that its field of fractions  $\operatorname{Frac}(\mathcal{A})$  is isomorphic to a field  $K(x_1, \ldots, x_m)$  of rational functions. Several classes of cluster algebras are known to be finitely generated, e.g. acyclic cluster algebras [BFZ, Corollary 1.21] and also a class of cluster algebras arising from Lie theory [GLS2, Theorem 3.2]. Berenstein,

Date: 13.10.2011.

<sup>2010</sup> Mathematics Subject Classification. Primary 13F60; Secondary 13F15,17B37.

Fomin and Zelevinsky gave an example of a cluster algebra which is not finitely generated. (One applies [BFZ, Theorem 1.24] to the example mentioned in [BFZ, Proposition 1.26].) Only very little is known on further ring theoretic properties of an arbitrary cluster algebra  $\mathcal{A}$ . Here are some basic questions we would like to address:

- Which elements in  $\mathcal{A}$  are invertible, irreducible or prime?
- When is  $\mathcal{A}$  a factorial ring?
- When is  $\mathcal{A}$  a polynomial ring?

In this paper, we work with cluster algebras of geometric type.

1.2. **Definition of a cluster algebra.** In this section we repeat Fomin and Zelevinsky's definition of a cluster algebra.

A matrix  $A = (a_{ij}) \in M_{n,n}(\mathbb{Z})$  is skew-symmetrizable (resp. symmetrizable) if there exists a diagonal matrix  $D = \text{Diag}(d_1, \ldots, d_n) \in M_{n,n}(\mathbb{Z})$  with positive diagonal entries  $d_1, \ldots, d_n$  such that DA is skew-symmetric (resp. symmetric), i.e.  $d_i a_{ij} = -d_j a_{ji}$  (resp.  $d_i a_{ij} = d_j a_{ji}$ ) for all i, j.

Let m, n and p be integers with

$$m \ge p \ge n \ge 1.$$

Let  $B = (b_{ij}) \in M_{m,n}(\mathbb{Z})$  be an  $(m \times n)$ -matrix with integer entries. By  $B^{\circ} \in M_{n,n}(\mathbb{Z})$  we denote the *principal part* of B, which is obtained from B by deleting the last m - n rows.

Let  $\Delta(B)$  be the graph with vertices  $1, \ldots, m$  and an edge between *i* and *j* provided  $b_{ij}$  or  $b_{ji}$  is non-zero. We call *B* connected if the graph  $\Delta(B)$  is connected.

For a field K of characteristic 0 let  $\mathcal{F} := K(X_1, \ldots, X_m)$  be the field of rational functions in m variables.

A seed of  $\mathcal{F}$  is a pair  $(\mathbf{x}, B)$  such that the following hold:

- (i)  $B \in M_{m,n}(\mathbb{Z}),$
- (ii) B is connected,
- (iii)  $B^{\circ}$  is skew-symmetrizable,
- (iv)  $\mathbf{x} = (x_1, \ldots, x_m)$  is an *m*-tuple of elements in  $\mathcal{F}$  such that  $x_1, \ldots, x_m$  are algebraically independent over K.

For a seed  $(\mathbf{x}, B)$ , the matrix B is the exchange matrix of  $(\mathbf{x}, B)$ . We say that B has maximal rank if rank(B) = n.

Given a seed  $(\mathbf{x}, B)$  and some  $1 \le k \le n$  we define the *mutation* of  $(\mathbf{x}, B)$  at k as

$$\mu_k(\mathbf{x}, B) := (\mathbf{x}', B')$$

where  $B' = (b'_{ij})$  is defined as

$$b'_{ij} := \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + \frac{|b_{ik}|b_{kj} + b_{ik}|b_{kj}|}{2} & \text{otherwise,} \end{cases}$$

and  $\mathbf{x}' = (x'_1, \dots, x'_m)$  is defined as

$$x'_{s} := \begin{cases} x_{k}^{-1} \prod_{b_{ik} > 0} x_{i}^{b_{ik}} + x_{k}^{-1} \prod_{b_{ik} < 0} x_{i}^{-b_{ik}} & \text{if } s = k, \\ x_{s} & \text{otherwise.} \end{cases}$$

The equality

(1) 
$$x_k x'_k = \prod_{b_{ik}>0} x_i^{b_{ik}} + \prod_{b_{ik}<0} x_i^{-b_{ik}}$$

is called a *mutation relation*. We write

$$\mu_{(\mathbf{x},B)}(x_k) := x'_k$$

and

$$\mu_k(B) := B'.$$

It is easy to check that  $(\mathbf{x}', B')$  is again a seed. Furthermore, we have  $\mu_k \mu_k(\mathbf{x}, B) = (\mathbf{x}, B)$ .

Two seeds  $(\mathbf{x}, B)$  and  $(\mathbf{y}, C)$  are *mutation equivalent* if there exists a sequence  $(i_1, \ldots, i_t)$  with  $1 \le i_j \le n$  for all j such that

$$\mu_{i_t}\cdots\mu_{i_2}\mu_{i_1}(\mathbf{x},B)=(\mathbf{y},C).$$

In this case, we write  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$ . This yields an equivalence relation on all seeds of  $\mathcal{F}$ . (By definition  $(\mathbf{x}, B)$  is also mutation equivalent to itself.)

For a seed  $(\mathbf{x}, B)$  of  $\mathcal{F}$  let

$$\mathcal{X}_{(\mathbf{x},B)} := \bigcup_{(\mathbf{y},C)\sim(\mathbf{x},B)} \{y_1,\ldots,y_n\}$$

where the union is over all seeds  $(\mathbf{y}, C)$  with  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$ . By definition, the *cluster algebra*  $\mathcal{A}(\mathbf{x}, B)$  associated to  $(\mathbf{x}, B)$  is the *L*-subalgebra of  $\mathcal{F}$  generated by  $\mathcal{X}_{(\mathbf{x},B)}$ , where

$$L := K[x_{n+1}^{\pm 1}, \dots, x_p^{\pm 1}, x_{p+1}, \dots, x_m]$$

is the localization of the polynomial ring  $K[x_{n+1}, \ldots, x_m]$  at  $x_{n+1} \cdots x_p$ . (For p = n we set  $x_{n+1} \cdots x_p := 1$ .) Thus  $\mathcal{A}(\mathbf{x}, B)$  is the K-subalgebra of  $\mathcal{F}$  generated by

$$\{x_{n+1}^{\pm 1}, \dots, x_p^{\pm 1}, x_{p+1}, \dots, x_m\} \cup \mathcal{X}_{(\mathbf{x},B)}.$$

The elements of  $\mathcal{X}_{(\mathbf{x},B)}$  are the *cluster variables* of  $\mathcal{A}(\mathbf{x},B)$ .

We call  $(\mathbf{y}, C)$  a seed of  $\mathcal{A}(\mathbf{x}, B)$  if  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$ . In this case, for any  $1 \leq k \leq n$ we call  $(y_k, \mu_{(\mathbf{y},C)}(y_k))$  an exchange pair of  $\mathcal{A}(\mathbf{x}, B)$ . Furthermore, the *m*-tuple  $\mathbf{y}$  is a cluster of  $\mathcal{A}(\mathbf{x}, B)$ , and monomials of the form  $y_1^{a_1}y_2^{a_2}\cdots y_m^{a_m}$  with  $a_i \geq 0$  for all *i* are called *cluster monomials* of  $\mathcal{A}(\mathbf{x}, B)$ .

Two clusters  $\mathbf{y}$  and  $\mathbf{z}$  of  $\mathcal{A}(\mathbf{x}, B)$  are *disjoint* if  $\{y_1, \ldots, y_n\} \cap \{z_1, \ldots, z_n\} = \emptyset$ . Note that for any cluster  $\mathbf{y}$  of  $\mathcal{A}(\mathbf{x}, B)$  we have  $y_i = x_i$  for all  $n+1 \leq i \leq m$ . These m-n elements are the *coefficients* of  $\mathcal{A}(\mathbf{x}, B)$ . There are no invertible coefficients if p = n. We call a cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  coefficient-free if m = n.

Clearly, for any two seeds of the form  $(\mathbf{x}, B)$  and  $(\mathbf{y}, B)$  there is an algebra isomorphism  $\eta: \mathcal{A}(\mathbf{x}, B) \to \mathcal{A}(\mathbf{y}, B)$  with  $\eta(x_i) = y_i$  for all  $1 \le i \le m$ , which respects the mutation relations. Furthermore, if  $(\mathbf{x}, B)$  and  $(\mathbf{y}, C)$  are mutation equivalent seeds, then  $\mathcal{A}(\mathbf{x}, B) = \mathcal{A}(\mathbf{y}, C)$  and we have  $K(x_1, \ldots, x_m) = K(y_1, \ldots, y_m)$ .

## 1.3. The Laurent phenomenon. For a seed $(\mathbf{x}, B)$ of $\mathcal{F}$ let

$$\mathscr{L}_{\mathbf{x}} := K[x_1^{\pm 1}, \dots, x_n^{\pm 1}, x_{n+1}^{\pm 1}, \dots, x_p^{\pm 1}, x_{p+1}, \dots, x_m]$$

be the localization of the polynomial ring  $K[x_1, \ldots, x_m]$  at  $x_1 x_2 \cdots x_p$ . We consider  $\mathscr{L}_{\mathbf{x}}$  as a subring of the field  $\mathcal{F}$ . The following result, known as the *Laurent phenomenon*, is due to Fomin and Zelevinsky and is our key tool to derive some ring theoretic properties of cluster algebras.

**Theorem 1.1** ([FZ1, Theorem 3.1]). For each seed  $(\mathbf{x}, B)$  of  $\mathcal{F}$  we have

$$\mathcal{A}(\mathbf{x},B) \subseteq \overline{\mathcal{A}}(\mathbf{x},B) := \bigcap_{(\mathbf{y},C) \sim (\mathbf{x},B)} \mathscr{L}_{\mathbf{y}}$$

The algebra  $\overline{\mathcal{A}}(\mathbf{x}, B)$  is called the *upper cluster algebra* associated to  $(\mathbf{x}, B)$ , compare [BFZ, Section 1].

1.4. Upper bounds. For a seed  $(\mathbf{x}, B)$  and  $1 \le k \le n$  let  $(\mathbf{x}_k, B_k) := \mu_k(\mathbf{x}, B)$ . Berenstein, Fomin and Zelevinsky [BFZ] called

$$U(\mathbf{x},B) := \mathscr{L}_{\mathbf{x}} \cap \bigcap_{k=1}^{n} \mathscr{L}_{\mathbf{x}_{k}}$$

the upper bound of  $\mathcal{A}(\mathbf{x}, B)$ . They prove the following:

**Theorem 1.2** ([BFZ, Corollary 1.9]). Let  $(\mathbf{x}, B)$  and  $(\mathbf{y}, C)$  be mutation equivalent seeds of  $\mathcal{F}$ . If B has maximal rank, then  $U(\mathbf{x}, B) = U(\mathbf{y}, C)$ . In particular, we have  $\overline{\mathcal{A}}(\mathbf{x}, B) = U(\mathbf{x}, B)$ .

For clusters **y** and **z** of  $\mathcal{A}(\mathbf{x}, B)$  define

$$U(\mathbf{y}, \mathbf{z}) := \mathscr{L}_{\mathbf{y}} \cap \mathscr{L}_{\mathbf{z}}.$$

1.5. Acyclic cluster algebras. Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$  with  $B = (b_{ij})$ . Let  $\Sigma(B)$  be the quiver with vertices  $1, \ldots, n$ , and arrows  $i \to j$  for all  $1 \leq i, j \leq n$  with  $b_{ij} > 0$ , compare [BFZ, Section 1.4]. So  $\Sigma(B)$  encodes the sign-pattern of the principal part  $B^{\circ}$  of B.

The seed  $(\mathbf{x}, B)$  and B are called *acyclic* if  $\Sigma(B)$  does not contain any oriented cycle. The cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is *acyclic* if there exists an acyclic seed  $(\mathbf{y}, C)$  with  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$ .

1.6. Skew-symmetric exchange matrices and quivers. Let  $B = (b_{ij})$  be a matrix in  $M_{m,n}(\mathbb{Z})$  such that  $B^{\circ}$  is skew-symmetric. Let  $\Gamma(B)$  be the quiver with vertices  $1, \ldots, m$  and  $b_{ij}$  arrows  $i \to j$  if  $b_{ij} > 0$ , and  $-b_{ij}$  arrows  $j \to i$  if  $b_{ij} < 0$ . Thus given  $\Gamma(B)$ , we can recover B. In the skew-symmetric case one often works with quivers and their mutations instead of exchange matrices.

1.7. Main results. For a ring R with 1, let  $R^{\times}$  be the set of invertible elements in R, and let  $R^* := R \setminus \{0\}$ . Non-zero rings without zero divisors are called *integral domains*. A non-invertible element a in an integral domain R is *irreducible* if it cannot be written as a product a = bc with  $b, c \in R$  both non-invertible. Cluster algebras are integral domains, since they are by definition subrings of fields.

**Theorem 1.3.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ . Then the following hold:

- (i) For m = 1 we have  $\mathcal{A}(\mathbf{x}, B)^{\times} = \{\lambda x_1^a \mid \lambda \in K^*, a \in \mathbb{Z}\}.$
- (ii) For m > 1 we have  $\mathcal{A}(\mathbf{x}, B)^{\times} = \left\{ \lambda x_{n+1}^{a_{n+1}} \cdots x_p^{a_p} \mid \lambda \in K^*, a_i \in \mathbb{Z} \right\}.$
- (iii) For m > 1 any cluster variable in  $\mathcal{A}(\mathbf{x}, B)$  is irreducible.

For elements a, b in an integral domain R we write a|b if there exists some  $c \in R$ with b = ac. A non-invertible element a in a commutative ring R is prime if whenever a|bc for some  $b, c \in R$ , then a|b or a|c. Every prime element is irreducible, but the converse is not true in general. However, if R is a polynomial ring, then each irreducible element in R is prime. Non-zero elements  $a, b \in R$  are associate if there is some unit  $c \in R^{\times}$  with a = bc. An integral domain R is factorial if the following hold:

- (i) Every non-zero element  $r \in R$  can be written as a product  $r = a_1 \cdots a_s$  of irreducible elements  $a_i \in R$ .
- (ii) If  $a_1 \cdots a_s = b_1 \cdots b_t$  with  $a_i, b_j \in R$  irreducible for all i and j, then s = tand there is a bijection  $\pi: \{1, \ldots, s\} \to \{1, \ldots, s\}$  such that  $a_i$  and  $b_{\pi(i)}$  are associate for all  $1 \leq i \leq s$ .

For example, any polynomial ring is factorial. In a factorial ring, all irreducible elements are prime.

The next result gives a useful criterion when a cluster algebra is a factorial ring.

**Theorem 1.4.** Let  $\mathbf{y}$  and  $\mathbf{z}$  be disjoint clusters of  $\mathcal{A}(\mathbf{x}, B)$ . If there is a subalgebra U of  $\mathcal{A}(\mathbf{x}, B)$ , such that U is factorial and

$$\{y_1, \ldots, y_n, z_1, \ldots, z_n, x_{n+1}^{\pm 1}, \ldots, x_p^{\pm 1}, x_{p+1}, \ldots, x_m\} \subset U,$$

then

$$U = \mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, \mathbf{z}).$$

In particular,  $\mathcal{A}(\mathbf{x}, B)$  is factorial and all cluster variables are prime.

We obtain the following corollary on upper bounds of factorial cluster algebras.

**Corollary 1.5.** Assume that  $\mathcal{A}(\mathbf{x}, B)$  is factorial.

- (i) If y and z are disjoint clusters of  $\mathcal{A}(\mathbf{x}, B)$ , then  $\mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, \mathbf{z})$ .
- (ii) For any  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$  we have  $\mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, C)$ .

In Section 7 we apply the above results to show that many cluster algebras are polynomial rings. In Section 8 we discuss some further applications concerning the dual of Lusztig's semicanonical basis and monoidal categorifications of cluster algebras.

1.8. Factoriality and maximal rank. In Section 6.1 we give examples of cluster algebras  $\mathcal{A}(\mathbf{x}, B)$ , which are not factorial. In these examples, B does not have maximal rank.

After we presented our results at the Abel Symposium in Balestrand in June 2011, Zelevinsky asked the following question:

**Problem 1.6.** Suppose  $(\mathbf{x}, B)$  is a seed of  $\mathcal{F}$  such that B has maximal rank. Does it follow that  $\mathcal{A}(\mathbf{x}, B)$  is factorial?

After we circulated a first version of this article, Philipp Lampe [La] discovered an example of a non-factorial cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  with B having maximal rank. With his permission we explain Lampe's example in Section 6.2.

### 2. Invertible elements in cluster algebras

In this section we prove Theorem 1.3(i),(ii), classifying the invertible elements of cluster algebras.

The following lemma is straightforward and well-known.

**Lemma 2.1.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ . Then  $\mathscr{L}_{\mathbf{x}}^{\times} = \{\lambda x_1^{a_1} \cdots x_p^{a_p} \mid \lambda \in K^*, a_i \in \mathbb{Z}\}.$ 

**Theorem 2.2.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ . Then the following hold:

- (i) For m = 1 we have  $\mathcal{A}(\mathbf{x}, B)^{\times} = \{\lambda x_1^a \mid \lambda \in K^*, a \in \mathbb{Z}\}.$
- (ii) For m > 1 we have  $\mathcal{A}(\mathbf{x}, B)^{\times} = \left\{ \lambda x_{n+1}^{a_{n+1}} \cdots x_p^{a_p} \mid \lambda \in K^*, a_i \in \mathbb{Z} \right\}.$

*Proof.* Assume m = 1. Then  $\mathcal{A}(\mathbf{x}, B)$  has exactly two cluster variables, namely  $x_1$  and  $2/x_1$ . It follows that  $\mathcal{A}(\mathbf{x}, B) = K[x_1^{\pm 1}]$  and by Lemma 2.1 we have  $\mathcal{A}(\mathbf{x}, B)^{\times} = \{\lambda x_1^a \mid \lambda \in K^*, a \in \mathbb{Z}\}.$ 

Next, assume that m > 1. Let u be an invertible element in  $\mathcal{A}(\mathbf{x}, B)$ , and let  $(\mathbf{y}, C)$  be any seed of  $\mathcal{A}(\mathbf{x}, B)$ . By the Laurent phenomenon Theorem 1.1 we know that  $\mathcal{A}(\mathbf{x}, B) \subseteq \mathscr{L}_{\mathbf{y}}$ . It follows that u is also invertible in  $\mathscr{L}_{\mathbf{y}}$ . Thus by Lemma 2.1 there are  $a_1, \ldots, a_p \in \mathbb{Z}$  and  $\lambda \in K^*$  such that  $u = \lambda M$  where

$$M = y_1^{a_1} \cdots y_k^{a_k} \cdots y_p^{a_p}.$$

If all  $a_i$  with  $1 \le i \le n$  are zero, we are done. To get a contradiction, assume that there is some  $1 \le k \le n$  with  $a_k \ne 0$ .

Let  $y_k^* := \mu_{(\mathbf{y},C)}(y_k)$ . Again the Laurent phenomenon yields  $b_1, \ldots, b_p \in \mathbb{Z}$  and  $\nu \in K^*$  such that

$$u = \nu y_1^{b_1} \cdots y_{k-1}^{b_{k-1}} (y_k^*)^{b_k} y_{k+1}^{b_{k+1}} \cdots y_p^{b_p}.$$

Without loss of generality let  $b_k \ge 0$ . (Otherwise we can work with  $u^{-1}$  instead of u.)

If  $b_k = 0$ , we get

$$\lambda y_1^{a_1} \cdots y_k^{a_k} \cdots y_p^{a_p} = \nu y_1^{b_1} \cdots y_{k-1}^{b_{k-1}} y_{k+1}^{b_{k+1}} \cdots y_p^{b_p},$$

where  $\lambda, \nu \in K^*$ . This is a contradiction, because  $a_k \neq 0$  and  $y_1, \ldots, y_m$  are algebraically independent, and therefore Laurent monomials in  $y_1, \ldots, y_m$  are linearly independent in  $\mathcal{F}$ .

Next, assume that  $b_k > 0$ . By definition we have

$$y_k^* = M_1 + M_2$$

with

$$M_1 = y_k^{-1} \prod_{c_{ik}>0} y_i^{c_{ik}}$$
 and  $M_2 = y_k^{-1} \prod_{c_{ik}<0} y_i^{-c_{ik}}$ 

where the products run over the positive, respectively negative, entries in the kth column of the matrix C.

Thus we get an equality of the form

(2) 
$$u = \lambda M = \nu (y_1^{b_1} \cdots y_{k-1}^{b_{k-1}}) (M_1 + M_2)^{b_k} (y_{k+1}^{b_{k+1}} \cdots y_p^{b_p}).$$

Since m > 1, we know that  $M_1 \neq M_2$ . (Here we use that exchange matrices are by definition connected. Otherwise, one could get exchange relations of the form  $x_k x'_k = 1 + 1$ .) Thus the right-hand side of Equation (2) is a non-trivial linear combination of  $b_k + 1 \geq 2$  pairwise different Laurent monomials in  $y_1, \ldots, y_m$ . This is again a contradiction, since  $y_1, \ldots, y_m$  are algebraically independent.  $\Box$ 

**Corollary 2.3.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$  with m > 1, and let y and z be nonzero elements in  $\mathcal{A}(\mathbf{x}, B)$ . Then y and z are associate if and only if there exist  $a_{n+1}, \ldots, a_p \in \mathbb{Z}$  and  $\lambda \in K^*$  with

$$\lambda x_{n+1}^{a_{n+1}} \cdots x_p^{a_p} y = z.$$

### 3. IRREDUCIBILITY OF CLUSTER VARIABLES

In this section we prove Theorem 1.3(iii). The proof is very similar to the proof of Theorem 2.2.

**Theorem 3.1.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$  with m > 1. Then any cluster variable in  $\mathcal{A}(\mathbf{x}, B)$  is irreducible.

*Proof.* Let  $(\mathbf{y}, C)$  be any seed of  $\mathcal{A}(\mathbf{x}, B)$ . Since m > 1, we know from Theorem 2.2(ii) that the cluster variables of  $\mathcal{A}(\mathbf{x}, B)$  are non-invertible in  $\mathcal{A}(\mathbf{x}, B)$ .

Assume that  $y_k$  is not irreducible for some  $1 \le k \le n$ . Thus  $y_k = y'_k y''_k$  for some non-invertible elements  $y'_k$  and  $y''_k$  in  $\mathcal{A}(\mathbf{x}, B)$ . Since  $y_k$  is invertible in  $\mathscr{L}_{\mathbf{y}}$ , we know

that  $y'_k$  and  $y''_k$  are both invertible in  $\mathscr{L}_{\mathbf{y}}$ . Thus by Lemma 2.1 there are  $a_i, b_i \in \mathbb{Z}$ and  $\lambda', \lambda'' \in K^*$  with

$$y'_k = \lambda' y_1^{a_1} \cdots y_s^{a_s} \cdots y_p^{a_p}$$
 and  $y''_k = \lambda'' y_1^{b_1} \cdots y_s^{b_s} \cdots y_p^{b_p}$ .

Since  $y_k = y'_k y''_k$ , we get  $a_s + b_s = 0$  for all  $s \neq k$  and  $a_k + b_k = 1$ .

Assume that  $a_s = 0$  for all  $1 \leq s \leq n$  with  $s \neq k$ . Then  $y'_k = \lambda' y^{a_k}_k y^{a_{n+1}}_{n+1} \cdots y^{a_p}_p$ and  $y''_k = \lambda'' y^{b_k}_k y^{b_{n+1}}_{n+1} \cdots y^{b_p}_p$ . If  $a_k \leq 0$ , then  $y'_k$  is invertible in  $\mathcal{A}(\mathbf{x}, B)$ , and if  $a_k > 0$ , then  $y''_k$  is invertible in  $\mathcal{A}(\mathbf{x}, B)$ . In both cases we get a contradiction.

Next assume  $a_s \neq 0$  for some  $1 \leq s \leq n$  with  $s \neq k$ . Let  $y_s^* := \mu_{(\mathbf{y},C)}(y_s)$ . Thus we have

$$y_s^* = M_1 + M_2$$

with

$$M_1 = y_s^{-1} \prod_{c_{is}>0} y_i^{c_{is}}$$
 and  $M_2 = y_s^{-1} \prod_{c_{is}<0} y_i^{-c_{is}}$ ,

where the products run over the positive, respectively negative, entries in the sth column of the matrix C.

Since  $s \neq k$ , we see that  $y_k$  and therefore also  $y'_k$  and  $y''_k$  are invertible in  $\mathscr{L}_{\mu_s(\mathbf{y},C)}$ . Thus by Lemma 2.1 there are  $c_i, d_i \in \mathbb{Z}$  and  $\nu', \nu'' \in K^*$  with

$$y'_{k} = \nu' y_{1}^{c_{1}} \cdots y_{s-1}^{c_{s-1}} (y_{s}^{*})^{c_{s}} y_{s+1}^{c_{s+1}} \cdots y_{p}^{c_{p}} \quad \text{and} \quad y''_{k} = \nu'' y_{1}^{d_{1}} \cdots y_{s-1}^{d_{s-1}} (y_{s}^{*})^{d_{s}} y_{s+1}^{d_{s+1}} \cdots y_{p}^{d_{p}}.$$

Note that  $c_s + d_s = 0$ . Without loss of generality we assume that  $c_s \ge 0$ . (If  $c_s < 0$ , we continue to work with  $y''_k$  instead of  $y'_k$ .) If  $c_s = 0$ , we get

$$y'_k = \lambda' y_1^{a_1} \cdots y_s^{a_s} \cdots y_p^{a_p} = \nu' y_1^{c_1} \cdots y_s^0 \cdots y_p^{c_p}.$$

This is a contradiction, since  $a_s \neq 0$  and  $y_1, \ldots, y_m$  are algebraically independent. If  $c_s > 0$ , then

$$y'_{k} = \lambda' y_{1}^{a_{1}} \cdots y_{s-1}^{a_{s-1}} y_{s}^{a_{s}} y_{s+1}^{a_{s+1}} \cdots y_{p}^{a_{p}}$$
  
=  $\nu' y_{1}^{c_{1}} \cdots y_{s-1}^{c_{s-1}} (y_{s}^{*})^{c_{s}} y_{s+1}^{c_{s+1}} \cdots y_{p}^{c_{p}}$   
=  $\nu' y_{1}^{c_{1}} \cdots y_{s-1}^{c_{s-1}} (M_{1} + M_{2})^{c_{s}} y_{s+1}^{c_{s+1}} \cdots y_{p}^{c_{p}}$ 

Since m > 1, we know that  $M_1 \neq M_2$ . Thus the Laurent monomial  $y'_k$  is a non-trivial linear combination of  $c_s + 1 \ge 2$  pairwise different Laurent monomials in  $y_1, \ldots, y_m$ , a contradiction.

Note that the coefficients  $x_{p+1}, \ldots, x_m$  of  $\mathcal{A}(\mathbf{x}, B)$  are obviously irreducible in  $\mathscr{L}_{\mathbf{x}}$ . Since  $\mathcal{A}(\mathbf{x}, B) \subseteq \mathscr{L}_{\mathbf{x}}$ , they are also irreducible in  $\mathcal{A}(\mathbf{x}, B)$ .

## 4. Factorial cluster algebras

4.1. A factoriality criterion. This section contains the proofs of Theorem 1.4 and Corollary 1.5.

**Lemma 4.1.** There is a cluster  $\mathbf{y}$  of  $\mathcal{A}(\mathbf{x}, B)$  with  $\mathcal{A}(\mathbf{x}, B) = \mathscr{L}_{\mathbf{y}}$  if and only if m = 1.

*Proof.* If m = 1, then there are exactly two clusters namely  $\mathbf{x} = (x_1)$  and  $\mathbf{y} = (2/x_1)$ , and therefore  $\mathcal{A}(\mathbf{x}, B) = \mathscr{L}_{\mathbf{x}} = \mathscr{L}_{\mathbf{y}}$ . Vice versa, if  $\mathcal{A}(\mathbf{x}, B) = \mathscr{L}_{\mathbf{y}}$ , we get an invertible cluster variable  $y_1$ . Then Theorem 2.2 yields m = 1.

**Lemma 4.2.** Let  $\mathbf{y}$  and  $\mathbf{z}$  be clusters of  $\mathcal{A}(\mathbf{x}, B)$  such that  $\mathbf{y}$  and  $\mathbf{z}$  are not disjoint. If m > 1, then we have  $\mathcal{A}(\mathbf{x}, B) \neq U(\mathbf{y}, \mathbf{z})$ .

*Proof.* Assume that m > 1 and  $\mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, \mathbf{z}) = \mathscr{L}_{\mathbf{y}} \cap \mathscr{L}_{\mathbf{z}}$  with  $\mathbf{y}$  and  $\mathbf{z}$  not disjoint. Then there exists an invertible cluster variable in  $\mathcal{A}(\mathbf{x}, B)$ . (Just take any element in the non-empty set  $\{y_1, \ldots, y_n\} \cap \{z_1, \ldots, z_n\}$ .) By Theorem 2.2 this implies m = 1, a contradiction.

**Theorem 4.3.** Let  $\mathbf{y}$  and  $\mathbf{z}$  be disjoint clusters of  $\mathcal{A}(\mathbf{x}, B)$ , and let U be a factorial subalgebra of  $\mathcal{A}(\mathbf{x}, B)$  such that

$$\{y_1, \ldots, y_n, z_1, \ldots, z_n, x_{n+1}^{\pm 1}, \ldots, x_p^{\pm 1}, x_{p+1}, \ldots, x_m\} \subset U.$$

Then we have

$$U = \mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, \mathbf{z}).$$

*Proof.* For m = 1 the statement follows from Lemma 4.1. Thus assume that m > 1. Let  $u \in U(\mathbf{y}, \mathbf{z}) = \mathscr{L}_{\mathbf{y}} \cap \mathscr{L}_{\mathbf{z}}$ . Thus we have

$$u = \frac{f}{y_1^{a_1} y_2^{a_2} \cdots y_p^{a_p}} = \frac{g}{z_1^{b_1} z_2^{b_2} \cdots z_p^{b_p}}$$

where f is a polynomial in  $y_1, \ldots, y_m$ , and g is a polynomial in  $z_1, \ldots, z_m$ , and  $a_i, b_i \ge 0$  for all  $1 \le i \le p$ . By the Laurent phenomenon it is enough to show that  $u \in U$ .

Since  $y_i, z_i \in U$  for all  $1 \leq i \leq m$ , we get the identity

$$fz_1^{b_1}z_2^{b_2}\cdots z_n^{b_n}z_{n+1}^{b_{n+1}}\cdots z_p^{b_p} = gy_1^{a_1}y_2^{a_2}\cdots y_n^{a_n}y_{n+1}^{a_{n+1}}\cdots y_p^{a_p}$$

in U.

By Theorem 3.1 the cluster variables  $y_i$  and  $z_i$  with  $1 \leq i \leq n$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . In particular, they are irreducible in the subalgebra U of  $\mathcal{A}(\mathbf{x}, B)$ . The elements  $y_{n+1}^{a_{n+1}} \cdots y_p^{a_p}$  and  $z_{n+1}^{b_{n+1}} \cdots z_p^{b_p}$  are units in U. (Recall that  $x_i = y_i = z_i$  for all  $n+1 \leq i \leq m$ .)

We have  $\{y_1, \ldots, y_n\} \cap \{z_1, \ldots, z_n\} = \emptyset$ , since **y** and **z** are disjoint. Thus, by the factoriality of U, the monomial  $y_1^{a_1} y_2^{a_2} \cdots y_n^{a_n}$  divides f in U. In other words there is some  $h \in U$  with  $f = hy_1^{a_1} y_2^{a_2} \cdots y_n^{a_n}$ . It follows that

$$u = \frac{f}{y_1^{a_1}y_2^{a_2}\cdots y_p^{a_p}} = \frac{hy_1^{a_1}y_2^{a_2}\cdots y_n^{a_n}}{y_1^{a_1}y_2^{a_2}\cdots y_p^{a_p}} = \frac{h}{y_{n+1}^{a_{n+1}}\cdots y_p^{a_p}} = hy_{n+1}^{-a_{n+1}}\cdots y_p^{-a_p}.$$

Since  $h \in U$  and  $y_{n+1}^{\pm 1}, \ldots, y_p^{\pm 1} \in U$ , we get  $u \in U$ . This finishes the proof.

- (i) If y and z are disjoint clusters of  $\mathcal{A}(\mathbf{x}, B)$ , then  $\mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, \mathbf{z})$ .
- (ii) For any  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$  we have  $\mathcal{A}(\mathbf{x}, B) = U(\mathbf{y}, C)$ .

**Corollary 4.4.** Assume that  $\mathcal{A}(\mathbf{x}, B)$  is factorial.

*Proof.* Part (i) follows directly from Theorem 1.4. To prove part (ii), assume  $(\mathbf{y}, C) \sim (\mathbf{x}, B)$  and let  $u \in U(\mathbf{y}, C)$ . For  $1 \leq k \leq n$  let  $(\mathbf{y}_k, C_k) := \mu_k(\mathbf{y}, C)$  and  $y_k^* := \mu_{(\mathbf{y}, C)}(y_k)$ . We get

$$u = \frac{f}{y_1^{a_1} \cdots y_k^{a_k} \cdots y_p^{a_p}} = \frac{f_k}{y_1^{b_1} \cdots (y_k^*)^{b_k} \cdots y_p^{b_p}}$$

for a polynomial f in  $y_1, \ldots, y_k, \ldots, y_m$ , a polynomial  $f_k$  in  $y_1, \ldots, y_k^*, \ldots, y_m$ , and  $a_i, b_i \ge 0$ . This yields an equality

(3) 
$$fy_1^{b_1}\cdots(y_k^*)^{b_k}\cdots y_p^{b_p} = f_k y_1^{a_1}\cdots y_k^{a_k}\cdots y_p^{a_p}$$

in  $\mathcal{A}(\mathbf{x}, B)$ . Now we argue similarly as in the proof of Theorem 4.3. By Theorem 3.1 the cluster variables  $y_1, \ldots, y_n$  and  $y_1^*, \ldots, y_n^*$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . By the factoriality of  $\mathcal{A}(\mathbf{x}, B)$ , Equation (3) implies that  $y_k^{a_k}$  divides f in  $\mathcal{A}(\mathbf{x}, B)$ . Since this holds for all  $1 \leq k \leq n$ , we get that  $y_1^{a_1} \cdots y_k^{a_k} \cdots y_n^{a_n}$  divides f in  $\mathcal{A}(\mathbf{x}, B)$ . It follows that  $u \in \mathcal{A}(\mathbf{x}, B)$ .

4.2. Existence of disjoint clusters. One assumption of Theorem 4.3 is the existence of disjoint clusters in  $\mathcal{A}(\mathbf{x}, B)$ . We can prove this under a mild assumption. But it should be true in general.

We say that a cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  satisfies the *separation property* if the following holds: If  $(y_k, y_k^*)$  is an exchange pair of  $\mathcal{A}(\mathbf{x}, B)$ , then there is no cluster  $\mathbf{z}$  of  $\mathcal{A}(\mathbf{x}, B)$  such that  $\{y_k, y_k^*\} \subseteq \{z_1, \ldots, z_m\}$ .

For example, if  $B^{\circ}$  is skew-symmetric, one can use the *E*-invariant introduced in [DWZ] to show that  $\mathcal{A}(\mathbf{x}, B)$  satisfies the separation property. Also, if all cluster monomials of  $\mathcal{A}(\mathbf{x}, B)$  are linearly independent, the separation property is obviously satisfied.

**Proposition 4.5.** Assume that  $\mathcal{A}(\mathbf{x}, B)$  satisfies the separation property. Let  $(\mathbf{y}, C)$  be a seed of  $\mathcal{A}(\mathbf{x}, B)$ , and let  $(\mathbf{z}, D) := \mu_n \cdots \mu_2 \mu_1(\mathbf{y}, C)$ . Then the clusters  $\mathbf{y}$  and  $\mathbf{z}$  are disjoint.

*Proof.* Set  $(\mathbf{y}[0], C[0]) := (\mathbf{y}, C)$ , and for  $1 \le k \le n$  let  $(\mathbf{y}[k], C[k]) := \mu_k(\mathbf{y}[k-1], C[k-1])$  and  $(y_1[k], \ldots, y_m[k]) := \mathbf{y}[k]$ . We claim that

$$\{y_1[k],\ldots,y_k[k]\}\cap\{y_1,\ldots,y_n\}=\emptyset.$$

For k = 1 this is straightforward. Thus let  $k \ge 2$ , and assume that our claim is true for k - 1. To get a contradiction, assume that  $y_k[k] \in \{y_1, \ldots, y_n\}$ . We have

$$\mathbf{y}[k] = (y_1[k], \dots, y_k[k], y_{k+1}, \dots, y_m).$$

Since  $y_1[k], \ldots, y_k[k], y_{k+1}, \ldots, y_m$  are algebraically independent and  $y_k[k] \neq y_k$ , we get  $y_k[k] \in \{y_1, \ldots, y_{k-1}\}$ , say  $y_k[k] = y_j$  for some  $1 \leq j \leq k-1$ . Since  $(\mathbf{y}[j], C[j]) = \mu_j(\mathbf{y}[j-1], C[j-1])$ , it follows that  $(y_j[j-1], y_j[j])$  is an exchange pair of  $\mathcal{A}(\mathbf{x}, B)$ . Next, observe that  $y_k[k] = y_j = y_j[j-1]$  and  $y_j[k] = y_j[j]$ . Thus  $y_j[j-1]$  and  $y_j[j]$  are both contained in  $\{y_1[k], \ldots, y_m[k]\}$ , a contradiction to the separation property.  $\Box$ 

### 5. The divisibility group of a cluster algebra

Let R be an integral domain, and let Frac(R) be the field of fractions of R. The abelian group

$$G(R) := (\operatorname{Frac}(R)^* / R^{\times}, \cdot)$$

is the *divisibility group* of R.

For  $g, h \in \operatorname{Frac}(R)^*$  let  $g \leq h$  provided  $hg^{-1} \in R$ . This relation is reflexive and transitive and it induces a partial ordering on G(R).

Let I be a set. The abelian group  $(\mathbb{Z}^{(I)}, +)$  is equipped with the following partial ordering: We set  $(x_i)_{i \in I} \leq (y_i)_{i \in I}$  if  $x_i \leq y_i$  for all i. (By definition, the elements in  $\mathbb{Z}^{(I)}$  are tuples  $(x_i)_{i \in I}$  of integers  $x_i$  such that only finitely many  $x_i$  are non-zero.)

There is the following well-known criterion for the factoriality of R, see for example [C, Section 2].

**Proposition 5.1.** For an integral domain R the following are equivalent:

- (i) R is factorial.
- (ii) There is a set I and a group isomorphism

 $\phi \colon G(R) \to \mathbb{Z}^{(I)}$ 

such that for all  $g, h \in G(R)$  we have  $g \leq h$  if and only if  $\phi(g) \leq \phi(h)$ .

Not all cluster algebras  $\mathcal{A}(\mathbf{x}, B)$  are factorial, but at least one part of the above factoriality criterion is satisfied:

**Proposition 5.2.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ . For m > 1 the divisibility group  $G(\mathcal{A}(\mathbf{x}, B))$  is isomorphic to  $\mathbb{Z}^{(I)}$ , where

 $I := \{f \in K[x_1, \dots, x_m] \mid f \text{ is irreducible and } f \neq x_i \text{ for } n+1 \leq i \leq p\} / K^*$ 

is the set of irreducible polynomials unequal to any  $x_{n+1}, \ldots, x_p$  in  $K[x_1, \ldots, x_m]$  up to non-zero scalar multiples.

*Proof.* By the Laurent phenomenon and the definition of a seed we get

 $\operatorname{Frac}(\mathcal{A}(\mathbf{x}, B)) = \operatorname{Frac}(\mathscr{L}_{\mathbf{x}}) = K(x_1, \dots, x_m).$ 

Furthermore, by Theorem 2.2 we have

$$\mathcal{A}(\mathbf{x}, B)^{\times} = \{\lambda x_{n+1}^{a_{n+1}} \cdots x_p^{a_p} \mid \lambda \in K^*, a_i \in \mathbb{Z}\}.$$

Any element in  $K(x_1, \ldots, x_m)$  is of the form  $f_1 \cdots f_s g_1^{-1} \cdots g_t^{-1}$  with  $f_i, g_j$  irreducible in  $K[x_1, \ldots, x_m]$ . Using that the polynomial ring  $K[x_1, \ldots, x_m]$  is factorial, and working modulo  $\mathcal{A}(\mathbf{x}, B)^{\times}$  yields the result.  $\Box$ 

6. Examples of non-factorial cluster algebras

6.1. For a matrix  $A \in M_{m,n}(\mathbb{Z})$  and  $1 \leq i \leq n$  let  $c_i(A)$  be the *i*th column of A.

**Proposition 6.1.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ . Assume that  $c_k(B) = c_s(B)$  or  $c_k(B) = -c_s(B)$  for some  $k \neq s$  with  $b_{ks} = 0$ . Then  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

*Proof.* Define  $(\mathbf{y}, C) := \mu_k(\mathbf{x}, B)$  and  $(\mathbf{z}, D) := \mu_s(\mathbf{y}, C)$ . We get  $y_k = z_k = x_k^{-1}(M_1 + M_2)$ 

where

$$M_1 := \prod_{b_{ik}>0} x_i^{b_{ik}}$$
 and  $M_2 := \prod_{b_{ik}<0} x_i^{-b_{ik}}$ 

By the mutation rule, we have  $c_k(C) = -c_k(B)$ , and since  $b_{ks} = 0$ , we get  $c_s(C) = c_s(B)$ . Since  $c_s(B) = c_k(B)$  or  $c_s(B) = -c_k(B)$ , this implies that

$$z_s = x_s^{-1}(M_1 + M_2).$$

Using Corollary 2.3 one easily checks that the cluster variables  $x_k, x_s, z_k, z_s$  are pairwise non-associate, and by Theorem 3.1 they are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . Obviously, we have

$$x_k z_k = x_s z_s$$

Thus  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

To give a concrete example of a cluster algebra, which is not factorial, assume m = n = p = 3, and let  $B \in M_{m,n}(\mathbb{Z})$  be the matrix

$$B = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

The matrix B obviously satisfies the assumptions of Proposition 6.1. Note that  $B = B^{\circ}$  is skew-symmetric, and that  $\Gamma(B)$  is the quiver

$$3 \longrightarrow 2 \longrightarrow 1.$$

Thus  $\mathcal{A}(\mathbf{x}, B)$  is a cluster algebra of Dynkin type  $\mathbb{A}_3$ . (Cluster algebras with finitely many cluster variables are classified via Dynkin types, for details see [FZ2].)

Define  $(\mathbf{z}, D) := \mu_3 \mu_1(\mathbf{x}, B)$ . We get  $z_1 = x_1^{-1}(1 + x_2)$ ,  $z_3 = x_3^{-1}(1 + x_2)$  and therefore  $x_1 z_1 = x_3 z_3$ . Using Corollary 2.3 we see that  $x_1, x_3, z_1, z_3$  are pairwise non-associate, and by Theorem 3.1 they are irreducible. Thus  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

6.2. The next example is due to Philipp Lampe. It gives a negative answer to Zelevinsky's Question 1.6.

**Proposition 6.2** ([La]). Let  $K = \mathbb{C}$ , m = n = 2 and

$$B = \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix}.$$

Then  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

*Proof.* Let  $y_1 := \mu_{(\mathbf{x},B)}(x_1)$ . The corresponding exchange relation is

$$x_1y_1 = 1 + x_2^2 = (x_2 + i)(x_2 - i),$$

where  $i \in \mathbb{C}$  with  $i^2 = -1$ . As polynomials in  $\mathbb{C}[x_2]$  the elements  $x_2 + i$  and  $x_2 - i$  are obviously both contained in  $\mathcal{A}(\mathbf{x}, B)$ . To get a contradiction, assume that  $\mathcal{A}(\mathbf{x}, B)$ is factorial. By Theorem 2.2 the elements  $x_2 + i$  and  $x_2 - i$  are not invertible in  $\mathcal{A}(\mathbf{x}, B)$ . By the factoriality of  $\mathcal{A}(\mathbf{x}, B)$ , each element in  $\mathcal{A}(\mathbf{x}, B)$  is a product of

irreducible elements. By Theorem 3.1 the cluster variables  $x_1$  and  $y_1$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . Now the identity  $x_1y_1 = (x_2 + i)(x_2 - i)$  and the factoriality of  $\mathcal{A}(\mathbf{x}, B)$ imply that  $x_2 + i$  and  $x_2 - i$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . Obviously, the elements  $x_1, y_1, x_2 + i, x_2 - i$  are pairwise non-associate, a contradiction to the factoriality of  $\mathcal{A}(\mathbf{x}, B)$ .

The following result is a straightforward generalization of Lampe's example.

**Proposition 6.3.** Let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$  with m > 1. Assume that there exists some  $1 \le k \le n$  such that the following hold:

- (i) For  $1 \leq i \leq m$  we have  $b_{ik} \geq 0$ .
- (ii) Let X be a variable. The polynomial  $1+X^d$  is not irreducible in K[X], where  $d := \gcd(b_{1k}, \ldots, b_{mk})$  is the greatest common divisor of  $b_{1k}, \ldots, b_{mk}$ .

Then  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

*Proof.* Let  $1 + X^d = f_1 \cdots f_t$ , where the  $f_j$  are irreducible polynomials in K[X]. Since  $1 + X^d$  is not irreducible in K[X], we have  $t \ge 2$ . Let  $y_k := \mu_{(\mathbf{x},B)}(x_k)$ . The corresponding exchange relation is

$$x_k y_k = 1 + \prod_{i=1}^m x_i^{b_{ik}} = 1 + M^d = \prod_{j=1}^t f_j(M),$$

where

$$M := \prod_{i=1}^m x_i^{b_{ik}/d}.$$

Clearly, each  $f_j(M)$  is contained in  $\mathcal{A}(\mathbf{x}, B)$ . To get a contradiction, assume that  $\mathcal{A}(\mathbf{x}, B)$  is factorial. By Theorem 2.2 none of the elements  $f_j(M)$  is invertible in  $\mathcal{A}(\mathbf{x}, B)$ . Since  $\mathcal{A}(\mathbf{x}, B)$  is factorial, each  $f_j(M)$  is equal to a product  $f_{1j} \cdots f_{a_jj}$ , where the  $f_{ij}$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$  and  $a_j \geq 1$ . By Theorem 3.1 the cluster variables  $x_k$  and  $y_k$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . It follows that  $a_1 + \cdots + a_t = 2$ , since  $\mathcal{A}(\mathbf{x}, B)$  is factorial. This implies t = 2 and  $a_1 = a_2 = 1$ . In particular,  $f_1(M)$  and  $f_2(M)$  are irreducible in  $\mathcal{A}(\mathbf{x}, B)$ , and we have  $x_k y_k = f_1(M) f_2(M)$ . For j = 1, 2 the elements  $x_k$  and  $f_j(M)$  cannot be associate, since  $f_j(M)$  is just a K-linear combination of powers of M and  $b_{kk} = 0$ . This is a contradiction to the factoriality of  $\mathcal{A}(\mathbf{x}, B)$ .

Corollary 6.4. Let  $K = \mathbb{C}$ , m = n = 2 and

$$B = \begin{pmatrix} 0 & -c \\ d & 0 \end{pmatrix}$$

with  $c \geq 1$  and  $d \geq 2$ . Then  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

*Proof.* For k = 1 assumption (i) of Proposition 6.3 holds, and we have gcd(0, d) = d. The polynomial  $1 + X^d$  is not irreducible in  $\mathbb{C}[X]$ . Thus assumption (ii) also holds.

Corollary 6.5. Let m = n = 2 and

$$B = \begin{pmatrix} 0 & -c \\ d & 0 \end{pmatrix}$$

with  $c \geq 1$  and  $d \geq 3$  an odd number. Then  $\mathcal{A}(\mathbf{x}, B)$  is not factorial.

*Proof.* For k = 1 assumption (i) of Proposition 6.3 holds, and we have gcd(0, d) = d. For odd d we have

$$1 + X^{d} = (X + 1) \left( \sum_{j=0}^{d-1} (-1)^{j} X^{j} \right)$$

Thus  $1 + X^d$  is not irreducible in K[X], and therefore assumption (ii) also holds.  $\Box$ 

## 7. Examples of factorial cluster algebras

7.1. Cluster algebras of Dynkin type A as polynomial rings. Assume m = n + 1 = p + 1, and let  $B \in M_{m,n}(\mathbb{Z})$  be the matrix

$$B = \begin{pmatrix} 0 & -1 & & & \\ 1 & 0 & -1 & & \\ & 1 & 0 & \ddots & \\ & & 1 & \ddots & -1 & \\ & & & \ddots & 0 & -1 \\ & & & & 1 & 0 \\ \hline & & & & & 1 \end{pmatrix}$$

Obviously,  $B^{\circ}$  is skew-symmetric,  $\Gamma(B)$  is the quiver

 $m \longrightarrow \cdots \longrightarrow 2 \longrightarrow 1,$ 

and  $\mathcal{A}(\mathbf{x}, B)$  is a cluster algebra of Dynkin type  $\mathbb{A}_n$ . Note that  $\mathcal{A}(\mathbf{x}, B)$  has exactly one coefficient, and that this coefficient is non-invertible.

Let  $(\mathbf{x}[0], B[0]) := (\mathbf{x}, B)$ . For each  $1 \le i \le m-1$  we define inductively a seed by  $(\mathbf{x}[i], B[i]) := \mu_{m-i} \cdots \mu_2 \mu_1 (\mathbf{x}[i-1], B[i-1]).$ 

For  $0 \le i \le m - 1$  set  $(x_1[i], \dots, x_m[i]) := \mathbf{x}[i]$ .

For simplicity we define  $x_0[i] := 1$  and  $x_{-1}[i] := 0$  for all i.

**Lemma 7.1.** For  $0 \le i \le m-2$ ,  $1 \le k \le m-1-i$  and  $0 \le j \le i$  we have

(4) 
$$\mu_{(\mathbf{x}[i],B[i])}(x_k[i]) = \frac{x_{k-1}[i] + x_{k+1}[i]}{x_k[i]} = \frac{x_{k-1+j}[i-j] + x_{k+1+j}[i-j]}{x_{k+j}[i-j]}$$

*Proof.* The first equality follows from the definition of  $(\mathbf{x}[i], B[i])$  and the mutation rule. The second equality is proved by induction on i.

Corollary 7.2. For  $0 \le i \le m - 2$  we have

(5) 
$$x_{i+2} = x_1[i+1]x_{i+1} - x_i,$$

(6)  $x_{i+1}[1] = x_1[i+1]x_i[1] - x_{i-1}[1].$ 

*Proof.* Equation (5) follows from (4) for k = 1 and j = i. The case k = 1 and j = i - 1 yields Equation (6).

**Proposition 7.3.** The elements  $x_1[0], x_1[1], \ldots, x_1[m-1]$  are algebraically independent and

$$K[x_1[0], x_1[1], \dots, x_1[m-1]] = \mathcal{A}(\mathbf{x}, B).$$

In particular, the cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is a polynomial ring in m variables.

*Proof.* It follows from Equation (5) that

$$x_1[i] \in K(x_1, \dots, x_{i+1}) \setminus K(x_1, \dots, x_i)$$

for all  $1 \leq i \leq m-1$ . Since  $x_1, \ldots, x_m$  are algebraically independent, this implies that  $x_1[0], x_1[1], \ldots, x_1[m-1]$  are algebraically independent as well. Thus

$$U := K[x_1[0], x_1[1], \dots, x_1[m-1]]$$

is a polynomial ring in m variables. In particular, U is factorial. Equation (5) implies that  $x_1, \ldots, x_m \in U$ , and Equation (6) yields that  $x_1[1], \ldots, x_m[1] \in U$ . Thus the assumptions of Theorem 4.3 are satisfied and we get  $U = \mathcal{A}(\mathbf{x}, B)$ .

The cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  as defined above has been studied by several people. It is related to a *T*-system of Dynkin type  $\mathbb{A}_1$  with a certain boundary condition, see [DK]. Furthermore, for  $K = \mathbb{C}$  the cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is naturally isomorphic to the complexified Grothendieck ring of the category  $\mathcal{C}_n$  of finite-dimensional modules of level *n* over the quantum loop algebra of Dynkin type  $\mathbb{A}_1$ , see [HL, N2]. It is well known, that  $\mathcal{A}(\mathbf{x}, B)$  is a polynomial ring. We just wanted to demonstrate how to use Theorem 4.3 in practise.

7.2. Acyclic cluster algebras as polynomial rings. Let  $C = (c_{ij}) \in M_{n,n}(\mathbb{Z})$  be a generalized Cartan matrix, i.e. C is symmetrizable,  $c_{ii} = 2$  for all i and  $c_{ij} \leq 0$  for all  $i \neq j$ .

Assume that m = 2n = 2p, and let  $(\mathbf{x}, B)$  be a seed of  $\mathcal{F}$ , where  $B = (b_{ij}) \in M_{2n,n}(\mathbb{Z})$  is defined as follows: For  $1 \leq i \leq 2n$  and  $1 \leq j \leq n$  let

$$b_{ij} := \begin{cases} 0 & \text{if } i = j, \\ -c_{ij} & \text{if } 1 \le i < j \le n, \\ c_{ij} & \text{if } 1 \le j < i \le n, \\ 1 & \text{if } i = n + j, \\ c_{i-n,j} & \text{if } n+1 \le i \le 2n \text{ and } i-n < j, \\ 0 & \text{if } n+1 \le i \le 2n \text{ and } i-n > j. \end{cases}$$

Thus we have

$$B = \begin{pmatrix} 0 & b_{12} & b_{13} & \cdots & b_{1n} \\ b_{21} & 0 & b_{23} & \cdots & b_{2n} \\ b_{31} & b_{32} & 0 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & b_{n-1,n} \\ b_{n1} & b_{n2} & \cdots & b_{n,n-1} & 0 \\ \hline 1 & -b_{12} & -b_{13} & \cdots & -b_{1n} \\ 0 & 1 & -b_{23} & \cdots & -b_{2n} \\ 0 & 0 & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & -b_{n-1,n} \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Clearly,  $(\mathbf{x}, B)$  is an acyclic seed. Namely, if  $i \to j$  is an arrow in  $\Sigma(B)$ , then i < j. Up to simultaneous reordering of columns and rows, each acyclic skew-symmetrizable matrix in  $M_{n,n}(\mathbb{Z})$  is of the form  $B^{\circ}$  with B defined as above. Note that  $\mathcal{A}(\mathbf{x}, B)$ has exactly n coefficients, and that all these coefficients are non-invertible.

For  $1 \leq i \leq n$  let

$$(\mathbf{x}[1], B[1]) := \mu_n \cdots \mu_2 \mu_1(\mathbf{x}, B)$$

and  $(x_1[1], \ldots, x_{2n}[1]) := \mathbf{x}[1]$ . Let  $B_0 := B$ , and for  $1 \le i \le n$  let  $B_i := \mu_i(B_{i-1})$ . Thus we have  $B_n = B[1]$ . It is easy to work out the matrices  $B_i$  explicitly: The matrix  $B_i$  is obtained from  $B_{i-1}$  by changing the sign in the *i*th row and the *i*th column of the principal part  $B_{i-1}^{\circ}$ . Furthermore, the (n+i)th row

$$(0,\ldots,0,1,-b_{i,i+1},-b_{i,i+2},\ldots,-b_{in})$$

of  $B_{i-1}$  gets replaced by

$$(-b_{i1}, -b_{i2}, \ldots, -b_{i,i-1}, -1, 0, \ldots, 0).$$

As an illustration, for n = 3 the matrices  $B_i$  look as follows:

$$B_{0} = \begin{pmatrix} 0 & b_{12} & b_{13} \\ b_{21} & 0 & b_{23} \\ b_{31} & b_{32} & 0 \\ \hline 1 & -b_{12} & -b_{13} \\ 0 & 1 & -b_{23} \\ 0 & 0 & 1 \end{pmatrix}, \qquad B_{1} = \begin{pmatrix} 0 & -b_{12} & -b_{13} \\ -b_{21} & 0 & b_{23} \\ \hline -b_{31} & b_{32} & 0 \\ \hline -1 & 0 & 0 \\ 0 & 1 & -b_{23} \\ 0 & 0 & 1 \end{pmatrix},$$
$$B_{2} = \begin{pmatrix} 0 & b_{12} & -b_{13} \\ b_{21} & 0 & -b_{23} \\ \hline -b_{31} & -b_{32} & 0 \\ \hline -1 & 0 & 0 \\ \hline -b_{21} & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad B_{3} = \begin{pmatrix} 0 & b_{12} & b_{13} \\ b_{21} & 0 & b_{23} \\ \hline b_{31} & b_{32} & 0 \\ \hline -1 & 0 & 0 \\ \hline -b_{21} & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

If we write  $N_+$  (resp.  $N_-$ ) for the upper (resp. lower) triangular part of  $B^{\circ}$ , we get

$$B = \begin{pmatrix} 1 & N_{+} \\ & \ddots & \\ N_{-} & 1 \\ \hline 1 & -N_{+} \\ & \ddots & \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad B[1] = \begin{pmatrix} 1 & N_{+} \\ & \ddots & \\ N_{-} & 1 \\ \hline -1 & 0 \\ & \ddots & \\ -N_{-} & -1 \end{pmatrix}.$$

In particular, the principal part  $B^{\circ}$  of B is equal to the principal part  $B[1]^{\circ}$  of B[1].

Now the definition of seed mutation yields

(7) 
$$x_k[1] = x_k^{-1} \left( x_{n+k} + \prod_{i=1}^{k-1} x_i[1]^{b_{ik}} \prod_{i=k+1}^n x_i^{-b_{ik}} \right)$$

for  $1 \leq k \leq n$ .

**Proposition 7.4.** The elements  $x_1, \ldots, x_n, x_1[1], \ldots, x_n[1]$  are algebraically independent and

$$K[x_1,\ldots,x_n,x_1[1],\ldots,x_n[1]] = \mathcal{A}(\mathbf{x},B).$$

In particular, the cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is a polynomial ring in 2n variables.

*Proof.* By Equation (7) and induction we have

$$x_k[1] \in K(x_1, \dots, x_{n+k}) \setminus K(x_1, \dots, x_{n+k-1})$$

for all  $1 \leq k \leq n$ . It follows that  $x_1, \ldots, x_n, x_1[1], \ldots, x_n[1]$  are algebraically independent. Let

$$U := K[x_1, \dots, x_n, x_1[1], \dots, x_n[1]].$$

Thus U is a polynomial ring in 2n variables. In particular, U is factorial. It follows from Equation (7) that

(8) 
$$x_{n+k} = x_k[1]x_k - \prod_{i=1}^{k-1} x_i[1]^{b_{ik}} \prod_{i=k+1}^n x_i^{-b_{ik}}.$$

This implies  $x_{n+k} \in U$  for all  $1 \leq k \leq n$ . Thus the assumptions of Theorem 4.3 are satisfied, and we can conclude that  $U = \mathcal{A}(\mathbf{x}, B)$ .

Proposition 7.4 is a special case of a much more general result proved in [GLS2]. But the proof presented here is new and more elementary.

Next, we compare the basis

$$\mathcal{P}_{\text{GLS}} := \left\{ x[\mathbf{a}] := x_1^{a_1} \cdots x_n^{a_n} x_1[1]^{a_{n+1}} \cdots x_n[1]^{a_{2n}} \mid \mathbf{a} = (a_1, \dots, a_{2n}) \in \mathbb{N}^{2n} \right\}$$

of  $\mathcal{A}(\mathbf{x}, B)$  resulting from Proposition 7.4 with a basis constructed by Berenstein, Fomin and Zelevinsky [BFZ]. For  $1 \leq k \leq n$  let

(9) 
$$x'_{k} := \mu_{(\mathbf{x},B)}(x_{k}) = x_{k}^{-1} \left( x_{n+k} \prod_{i=1}^{k-1} x_{i}^{b_{ik}} + \prod_{i=k+1}^{n} x_{i}^{-b_{ik}} \prod_{i=1}^{k-1} x_{n+i}^{b_{ik}} \right)$$

and set

$$\mathcal{P}_{BFZ} := \{ x'[\mathbf{a}] := x_1^{a_1} \cdots x_n^{a_n} x_{n+1}^{a_{n+1}} \cdots x_{2n}^{a_{2n}} (x'_1)^{a_{2n+1}} \cdots (x'_n)^{a_{3n}} \mid \mathbf{a} = (a_1, \dots, a_{3n}) \in \mathbb{N}^{3n}, a_k a_{2n+k} = 0 \text{ for } 1 \le k \le n \}.$$

**Proposition 7.5** ([BFZ, Corollary 1.21]). The set  $\mathcal{P}_{BFZ}$  is a basis of  $\mathcal{A}(\mathbf{x}, B)$ .

Note that the basis  $\mathcal{P}_{\text{GLS}}$  is constructed by using cluster variables from two seeds, namely  $(\mathbf{x}, B)$  and  $\mu_n \cdots \mu_1(\mathbf{x}, B)$ , whereas  $\mathcal{P}_{\text{BFZ}}$  uses cluster variables from n + 1 seeds, namely  $(\mathbf{x}, B)$  and  $\mu_k(\mathbf{x}, B)$ , where  $1 \le k \le n$ .

Now we insert Equation (8) into Equation (9) and obtain

(10) 
$$x_{k}x_{k}' = \left(x_{k}[1]x_{k} - \prod_{i=1}^{k-1} x_{i}[1]^{b_{ik}} \prod_{i=k+1}^{n} x_{i}^{-b_{ik}}\right) \prod_{i=1}^{k-1} x_{i}^{b_{ik}} + \prod_{i=k+1}^{n} x_{i}^{-b_{ik}} \prod_{i=1}^{k-1} \left(x_{i}[1]x_{i} - \prod_{j=1}^{i-1} x_{j}[1]^{b_{ji}} \prod_{j=i+1}^{n} x_{j}^{-b_{ji}}\right)^{b_{ik}}$$

Then we observe that the right-hand side of Equation (10) is divisible by  $x_k$  and that  $x'_k$  is a polynomial in  $x_1, \ldots, x_n, x_1[1], \ldots, x_n[1]$ . Thus we can express every element of the basis  $\mathcal{P}_{BFZ}$  explicitly as a linear combination of vectors from the basis  $\mathcal{P}_{GLS}$ .

One could use Equation (10) to get an alternative proof of Proposition 7.4 as pointed out by Zelevinsky [Z]. Vice versa, using Proposition 7.4 yields another proof that  $\mathcal{P}_{BFZ}$  is a basis.

7.3. Cluster algebras arising in Lie theory as polynomial rings. The next class of examples can be seen as a fusion of the examples discussed in Sections 7.1 and 7.2. In the following we use the same notation as in [GLS2].

Let  $C \in M_{n,n}(\mathbb{Z})$  be a symmetric generalized Cartan matrix, and let  $\mathfrak{g}$  be the associated Kac-Moody Lie algebra over  $K = \mathbb{C}$  with triangular decomposition  $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}$ , see [K].

Let  $U(\mathfrak{n})_{\mathrm{gr}}^*$  be the graded dual of the enveloping algebra  $U(\mathfrak{n})$  of  $\mathfrak{n}$ . To each element w in the Weyl group W of  $\mathfrak{g}$  one can associate a subalgebra  $\mathcal{R}(\mathcal{C}_w)$  of  $U(\mathfrak{n})_{\mathrm{gr}}^*$  and a cluster algebra  $\mathcal{A}(\mathcal{C}_w)$ , see [GLS2]. Here  $\mathcal{C}_w$  denotes a Frobenius category associated to w, see [BIRS, GLS2].

In [GLS2] we constructed a natural algebra isomorphism

$$\mathcal{A}(\mathcal{C}_w) \to \mathcal{R}(\mathcal{C}_w).$$

This yields a cluster algebra structure on  $\mathcal{R}(\mathcal{C}_w)$ .

Let  $\mathbf{i} = (i_r, \ldots, i_1)$  be a reduced expression of w. In [GLS2] we studied two clustertilting modules  $V_{\mathbf{i}} = V_1 \oplus \cdots \oplus V_r$  and  $T_{\mathbf{i}} = T_1 \oplus \cdots \oplus T_r$  in  $\mathcal{C}_w$ , which are associated to  $\mathbf{i}$ . These modules yield two disjoint clusters  $(\delta_{V_1}, \ldots, \delta_{V_r})$  and  $(\delta_{T_1}, \ldots, \delta_{T_r})$  of  $\mathcal{R}(\mathcal{C}_w)$ . The exchanges matrices are of size  $r \times (r - n)$ . In contrast to our conventions in this article, the n coefficients are  $\delta_{V_k} = \delta_{T_k}$  with  $k^+ = r + 1$ , where  $k^+$  is defined as in [GLS2], and none of these coefficients is invertible. Furthermore, we studied a module  $M_{\mathbf{i}} = M_1 \oplus \cdots \oplus M_r$  in  $\mathcal{C}_w$ , which yields cluster variables  $\delta_{M_1}, \ldots, \delta_{M_r}$  of  $\mathcal{R}(\mathcal{C}_w)$ . (These do not form a cluster.) Using methods from Lie theory we obtained the following result.

**Theorem 7.6** ([GLS2, Theorem 3.2]). The cluster algebra  $\mathcal{R}(\mathcal{C}_w)$  is a polynomial ring in the variables  $\delta_{M_1}, \ldots, \delta_{M_r}$ .

To obtain an alternative proof of Theorem 7.6, one can proceed as follows:

- (i) Show that the cluster variables  $\delta_{M_1}, \ldots, \delta_{M_r}$  are algebraically independent.
- (ii) Show that for  $1 \leq k \leq r$  the cluster variables  $\delta_{V_k}$  and  $\delta_{T_k}$  are polynomials in  $\delta_{M_1}, \ldots, \delta_{M_r}$ .
- (iii) Apply Theorem 4.3.

Part (i) can be done easily using induction and the mutation sequence in [GLS2, Section 13]. Part (ii) is not at all straightforward.

Let us give a concrete example illustrating Theorem 7.6. Let  $\mathfrak{g}$  be the Kac-Moody Lie algebra associated to the generalized Cartan matrix

$$C = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix},$$

and let  $\mathbf{i} = (2, 1, 2, 1, 2, 1, 2, 1)$ . Then  $\mathcal{A}(\mathcal{C}_w) = \mathcal{A}(\mathbf{x}, B_{\mathbf{i}})$ , where r = n + 2 = 8,  $x_7$  and  $x_8$  are the (non-invertible) coefficients, and

$$B_{\mathbf{i}} = \begin{pmatrix} 0 & 2 & -1 & & \\ -2 & 0 & 2 & -1 & & \\ 1 & -2 & 0 & 2 & -1 & \\ & 1 & -2 & 0 & 2 & -1 \\ & & 1 & -2 & 0 & 2 \\ & & & 1 & -2 & 0 \\ \hline & & & & 1 & -2 \\ & & & & & 1 \end{pmatrix}$$

The principal part  $B_{\mathbf{i}}^{\circ}$  of  $B_{\mathbf{i}}$  is skew-symmetric, and the quiver  $\Gamma(B_{\mathbf{i}})$  looks as follows:



Define

$$\begin{aligned} &(\mathbf{x}[0], B[0]) := (\mathbf{x}, B_{\mathbf{i}}), \\ &(\mathbf{x}[1], B[1]) := \mu_{5}\mu_{3}\mu_{1}(\mathbf{x}[0], B[0]), \\ &(\mathbf{x}[3], B[3]) := \mu_{3}\mu_{1}(\mathbf{x}[2], B[2]), \\ &(\mathbf{x}[5], B[5]) := \mu_{1}(\mathbf{x}[4], B[4]), \end{aligned}$$

$$\begin{aligned} &(\mathbf{x}[2], B[2]) := \mu_{6}\mu_{4}\mu_{2}(\mathbf{x}[1], B[1]), \\ &(\mathbf{x}[4], B[4]) := \mu_{4}\mu_{2}(\mathbf{x}[3], B[3]), \\ &(\mathbf{x}[5], B[5]) := \mu_{1}(\mathbf{x}[4], B[4]), \end{aligned}$$

and for  $0 \le k \le 6$  let  $(x_1[k], \ldots, x_8[k]) := \mathbf{x}[k]$ .

Under the isomorphism  $\mathcal{A}(\mathcal{C}_w) \to \mathcal{R}(\mathcal{C}_w)$  the cluster  $\mathbf{x}[0]$  of  $\mathcal{A}(\mathcal{C}_w) = \mathcal{A}(\mathbf{x}, B_{\mathbf{i}})$ corresponds to the cluster  $(\delta_{V_1}, \ldots, \delta_{V_8})$  of  $\mathcal{R}(\mathcal{C}_w)$ , the cluster  $\mathbf{x}[6]$  corresponds to  $(\delta_{T_1}, \ldots, \delta_{T_8})$ , and we have

$$\begin{aligned} x_1[0] \mapsto \delta_{M_1}, & x_2[0] \mapsto \delta_{M_2}, & x_1[2] \mapsto \delta_{M_3}, & x_2[2] \mapsto \delta_{M_4}, \\ x_1[4] \mapsto \delta_{M_5}, & x_2[4] \mapsto \delta_{M_6}, & x_1[6] \mapsto \delta_{M_7}, & x_2[6] \mapsto \delta_{M_8}. \end{aligned}$$

By Theorem 7.6 we know that the cluster algebra  $\mathcal{A}(\mathbf{x}, B_{\mathbf{i}})$  is a polynomial ring in the variables  $x_1[0], x_2[0], x_1[2], x_2[2], x_1[4], x_2[4], x_1[6], x_2[6]$ .

### 8. Applications

8.1. Prime elements in the dual semicanonical basis. As in Section 7.3 let  $C \in M_{n,n}(\mathbb{Z})$  be a symmetric generalized Cartan matrix, and let  $\mathfrak{g} = \mathfrak{n}_{-} \oplus \mathfrak{h} \oplus \mathfrak{n}$  be the associated Lie algebra.

As before let W be the Weyl group of  $\mathfrak{g}$ . To C one can also associate a preprojective algebra  $\Lambda$  over  $\mathbb{C}$ , see for example [GLS2, R]

Lusztig [Lu] realized the universal enveloping algebra  $U(\mathfrak{n})$  of  $\mathfrak{n}$  as an algebra of constructible functions on the varieties  $\Lambda_d$  of nilpotent  $\Lambda$ -modules with dimension vector  $d \in \mathbb{N}^n$ . He also constructed the semicanonical basis S of  $U(\mathfrak{n})$ . The elements of S are naturally parametrized by the irreducible components of the varieties  $\Lambda_d$ .

An irreducible component Z of  $\Lambda_d$  is called *indecomposable* if it contains a Zariski dense subset of indecomposable  $\Lambda$ -modules, and Z is *rigid* if it contains a rigid  $\Lambda$ -module M, i.e. M is a module with  $\operatorname{Ext}^1_{\Lambda}(M, M) = 0$ .

Let  $\mathcal{S}^*$  be the dual semicanonical basis of the graded dual  $U(\mathfrak{n})^*_{gr}$  of  $U(\mathfrak{n})$ . The elements  $\rho_Z$  in  $\mathcal{S}^*$  are also parametrized by irreducible components Z of the varieties  $\Lambda_d$ . We call  $\rho_Z$  indecomposable (resp. rigid) if Z is indecomposable (resp. rigid). An element  $b \in \mathcal{S}^*$  is called *primitive* if it cannot be written as a product  $b = b_1b_2$ with  $b_1, b_2 \in \mathcal{S}^* \setminus \{1\}$ .

**Theorem 8.1** ([GLS1, Theorem 1.1]). If  $\rho_Z$  is primitive, then Z is indecomposable.

**Theorem 8.2** ([GLS2, Theorem 3.1]). For  $w \in W$  all cluster monomials of the cluster algebra  $\mathcal{R}(\mathcal{C}_w)$  belong to the dual semicanonical basis  $\mathcal{S}^*$  of  $U(\mathfrak{n})^*_{gr}$ . More precisely, we have

{cluster variables of  $\mathcal{R}(\mathcal{C}_w)$ }  $\subseteq$  { $\rho_Z \in \mathcal{S}^* \mid Z$  is indecomposable and rigid}, {cluster monomials of  $\mathcal{R}(\mathcal{C}_w)$ }  $\subseteq$  { $\rho_Z \in \mathcal{S}^* \mid Z$  is rigid}.

Combining Theorems 3.1, 7.6 and 8.2 we obtain a partial converse of Theorem 8.1.

**Theorem 8.3.** The cluster variables in  $\mathcal{R}(\mathcal{C}_w)$  are prime, and they are primitive elements of  $\mathcal{S}^*$ .

**Conjecture 8.4.** If  $\rho_Z \in S^*$  is indecomposable and rigid, then  $\rho_Z$  is prime in  $U(\mathfrak{n})^*_{gr}$ .

8.2. Monoidal categorifications of cluster algebras. Let  $\mathcal{C}$  be an abelian tensor category with unit object  $I_{\mathcal{C}}$ . We assume that  $\mathcal{C}$  is a Krull-Schmidt category, and that all objects in  $\mathcal{C}$  are of finite length. Let  $\mathcal{M}(\mathcal{C}) := \mathcal{K}_0(\mathcal{C})$  be the Grothendieck ring of  $\mathcal{C}$ . The class of an object  $M \in \mathcal{C}$  is denoted by [M]. The addition in  $\mathcal{M}(\mathcal{C})$  is given by  $[M]+[N] := [M \oplus N]$  and the multiplication is defined by  $[M][N] := [M \otimes N]$ . We assume that  $[M \otimes N] = [N \otimes M]$ . (In general this does not imply  $M \otimes N \cong N \otimes M$ .) Thus  $\mathcal{M}(\mathcal{C})$  is a commutative ring.

Tensoring with K over  $\mathbb{Z}$  yields a K-algebra  $\mathcal{M}_K(\mathcal{C}) := K \otimes_{\mathbb{Z}} \mathcal{K}_0(\mathcal{C})$  with K-basis the classes of simple objects in  $\mathcal{C}$ . Note that the unit object  $I_{\mathcal{C}}$  is simple.

A monoidal categorification of a cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is an algebra isomorphism

$$\Phi\colon \mathcal{A}(\mathbf{x},B)\to \mathcal{M}_K(\mathcal{C}),$$

where  $\mathcal{C}$  is a tensor category as above, such that each cluster monomial  $y = y_1^{a_1} \cdots y_m^{a_m}$  of  $\mathcal{A}(\mathbf{x}, B)$  is mapped to a class  $[S_y]$  of some simple object  $S_y \in \mathcal{C}$ . In particular, we have

$$[S_y] = [S_{y_1}]^{a_1} \cdots [S_{y_m}]^{a_m} = [S_{y_1}^{\otimes a_1} \otimes \cdots \otimes S_{y_m}^{\otimes a_m}].$$

For an object  $M \in \mathcal{C}$  let  $x_M$  be the element in  $\mathcal{A}(\mathbf{x}, B)$  with  $\Phi(x_M) = [M]$ .

The concept of a monoidal categorification of a cluster algebra was introduced in [HL, Definition 2.1]. But note that our definition uses weaker conditions than in [HL].

An object  $M \in \mathcal{C}$  is called *invertible* if [M] is invertible in  $\mathcal{M}_K(\mathcal{C})$ . An object  $M \in \mathcal{C}$  is *primitive* if there are no non-invertible objects  $M_1$  and  $M_2$  in  $\mathcal{C}$  with  $M \cong M_1 \otimes M_2$ .

**Proposition 8.5.** Let  $\Phi: \mathcal{A}(\mathbf{x}, B) \to \mathcal{M}_K(\mathcal{C})$  be a monoidal categorification of a cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  with m > 1. Then the following hold:

(i) The invertible elements in  $\mathcal{M}_K(\mathcal{C})$  are

$$\mathcal{M}_K(\mathcal{C})^{\times} = \left\{ \lambda[I_{\mathcal{C}}][S_{x_{n+1}}]^{a_{n+1}} \cdots [S_{x_p}]^{a_p} \mid \lambda \in K^*, a_i \in \mathbb{Z} \right\}.$$

(ii) Let M be an object in C such that the element  $x_M$  is irreducible in  $\mathcal{A}(\mathbf{x}, B)$ . Then M is primitive.

Proof. Part (i) follows directly from Theorem 2.2. To prove (ii), assume that M is not primitive. Thus there are non-invertible objects  $M_1$  and  $M_2$  in  $\mathcal{C}$  with  $M \cong M_1 \otimes$  $M_2$ . Thus in  $\mathcal{M}_K(\mathcal{C})$  we have  $[M] = [M_1][M_2]$ . Since  $\Phi$  is an algebra isomorphism, we get  $x_M = x_{M_1} x_{M_2}$  with  $x_{M_1}$  and  $x_{M_2}$  non-invertible in  $\mathcal{A}(\mathbf{x}, B)$ . Since  $x_M$  is irreducible, we have a contradiction.  $\Box$ 

Combining Proposition 8.5 with Theorem 3.1 we get the following result.

**Corollary 8.6.** Let  $\Phi: \mathcal{A}(\mathbf{x}, B) \to \mathcal{M}_K(\mathcal{C})$  be a monoidal categorification of a cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  with m > 1. For each cluster variable y of  $\mathcal{A}(\mathbf{x}, B)$ , the simple object  $S_y$  is primitive.

Examples of monoidal categorifications of cluster algebras can be found in [HL, N1], see also [Le].

Acknowledgements. We thank Giovanni Cerulli Irelli, Sergey Fomin, Daniel Labardini Fragoso, Philipp Lampe and Andrei Zelevinsky for helpful discussions. We thank Giovanni Cerulli Irelli for carefully reading several preliminary versions of this article. The first author likes to thank the Max-Planck Institute for Mathematics in Bonn for a one year research stay in 2010/2011. The third author thanks the Sonderforschungsbereich/Transregio SFB 45 for financial support, and all three authors thank the Hausdorff Institute for Mathematics in Bonn for support and hospitality.

### References

- [BFZ] A. Berenstein, S. Fomin, A. Zelevinsky, Cluster algebras III: Upper bounds and double Bruhat cells, Duke Math. J. 126 (2005), no. 1, 1–52.
- [BIRS] A. Buan, O. Iyama, I. Reiten, J. Scott, Cluster structures for 2-Calabi-Yau categories and unipotent groups, Compos. Math. 145 (2009), no. 4, 1035–1079.
- [C] P.M. Cohn, Unique factorization domains, Amer. Math. Monthly 80 (1973), 1–18.
- [DWZ] H. Derksen, J. Weyman, A. Zelevinsky, Quivers with potentials and their representations II: applications to cluster algebras, J. Amer. Math. Soc. 23 (2010), no. 3, 749–790.
- [DK] P. Di Francesco, R. Kedem, Q-systems as cluster algebras. II. Cartan matrix of finite type and the polynomial property, Lett. Math. Phys. 89 (2009), no. 3, 183–216.
- [FZ1] S. Fomin, A. Zelevinsky, Cluster algebras. I. Foundations, J. Amer. Math. Soc. 15 (2002), no. 2, 497–529.
- [FZ2] S. Fomin, A. Zelevinsky, Cluster algebras. II. Finite type classification, Invent. Math. 154 (2003), no. 1, 63–121.
- [GLS1] C. Geiß, B. Leclerc, J. Schröer, Semicanonical bases and preprojective algebras, Ann. Sci. École Norm. Sup. (4) 38 (2005), no. 2, 193–253.
- [GLS2] C. Geiß, B. Leclerc, J. Schröer, Kac-Moody groups and cluster algebras, Adv. Math. 228 (2011), 329–433.
- [HL] D. Hernandez, B. Leclerc, Cluster algebras and quantum affine algebras, Duke Math. J. 154 (2010), no. 2, 265–341.
- [K] V. Kac, Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990. xxii+400pp.
- [La] P. Lampe, Email from October 8th, 2011.
- [Le] B. Leclerc, Quantum loop algebras, quiver varieties, and cluster algebras, To appear in the Proceedings of ICRA XIV "Representations of Algebras and Related Topics", published by the European Mathematical Society, Preprint (2011), 34pp, arXiv:1102.1076
- [Lu] G. Lusztig, Semicanonical bases arising from enveloping algebras, Adv. Math. 151 (2000), no. 2, 129–139.
- [N1] H. Nakajima, Quiver varieties and cluster algebras, Kyoto J. Math. 51 (2011), no. 1, 71–126.
- [N2] H. Nakajima, Lecture at the workshop "Cluster algebras, representation theory, and Poisson geometry", Banff, September 2011.
- [R] C.M. Ringel, The preprojective algebra of a quiver. Algebras and modules, II (Geiranger, 1996), 467–480, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.
- [Z] A. Zelevinsky, Private communication, Balestrand, June 2011, and Email exchange, July 2011.

Christof Geiss Instituto de Matemáticas Universidad Nacional Autónoma de México Ciudad Universitaria 04510 México D.F. México

*E-mail address*: christof@math.unam.mx

Bernard Leclerc Université de Caen LMNO, CNRS UMR 6139 Institut Universitaire de France 14032 Caen cedex France

*E-mail address*: bernard.leclerc@unicaen.fr

Jan Schröer Mathematisches Institut Universität Bonn Endenicher Allee 60 53115 Bonn Germany

*E-mail address*: schroer@math.uni-bonn.de