Excluding Degree Sequences of Cycles

Christian Joseph Altomare¹

The Ohio State University, 231, West 18th Avenue, Columbus, Ohio, United States

Abstract

In this paper, we characterize the degree sequences excluding the degree sequence of a square in terms of forcibly chordal graphs, and we prove several related results.

1. Introduction

Let G be a finite, simple graph and let $D(G) = (d_1, \ldots, d_n)$ be its sequence of vertex degrees listed in decreasing order. The sequence D(G) is known as the *degree sequence* of G, the graph G is said to *realize* D(G), and G is said to be a *realization* of D(G). We call a sequence (d_1, \ldots, d_n) of nonnegative integers a *degree sequence* if it is realized by some graph. We make the convention that if D is used without comment, it denotes a degree sequence. Similarly for D_1 , D_2 , and so on.

Given D_1 and D_2 , we define $D_1 \leq D_2$ to mean there is G_1 realizing D_1 and G_2 realizing D_2 such that $G_1 \sqsubseteq G_2$, where \sqsubseteq is the induced subgraph relation. The reader may check that \preceq is a transitive relation on degree sequences. We say that G_2 excludes G_1 if $G_1 \not\sqsubseteq G_2$, that D_2 excludes D_1 if $D_1 \not\preceq D_2$, that D excludes G if D excludes D(G), and that G excludes D if D(G) excludes D.

A set X of vertices in a graph is *complete* (*anticomplete*) if every (no) two distinct vertices in X are adjacent. Given disjoint sets X and Y, we say that X is *complete to* (*anticomplete to*) Y if each (no) x in X is adjacent to each (any) y in Y. A graph S is *split* if its vertex set has a partition (A, B) with A

Email address: altomare@math.ohio-state.edu (Christian Joseph Altomare)

complete and B anticomplete. Such partitions are called *split partitions*. Split partitions of a split graph are not in general unique. In writing a split partition (A, B), we understand that A is complete and B is anticomplete.

A graph is *chordal* if every induced cycle is a triangle. For a graph G and graph property \mathcal{P} , we say G and D(G) are *forcibly*- \mathcal{P} if every realization of D(G)has property \mathcal{P} . We let $G_1 \coprod G_2$ denote the disjoint union of G_1 and G_2 . We let G[X] be the induced subgraph of G with vertex set X. We call a two edge matching M_2 . For other basic graph theoretic definitions and terminology, we refer the reader to [3].

In [2], Chudnovsky and Seymour prove Rao's Conjecture; given infinitely many degree sequences $D_1, D_2, \ldots, D_n, \ldots$, there are positive integers i < jsuch that $D_i \leq D_j$. To prove this they essentially give, for an arbitrary degree sequence D, an approximate structure theorem for those graphs excluding D. This general theorem is very powerful, allowing them to resolve a nearly thirty year old conjecture. It further suggests and leaves open a problem of independent interest: to give *exact* structure theorems for *specific* degree sequence exclusions. That is the focus of the current paper. Our main result is a structure theorem characterizing degree sequences excluding the square C_4 .

2. Technical Lemmas

We recall the following folklore theorem, whose simple proof we omit.

Proposition 1. A graph G is a split graph iff G excludes M_2 and all holes. Equivalently, G is split iff G excludes M_2 , C_4 , and C_5 .

In particular split graphs are chordal. We have the following corollary.

Corollary 2. A degree sequence D is the degree sequence of a split graph iff D excludes M_2 , C_4 , and C_5 , or equivalently, iff D excludes M_2 and all cycles on at least 4 vertices.

By the well known characterization [5] of split graphs as those graphs for which some Erdös-Gallai inequality [4] is equality, we see that every realization of a split graph is also split; every split graph is thus forcibly split. Let S be split with split partition (A, B). Let H be an arbitrary graph. We define $(S, A, B) \circ H$ as the graph G with vertex set $V(S) \cup V(H)$ formed by joining H completely to A and anticompletely to B. This operation is defined by R. Tyshkevish in [6], where she states and proves a unique decomposition theorem for finite graphs with respect to \circ .

Lemma 3. Let $n \ge 4$. Let S be a split graph with split partition (A, B) and let H be an arbitrary graph. If C_n is an induced subgraph of $(S, A, B) \circ H$, then C_n is an induced subgraph of S or H.

PROOF. Choose an induced n point cyle C in $(S, A, B) \circ H$. The set $V(C) \cap B$ is either empty or nonempty. We consider these cases.

Suppose there is x in $V(C) \cap B$. Since $d_C(x) = 2$ and x is only adjacent to vertices in A, we see that $|V(C) \cap A| \ge 2$, so choose distinct vertices y, z in $V(C) \cap A$. If $V(C) \cap H$ contains a vertex w, then $\{y, z, w\}$ is the vertex set of an induced triangle in C since A is both complete and complete to H. By hypothesis, C is a cycle on $n \ge 4$ vertices and thus has no induced triangle. Therefore $V(C) \cap H$ is empty, showing C is an induced subgraph of S.

We now suppose $V(C) \cap B$ is empty. If $V(C) \cap A$ is also empty, then $C \sqsubseteq H$, which proves the lemma. If $|V(C) \cap A| = 1$, then C has at least three remaining elements, which are all contained in H as $V(C) \cap B$ is empty by hypothesis. Since A is complete to H it thus follows that $d_C(x) \ge 3$, a contradiction. If $|V(C) \cap A| = 2$, then $V(C) \cap H$ must be nonempty so the two vertices of $V(C) \cap A$ together with a vertex of $V(C) \cap H$ comprise an induced triangle in C, a contradiction. If $|V(C) \cap A| \ge 3$, then any three vertices of $V(C) \cap A$ comprise an induced triangle in C since A is complete, a contradiction.

In any case, we have either that C is contained in S, H, or a contradiction, completing the proof.

Lemma 4. Let $k \ge 5$. Suppose D excludes C_{k-1} , but D does not exclude C_k . Let G be a realization of D containing a cycle C isomorphic to C_k , and let Aand B be the sets of vertices of G - C that are complete and anticomplete to C, respectively. Then $G = (G[A \cup B], A, B) \circ C$. PROOF. We have only to show that every vertex of G - C is complete or anticomplete to C, that vertices complete to C are pairwise adjacent, and that vertices anticomplete to C are pairwise nonadjacent.

First we show that every vertex of G - C is complete or anticomplete to C. Assume not. Then there is a vertex x outside of C adjacent to some vertex y of C and nonadjacent to some other vertex z of C. Let v be a neighbor of z in C distinct from y. Let $K = G[C \cup x]$. Define K' as the graph obtained from $K/\{v, z\}$ by subdividing the edge xy with a new vertex t. Simple checking shows that K and K' have the same degree sequence. But $K' - \{x, t\}$ is isomorphic to C_{k-1} . Therefore K' contains C_{k-1} as an induced subgraph. Therefore K does not exclude $D(C_{k-1})$, and hence G does not exclude $D(C_{k-1})$ either. This implies that $D(C_{k-1}) \leq D$, contrary to hypothesis. This contradiction shows that every vertex outside C is complete or anticomplete to C as claimed.

Next, assume there are nonadjacent vertices x and y, both complete to C. Write C in cyclic order as c_1, c_2, \ldots, c_k . Let $G' = G + c_1c_3 - c_3x + xy - yc_1$. One may check that D(G) = D(G') and that $G'[c_1, c_3, c_4, \ldots, c_k]$ is a cycle in that cyclic order. Therefore G' contains an induced C_{k-1} . We thus see that D does not exclude C_{k-1} , contrary to hypothesis. This contradiction shows x and ymust be adjacent. Since x and y are arbitrary elements of A, it follows that Ais complete as claimed.

Finally, let x and y be distinct vertices in B. It is enough to show x and y are not adjacent. Suppose they are adjacent. Then $G[C \cup \{x, y\}]$ is isomorphic to $C_k \coprod P_2$, which has the same degree sequence as $C_{k-1} \coprod P_3$. Therefore D does not exclude C_{k-1} , contrary to assumption. This completes the proof.

3. The Main Results

We now state our main theorem, from which we derive our other main results as corollaries. A certain abuse of notation makes the statements of these results more concise, so we make the convention that $D = D(\text{SPLIT} \circ G)$ means that there is some split graph S with split partition (A, B) such that D is the degree sequence of $(S, A, B) \circ G$, and D = D(SPLIT) means D is the degree sequence of a split graph.

Theorem 5. Let $n \ge 4$. A degree sequence D excludes C_n iff either $D = D(\text{SPLIT} \circ C_{n+1}), D = D(\text{SPLIT} \circ C_{n+2}), \text{ or } D$ forcibly excludes each chordless cycle on at least n vertices.

PROOF. First, if D excludes all chordless cycles on n or more vertices, then in particular D excludes C_n . If $D = D(\text{SPLIT} \circ C_{n+1})$, then by Lemma 3, if $D(C_n) \preceq D$ then $C_n \sqsubseteq C_{n+1}$ or $C_n \sqsubseteq S$ for some split graph S, a contradiction. Therefore D excludes C_n . Similarly if $D = D(\text{SPLIT} \circ C_{n+2})$ then D excludes C_n . One direction of the theorem is thus proved.

We now prove the converse. So, let D exclude C_n . We must show D falls into one of the above three classes as claimed.

First, note D excludes C_{n+k} for all $k \ge 3$. To see this, assume not. Note that $D(C_{n+k}) = D(C_n \coprod C_k)$, as C_k exists since $k \ge 3$ by assumption. Therefore $D(C_n \coprod C_k) \preceq D$, so that D has a realization G such that $C_n \coprod C_k \sqsubseteq G$. In particular $C_n \sqsubseteq G$, contrary to assumption that D excludes C_n . This contradiction proves our claim.

Next, we break into cases. The first case we consider is that D excludes C_{n+1} and C_{n+2} . D excludes C_n by hypothesis, and by the previous paragraph, D excludes C_{n+k} for k at least three. Therefore D excludes all cycles on at least n vertices. Therefore, as claimed, no realization has a chordless cycle on n or more vertices.

The other case is that D does not exclude both C_{n+1} and C_{n+2} . Then D has a realization G containing either C_{n+1} or C_{n+2} as an induced subgraph. Let k = n + 1 if G contains an induced C_{n+1} and let k = n + 2 otherwise. In either case, D excludes C_{k-1} but not C_k . Let $C_k = C$. Then it follows by Lemma 4 that $G = (G[A \cup B], A, B) \circ C)$, thus completing the proof.

Applying Theorem 5 with n = 4, we obtain the following theorem as a corollary.

Theorem 6. A degree sequence D excludes C_4 iff either D is forcibly chordal, $D = D(\text{SPLIT} \circ C_5)$, or $D = D(\text{SPLIT} \circ C_6)$.

Taking complements yields the following theorem as well.

Theorem 7. A degree sequence D excludes M_2 iff either D is forcibly antichordal, $D = D(\text{SPLIT} \circ C_5)$, or $D = D(\text{SPLIT} \circ K_{3,3})$.

PROOF. Just take complements, use Theorem 6, and note antichordal graphs are the complements of chordal graphs by definition, the pentagon is selfcomplementary, and the complement of a hexagon has the same degree sequence as $K_{3,3}$.

We omit proofs of the following corollaries. Details may be found in [1].

Corollary 8. A degree sequence D excludes C_4 and C_5 iff D is forcibly chordal or $D = D(\text{SPLIT} \circ C_6)$.

Corollary 9. A degree sequence D excludes M_2 and C_4 iff $D = D(SPLIT \circ C_5)$ or D = D(SPLIT).

In fact, it is proved in [1] that Corollary 9 holds not only for degree sequences, but in fact for graphs as well. More precisely, the following proposition holds.

Proposition 10. A graph G excludes M_2 and C_4 iff G is a split graph or $G = (S, A, B) \circ C_5$ for some split graph S with split partition (A, B).

References

- Christian Altomare. Degree Sequences, Forcibly Chordal Graphs and Combinatorial Proof Systems. PhD thesis, The Ohio State University, December 2009. http://etd.ohiolink.edu.
- [2] Maria Chudnovsky and Paul Seymour. The proof of Rao's Conjecture on degree sequences. In Preparation.

- [3] Reinhard Diestel. *Graph Theory*. Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 2000.
- [4] Paul Erdös and Tibor Gallai. Gráfok előírt fokú pontokkal (Graphs with Prescribed Degrees of Vertices). Mat. Lapok, 11:264–274, 1960. In Hungarian.
- [5] Peter L. Hammer and Bruno Simeone. The splittance of a graph. Combinatorica, 1(3):275–284, 1981.
- [6] Regina Tyshkevich. Decomposition of graphical sequences and unigraphs. Discrete Math, 220(1-3):201-238, 2000.