

An adjustable law of motion for relativistic spherical shells

Research Article

L. Zaninetti ^{1*},

Dipartimento di Fisica Generale,
Università degli Studi di Torino
via P.Giuria 1, 10125 Torino, Italy

Abstract: A classical and a relativistic law of motion for an advancing shell are deduced applying the thin layer approximation. A new parameter connected with the quantity of absorbed matter in the expansion is introduced; this allows of matching theory and observation.

PACS (2008): 47.75.+f, 52.27.Ny, 98.38.Mz

Keywords: Relativistic fluid dynamics, Relativistic plasmas, Supernova remnants
© Versita Warsaw and Springer-Verlag Berlin Heidelberg.

1. Introduction

The first law of motion to be analyzed is the Sedov–Taylor solution

$$R(t) = \left(\frac{25}{4} \frac{E t^2}{\pi \rho} \right)^{1/5}, \quad (1)$$

where E is the energy injected into the process, ρ is the density of the surrounding medium, and t is the time, see [1–3]. This equation allows of deducing the energy of the first explosion of the atomic bomb, the Trinity test in New Mexico in 1945, which is $E \approx 10^{21}$ ergs. A second application of this equation belongs to astrophysics and is the supernova explosion (SN) in which $E \approx 10^{51}$ ergs, see [4]. A different approach as given by [5] and [6] analyzes two self-similar solutions with varying inverse power law exponents for the density profile of the advancing matter, R^{-n} , and ambient medium, R^{-s} . The previous assumptions give a law of motion $R \propto t^{\frac{n-3}{n-s}}$ when $n > 5$. Another example is an analytical solution suggested by [7] where the radius–time relationship is regulated by the decrease in density: as an example, a density proportional to R^{-9} gives $R \propto t^{2/3}$: this mechanism has been applied to young supernova remnants, see [8]. The case of spherically symmetric thin shells has been recently analyzed both

* E-mail: E-mail: zaninetti@ph.unito.it

in special relativity, see [9] with application to the mass inflation phenomenon, and in general relativity, see [10] where the solutions are split into “ordinary” matter and “exotic” matter. The actual theoretical situation leaves a series of questions unanswered or only partially answered:

- Is it possible to deduce an equation of motion for an expanding shell by assuming that only a fraction of the mass enclosed in the advancing sphere is absorbed in the thin layer?
- Can this new adjustable law of motion for an expanding shell be calibrated on the observed data?
- Can we deduce an asymptotic behavior for the advancing radius of the thin layer of the type $\propto t^\alpha$, where t is the time and α an observational exponent?
- Can we model the momentum carried away by the photons?
- Can we derive a relativistic law of motion for the expanding shell under the previously outlined hypothesis?

In order to answer these questions, Section 2 reviews the existing situation and reports a new classical law of motion which introduces the two concepts of porosity and photon’s losses. Section 3 reviews a first relativistic law of motion and introduces a new relativistic law of motion which includes the porosity.

2. Classical thin layer approximation

This Section reviews the general purpose power law model, the standard thin layer approximation with constant density and the thin layer approximation in a medium with density regulated by a power law. A first new equation of motion is introduced which works in a medium with constant density but swept mass regulated by a parameter called porosity. A second new equation of motion includes porosity and photon’s losses.

2.1. The power law solution

The equation of the expansion of a shell can be modeled by a power law of the type

$$R(t) = R_0 \left(\frac{t}{t_0} \right)^\alpha \quad , \quad (2)$$

where R is the radius of the expansion, t is the time, R_0 is the radius at $t = t_0$, and α is an exponent which can be found from a numerical analysis.

The velocity is

$$v(t) = \alpha R_0 \left(\frac{1}{t_0} \right)^\alpha t^{\alpha-1} \quad . \quad (3)$$

As an example, Fig. 1 reports the fit of SN 1993J where the radius is growing more slowly than the free expansion, $R \propto t$, and more quickly than the Sedov–Taylor solution, $R \propto t^{0.4}$, see equation (1).




Figure 1. Theoretical radius as given by the power law fit represented by equation (2) with $\alpha = 0.828$, $R_0 = 0.0087$ pc and $t_0 = 0.498$ yr. The astronomical data of SN1993J are represented by vertical error bars and are extracted from Table 1 in [11].

2.2. Standard case

The thin layer approximation assumes that all the swept-up gas accumulates infinitely in a thin shell just after the shock front. The conservation of the radial momentum requires that

$$\frac{4}{3}\pi R^3 \rho \dot{R} = \frac{4}{3}\pi R_0^3 \rho \dot{R}_0 \quad , \quad (4)$$

where R and \dot{R} are the radius and the velocity of the advancing shock, ρ the density of the ambient medium, R_0 the initial radius evaluated at $t = t_0$, and \dot{R}_0 the initial velocity, see [12, 13]. The law of motion is

$$R = R_0 \left(1 + 4 \frac{\dot{R}_0}{R_0} (t - t_0) \right)^{\frac{1}{4}} \quad . \quad (5)$$

and the velocity

$$\dot{R} = \dot{R}_0 \left(1 + 4 \frac{\dot{R}_0}{R_0} (t - t_0) \right)^{-\frac{3}{4}} \quad . \quad (6)$$

From equation (5) we can extract \dot{R}_0 and insert it in equation (6)

$$\dot{R} = \frac{1}{4(t - t_0)} \frac{R^4 - R_0^4}{R_0^3} \left(1 + \frac{R^4 - R_0^4}{R_0^4} \right)^{-\frac{3}{4}} \quad . \quad (7)$$

The astrophysical radius in pc as a function of time is

$$R(t) = \sqrt[4]{R_0^3 (4.08 \cdot 10^{-6} v_1 (t_1 - t_0) + R_0)} \text{ pc} \quad , \quad (8)$$

where t_1 and t_0 are times in years, R_0 is the radius in pc when $t_1 = t_0$ and v_1 is the velocity in $\frac{1 \text{ km}}{\text{s}}$ units when $t_1 = t_0$. The momentum solution in the presence of a constant density medium scales as $t^{0.25}$. More details can be found in [14].

2.3. Classical medium with power law

We assume that around the SNR the density of the interstellar medium (ISM) has the following two piecewise dependencies

$$\rho(R) = \begin{cases} \rho_0 & \text{if } R \leq R_0 \\ \rho_0 \left(\frac{R_0}{R}\right)^d & \text{if } R > R_0 \end{cases} . \quad (9)$$

In this framework, the density decreases as an inverse power law with an exponent d which can be fixed from the observed temporal evolution of the radius. The mass swept, M_0 , in the interval $0 \leq r \leq R_0$ is

$$M_0 = \frac{4}{3} \rho_0 \pi R_0^3 . \quad (10)$$

The mass swept, M , in the interval $0 \leq r \leq R$ with $r \geq R_0$ is

$$\begin{aligned} M &= -4r^3 \rho_0 \pi \left(\frac{R_0}{r}\right)^d (d-3)^{-1} \\ &\quad + 4 \frac{\rho_0 \pi R_0^3}{d-3} + \frac{4}{3} \rho_0 \pi R_0^3 . \end{aligned} \quad (11)$$

Momentum conservation in the thin layer approximation requires that

$$Mv = M_0 v_0 , \quad (12)$$

where v is the velocity at t and v_0 is the velocity at $t = t_0$. The previous expression as a function of the radius is

$$v = \frac{r_0^3 v_0 (3-d)}{3 r_0^d R^{3-d} - r_0^3 d} . \quad (13)$$

In this differential equation of first order in R , the variables can be separated and an integration term-by-term gives the following nonlinear equation \mathcal{F}_{NL}

$$\begin{aligned} \mathcal{F}_{NL} &= (4R_0^3 d - R_0^3 d^2) R - 3R_0^d R^{4-d} + R_0^4 d^2 \\ &\quad + 12R_0^3 v_0 t + 3R_0^4 - 4R_0^4 d \\ &\quad + 7R_0^3 v_0 dt_0 + R_0^3 v_0 d^2 t \\ &\quad - 7R_0^3 v_0 dt - 12R_0^3 v_0 t_0 - R_0^3 v_0 d^2 t_0 = 0 . \end{aligned} \quad (14)$$

An approximate solution of $\mathcal{F}_{NL}(r)$ can be obtained assuming that

$$3R_0^d R^{4-d} \gg -(4R_0^3 d - R_0^3 d^2)R$$

$$\begin{aligned} R(t) = & \\ & (R_0^{4-d} - \frac{1}{3}dR_0^{4-d}(4-d) \\ & + \frac{1}{3}(4-d)v_0 R_0^{3-d}(3-d)(t-t_0))^{\frac{1}{4-d}} . \end{aligned} \quad (15)$$

Up to now, the physical units have not been specified: pc for length and yr for time are the hybrid units usually adopted for SNRs. With these units, the initial velocity v_0 is expressed in $\frac{pc}{yr}$ and should be converted into $\frac{km}{s}$. This means that $v_0 = 1.02 \cdot 10^{-6} v_{0,1}$ where $v_{0,1}$ is the initial velocity expressed in $\frac{km}{s}$; in this transformation we have used $1yr = 3.15 \times 10^7 sec$ and $1 pc = 3.086 \times 10^{18} cm$. The astrophysical version of the above equation in pc is

$$\begin{aligned} R(t) = & \\ & (R_0^{4-d} - \frac{1}{3}dR_0^{4-d}(4-d) + 3.402 \cdot 10^{-7} \times \\ & \times (4-d)v_1 R_0^{3-d}(3-d)(t_1-t_0))^{\frac{1}{4-d}} pc , \end{aligned} \quad (16)$$

where t_1 and t_0 are times in years, R_0 is the radius in pc at $t_1 = t_0$ and v_1 is the velocity at $t_1 = t_0$ in $\frac{km}{s}$. The approximate solution (15) has the following limit as $t \rightarrow \infty$

$$R(t) = C_{th} t^{\frac{1}{4-d}} , \quad (17)$$

where

$$C_{th} = \left(\frac{1}{3} \frac{R_0^3 v_0 (-3+d)(-4+d)}{R_0^d} \right)^{\frac{1}{(4-d)}} . \quad (18)$$

On imposing

$$\alpha = \frac{1}{(4-d)} , \quad (19)$$

we obtain

$$d = \frac{4\alpha - 1}{\alpha} . \quad (20)$$

where α is an observable parameter defined in Section 2.1. This means that the unknown parameter d can be deduced from the observed parameter α . The numerical result is reported in Fig. 2. More details can be found in [15].




Figure 2. Theoretical radius in pc as function of the time in yr as obtained by the solution of the non linear equation (14) (full line) which models the medium with variable density. The parameters of the simulation are $d = 2.24$, $R_0 = 0.031$ pc, $t_0 = 0.049$ yr and $v_{0,1} = 50\,000$. The astronomical data of SN 1993J at 6 cm are represented through the error bars and are extracted from Table 1 in [11]

2.4. Classical porosity

The thin layer approximation with porosity p in classical physics assumes that only a fraction of the total mass enclosed in the volume of the expansion accumulates in a thin shell just after the shock front. The global mass between 0 and R_0 is $\frac{4}{3}\pi\rho R_0^3$ where ρ is the density of the ambient medium. The mass that resides beyond the expansion is

$$M_0 = \left(\frac{4}{3}\pi\rho R_0^3\right)^{\frac{1}{p}} . \quad (21)$$

The mass swept between 0 and R is

$$M = \left(\frac{4}{3}\pi\rho R^3\right)^{\frac{1}{p}} . \quad (22)$$

The parameter p defined in the interval $[0, 1]$ has been introduced because the observed radius time relationship in SN 1993J can be approximated by a power law dependence of the type $R \propto t^{0.82}$ [11]. The theory suggests $R \propto R^{2/5}$ for the Sedov solution, see equation (1) or $R \propto R^{1/4}$ for the thin layer approximation, see equation (5); therefore these two models does not match with the observations. We will see later on how is possible deduce p from the observed data. The conservation of radial momentum requires that, after the initial radius R_0 ,

$$Mv = M_0v_0 \quad , \quad (23)$$

where R and v are the radius and velocity of the advancing shock. In classical physics, the velocity as a function of radius is

$$v = v_0 \left(\frac{R_0}{R} \right)^{\frac{3}{p}} , \quad (24)$$

and introducing $\beta_0 = \frac{v_0}{c}$ and $\beta = \frac{v}{c}$ we obtain

$$\beta = \beta_0 \left(\frac{R_0}{R} \right)^{\frac{3}{p}} . \quad (25)$$

The law of motion is

$$R(t) = \left(R_0^{1+3p^{-1}} + (3+p) v_0 R_0^{3p^{-1}} (t-t_0) p^{-1} \right)^{\frac{p}{3+p}} , \quad (26)$$

where t is the time and t_0 is the initial time. In classical physics, the velocity as a function of time is

$$v = \frac{N}{D} \quad (27)$$

where

$$\begin{aligned} N &= \left(R_0^{1+3p^{-1}} + (3+p) v_0 R_0^{3p^{-1}} (t-t_0) p^{-1} \right)^{\frac{p}{3+p}} v_0 R_0^{3p^{-1}} p \\ D &= R_0^{1+3p^{-1}} p + 3 v_0 R_0^{3p^{-1}} t + v_0 R_0^{3p^{-1}} t p \\ &\quad - 3 v_0 R_0^{3p^{-1}} t_0 - v_0 R_0^{3p^{-1}} t_0 p . \end{aligned} \quad (28)$$

Equation (26) can also be solved with a similar solution of type $R = K(t-t_0)^\alpha$, k being a constant, and the classical result is

$$R(t) = \left(\frac{(3+p) v_0 R_0^{3/p} (t-t_0)}{p} \right)^{\frac{p}{3+p}} . \quad (29)$$

The similar solution for the velocity is

$$v(t) = (3+p)^{-\frac{3}{3+p}} p^{\frac{3}{3+p}} v_0^{\frac{p}{3+p}} R_0^{\frac{3}{3+p}} (t-t_0)^{-\frac{3}{3+p}} . \quad (30)$$

The similar formula (29) for the radius can be compared with the observed radius–time relationship reported as

$$R(t) = r_{obs} t^{\alpha_{obs}} , \quad (31)$$

where the two parameters r_{obs} and α_{obs} are found from the numerical analysis of the observational data, see Section 2.1. In this case the velocity is

$$v(t) = r_{obs} \alpha_{obs} t^{(\alpha_{obs}-1)} . \quad (32)$$

The comparison between theory and observation allows of deducing p :

$$p = \frac{3\alpha_{obs}}{1-\alpha_{obs}} . \quad (33)$$

The theoretical solution as given by Equation (26) can be found through the Levenberg–Marquardt method (subroutine MRQMIN in [16]) and Fig. 3 reports a numerical example.




Figure 3. Theoretical radius in pc as function of the time in yr as obtained by the solution of Equation (26) (full line) which models the medium with constant density and porosity. The parameters of the simulation are $p=18.8$, $R_0 = 5 \cdot 10^{-5}$ pc, $t_0 = 4.98 \cdot 10^{-6}$ yr and $v_{0,1} = 30\,000$. The astronomical data of SN 1993J at 6 cm are represented through the error bars and are extracted from Table 1 in [11]

2.5. Classical porosity and photon losses

The emission of photons produces losses in the momentum carried by the advancing shell. This effect can be parametrized by the following conservation law

$$(4/3 \rho \pi R^3)^{p-1} v + C_l (4/3 \rho \pi)^{p-1} (R^2 - R_0^2) v = M_0 v_0 \quad , \quad (34)$$

where $C_l (4/3 \rho \pi)^{p-1} (R^2 - R_0^2)$ takes account of the losses and C_l is a constant. At the moment of writing is not possible to deduce the parameter C_l which controls the quantity of momentum carried away by the photons due to the presence of another parameter p which controls the quantity of swept mass during the expansion. The expression for the velocity in presence of porosity and momentum losses by a radiative process is

$$\beta = \frac{(R_0^{p-1})^3 \beta_0}{(R^{p-1})^3 + C_l R^2 - C_l R_0^2} \quad . \quad (35)$$

Fig. 4 reports how a given value of C_l modifies the velocity. The non linear equation for the trajectory in presence of porosity and losses is

$$\begin{aligned} & - \frac{(R_0^{p-1})^3 R_0 p}{3+p} + 2 \frac{C_l R_0^3}{3+p} + 2/3 \frac{C_l R_0^3 p}{3+p} + \frac{(R^{p-1})^3 R p}{3+p} + 1/3 \frac{C_l R^3 p}{3+p} \\ & + \frac{C_l R^3}{3+p} - \frac{C_l R_0^2 R p}{3+p} - 3 \frac{C_l R_0^2 R}{3+p} - (R_0^{p-1})^3 \beta_0 c (t - t_0) = 0 \quad . \end{aligned} \quad (36)$$




Figure 4. Theoretical classical velocity with porosity as given by Equation (24) (full line), velocity as given by the power law dependence as in Equation (32) (presence of porosity) (dash-dash line) and velocity with porosity and photon losses as given by equation (35) when $C_l=50$ (dash-dot-dash line). The parameters of the simulation are $p=18.8$, $R_0 = 5 \cdot 10^{-5}$ pc, $t_0 = 4.98 \cdot 10^{-6}$ yr and $v_{0,1} = 30\,000$.

An example of the modification of the trajectory as given by the parameter C_l is reported in Fig. 5.

3. Relativistic thin layer approximation

The Newton's second law in special relativity is:

$$F = \frac{dP}{dt} = \frac{d}{dt}(mv) = \frac{d}{dt} \left(\frac{m_0 v}{\sqrt{1 - \frac{v^2}{c^2}}} \right) , \quad (37)$$

where F is the force, P is the relativistic momentum, m is the relativistic mass, m_0 is the mass at rest and v is the velocity, see equation (7.16) in [17]. This Section reviews the relativistic thin layer approximation in a medium with density regulated by a power law. A new relativistic equation of motion is introduced which works in a medium with constant density but adjustable swept mass.

3.1. Relativistic power law

In the case of the relativistic expansion of a shell in which all the swept material resides at two different points, denoted by radius R and radius R_0 , the previous equation (37) gives:

$$M \frac{\beta}{\sqrt{1 - \beta^2}} = M_0 \frac{\beta_0}{\sqrt{1 - \beta_0^2}} , \quad (38)$$




Figure 5. Theoretical radius in pc as function of the time in yr as obtained by the solution of Equation (26) with porosity (full line) and radius with with porosity and losses as given by Equation (36) (dashed line) when $C_l=50$. The parameters of the simulation are $\rho=18.8$, $R_0 = 5 \cdot 10^{-5}$ pc, $t_0 = 4.98 \cdot 10^{-6}$ yr and $v_{0,1} = 30\,000$.

where $\beta_0=v_0/c$, $\beta=v/c$, M is the rest mass swept between 0 and R and M_0 is the rest mass swept between 0 and R_0 . This formula is invariant under Lorentz transformations and the initial velocity, v_0 , cannot be greater than the velocity of light. Assuming a spatial dependence of the ISM as given by formula (9), the relativistic conservation of momentum gives

$$\frac{-4 \rho \pi \left(3 r^3 \left(\frac{r_0}{r} \right)^d - r_0^3 d \right) \beta}{3 (-3 + d) \sqrt{1 - \beta^2}} = \frac{4 \rho \pi r_0^3 \beta_0}{3 \sqrt{1 - \beta_0^2}} . \quad (39)$$

According to the previous equation, β is

$$\beta = \frac{-(-3 + d) \beta_0 r_0^3}{\sqrt{D_\beta}} \quad (40)$$

$$\text{with } D_\beta = 9 R_0^6 \beta_0^2 - 6 R_0^6 \beta_0^2 d$$

$$+ 9 r^{6-2d} R_0^{2d}$$

$$- 6 r^{3-d} R_0^{d+3} d + R_0^6 d^2$$

$$- 9 \beta_0^2 r^{6-2d} R_0^{2d} + 6 \beta_0^2 r^{3-d} R_0^{d+3} d .$$

In this differential equation of first order in r , the variables can be separated and the integration can be expressed as

$$\int_{R_0}^R \sqrt{D_\beta} dR = c (3 - d) \beta_0 R_0^3 (t - t_0) . \quad (41)$$




Figure 6. Theoretical radius as obtained by the solution of the relativistic equation (41) which models the medium with variable density (full line). The parameters of the simulation are $d = 2.52$, $R_0 = 0.0068$ pc, $t_0 = 0.14$ yr; and $\beta_0 = 0.30$. The astronomical data of SN 1993J are represented with vertical error bars.

The integral of the previous equation can be performed analytically only in the cases $d = 0$, $d = 1$ and $d = 2$, but we do not report the result because we are interested in a variable value of d . The integral can be easily evaluated from a theoretical point of view using the subroutine QROMB from [16]. The numerical result is reported in Fig. 6. More details can be found in [15, 18].

3.2. Relativistic porosity

According to equation (37) the Newton's law in special relativity is

$$F = \frac{dp}{dt} = \frac{d}{dt}(Mv) = \frac{d}{dt}\left(\frac{M_r v}{\sqrt{1 - \frac{v^2}{c^2}}}\right) = \frac{d}{dt}\left(\frac{M_r v}{\sqrt{1 - \beta^2}}\right) , \quad (42)$$

where F is the force, p is the relativistic momentum, M is the relativistic mass, M_r is the mass at rest, v is the velocity, $\beta = \frac{v}{c}$, and c is the speed of light. In the case of the relativistic expansion of a shell in which the swept material is part of the mass contained in the advancing layer, the equation of motion is

$$\frac{(\rho \frac{4}{3} \pi R^3)^{1/p} \beta}{\sqrt{1 - \beta^2}} = \frac{(\rho \frac{4}{3} \pi R_0^3)^{1/p} \beta_0}{\sqrt{1 - \beta_0^2}} , \quad (43)$$

where $\beta_0 = v_0/c$, R_0 is the initial radius, and R is the temporary radius. The velocity of a relativistic expanding shell is

$$\beta = \frac{\beta_0 R_0^{3/p}}{\sqrt{R_0^{6/p} \beta_0^2 + R^{6/p} - R^{6/p} \beta_0^2}} . \quad (44)$$




Figure 7. Theoretical relativistic velocity for SN 1993J as given by Equation (44) (relativistic porosity-dashed line), theoretical classical velocity as given by Equation (24) (classical porosity -full line), velocity as given by the power law dependence as in Equation (32) (dot-dash-dot-dash line).

Fig. 7 shows the classical and relativistic behaviors of the velocity as a function of the radius R . In this differential equation of the first order in r the variables can be separated and the integration can be expressed as

$$\int_{R_0}^R \sqrt{R_0^{6/p} \beta_0^2 + R^{6/p} - R^{6/p} \beta_0^2} dr = \beta_0 R_0^{3/p} (t - t_0) dt \quad . \quad (45)$$

The integral of the previous equation can be expressed analytically but the result is complicated. The integral can be easily evaluated numerically using the subroutine QROMB from [16]. The numerical result is reported in Fig. 8. An interesting property of the Gamma-ray burst (GRB) is that the frequency of observation at which it starts to be visible decreases with time. An example of such behavior for GRB 050904 in the BAT, XRT, J, and I bands, can be found in Table 1 of [19]. The numerical analysis of the data as reported in Fig. 9 gives

$$\nu = 1.0310^{25} t(s)^{-2.27} \text{ Hz} \quad (46)$$

where t has been chosen as the value of time at which the flux at the chosen frequency is maximum. The interval in time at which the GRB becomes visible in different astronomical bands is here correlated with the velocity of the shell's expansion, see Fig. 10.

4. Conclusions

We have deduced two new laws of motion for an advancing shell assuming that only a fraction of the mass which resides in the surrounding medium is accumulated in the advancing layer. According to the present laws of




Figure 8. Theoretical radius in pc as function of the time in yr as obtained by the solution of the relativistic Equation (45) (relativistic porosity) (full line). The parameters of the simulation are $p=18.3$, $R_0 = 6 \cdot 10^{-5}$ pc, $t_0 = 5 \cdot 10^{-8}$ yr, and $\beta_0 = 0.1$. The astronomical data of SN 1993J are represented through error bars and are extracted from Table 1 in [11]




Figure 9. Bands from the radio through the IR/optical to X-ray and BAT at which the expansion becomes optically thin. The energy bands are expressed in Hz and the y-axis as the logarithm of the frequency. The x-axis reports the logarithm of the time in s. The data are extracted by the author from Table 1 in [19].




Figure 10. Theoretical relativistic velocity for SN 1993J as given by Equation (44) as function of Log(time) in seconds. The parameters of the simulation are $p=18.3$, $R_0 = 2.25 \cdot 10^{-8}$ pc, $t_0 = 5 \cdot 10^{-8}$ yr, and $\beta_0 = 0.33$.

mechanics the analysis has been split into classical and relativistic cases, see Equations (26) and (45). In the classical case a similar solution has been found, see Equation (29). The momentum carried away by the photons can be modeled in the classical case by introducing a reasonable form for the photon's losses, see (36).

Acknowledgements

I would like to thank the two anonymous referees for constructive comments on the text.

References

-
- [1] G. Taylor, Royal Society of London Proceedings Series A 201, 159 (1950).
 - [2] G. Taylor, Royal Society of London Proceedings Series A 201, 175 (1950).
 - [3] L. I. Sedov, Similarity and Dimensional Methods in Mechanics (Academic Press, New York, 1959).
 - [4] A. McCray, R. In: Dalgarno and D. Layzer, eds., Spectroscopy of astrophysical plasmas (Cambridge University Press, Cambridge, 1987).
 - [5] R. A. Chevalier, ApJ 258, 790 (1982).
 - [6] R. A. Chevalier, ApJ 259, 302 (1982).
 - [7] J. K. Truelove and C. F. McKee, ApJS 120, 299 (1999).

- [8] C. D. Vigh, P. F. Velázquez, D. O. Gómez, E. M. Reynoso, A. Esquivel, and E. Matias Schneiter, *ApJ* 727, 32 (2011).
- [9] M. E. Gáspár and I. Rácz, *Classical and Quantum Gravity* 28, 085005 (2011).
- [10] J. Kijowski, G. Magli, and D. Malafarina, *International Journal of Modern Physics D* 18, 1801 (2009), arXiv:gr-qc/0512089.
- [11] J. M. Marcaide, I. Martí-Vidal, A. Alberdi, and M. A. Pérez-Torres, *A&A* 505, 927 (2009), 0903.3833.
- [12] Dyson, J. E. and Williams, D. A., *The physics of the interstellar medium* (Institute of Physics Publishing, Bristol, 1997).
- [13] P. Padmanabhan, *Theoretical astrophysics. Vol. II: Stars and Stellar Systems* (Cambridge University Press, Cambridge, MA, 2001).
- [14] L. Zaninetti, *MNRAS* 395, 667 (2009).
- [15] L. Zaninetti, *Astrophysics and Space Science* 333, 99 (2011).
- [16] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in FORTRAN. The Art of Scientific Computing* (Cambridge University Press, Cambridge, 1992).
- [17] French, A.P., *Special Relativity* (CRC, New York, 1968).
- [18] L. Zaninetti, *Adv. Stud. Theor. Phys.* 4, 525 (2010).
- [19] L. Gou, D. B. Fox, and P. Mészáros, *ApJ* 668, 1083 (2007), arXiv:astro-ph/0612256.